



This is a repository copy of *First search for light dark matter in the neutrino fog with XENONnT*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/226430/>

Version: Published Version

Article:

Aprile, E. orcid.org/0000-0001-6595-7098, Aalbers, J. orcid.org/0000-0003-0030-0030, Abe, K. orcid.org/0009-0000-9620-788X et al. (165 more authors) (2025) First search for light dark matter in the neutrino fog with XENONnT. Physical Review Letters, 134 (11). 111802. ISSN 0031-9007

<https://doi.org/10.1103/physrevlett.134.111802>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

First Search for Light Dark Matter in the Neutrino Fog with XENONnT

E. Aprile¹, J. Aalbers², K. Abe³, S. Ahmed Maouloud⁴, L. Althueser⁵, B. Andrieu⁴, E. Angelino^{6,7}, D. Antón Martín⁸, F. Arneodo⁹, L. Baudis¹⁰, M. Bazyk¹¹, L. Bellagamba¹², R. Biondi¹³, A. Bismark¹⁰, K. Boese¹³, A. Brown¹⁴, G. Bruno¹¹, R. Budnik¹⁵, C. Cai¹⁶, C. Capelli¹⁰, J. M. R. Cardoso¹⁷, A. P. Cimental Chávez¹⁰, A. P. Colijn¹⁸, J. Conrad¹⁹, J. J. Cuenca-García¹⁰, V. D'Andrea^{7,*}, L. C. Daniel Garcia⁴, M. P. Decowski¹⁸, A. Deisting²⁰, C. Di Donato^{21,7}, P. Di Gangi¹², S. Diglio¹¹, K. Eitel²², S. el Morabit¹⁸, A. Elykov²², A. D. Ferella^{21,7}, C. Ferrari⁷, H. Fischer¹⁴, T. Flehmke¹⁹, M. Flierman¹⁸, W. Fulgione^{6,7}, C. Fuselli¹⁸, P. Gaemers¹⁸, R. Gaior⁴, M. Galloway¹⁰, F. Gao¹⁶, S. Ghosh²³, R. Giacomobono²⁴, R. Glade-Beucke¹⁴, L. Grandi⁸, J. Grigat¹⁴, H. Guan²³, M. Guida¹³, P. Gyorgy²⁰, R. Hammann¹³, A. Higuera²⁵, C. Hils²⁰, L. Hoetzsch¹³, N. F. Hood²⁶, M. Iacovacci²⁴, Y. Itow²⁷, J. Jakob⁵, F. Joerg^{13,10}, Y. Kaminaga³, M. Kara²², P. Kavargin¹⁵, S. Kazama²⁷, M. Kobayashi²⁷, D. Koke⁵, A. Kopec^{26,‡}, H. Landsman¹⁵, R. F. Lang²³, L. Levinson¹⁵, I. Li²⁵, S. Li²⁸, S. Liang²⁵, Y.-T. Lin¹³, S. Lindemann¹⁴, M. Lindner¹³, K. Liu¹⁶, M. Liu^{1,16}, J. Loizeau¹¹, F. Lombardi²⁰, J. Long⁸, J. A. M. Lopes^{17,†}, T. Luce¹⁴, Y. Ma²⁶, C. Macolino^{21,7}, J. Mahlstedt¹⁹, A. Mancuso¹², L. Manenti⁹, F. Marignetti²⁴, T. Marrodán Undagoitia¹³, K. Martens³, J. Masbou¹¹, E. Masson⁴, S. Mastroianni²⁴, A. Melchiorre^{21,7}, J. Merz²⁰, M. Messina⁷, A. Michael⁵, K. Miuchi²⁹, A. Molinaro⁶, S. Moriyama³, K. Morā¹, Y. Mosbacher¹⁵, M. Murra¹, J. Müller¹⁴, K. Ni²⁶, U. Oberlack²⁰, B. Paetsch¹⁵, Y. Pan⁴, Q. Pellegrini⁴, R. Peres¹⁰, C. Peters²⁵, J. Pienaar^{8,15}, M. Pierre¹⁸, G. Plante¹, T. R. Pollmann¹⁸, L. Principe¹¹, J. Qi²⁶, J. Qin²⁵, D. Ramírez García¹⁰, M. Rajado¹⁰, R. Singh²³, L. Sanchez²⁵, J. M. F. dos Santos¹⁷, I. Sarnoff⁹, G. Sartorelli¹², J. Schreiner¹³, P. Schulte⁵, H. Schulze EiBing⁵, M. Schumann¹⁴, L. Scotto Lavina⁴, M. Selvi¹², F. Semeria¹², P. Shagin²⁰, S. Shi^{1,§}, J. Shi¹⁶, M. Silva¹⁷, H. Simgen¹³, C. Szyska²⁰, A. Takeda³, P.-L. Tan¹⁹, D. Thers¹¹, F. Toschi²², G. Trincherio⁶, C. D. Tunnell²⁵, F. Tönnies¹⁴, K. Valerius²², S. Vecchi³⁰, S. Vetter²², F. I. Villazon Solar²⁰, G. Volta¹³, C. Weinheimer⁵, M. Weiss¹⁵, D. Wenz⁵, C. Wittweg¹⁰, V. H. S. Wu²², Y. Xing¹¹, D. Xu¹, Z. Xu¹, M. Yamashita³, L. Yang²⁶, J. Ye³¹, L. Yuan^{8,||}, G. Zavattini³⁰ and M. Zhong²⁶

(XENON Collaboration)

¹Physics Department, Columbia University, New York, New York 10027, USA²Nikhef and the University of Groningen, Van Swinderen Institute, 9747AG Groningen, Netherlands³Kamioka Observatory, Institute for Cosmic Ray Research, and Kavli Institute for the Physics and Mathematics of the Universe (WPI), University of Tokyo, Higashi-Mozumi, Kamioka, Hida, Gifu 506-1205, Japan⁴LPNHE, Sorbonne Université, CNRS/IN2P3, 75005 Paris, France⁵Institut für Kernphysik, University of Münster, 48149 Münster, Germany⁶INAF-Astrophysical Observatory of Torino, Department of Physics, University of Torino and INFN-Torino, 10125 Torino, Italy⁷INFN-Laboratori Nazionali del Gran Sasso and Gran Sasso Science Institute, 67100 L'Aquila, Italy⁸Department of Physics, Enrico Fermi Institute and Kavli Institute for Cosmological Physics, University of Chicago, Chicago, Illinois 60637, USA⁹New York University Abu Dhabi - Center for Astro, Particle and Planetary Physics, Abu Dhabi, United Arab Emirates¹⁰Physik-Institut, University of Zürich, 8057 Zürich, Switzerland¹¹SUBATECH, IMT Atlantique, CNRS/IN2P3, Nantes Université, Nantes 44307, France¹²Department of Physics and Astronomy, University of Bologna and INFN-Bologna, 40126 Bologna, Italy¹³Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany¹⁴Physikalisches Institut, Universität Freiburg, 79104 Freiburg, Germany¹⁵Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot 7610001, Israel¹⁶Department of Physics and Center for High Energy Physics, Tsinghua University, Beijing 100084, Peoples Republic of China¹⁷LIBPhys, Department of Physics, University of Coimbra, 3004-516 Coimbra, Portugal¹⁸Nikhef and the University of Amsterdam, Science Park, 1098XG Amsterdam, Netherlands¹⁹Oskar Klein Centre, Department of Physics, Stockholm University, AlbaNova, Stockholm SE-10691, Sweden²⁰Institut für Physik & Exzellenzcluster PRISMA⁺, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany²¹Department of Physics and Chemistry, University of L'Aquila, 67100 L'Aquila, Italy²²Institute for Astroparticle Physics, Karlsruhe Institute of Technology, 76021 Karlsruhe, Germany²³Department of Physics and Astronomy, Purdue University, West Lafayette, Indiana 47907, USA²⁴Department of Physics "Ettore Pancini," University of Napoli and INFN-Napoli, 80126 Napoli, Italy

²⁵Department of Physics and Astronomy, Rice University, Houston, Texas 77005, USA

²⁶Department of Physics, University of California San Diego, La Jolla, California 92093, USA

²⁷Kobayashi-Maskawa Institute for the Origin of Particles and the Universe,
and Institute for Space-Earth Environmental Research, Nagoya University,
Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan

²⁸Department of Physics, School of Science, Westlake University, Hangzhou 310030, Peoples Republic of China

²⁹Department of Physics, Kobe University, Kobe, Hyogo 657-8501, Japan

³⁰INFN-Ferrara and Dipartimento di Fisica e Scienze della Terra, Università di Ferrara, 44122 Ferrara, Italy

³¹School of Science and Engineering, The Chinese University of Hong Kong,
Shenzhen, Guangdong 518172, Peoples Republic of China



(Received 26 September 2024; accepted 21 January 2025; published 20 March 2025)

We search for dark matter (DM) with a mass $[3, 12] \text{ GeV}/c^2$ using an exposure of 3.51 tonne year with the XENONnT experiment. We consider spin-independent DM-nucleon interactions mediated by a heavy or light mediator, spin-dependent DM-neutron interactions, momentum-dependent DM scattering, and mirror DM. Using a lowered energy threshold compared to the previous weakly interacting massive particle search, a blind analysis of $[0.5, 5.0] \text{ keV}$ nuclear recoil events reveals no significant signal excess over the background. XENONnT excludes spin-independent DM-nucleon cross sections $> 2.5 \times 10^{-45} \text{ cm}^2$ at 90% confidence level for $6 \text{ GeV}/c^2$ DM. In the considered mass range, the DM sensitivity approaches the “neutrino fog,” the limitation where neutrinos produce a signal that is indistinguishable from that of light DM-xenon nucleus scattering.

DOI: [10.1103/PhysRevLett.134.111802](https://doi.org/10.1103/PhysRevLett.134.111802)

Introduction—The XENONnT experiment [1], operated at INFN Laboratori Nazionali del Gran Sasso, is primarily designed to search for weakly interacting massive particles (WIMPs) [2–4] with a mass scale of GeV/c^2 – TeV/c^2 . By lowering the energy threshold compared to our previous WIMP search [5], we are able to improve the sensitivity to light dark matter (DM). However, as we increase the exposure, the solar ^8B neutrino coherent elastic neutrino-nucleus scattering (CE ν NS) background begins to affect the sensitivity gain. This gradual slowdown in sensitivity improvement for light DM with additional exposure is known as the neutrino fog [6–10]. In this Letter, we report on the first search for light DM with a mass of $[3, 12] \text{ GeV}/c^2$, achieving DM sensitivities into the neutrino fog. This search utilizes the same dataset and analysis techniques as in our recently reported search for solar ^8B neutrino via CE ν NS interaction [11].

The XENONnT experiment comprises a central detector, operating as a time projection chamber (TPC), with an active target of 5.9 tonnes of cryogenic liquid xenon, enclosed by a Cherenkov neutron veto and muon veto [12] detector to mitigate radiogenic neutrons and cosmogenic muon-induced backgrounds. The operation of XENONnT is supported by various subsystems, including a gas-phase and a liquid-phase xenon circulation and purification system, a radon removal system [13], and a krypton distillation column [14]. A particle interaction in the xenon target may produce a nuclear recoil (NR) with the xenon nucleus, or an electronic recoil (ER) with the atomic electrons. Two arrays of photomultiplier tubes (PMTs) at the top and bottom of the TPC collect the light from the scintillation signal (S1) and the ionization signal (S2), converting them to photoelectrons (PEs). Digitizers convert the PE waveforms that cross the digitization threshold into PMT hits, which are clustered into peaks by reconstruction software [15,16]. The detector wall consists of polytetrafluoroethylene (PTFE) panels to enhance photon gain (g1). The cathode electrode near the bottom of the TPC and the gate electrode below the liquid-gas interface establish a vertical electric field of approximately 23 V/cm, which drifts the ionization electrons to the liquid-gas interface with a maximum drift time of 2.2 ms. An extraction field is generated by the gate and anode electrodes, leading to secondary scintillation of the extracted electrons. The extraction efficiency and single electron scintillation gain contribute to the electron gain (g2). Further details on XENONnT can be found in Ref. [1].

* Also at INFN-Roma Tre, 00146 Roma, Italy.

† Also at Coimbra Polytechnic—ISEC, 3030-199 Coimbra, Portugal.

‡ Present address: Department of Physics and Astronomy, Bucknell University, Lewisburg, Pennsylvania, USA.

§ Contact author: shenyang.shi@columbia.edu

|| Contact author: yuanlq@uchicago.edu

Dataset—This DM search combines data from two science runs spanning from May 1, 2021, to December 10, 2021 (SR0, live time 108.0 days), and from May 19, 2022, to August 8, 2023 (SR1, live time 208.5 days). Between SR0 and SR1, the radon distillation system was upgraded to operate in both gas xenon (GXe) and liquid xenon (LXe) modes. On July 15, 2022, the liquid level was decreased by 0.2 mm from (5.0 ± 0.2) mm and the anode voltage was increased from 4.90 to 4.95 kV to mitigate clustered electron emissions [17,18], while keeping a constant electron gain. In SR0 (SR1), the average electron lifetime is 14.2 ms (20.1 ms), g_1 is 0.1515 ± 0.0014 PE/photon (0.1367 ± 0.0010 PE/photon), and g_2 is 16.45 ± 0.64 PE/e $^-$ (16.85 ± 0.46 PE/e $^-$).

Signal—We consider DM models involving nuclear recoils from the elastic scattering of DM particles on xenon nuclei, focusing on a mass ranging from 3 to 12 GeV/ c^2 . This mass range is favored by various DM models such as WIMP [2–4], asymmetric DM [19,20], and self-interacting DM [21–24]. We examine spin-independent (SI) DM-nucleon [25] and spin-dependent (SD) DM-neutron [26] interactions mediated by a heavy mediator with mass $m_\phi \gg q_0/c \equiv 10$ MeV/ c^2 , where q_0 represents the typical momentum transfer in this search. We also consider DM with SI interaction mediated by a scalar or vector light mediator (LM) with mass $m_\phi \ll q_0/c$ (SI-LM) [24,27,28]. In this case, the differential rate is proportional to $\sigma m_\phi^4 / (q^2/c^2 + m_\phi^2)^2 \approx \sigma m_\phi^4 c^4 / q^4$, where q denotes the momentum transfer, and σ is the DM-nucleon cross section [29,30]. Momentum-dependent DM scattering (MDDM) [31] is also considered, characterized by a modified SI interaction cross section of the form $\sigma_{\chi N} = (q/q_0)^{2n} \sigma_0$, where $n \in \{1, 2\}$ and σ_0 is the DM-nucleon cross section. Additionally, we search for mirror dark matter, characterized by a differential cross section resembling Rutherford scattering [32,33]. The recoil energy spectra are evaluated using Ref. [34], and the spectra for SI interaction are shown in Fig. 1 (top). For astrophysical models and nuclear form factors, we follow Ref. [5] with the exception of mirror DM, where we adopt a DM-mass-dependent Boltzmann velocity v_0 as assumed in Ref. [33] beyond the standard halo model recommended in Ref. [35].

The average number of scintillation photons and ionization electrons from the energy of an NR interaction (E_R) is quantified by the light yield model L_y and the charge yield model Q_y . We model detector responses for low-energy NR events [36] with ^{88}YBe neutron calibration data [37] (with a mean $\bar{E}_R \approx 2$ keV per scatter), and fit L_y and Q_y following the methodology described in Ref. [38]. Yields for $E_R < 0.5$ keV are assumed to be zero, and light DM signals with $E_R > 5.0$ keV are neglected. For $E_R < 5.0$ keV, the ^{88}YBe calibrated yield model aligns with the Q_y of the NEST v2.3.6 [39] yield model within systematic uncertainty, while the L_y uncertainties generally encompass the median




FIG. 1. Top: SI-DM signal spectrum with a cross section of 4.4×10^{-45} cm 2 before (dotted) and after (solid) accounting for the signal efficiency. The ^{88}B CEvNS spectrum (gray) is shown for comparison. Bottom: signal efficiency (black) as a combination of individual efficiencies of S1 ROI (blue), S1 reconstruction (light blue), S2 ROI (dark green), S2 quality selection (light green), and event efficiency (red). These efficiencies are evaluated per science run and weighted by live time.

of the NEST model. The uncertainties of L_y and Q_y are modeled using shape parameters t_{L_y} and t_{Q_y} , respectively. t_{L_y} and t_{Q_y} determine the relative shift of L_y and Q_y from the median toward the upper and lower bound of their 68% credible intervals [36].

A simulation-based detector response model converts the produced photons and electrons to detector observables [40]. Using the depth (Z) and radius (R) position of the event, the fiducial volume is defined as -142.0 cm $\leq Z \leq -13.0$ cm, $R \leq 60.2$ cm for SR0 and $R \leq 59.6$ cm for SR1. The region of interest (ROI) is defined as two or three hits for S1 and [120, 500] PE for S2, corresponding to approximately [4, 16] extracted electrons. Events reconstructed with $R \leq 63.0$ cm in the ROI were blinded before finalizing the analysis. Compared to the previous XENONnT NR search [5], the ROI in this Letter leads to a 17 times higher signal rate expectation for a 6 GeV/ c^2 SI-DM. The overlap of ROI between this Letter and the previously unblinded SR0 WIMP search [5] was not used in the data selection development, and is used in this search.

Three types of selection are implemented, following the ^{88}B CEvNS analysis [11]. For the S1 peaks, the reconstruction requires at least two hits from two different PMTs within 50 ns, which is referred to as the twofold PMT coincidence. For the S2 peaks, other than the waveform width and PMT distribution quality selections

inherited from Ref. [5], an S2 boosted-decision-tree machine is developed as detailed in the background section. To distinguish physical events from detector artifacts, we implement selections that consider temporal and spatial correlations with preceding high-energy events [11], including $S2_{\text{pre}}/\Delta t_{\text{pre}}$, defined as the ratio between the S2 peak area of high-energy events and the time difference from candidate events. Following Ref. [5], a single-scatter selection is applied. A selection based on the single electron rate within a preceding 5 s time window is implemented to mitigate the impact of single electron pileup [17,18].

The signal efficiency consists of the ROI efficiency, the S1 reconstruction efficiency, the S2 quality selection efficiency, and the event efficiency, as illustrated in Fig. 1 (bottom). Excluding yield model uncertainties, the systematic uncertainty of the combined efficiency is estimated to be 25%, predominantly due to the uncertainties in the S1 reconstruction efficiency and the S2 quality selection efficiency.

The ROI efficiency is assessed through detector reconstruction simulation and is sensitive to the yield model. For a 6 GeV/ c^2 SI-DM, the signal efficiency uncertainty due to the yield model alone is estimated to be 34%, resulting from the propagation of t_{Ly} and t_{Qy} to the signal rate computation.

The S1 reconstruction efficiency originates from the twofold PMT coincidence requirement. We employ a data-driven method [40] that simulates low-energy S1 waveforms by sampling a subset of photon waveforms from $^{83\text{m}}\text{Kr}$ 32.1 keV and 9.4 keV conversion electrons and ^{37}Ar 2.8 keV electron capture calibration data. We incorporate a conservative systematic uncertainty of 10% to account for the systematic uncertainty in the shape of the S1 waveform. The S1 reconstruction efficiency is 65% (85%) for S1 signals with 2 (3) detected photons, while the other S1 quality selections achieve acceptances over 99% and are thus neglected in efficiency evaluation.

The S2 quality selection efficiency is evaluated by studying the selection survival ratio of evaluation datasets within the ROI, including waveform simulation [41], surface ^{210}Pb β events from PTFE, ^{214}Pb β events from the cathode, ^{37}Ar 0.27 keV electron capture, and neutron-veto-tagged low energy $^{241}\text{AmBe}$ neutrons. A systematic uncertainty of 10% is adopted based on the variation among the evaluation datasets.

The event reconstruction algorithm pairs one S1 to one S2 peak to build an event. The algorithm may fail when S1 and S2 peaks of areas similar to candidate events occur within a time window of $\mathcal{O}(10)$ ms. To account for this effect, along with the aforementioned event selections, the event efficiency is evaluated per science run by tracking the survival ratio of simulated events randomly inserted into real data assuming a spatially and temporally uniform signal distribution [42]. The event efficiency is 77% (73%) for two-hit S1, and 82% (78%) for three-hit S1 in SR0 (SR1) within the ROI.

Backgrounds—This analysis considers the background described in Refs. [5,38,43]. The expected rate and systematic uncertainties, both in the ROI and in the signal-like region, which is defined as the region containing 50% of the signal for bins ranked by signal-to-background ratio, are summarized in Table I.

The solar ^8B CE ν NS interaction, whose recoil spectrum is nearly indistinguishable from a 5.5 GeV/ c^2 DM-nucleon scattering at cross section of 4.4×10^{-45} cm 2 , is one of the dominant backgrounds of this search. Following the recommended convention [35], the predicted solar ^8B CE ν NS rate after all selections is (3.6 ± 0.9) events $\text{t}^{-1} \text{y}^{-1}$ in SR0 and (3.3 ± 0.8) events $\text{t}^{-1} \text{y}^{-1}$ in SR1, based on the standard solar model ^8B neutrino spectrum [45] and the measured flux of $(5.25 \pm 0.20) \times 10^6$ cm 2 s $^{-1}$ from Sudbury Neutrino Observatory [44], using the nominal NR yield model.

TABLE I. For each model component, we show the background expectation and best-fit values for the background-only hypothesis, background, and SI-DM signal hypothesis with a 4, 6, 10 GeV/ c^2 mass DM. The expectation uncertainties for ^8B CE ν NS background consider both signal efficiency uncertainty and yield uncertainties while neglecting the subdominant 3% ^8B neutrino flux uncertainty [44]. All uncertainties in the last four columns are calculated using the 68% confidence interval constructed by the profile likelihood method. In addition to the event number in the full ROI, the event number in the most signal-like region is shown in parentheses. The best fit and expectation agree within the uncertainties for all components.

Component	Expectation	Background only	4 GeV/ c^2		6 GeV/ c^2		10 GeV/ c^2	
SI-DM	3.2	(1.6)	0.0	(0.0)	0.0	(0.0)
^8B CE ν NS	$11.9^{+4.5}_{-4.2}$	$11.4^{+2.0}_{-3.6}$	10.2 ± 2.7	(2.5 ± 0.7)	$11.4^{+2.7}_{-2.6}$	(6.0 ± 1.4)	$11.4^{+2.7}_{-2.6}$	$(5.4^{+1.3}_{-1.2})$
AC	25.3 ± 1.2	$25.3^{+1.1}_{-1.3}$	25.1 ± 1.2	(3.7 ± 0.1)	25.3 ± 1.2	(4.1 ± 0.2)	25.3 ± 1.2	(3.3 ± 0.1)
ER	0.7 ± 0.7	$0.5^{+0.7}_{-0.6}$	$0.6^{+0.7}_{-0.6}$	(0.0 ± 0.0)	$0.5^{+0.6}_{-0.5}$	(0.3 ± 0.3)	$0.5^{+0.7}_{-0.5}$	$(0.4^{+0.5}_{-0.4})$
Neutron	$0.5^{+0.2}_{-0.3}$	0.5 ± 0.3	0.5 ± 0.3	(0.1 ± 0.1)	0.5 ± 0.3	(0.2 ± 0.1)	0.5 ± 0.3	(0.2 ± 0.1)
Total background	$38.3^{+4.7}_{-4.4}$	$37.7^{+2.5}_{-3.9}$	$36.4^{+3.0}_{-3.0}$	$(6.3^{+0.7}_{-0.7})$	$37.7^{+3.0}_{-2.9}$	$(10.6^{+1.5}_{-1.4})$	$37.7^{+3.0}_{-2.9}$	$(9.2^{+1.4}_{-1.3})$
Observed	...	37	37	(10)	37	(10)	37	(4)

The accidental coincidence (AC) background arises from the accidental pairing of isolated S1 and isolated S2 peaks, where “isolated” refers to peaks that lack a physically corresponding S1 or S2 [11,42]. The quality and reconstruction selections outlined in the signal section reduce the isolated S1 and S2 peak rates to 2.3 (2.2) Hz for S1 and 18 (26) mHz for S2 in SR0 (SR1). Utilizing a synthesized data-driven AC dataset [38,43], two boosted decision tree (BDT) classifiers, S1 BDT and S2 BDT, are trained on an equivalent 6 GeV/ c^2 DM signal, each targeting the AC of different origins. The S1 BDT distinguishes isolated S1 signals originating from random PMT hit clustering by incorporating S1 hit counts, PMT channel distribution, double photoelectron emission [46], and S1 pulse shape [47] into the S1 BDT score. The S2 BDT evaluates whether the S2 pulse shape of the candidate event is compatible with the correlation between electron diffusion and the drift time [48], converting this assessment into an S2 BDT score. These waveform-feature-based BDT scores are used as an inference dimension to enhance signal-to-background discrimination power, as shown in Fig. 2. An AC sideband, primarily consisting of events rejected by the S2 BDT selection, was unblinded before the ROI unblinding to examine the validity of the AC modeling. Subsequently, the S2 area threshold of the ROI was raised from 100 to 120 PE due to an observed tension between the observation and the AC background model. The discrepancy in the AC rate between the unblinding

result of the AC sideband and the prediction is then less than 5%, and the statistical uncertainty of the AC sideband data is conservatively chosen to be the systematic uncertainty of the AC background. The details of sideband validation can be found in Ref. [11]. The AC background prediction is (6.4 ± 0.6) events $t^{-1} y^{-1}$ in SR0 and (7.6 ± 0.4) events $t^{-1} y^{-1}$ in SR1.

The surface background originates from ^{222}Rn progeny plate-out on the PTFE panels, such as β decays from ^{210}Pb . This can result in a higher background rate in the ROI due to the presence of S2 charge-insensitive volumes and the worse position reconstruction resolution of small S2 signals ($\Delta R \approx 2$ cm) near the detector wall. The surface background is modeled in a data-driven manner using nonblinded events. The fiducial volume is selected to mitigate the surface background contribution, yielding a negligible rate of less than 0.10 events $t^{-1} y^{-1}$ at 90% CL. The prediction is validated by a sideband unblinding of events within $61.4 \text{ cm} < R < 63.0 \text{ cm}$. Because of the negligible rate, the surface background is not included in the statistical model.

The radiogenic neutron background originates from nuclear fission and the (α, n) process in the uranium and thorium decay chains. It is modeled using full-chain simulations as described in Ref. [38]. The levels of the radioactive contamination, as determined from the radio assay [5], are further adjusted based on the comparison between observed and simulated high-energy γ ray activity. The prediction of the neutron background is




FIG. 2. DM candidate events in the search space, distributed in quantiles of (S1 BDT, cS2) (left), (S2 BDT, cS2) (center), and ($S2_{\text{pre}}/\Delta t_{\text{pre}}$, cS2) (right). The quantiles are defined to achieve an equal-probable binning of the AC background. Events after selections are shown as pie charts, with each wedge representing the contribution of the components in the best-fit model assuming a 6 GeV/ c^2 DM. The size of the pie charts is proportional to the DM signal model at that position. The contributions of DM, ER, and neutron events in the wedges and histograms are barely visible since the best-fit rates are negligible. The shaded contours show the 1σ and 2σ distributions of the AC background (purple) and solar ^8B CE ν NS background (green). The solid and dashed line contours show the 1σ and 2σ distribution of a 6 GeV/ c^2 DM signal (orange), showing its similarity with ^8B CE ν NS. The equivalent nuclear recoil energy based solely on cS2 is indicated by the gray lines. The one-dimensional projection of best-fit and observed data (black) is shown on the side.

(0.11 ± 0.06) events $\text{t}^{-1} \text{y}^{-1}$ in SR0 and (0.14 ± 0.08) events $\text{t}^{-1} \text{y}^{-1}$ in SR1, considering an effective neutron veto tagging efficiency of 49% (SR0) and 52% (SR1).

The electronic recoil background originates from internal radioactive contaminations, including the β decays of ^{214}Pb , ^{85}Kr , material γ rays, and the neutrino-electron weak interaction of solar pp neutrinos [49]. The ER rates per SR are determined by fitting the already unblinded events in the $[1, 140]$ keV range in SR0 [49] and the nonblinded events in the $[10, 140]$ keV range in SR1. Because of the uncertainty of the shape of the ER background energy spectrum, a systematic uncertainty of 100% is assigned to the ER background, under the assumption of a flat ER energy spectrum in the ROI and utilizing the combined fit yield model from SR0 ^{220}Rn and ^{37}Ar calibration data [38]. The conservative uncertainty has minimal impact on our results due to the low rate of ER background. The prediction of ER background is (0.11 ± 0.11) events $\text{t}^{-1} \text{y}^{-1}$ in SR0 and (0.24 ± 0.24) events $\text{t}^{-1} \text{y}^{-1}$ in SR1, and the difference

between SRs is due to an increased concentration of ^{85}Kr at the start of SR1, which was subsequently removed by cryogenic distillation.

Inference—An extended binned likelihood function is constructed for the SR0 and SR1 data using four analysis dimensions: corrected S2 (cS2) [50], S1 BDT score, S2 BDT score, and $\text{S2}_{\text{pre}}/\Delta t_{\text{pre}}$. The bin edges are defined by binning the AC background in each dimension with equal probability, as shown in Fig. 2, resulting in a total of $3^4 = 81$ bins in the 4D histogram. The likelihood function is expressed as

$$\mathcal{L}(\sigma, \theta) = \mathcal{L}_{\text{SR0}}(\sigma, \theta) \times \mathcal{L}_{\text{SR1}}(\sigma, \theta) \times \prod_m \mathcal{L}_m^{\text{anc}}(\theta_m), \quad (1)$$

where σ represents the DM-nucleon interaction cross section, and θ are the Gaussian ancillary likelihood-constrained nuisance parameters. The index m iterates through the nuisance parameters, including the expected




FIG. 3. DM cross-section upper limit at 90% confidence level with the 1σ (green) and 2σ (yellow) sensitivity band. The results with and without sensitivity -1σ PCL are shown by solid and dashed lines, respectively. Previous results are shown for comparison [28,30,43,54–58], where limits from other liquid-xenon-based experiments are power-constrained to sensitivity -1σ . (a) Shows the constraint on SI-DM–nucleon interaction. The neutrino fog [6] is defined as $n = \max_{\sigma \in [\sigma, \infty]} [-(d \ln \sigma / d \ln MT)^{-1}]$, in which $n = 2, 2.5, 3$ from lighter to darker gray. The ^8B CE ν NS equivalent DM of $5.5 \text{ GeV}/c^2$ with cross section of $4.4 \times 10^{-45} \text{ cm}^2$ is shown. (b) Shows the constraints on SI-LM, where the differential rate is proportional to σm_ϕ^4 assuming $m_\phi \ll 10 \text{ MeV}/c^2$ [28,60]. (c) Shows the constraint on SD-DM–neutron interaction assuming the median of nuclear form factors from Ref. [59]. (d) Shows the constraint on MDDM with momentum dependence $n = 1, 2$ (red, blue), where σ_0 is the SI cross section. The 1 and 2σ sensitivity bands are depicted using shadow and dots, respectively.

rate of each background with their uncertainties shown in Table I, the signal efficiency with 25% uncertainty, and the NR yield model shape parameters (t_{Ly} , t_{Qy}).

Prior to unblinding, the sensitivity band was determined by computing the profile-likelihood-ratio test statistics between the signal-plus-background hypothesis and the background-only hypothesis using toy Monte Carlo datasets [35,51,52]. We adopt a power-constrained limit (PCL) [53] with a power threshold of -1σ to report the final limit. The power threshold is changed from our previous 50% threshold [5] to be in line with the limits reported by other experiments. A future publication will discuss the difference between PCL thresholds and further motivate this choice.

Results—After unblinding, SR0 and SR1 reveal 9 and 28 events, respectively. The discovery p value for the DM of SI interaction indicates no significant excess, with a minimum p value of 0.18 at 3 GeV/ c^2 . Their distributions in the search space are shown in Fig. 2. The background-only and signal-plus-background best fits for 4, 6, 10 GeV/ c^2 DMs are presented in Table I. Two goodness-of-fit tests, with a reporting p -value threshold of 0.025 determined prior to unblinding, are performed on the spatial X - Y dimension. The resulting p values, 0.018 and 0.037, suggest potential spatial clustering of events. However, the positions are not used in the inference framework defined prior to unblinding. Postunblinding checks that include inspection of the distribution in data selection space as well as the event waveforms show no indication of mismodeling.

The results for SI, SD, MDDM, and SI-LM, along with previous limits [28,30,43,54–58], are presented in Fig. 3 for comparison. Our results tighten the constraints across the parameter spaces for all listed DM models, with the SI limit at the benchmark 6 GeV/ c^2 DM mass reaching 2.5×10^{-45} cm 2 . An upward fluctuation is observed below 5 GeV/ c^2 and a downward fluctuation above this mass. The SD DM-neutron limit is susceptible to the current $\sim 10\%$ uncertainty of the nuclear form factors in Ref. [59]. As the uncertainty predominately changes the expected recoil rate, the results presented here can be rescaled. Additionally, we exclude all metallic components of mirror DM proposed by Refs. [32,33], specifically ruling out mirror oxygen with a 90% upper limit on the rate parameter $\epsilon\sqrt{\xi_{O'}}$ of 1.3×10^{-12} , with the theoretically favored region $10^{-11} \lesssim \epsilon\sqrt{\xi_{O'}} \lesssim 4 \times 10^{-10}$.

XENONnT achieves the world-leading sensitivity for light DM candidates within the ^8B CE ν NS neutrino fog, as illustrated by Fig. 3(a). The background-only model yields a best fit of $11.4^{+2.0}_{-3.6}$ ^8B CE ν NS events. In the 6 GeV/ c^2 model, ^8B CE ν NS emerges as the most significant background, accounting for 60% of the total background in the signal-like region, marking the first step into the neutrino fog. We have prepared a

dedicated data release to allow further interpretation of this work with different DM models and yield models in Ref. [61].

Acknowledgments—We gratefully acknowledge support from the National Science Foundation, Swiss National Science Foundation, German Ministry for Education and Research, Max Planck Gesellschaft, Deutsche Forschungsgemeinschaft, Helmholtz Association, Dutch Research Council (NWO), Fundacao para a Ciencia e Tecnologia, Weizmann Institute of Science, Binational Science Foundation, Région des Pays de la Loire, Knut and Alice Wallenberg Foundation, Kavli Foundation, JSPS Kakenhi, JST FOREST Program, and ERAN in Japan, Tsinghua University Initiative Scientific Research Program, DIM-ACAV+ Région Ile-de-France, and Istituto Nazionale di Fisica Nucleare. This project has received funding and support from the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie Grant Agreement No. 860881-HIDDeN. We gratefully acknowledge support for providing computing and data-processing resources of the Open Science Pool and the European Grid Initiative, at the following computing centers: the CNRS/IN2P3 (Lyon, France), the Dutch national e-infrastructure with the support of SURF Cooperative, the Nikhef Data-Processing Facility (Amsterdam, Netherlands), the INFN-CNAF (Bologna, Italy), the San Diego Supercomputer Center (San Diego, USA) and the Enrico Fermi Institute (Chicago, USA). We acknowledge the support of the Research Computing Center (RCC) at The University of Chicago for providing computing resources for data analysis. We thank the INFN Laboratori Nazionali del Gran Sasso for hosting and supporting the XENON project.

-
- [1] E. Aprile *et al.* (XENON Collaboration), The XENONnT dark matter experiment, *Eur. Phys. J. C* **84**, 784 (2024).
 - [2] L. Roszkowski, E. M. Sessolo, and S. Trojanowski, WIMP dark matter candidates and searches—current status and future prospects, *Rep. Prog. Phys.* **81**, 066201 (2018).
 - [3] G. Jungman, M. Kamionkowski, and K. Griest, Supersymmetric dark matter, *Phys. Rep.* **267**, 195 (1996).
 - [4] G. Bertone and D. Hooper, History of dark matter, *Rev. Mod. Phys.* **90**, 045002 (2018).
 - [5] E. Aprile *et al.* (XENON Collaboration), First dark matter search with nuclear recoils from the XENONnT experiment, *Phys. Rev. Lett.* **131**, 041003 (2023).
 - [6] C. A. J. O’Hare, New definition of the neutrino floor for direct dark matter searches, *Phys. Rev. Lett.* **127**, 251802 (2021).
 - [7] J. Billard, L. Strigari, and E. Figueroa-Feliciano, Implication of neutrino backgrounds on the reach of next generation dark matter direct detection experiments, *Phys. Rev. D* **89**, 023524 (2014).
 - [8] F. Ruppin, J. Billard, E. Figueroa-Feliciano, and L. Strigari, Complementarity of dark matter detectors in light of the neutrino background, *Phys. Rev. D* **90**, 083510 (2014).

- [9] D. S. Akerib *et al.*, Snowmass2021 cosmic frontier dark matter direct detection to the neutrino fog, in *Snowmass 2021* (2022), [arXiv:2203.08084](#).
- [10] J. Billard *et al.*, Direct detection of dark matter—APPEC committee report, *Rep. Prog. Phys.* **85**, 056201 (2022).
- [11] E. Aprile *et al.* (XENON Collaboration), First indication of solar ^8B neutrinos via coherent elastic neutrino-nucleus scattering with XENONnT, *Phys. Rev. Lett.* **133**, 191002 (2024).
- [12] E. Aprile *et al.* (XENON1T Collaboration), Conceptual design and simulation of a water Cherenkov muon veto for the XENON1T experiment, *J. Instrum.* **9**, P11006 (2014).
- [13] M. Murra, D. Schulte, C. Huhmann, and C. Weinheimer, Design, construction and commissioning of a high-flow radon removal system for XENONnT, *Eur. Phys. J. C* **82**, 1104 (2022).
- [14] E. Aprile *et al.* (XENON Collaboration), Removing krypton from xenon by cryogenic distillation to the ppq level, *Eur. Phys. J. C* **77**, 275 (2017).
- [15] J. Aalbers *et al.*, AxFoundation/strax: Stream analysis for xenon tpcs, [10.5281/zenodo.11355772](#) (2024).
- [16] XENON Collaboration, XENONnT/straxen: Streaming analysis for xenon, [10.5281/zenodo.12608732](#) (2024).
- [17] A. Tomás, H. M. Araújo, A. J. Bailey, A. Bayer, E. Chen, B. López Paredes, and T. J. Sumner, Study and mitigation of spurious electron emission from cathodic wires in noble liquid time projection chambers, *Astropart. Phys.* **103**, 49 (2018).
- [18] D. S. Akerib *et al.* (LUX Collaboration), Investigation of background electron emission in the LUX detector, *Phys. Rev. D* **102**, 092004 (2020).
- [19] K. Petraki and R. R. Volkas, Review of asymmetric dark matter, *Int. J. Mod. Phys. A* **28**, 1330028 (2013).
- [20] D. E. Kaplan, M. A. Luty, and K. M. Zurek, Asymmetric dark matter, *Phys. Rev. D* **79**, 115016 (2009).
- [21] D. N. Spergel and P. J. Steinhardt, Observational evidence for selfinteracting cold dark matter, *Phys. Rev. Lett.* **84**, 3760 (2000).
- [22] S. Tulin and H.-B. Yu, Dark matter self-interactions and small scale structure, *Phys. Rep.* **730**, 1 (2018).
- [23] K. M. Zurek, Multi-component dark matter, *Phys. Rev. D* **79**, 115002 (2009).
- [24] N. Fornengo, P. Panci, and M. Regis, Long-range forces in direct dark matter searches, *Phys. Rev. D* **84**, 115002 (2011).
- [25] J. D. Lewin and P. F. Smith, Review of mathematics, numerical factors, and corrections for dark matter experiments based on elastic nuclear recoil, *Astropart. Phys.* **6**, 87 (1996).
- [26] J. Menendez, D. Gazit, and A. Schwenk, Spin-dependent WIMP scattering off nuclei, *Phys. Rev. D* **86**, 103511 (2012).
- [27] E. Del Nobile, M. Kaplinghat, and H.-B. Yu, Direct detection signatures of self-interacting dark matter with a light mediator, *J. Cosmol. Astropart. Phys.* **10** (2015) 055.
- [28] E. Aprile *et al.* (XENON Collaboration), Light dark matter search with ionization signals in XENON1T, *Phys. Rev. Lett.* **123**, 251801 (2019).
- [29] X. Ren *et al.* (PandaX-II Collaboration), Constraining dark matter models with a light mediator at the PandaX-II experiment, *Phys. Rev. Lett.* **121**, 021304 (2018).
- [30] S. Li *et al.* (PandaX Collaboration), Search for light dark matter with ionization signals in the PandaX-4T experiment, *Phys. Rev. Lett.* **130**, 261001 (2023).
- [31] S. Chang, A. Pierce, and N. Weiner, Momentum dependent dark matter scattering, *J. Cosmol. Astropart. Phys.* **01** (2010) 006.
- [32] R. Foot, Mirror dark matter: Cosmology, galaxy structure and direct detection, *Int. J. Mod. Phys. A* **29**, 1430013 (2014).
- [33] J. D. Clarke and R. Foot, Mirror dark matter will be confirmed or excluded by XENON1T, *Phys. Lett. B* **766**, 29 (2017).
- [34] J. Aalbers, B. Pelssers, J. R. Angevaere, and K. D. Morá, Jelleaalbers/wimprates: v0.5.0, [10.5281/zenodo.7636982](#) (2023).
- [35] D. Baxter *et al.*, Recommended conventions for reporting results from direct dark matter searches, *Eur. Phys. J. C* **81**, 907 (2021).
- [36] E. Aprile *et al.*, Low-energy nuclear recoil calibration of XENONnT with a ^{88}Y Be photoneutron source, [arXiv:2412.10451](#).
- [37] J. I. Collar, Applications of an ^{88}Y /Be photo-neutron calibration source to dark matter and neutrino experiments, *Phys. Rev. Lett.* **110**, 211101 (2013).
- [38] E. Aprile *et al.* (XENON Collaboration), XENONnT WIMP search: Signal & background modeling and statistical inference, [arXiv:2406.13638](#).
- [39] M. Szydagis, N. Barry, K. Kazkaz, J. Mock, D. Stolp, M. Sweany, M. Tripathi, S. Uvarov, N. Walsh, and M. Woods, NEST: A comprehensive model for scintillation yield in liquid xenon, *J. Instrum.* **6**, P10002 (2011).
- [40] E. Aprile *et al.*, XENONnT analysis: Signal reconstruction, calibration, and event selection, [arXiv:2409.08778](#).
- [41] XENON Collaboration, XENONnT/fuse: Refactor xenonnt epix and wfsim code, [10.5281/zenodo.11551366](#) (2024).
- [42] XENON Collaboration, XENONnT/axidence: Strax-based data-driven accidental coincidence background simulation and peak-level salting, [10.5281/zenodo.12791105](#) (2024).
- [43] E. Aprile *et al.* (XENON Collaboration), Search for coherent elastic scattering of solar ^8B neutrinos in the XENON1T dark matter experiment, *Phys. Rev. Lett.* **126**, 091301 (2021).
- [44] B. Aharmim *et al.* (SNO Collaboration), Combined analysis of all three phases of solar neutrino data from the sudbury neutrino observatory, *Phys. Rev. C* **88**, 025501 (2013).
- [45] J. N. Bahcall, E. Lisi, D. E. Alburger, L. De Braekeleer, S. J. Freedman, and J. Napolitano, Standard neutrino spectrum from B-8 decay, *Phys. Rev. C* **54**, 411 (1996).
- [46] C. H. Faham, V. M. Gehman, A. Currie, A. Dobi, P. Sorensen, and R. J. Gaitskell, Measurements of wavelength-dependent double photoelectron emission from single photons in VUV-sensitive photomultiplier tubes, *J. Instrum.* **10**, P09010 (2015).
- [47] D. S. Akerib *et al.* (LZ Collaboration), Enhancing the sensitivity of the LUX-ZEPLIN (LZ) dark matter experiment to low energy signals, [arXiv:2101.08753](#).
- [48] P. Sorensen, Anisotropic diffusion of electrons in liquid xenon with application to improving the sensitivity of direct

- dark matter searches, *Nucl. Instrum. Methods Phys. Res., Sect. A* **635**, 41 (2011).
- [49] E. Aprile *et al.* (XENON Collaboration), Search for new physics in electronic recoil data from XENONnT, *Phys. Rev. Lett.* **129**, 161805 (2022).
- [50] E. Aprile *et al.* (XENON Collaboration), XENONnT analysis: Signal reconstruction, calibration and event selection, [arXiv:2409.08778](#).
- [51] G. J. Feldman and R. D. Cousins, A unified approach to the classical statistical analysis of small signals, *Phys. Rev. D* **57**, 3873 (1998).
- [52] XENON Collaboration, XENONnT/alea: A tool to perform toymc-based inference constructions, [10.5281/zenodo.10829030](#) (2024).
- [53] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, Power-constrained limits, [arXiv:1105.3166](#).
- [54] J. Aalbers *et al.* (LZ Collaboration), First dark matter search results from the LUX-ZEPLIN (LZ) experiment, *Phys. Rev. Lett.* **131**, 041002 (2023).
- [55] W. Ma *et al.* (PandaX Collaboration), Search for solar B8 neutrinos in the PandaX-4T experiment using neutrino-nucleus coherent scattering, *Phys. Rev. Lett.* **130**, 021802 (2023).
- [56] G. Angloher *et al.* (CRESST Collaboration), Limits on momentum-dependent asymmetric dark matter with CRESST-II, *Phys. Rev. Lett.* **117**, 021303 (2016).
- [57] P. Agnes *et al.* (DarkSide-50 Collaboration), Search for low-mass dark matter WIMPs with 12 ton-day exposure of DarkSide-50, *Phys. Rev. D* **107**, 063001 (2023).
- [58] D. Huang *et al.* (PandaX Collaboration), Search for dark-matter–nucleon interactions with a dark mediator in PandaX-4T, *Phys. Rev. Lett.* **131**, 191002 (2023).
- [59] P. Klos, J. Menéndez, D. Gazit, and A. Schwenk, Large-scale nuclear structure calculations for spin-dependent WIMP scattering with chiral effective field theory currents, *Phys. Rev. D* **88**, 083516 (2013).
- [60] E. D. Nobile, M. Kaplinghat, and H.-B. Yu, Direct detection signatures of self-interacting dark matter with a light mediator, *J. Cosmol. Astropart. Phys.* **10** (2015) 055.
- [61] L. Yuan and S. Shi, XENONnT/light_wimp_data_release: v0.0.1, Zenodo v0.0.1 (2024), [10.5281/zenodo.14522707](#).