

This is a repository copy of *Long-term* exposure to residential greenness and cardiovascular disease and all-cause mortality in China.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/226654/</u>

Version: Published Version

Article:

Cao, X., Tuerdi, N., Tang, H. orcid.org/0000-0002-2924-0126 et al. (8 more authors) (2025) Long-term exposure to residential greenness and cardiovascular disease and all-cause mortality in China. BMC Public Health, 25 (1). 1645. ISSN 1472-698X

https://doi.org/10.1186/s12889-025-22899-7

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk https://eprints.whiterose.ac.uk/

RESEARCH





Long-term exposure to residential greenness and cardiovascular disease and all-cause mortality in China

Xue Cao^{1†}, Nuerguli Tuerdi^{1†}, Haosu Tang^{2,3}, Yujie Zhang¹, Xin Wang¹, Congyi Zheng¹, Yixin Tian¹, Xue Yu¹, Xuyan Pei¹, Gang Huang^{2,3} and Zengwu Wang^{1,4*}

Abstract

Background Influence of residential greenness on CVD risk has garnered increasing attention, however, evidence from large-scale cohort studies in developing nations, such as China, remains sparse. This study aimed to evaluate the association of residential greenness with CVD and mortality, and explore the potential mediating role of modifiable risk factors in the associations.

Methods A total of 22,702 participants aged 35 years and above were enrolled between October 2012 and December 2015. Residential greenness was assessed using Normalized Difference Vegetation Index (NDVI) within radii of 300 m, 500 m, and 1000 m from participants' residential address (NDVI_{300 m}, NDVI_{500 m}, and NDVI_{1000 m}). Primary outcomes comprised CVD events and all-cause mortality, with follow-up from 2018 to 2019. Multivariable Cox regression models were employed to estimate hazard ratios (HRs), and causal mediation analysis was conducted to assess the role of modifiable risk factors in the observed associations.

Results Residential greenness demonstrated a significant association with the risk of CVD, with HRs per tertile increment of 0.84 (95% confidential interval [CI]: 0.77–0.92) for NDVI_{300 m}, 0.86 (95% CI: 0.79–0.94) for NDVI_{500 m}, and 0.90 (95% CI: 0.82–0.98) for NDVI_{1000 m}, separately. Compared to areas with the lowest NDVI_{500 m}, the HR for CVD incidence in areas with medium and high NDVI_{500 m} were 0.89 (95% CI: 0.76–1.06), and 0.74 (95% CI: 0.62–0.89), respectively. Utilizing a newly proposed two-stage regression method in mediation analysis, approximately 16.18%, 5.34%, 4.04%, and 2.45% of the total effect of NDVI_{500 m} on CVD risk were mediated by high-density lipoprotein cholesterol, physical activity, body mass index, and diabetes mellitus, respectively.

Conclusion This study provides compelling evidence that higher residential greenness is associated with a reduced risk of CVD among the adult Chinese population, with specific modifiable risk factors playing a mediating role. These findings underscore the significance of incorporating green space interventions into CVD prevention strategies.

Keywords Cardiovascular disease, Greenness, Mediation analysis, Epidemiology

[†]Xue Cao and Nuerguli Tuerdi contributed equally to this work.

*Correspondence: Zengwu Wang wangzengwu@foxmail.com Full list of author information is available at the end of the article



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by-nc-nd/4.0/.

Introduction

Cardiovascular diseases(CVD) remain a major causes of death in the world, particularly in low- and middleincome countries [1]. In 2020, there were 4.58 million deaths attributed to CVD in China [2], with a predicted growth rate of CVD events exceeding 50% between 2010 and 2030, solely based on population aging and growth, assuming that risk factors remain unchanged [3]. Moreover, prior studies have already demonstrated that some common and well known risk factors, such as smoking, obesity and hypertension, are associated with the risk of CVD incidence and deaths [4], and effective intervention concerning lifestyle modifications can significantly improve the control rate of blood pressure [5, 6]. At the same time, there is an increasing recognition that environmental elements, such as built environment and air pollution, can influence individual behavioral and metabolic factors, as well as the CVD incidence, also potentially serving as significant factors in accurate CVD risk assessment and prediction.

In recent years, residential greenness has attracted significant global attention for its potential to enhance public health. Scientific study showed that living in greener areas, or even brief exposure to vegetation, can improve both physical and mental well-being [7]. These benefits stem from several factors, including reducing stress and air pollution, increasing convenience of engaging in physical activities, and enhancing social cohesion [8–10]. Collectively, these effects contribute to a lower risk of chronic diseases, particularly cardiovascular conditions. Previous research has indicated that individuals residing in areas characterized by higher residential greenness face a reduced risk of myocardial infarction, heart failure, ischemic stroke, high blood pressure, and dyslipidemia [11–18]. However, most investigations were conducted in middle- and high-income developed countries, resulting in a scarcity of evidence from large-scale population-based cohort studies in developing nations, such as China, where the prevalence of CVD and cardiometabolic disorders have surged. Furthermore, to our knowledge, only a limited number of studies within the Chinese population has explored the association between residential greenness and CVD risk [19, 20]. Nonetheless, those investigations adopted cross-sectional designs. Moreover, only a few studies have explored the potential mediation effect of hypertension, diabetes, lipid levels, body mass index (BMI), physical activity, and air pollution on the association between greenness and CVD risk, yielding inconsistent findings [19-22]. In the context of China's rapid economic development and the accelerated surge in certain modifiable risk factors and the associated burden of CVD, it becomes particularly imperative to scrutinize the association between greenness and both CVD incidence and mortality risk. This necessitates a comprehensive investigation based on a large-scale population-based cohort study, focusing on discerning potential factors that may influence this association.

Therefore, in this present study, we aimed to evaluate the association of residential greenness with CVD and mortality among Chinese adults based on a large-scale longitudinal study, and explore whether modifiable risk factors could mediate the relationship between residential greenness and CVD.

Methods

Data source and study population

In this study, individuals were selected from the China Hypertension Survey (CHS), a nationally representative population-based survey conducted from October 2012 to December 2015. Briefly, a stratified multistage random sampling method was used to recruit nearly 0.5 million participants aged 15 years and older from 31 provinces in mainland China, more information was provided elsewhere [23]. For this sub-study, all selected urban and rural areas were further stratified into eastern, central, and western regions based on geographical location and economic levels. Then a total of 16 cities and 17 counties were selected using simple random sampling (SRS), and at least three communities or villages were randomly selected from each region [24, 25]. Ultimately, 30,036 individuals aged \geq 35 years, with essential baseline information at baseline (including gender, name, and valid blood samples), were enrolled initially. The followed up was conducted in 2018 and in 2019, and the outcomes were identified from baseline (2012-2015) to the followup survey (2018–2019).

In this current study, we first excluded participants lost to follow-up (n = 4,726). Next, we further removed 2,135 subjects with a history of CVD and 473 participants with incomplete baseline covariate data. Finally, 24.4% participants were excluded from the original population, and 22,702 participants with no CVD and had complete baseline information on hypertension, diabetes, and blood lipid levels were included in the final analysis. (Figure S1 in supplementary materials 1). This study was approved by the ethics committee of Fuwai Hospital, Beijing, China, and all participants provided written informed consent.

Assessment of residential greenness levels

Normalized Difference Vegetation Index (NDVI) was used to quantify residential greenness in the 300 m, 500 m and 1000 m radius around the participant's residential addresses. The NDVI is a vegetation indicator, which is defined as (NIR—RED)/(NIR + RED), where NIR represents the near-infrared reflectance and RED signifies red reflectance. The NDVI ranges from -1 to 1, with positive values indicating vegetation, zero indicating bare soil, and negative values representing non-vegetated surfaces, such as blue spaces or water bodies. The higher the positive NDVI values, the greater the level of green vegetation. Monthly satellite-derived NDVI datasets across China, archived in the SPOT/VEGETATION PROBA-V 100 m products [26], were used in this study. These datasets have a spatial resolution of 100 m \times 100 m and were derived by calculating the monthly maximum values. We assessed the residential greenness levels via averaging the values within each participant's boundaries. Furthermore, regarding that water surfaces or blue spaces may have independent beneficial effects on health, negative NDVI values were removed in our analysis [27]. We calculated the three-year average NDVI before baseline survey to reflect their long-term exposure to green environments. To take into account seasonal variability in greenspace, in the sensitivity analysis, the average NDVI values between May and September were also computed to maximise the contrast in estimated exposure since the summer months are the greenest months in China. Additionally, NDVI of 500 m radii around participant's residential addresses were stratified into tertiles based on established research practice [27], with ranges of < 0.33 (Q1, Low), 0.33-0.44 (Q2, Medium), and >0.44 (Q3, High), respectively.

Mediating factors and covariates

Hypertension, diabetes mellitus, overweight/obesity, dyslipidemia, physical activity and fine particulate matter concentration ($PM_{2.5}$, particles with an aerodynamic diameter of $\leq 2.5 \,\mu\text{m}$) were considered as potential mediators of greenness and CVD incidence. In this study, hypertension was defined as a systolic blood pressure (SBP) \geq 140 mmHg, and/or a diastolic blood pressure (DBP) \geq 90 mmHg, and/or self-reported taking antihypertensive medication at baseline. According to the 2017 standards of the Chinese Medical Association's Diabetes Study Group, diabetes was defined as a fasting plasma glucose (FPG) level reaching or exceeding 7.0 mmol/L (126 mg/dL), or a self-reported diagnosis of type 2 diabetes by a healthcare professional, or being on anti-diabetic medication [28]. Overweight and obesity were defined as a body mass index (BMI) of 24.0 and \geq 28.0 kg/m², respectively. Participants with total cholesterol (TC) \geq 6.22 mmol/L, triglyceride (TG) \geq 2.26 mmol/L, low-density lipoprotein cholesterol (LDL-C) >4.14 mmol/L, or high-density lipoprotein cholesterol (HDL-C) <1.04 mmol/L were considered as having dyslipidemia based on the Chinese Guidelines for Lipid Management (2023) [29]. Additionally, physical activity was defined by metabolic equivalent tasks (METs) based on the participants'occupation/housework, commuting,

and leisure time activities, and low physical activity was defined as <600 METs/week. Average $PM_{2.5}$ concentrations in the year before CVD incidence or death were also included. By linking each participant's residence by residential counties/districts to average $PM_{2.5}$ concentrations in a 1 km ×1 km grid, which has been cross-validated [30]. Drinker was defined as consuming at least 1 alcoholic beverage per week in the past month. Daily maximum temperature data was obtained from meteorological station within each district or county. Heatwaves were defined as daily maximum temperatures exceeding the 92.5 th percentile of the warm season for at least three consecutive days.

We adjusted the following variables that were likely to confound the association based on the questionnaire, including age, sex, level of education (junior high school or below and high school or above), urbanization (urban/ rural), and geographic region (eastern, central, or western), alcohol consumption (drinker or non-drinker), smoking status (never smoked, former smoker, or current smoker), family history of CVD, and anti-dyslipidemia medication receipt. Moreover, we also included per capita Gross Domestic Product (GDP) to represent socioeconomic status (SES) at district/county levels, which were collected from statistic yearbooks, national bureau of statistics and China national knowledge infrastructure.

Outcome measures

The primary outcomes were composite of CVD events and mortality at any point before the follow-up survey conducted in 2018-2019, including coronary heart disease (CHD, International Classification of Diseases 10 th Revision [ICD-10] codes I20-I25), stroke (I60-I61 and I63-I64), chronic heart failure (I50), and deaths attributable to CVD (I00-I25, I27-I88, and I95-99). Specifically, CHD was defined as having one of the following conditions or treatments: myocardial infarction, coronary artery bypass graft surgery, or percutaneous coronary intervention coronary heart disease-related deaths, or CHD-related deaths. Stroke encompassed both non-fatal and fatal stroke, including subarachnoid hemorrhage, intracerebral hemorrhage, ischemic stroke, and unspecified stroke. During the follow-up, we tracked CVD events by contacting participants or their proxies through faceto-face interviews, telephone calls, or mailed questionnaires. and further verified these events by reviewing medical records or death certification. More detailed information on the questionnaires is in the supplementary materials 2.

Statistical analysis

In this study, descriptive statistics were carried out for all variables of interest. Continuous variables were represented by means with standard deviations (SDs), and categorical data by frequency and percentages. The association of greenness levels with CVD and mortality was estimated by multivariable Cox proportional hazards models, adjusting for age, sex, level of education, geographical region, urbanization, alcohol consumption, smoking status, family history of CVD, and per GDP. Hazard ratios (HRs) and 95% confidential interval (CI) were computed. Moreover, the Cox proportional hazard assumption was tested by weighted Schoenfeld residuals and P-values were greater than 0.05 for all outcomes. The survival time was measured in years from the date of initial interview administration to the occurrence date of study endpoint events. Subjects meeting any of the following criteria were treated as censored observations: (1) documented history of CVD at baseline assessment; (2) loss to follow-up during the study period; or (3) incomplete baseline covariate measurements essential for analysis. Median values were earmarked to each group of NDVI levels and were treated as continuous variables when performing trend testing. Exposureresponse curves between $\mathrm{NDVI}_{500\,\mathrm{m}}$ and incident CVD and mortality risk were plotted using penalized splines with two degrees of freedom. Furthermore, stratified analyses were conducted to explore variation in the association of greenness levels per tertile increment with CVD by sex, age and urbanization, and we also added a cross-product term between green and stratified variables into the fully adjusted model to test for interaction on the multiplicative scale.

Finally, we conducted mediation analysis to evaluate the effect of NDVI levels in 500 m radii on outcomes (expressed as the HR per tertile increase in NDVI) described by VanderWeele for Cox proportional hazards regression, keeping cardiometabolic risk factors, physical activity, PM₂₅ concentration, and heatwave days as mediator variables [31, 32]. Regarding that the 500 m buffer can cover more green space the participants may use, we used it as the main exposure to explore the mediation analyses. Briefly, the total effect was decomposed into the natural direct effect (NDE) and natural indirect effect (NIE), where NDE represents the effect size of NDVI unexplained through the mediator, while NIE is interpreted as the effect size of NDVI resulting from mediation through the specified mediators. The proportion of the association of NDVI with CVD events and all-cause mortality mediated via the designated mediator, was computed as a metric of the contribution of the natural indirect association to the overall association, employing the following formula: [32]

Moreover, all of the 95% CIs for the total effect, NDE, NIE and the proportion mediated were calculated using the delta method, and the mediation effects of each potential mediator were assessed separately in this current study.

Penalized splines analysis was conducted using the *survival* package, version 3.4.2 for R software (R Foundation for Statistical Computing, Vienna, Austria). All other analyses were performed with SAS version 9.4 (SAS Institute Inc., Cary, NC USA). *Mediation* macro for SAS was applied to perform the mediation analysis. All statistical tests were 2-sided at a significance level of P < 0.05.

Results

Baseline characteristics

The baseline characteristics according to different levels of NDVI (low, medium, high) are presented in Table 1. Among 22 702 adult participants initially free of CVD, the mean baseline age was 56.1 \pm 13.1 years. During the follow-up, 993 participants developed fatal or nonfatal CVD and 1096 died from all causes (Figure S1 in supplementary materials 1). The cumulative incidence of CVD and all-cause mortality were 9.45 and 10.81 per 1000 person-years, respectively (Table 2). Participants resided in areas with higher greenness were more likely to lower BMI levels, and from rural areas. The median (inter-quartile range) of average 3-year NDVI exposure in circular buffers of 300 m, 500 m and 1000 m around each address's centroid were 0.39 (0.18), 0.39 (0.19), and 0.38 (0.20), respectively. Figure 1 displays the geographical distributions of average NDVI levels from 2012-2015 in China, with substantial variations in greening across the entire surveyed region.

Association of greenness with CVD and all-cause mortality

Table 2 presents the association between different levels of NDVI and CVD incidence as well as all-cause mortality. In the multivariate adjustment model, we found that higher levels of greenness (measured by NDVI) were associated with greater protective effects (i.e. beneficial effect) for CVD. Specifically, for per tertile increment in NDVI_{300 m}, NDVI_{500 m}, and NDVI_{1000 m}, the risk of total CVD reduced by 16% (HR =0.84 [95% CI: 0.77–0.92]), 14% (HR =0.86 [95% CI: 0.79–0.94)), and 10% (HR =0.90 [95% CI: 0.82–0.98]), respectively. This trend towards reducing the CVD risk was significant for the classes with elevated levels of NDVI ($P_{trend} < 0.05$).

 $Proportion mediated = HR^{NDE} \times (HR^{NIE} - 1)/(HR^{NDE} \times HR^{NIE} - 1)$

Table 1 Characteristics of study participants

Characteristic	Total (N = 22,702)	NDVI _{500 m}		P value	
		Low (N= 7458)	Medium (N = 7712)	High (N= 7532)	
Socio-demographics					
Age, years	56.1 ± 13.1	55.7 ± 13.3	55.3 ± 12.7	57.4 ± 13.2	<.001
Gender, %					0.558
Male	10,505 (46.3)	3489 (46.8)	3546 (46.0)	3470 (46.1)	
Female	12,197 (53.7)	3969 (53.2)	4166 (54.0)	4062 (53.9)	
Educational level, %					<.001
Elementary middle School or lower	11,593 (51.1)	2537 (34.0)	4246 (55.1)	4810 (63.9)	
High school or above	11,109 (48.9)	4921 (66.0)	3466 (44.9)	2722 (36.1)	
Residence, %					<.001
Urban	10,130 (44.6)	5300 (71.1)	2556 (33.1)	2274 (30.2)	
Rural	12,572 (55.4)	2158 (28.9)	5156 (66.9)	5258 (69.8)	
Region, %					<.001
East	9263 (40.8)	3149 (42.2)	3313 (43.0)	2801 (37.2)	
Central	9460 (41.7)	2322 (31.1)	3426 (44.4)	3712 (49.3)	
West	3979 (17.5)	1987 (26.6)	973 (12.6)	1019 (13.5)	
Lifestyle behaviors					
Consumption of Alcohol, %	6318 (27.8)	2081 (27.9)	2123 (27.5)	2114 (28.1)	0.749
Smoking types, %					<.001
Current	5731 (25.2)	1599 (21.4)	2068 (26.8)	2064 (27.4)	
Former	1216 (5.4)	379 (5.1)	448 (5.8)	389 (5.2)	
Never	15,755 (69.4)	5480 (73.5)	5196 (67.4)	5079 (67.4)	
Mediators					
Low physical activity (< 600 METs/week)	4860 (21.4)	1951 (26.2)	1338 (17.3)	1571 (20.9)	<.001
Hypertension	8957 (39.5)	3037 (40.7)	2931 (38.0)	2989 (39.7)	0.003
Diabetes Mellitus	2286 (10.1)	908 (12.2)	707 (9.2)	671 (8.9)	<.001
BMI, Kg/m ²	24.5 ± 3.5	25.0 ± 3.5	24.4 ± 3.5	24.1 ± 3.4	<.001
Triglycerides, mmol/L	1.4 ± 1.0	1.5 ± 1.0	1.5 ± 1.0	1.4 ± 1.0	
Cholesterol, mmol/L					
Total	4.8 ± 1.0	4.7 ± 1.0	4.8 ± 1.0	4.8 ± 1.0	<.001
HDL-C	1.4 ± 0.3	1.3 ±0.3	1.4 ±0.3	1.4 ± 0.4	<.001
LDL-C	2.8 ± 0.8	2.7 ±0.8	2.8 ± 0.8	2.8 ± 0.8	<.001
FPG, mmol/L	5.6 ± 1.5	5.6 ± 1.7	5.6 ± 1.4	5.5 ± 1.4	<.001
PM _{2.5} , μg/m ³	53.4 ± 13.2	54.0 ± 12.7	50.8 ± 12.7	55.3 ± 13.8	<.001
GDP per capita, yuan	65,339.0 ± 36,929.8	70,443.6 ± 43,915.4	66,955.6 ± 36,939.3	58,629.2 ± 27,032.1	<.001

NDVI normalized difference vegetation index, BMI body mass index, HDL-C high-density lipoprotein cholesterol, LDL-C low density lipoprotein cholesterol, FPG fasting plasma glucose, MET metabolic equivalent tasks, PM particulate matter, GDP gross domestic product

Furthermore, compared to areas with the lowest of NDVI per 500 m, the HR for the incidence of total CVD in areas with medium and high NDVI_{500 m} were 0.89 (95% CI: 0.76–1.06), and 0.74 (95% CI: 0.62–0.89), respectively. For CHD outcome, we also found significant association, with estimated HRs of 0.55 (95% CI: 0.40–0.75) and 0.67 (95% CI: 0.49–0.90) for participant living in the medium and high NDVI in 500 m radii. Similar results were observed for NDVI levels per 300 m and 1000 m. Moreover, the correlation between greenness and stroke

was interesting, because we found that residential greenness in the 300 m radius could reduce the risk of stroke, but no statistically significant difference was observed for NDVI_{1000 m}. Also, no important associations were observed between overall greenness and all-cause mortality. The exposure–response curves based on penalized spline models revealed that the risk of CVD decreased with the increment of NDVI_{500 m}, and the HR for the association between NDVI_{500 m} and total CVD, and

	CVD		CHD		Stroke		All-cause mortality		
Exposure		Incidence/1000 person-years	HR (95%CI)						
NDVI _{300 m}	Low	10.39	[Reference]	3.91	[Reference]	5.40	[Reference]	9.92	[Reference]
	Medium	9.00	0.79(0.66– 0.93)	1.88	0.45(0.32– 0.62)	6.20	1.02(0.81– 1.27)	10.35	0.97(0.82– 1.15)
	High	9.12	0.70(0.59– 0.84)	2.84	0.60(0.44- 0.81)	5.33	0.77(0.61– 0.98)	12.09	0.88(0.74– 1.04)
	Per tertile increment ^a	9.45	0.84(0.77– 0.92)	2.86	0.79(0.67– 0.92)	5.64	0.87(0.77– 0.98)	10.81	0.93(0.86– 1.01)
	P for trend		<.001		0.002		0.020		0.100
NDVI _{500 m}	Low	10.02	[Reference]	3.70	[Reference]	5.20	[Reference]	9.73	[Reference]
	Medium	9.51	0.89(0.76– 1.06)	2.07	0.55(0.40– 0.75)	6.62	1.15(0.92– 1.43)	10.92	1.05(0.89– 1.25)
	High	8.84	0.74(0.62– 0.89)	2.85	0.67(0.49– 0.90)	5.06	0.80(0.63– 1.02)	11.76	0.90(0.76– 1.07)
	Per tertile increment ^a		0.86(0.79– 0.94)		0.83(0.70– 0.97)		0.88(0.79– 0.99)		0.94(0.86– 1.02)
	P for trend		<.001		0.016		0.038		0.125
NDVI _{1000 m}	Low	9.97	[Reference]	3.63	[Reference]	5.24	[Reference]	9.68	[Reference]
	Medium	9.30	0.83(0.70– 0.98)	1.93	0.50(0.36– 0.68)	6.53	1.05(0.84– 1.30)	10.73	1.01(0.86– 1.20)
	High	9.10	0.80(0.66– 0.96)	3.17	0.79(0.58– 1.08)	5.00	0.82(0.64– 1.04)	12.07	0.94(0.79– 1.12)
	Per tertile increment ^a		0.90(0.82– 0.98)		0.91(0.77– 1.07)		0.90(0.80- 1.01)		0.97(0.89– 1.05)
	P for trend		0.017		0.217		0.077		0.426

	Table 2	Hazard rati	io for CVE) and all-	-cause mortality	/ associated	with	residential	greenness
--	---------	-------------	------------	------------	------------------	--------------	------	-------------	-----------

Notes: All models were adjusted for age, sex, alcohol consumption, urbanization and geographical region, smoking, educational level, family history of CVD, CVD medication history, and per capita GDP

Abbreviations: HR hazard ratio, CVD cardiovascular disease, NDVI normalized difference vegetation index, CHD coronary heart disease, HR hazard ratio, CI confidential interval

^a NDVI level per tertile increment



Colors: Annual mean NDVI across China for 2012-2015

Fig. 1 Annual mean NDVI (colors) at baseline (2012–2015) and 38 China Hypertension Survey sites (red dots) across China. Insets: South China Sea. Abbreviations: NDVI, Normalized Difference Vegetation Index



Fig. 2 Exposure–response relations between long term exposure to NDVI within the 500 m circular buffers around the participants'residential addresses and incidence of CVD and all-cause mortality. Solid line represents point estimate and dashed lines represent 95% confidence intervals. Boxplot represents distribution of NDVI exposure levels for participants. NDVI, normalized difference vegetation index; CVD, cardiovascular disease

all-cause mortality were non-linear (*P*-nonlinear < 0.05), respectively (Fig. 2).

Stratified analysis

In the stratified analyses (Fig. 3), the association of a tertile increase in NDVI_{300 m} and NDVI_{500 m} with the risk of CVD did not differ substantially by age and sex. However, the association differed according to urbanization. In particular, we observed a different association between NDVI_{300 m} and CVD risk in the rural areas (HR = 0.71; 95%CI:0.63, 0.80) and in urban area (HR = 0.97;95%CI:0.85,1.11), and the *P* interaction lower than 0.001. Similar results were observed for NDVI_{500 m} and NDVI_{1000 m}. Additionally, participants younger than 60 had a greater significant modification effect of NDVI_{1000 m}, and the *P* value of multiplicative interaction was 0.034.

Mediation analysis

Mediation analysis showed that diabetes mellitus, BMI continuous, HDL-C, and physical activity played mediation effects on the association between NDVI_{500 m} and CVD. In contrast, no significant mediation effect was found for hypertension and $PM_{2.5}$ (Table 3). Specifically, an increase in per tertile for NDVI_{500 m} reduced the risk of CVD by 13% (HR [total association], 0.87; 95%CI, 0.79–0.95), and that approximately 16.18% of the total association], 0.98; 95%CI, 0.96–0.99). Also, Table 3 showed that an increase in NDVI_{500 m} decreased the risk of CVD by 15% (HR [total association], 0.85; 95% CI, 0.78–0.93), and that 5.34% of the total association (95%)

CI, 2.01%–13.55%) was mediated through the physical activity (HR [indirect association], 0.99; 95% CI, 0.99–1.00, P= 0.001). Moreover, proportions mediated were 4.04% for BMI, and 2.45%(95%CI, 35.5%–64.2%) for diabetes mellitus. Additionally, no mediation effects of hypertension, diabetes mellitus, and lipids were found to play a significant role in the association of NDVI_{500 m} with CHD and all-cause mortality (P> 0.05) (Table S1–2 in supplementary materials 1). However, in the mediation analyses of NDVI and stroke risk, physical activity showed the most significant mediation effects (Table S1 in supplementary materials 1).

Discussion

In this prospective cohort study in China, we observed that individuals residing in higher residential greenness, as measured by satellite-derived NDVI_{300 m} and NDVI₅₀₀ m, had a reduced risk of CVD incidence, whereas non-significant relationship was found between residential greenness and the risk of all-cause mortality. Additionally, the association between residential greenness and the risk of CVD was partially mediated by diabetes mellitus, BMI, HDL-C, and physical activity.

Findings in this study of negative associations between greenness level and the risk of CVD incidence were supported by previous research [11–13, 17, 18]. The longitudinal study conducted in Korea, enrolling 351 409 participant aged over 20 years, demonstrated that living in urban area with higher level of greenness may beneficially affect the risk of total CVD [18]. Another study combining four large population-based cohorts in

Subgroup	HR (95% CI)		P value
NDVI _{300m}			
Age, vears			0.066
< 60	0.70 (0.57-0.85)	⊢∎(
≥ 60	0.90 (0.81-0.99)	⊢∎ -1	
Gender			0.156
Male	0.81 (0.72-0.91)	⊢∎ -4	
Female	0.89 (0.78-1.02)	⊢∎⊣	
Residence			<.001
Urban	0.97 (0.85-1.11)	⊢ ∎	
Rural	0.71 (0.63-0.80)	⊨∎⊣	
NDVI _{500m}			
Age, years			0.131
< 60	0.73 (0.60-0.89)	┝━╋━┥	
≥ 60	0.90 (0.81-0.99)	⊢∎⊣	
Gender			0.104
Male	0.82 (0.73-0.92)	+∎-1	
Female	0.93 (0.81-1.06)	⊢∎⊣	
Residence			<.001
Urban	1.02 (0.89-1.17)	⊢ ∎1	
Rural	0.71 (0.62-0.80)		
NDVI _{1000m}			
Age, years			0.034
< 60	0.71 (0.58-0.87)	┝━╋━┥	
≥ 60	0.95 (0.86-1.06)	⊨∎⊣	
Gender			0.068
Male	0.84 (0.74-0.94)	┝╼╋╾┥	
Female	0.98 (0.85-1.13)	⊢ ∎1	
Residence			<.001
Urban	1.06 (0.92-1.21)	₽ <mark>₩</mark> ₽₩	
Rural	0.74 (0.65-0.83) 「		
	0.4	4 0.6 0.8 1 1.2 1.4	

Fig. 3 Stratified analyses of association between greenness levels per tertile increment and the risk of cardiovascular events. NDVI, normalized difference vegetation index

Ontario revealed that the incidence risk of acute myocardial infarction and heart failure decreased by 6% and 7%, respectively, as an interquartile range increased in residential greenness (measured by NDVI level) [17]. To our knowledge, only a few studies conducted in in developing nations, such as China, that have examined the association between greenness exposure and incidence risk of CVD [19, 20]. For example, the 33 Communities Chinese Health Study, based on population-based cross-sectional study, and recruiting 24,845 participants aged from 18 to 74 years, suggested that an interquartile range increase in NDVI_{500 m}of a community was associated with a 27% lower likelihood of CVD prevalence, with an OR being 0.73 (95% CI:0.65–0.83), which was similar to our findings [20]. Insights derived from these studies, combined with this current study, can offer valuable guidance to decision-makers involved in shaping development strategies aimed at improving cardiovascular health among

associations mediate	associations mediated by cardiometabolic risk factors								
Variable	Associations ^a	Proportion							
	Total	P value	Indirect		Direct		CI)		
	HR (95% CI)		HR (95% CI)	P value	HR (95% CI)	P value			
Hypertension	0.86(0.78-0.94)	0.001	1.00(0.99–1.00)	0.129	0.86(0.79–0.94)	0.001	2.52(-0.80,7.91)		
Diabetes mellitus	0.86(0.79-0.94)	0.001	1.00(0.99–1.00)	0.020	0.86(0.79-0.94)	0.001	2.45(0.33,7.17)*		
Overweight/Obesity	0.92(0.85-0.99)	0.036	1.00(0.99-1.01)	0.717	0.92(0.85-0.99)	0.034	-1.19(-28.24,9.43)		
BMI continuous	0.86(0.79-0.94)	0.001	0.99(0.99–1.00)	0.024	0.87(0.79–0.95)	0.002	4.04(0.45,11.95)*		
Dyslipidemia	0.86(0.79–0.94)	0.001	1.00(0.99–1.00)	0.302	0.86(0.79–0.94)	0.001	1.90(-2.02,7.35)		
TC	0.86(0.79-0.94)	0.001	1.00(0.99-1.00)	0.048	0.87(0.79-0.95)	0.001	2.75(-0.07,8.46)		

0.507

<.001

0.283

0.001

0.058

0.017

0.87(0.80-0.96)

0.89(0.81-0.97)

0.86(0.79-0.94)

0.86(0.79-0.94)

0.86(0.79-0.94)

0.84(0.77-0.92)

Table 3 Decomposition of the total association between NDVI within 500 m buffer and the risk of total CVD direct and indirect associations mediated by cardiometabolic risk factors

Abbreviations: CVD cardiovascular disease, NDVI normalized difference vegetation index, HR hazard ratio, CI confidential interval, BMI body mass index, TC total cholesterol, TG triglyceride, HDL-C high-density lipoprotein cholesterol, LDL-C low-density lipoprotein cholesterol, PM particulate matter

1.00(0.99-1.00)

0.98(0.96-0.99)

1.00(1.00 - 1.00)

0.99(0.99-1.00)

0.99(0.99-1.00)

1.02(1.00 - 1.04)

^a Hazard ratios given per tertile increase for NDVI level

0.87(0.79-0.95)

0.87(0.79-0.95)

0.86(0.79-0.94)

0.85(0.78-0.93)

0.85(0.78-0.93)

0.86(0.78-0.94)

0.003

0.002

0.001

0.001

0.001

< 0.01

^{*} P < 0.05

ΤG

HDL-C

LDL-C

PM_{2.5}

Physical activity

Heatwave days

citizens. This includes acknowledging the pivotal influence of residential greenness and contemplating initiatives such as the expansion of green spaces. Most importantly, our findings provide evidence of the association between greenness levels and CVD in developing country, and highlight the importance of implementing greenness interventions at population level for primary prevention of CVD and the comprehensive management of modifiable risk factors. Moreover, our initiatives should not be exclusively focus on expanding the quantity of green spaces, rather, equal attention should be given to enhancing public perception of these environments to optimize health benefits.

Stratified analyses revealed that rural residents derived greater health benefits from green space exposure compared to their urban counterparts. While prior studies have predominantly focused on urban settings in highincome countries [18, 33-35], our findings align with a recent Chinese cohort study documenting stronger protective effects of rural greenness [36]. Three key factors may explain this disparity: First, rural regions typically exhibit higher vegetation density and structurally distinct green spaces (e.g., forests, farmland) compared to fragmented urban greenery, which may limit urban residents'exposure to meaningful vegetation [8, 37]. Second, accelerated urbanization in Chinese cities correlates with elevated air pollution levels, potentially counteracting green space benefits [38]. Third, urban parks-often prioritized in green space metrics-frequently lack dense vegetation; studies show minimal correlation between urban park coverage and actual greenness exposure [39]. These findings underscore the need to contextualize green space research within regional urbanization patterns and ecological characteristics.

0.004

0.011

0.001

0.001

0.001

< 0.01

0.92(-2.33,4.89)

16.18(7.37,47.96)*

1.02(-0.98,3.98)

5.34(2.01,13.55)*

4.16(-0.31.12.05)

-12.82(-48.79,-2.47)*

Additionally, the evidence concerning the impact of residential greenness on specific CVD subgroups, particularly stroke was sparse. Several studies have investigated the relationship between residential greenness and the risk of stroke, however, they have yield inconsistent findings [18, 40]. Possible explanations may be the difference in exposure definitions and population selection. In this current large-scale cohort study, we found that residential greenness in the 300 m radius around the participant's residential addresses could reduce the risk of stroke, but no statistically significant difference between NDVI per 1000 m and stroke risk. It seems possible that vegetation appears decrease the risk of stroke via directly and efficiently reducing traffic related air pollution and noise, particularly within a small buffer size surrounding individual's residence address. Undoubtedly, future studies should shed light on whether these associations are sensitive to different buffer size around individual's neighbourhood, a factor which may be related to the accessibility of the greenspace.

Although some studies have suggested that greenspaces may enhance the cardiovascular health by diminishing exposure to air pollution and traffic noise, alleviating the heat, and promoting physical activity among nearby residents [19, 41, 42], the exact mechanisms remain incompletely delineated. A report has provided three potential functions of greenspace on health, including reducing harm, restoring capacities, and building capacities [42]. In this context, some studies have explored the association of residential greenness with CVD, focusing on the mediators such as physical activity, road proximity and air pollution [11, 12]. A recent analysis of data from five southwestern provinces of China based on a cross-sectional design indicated that higher physical activity level and lower BMI may partially mediate the benefit effect of greenspace on the risk of 10-year atherosclerosis cardiovascular disease (ASCVD), with physical activity and BMI mediating 44.73% and 35.15% of the total effect for Enhanced Vegetation Index per 500 m, respectively, in the medium-risk and low-risk groups compared [19]. These also accord with our finding, which showed that physical activity and BMI continuous mediated 5.34% and 4.04% of the total effects for NDVI_{500 m}on CVD. A community-based study in eastern China also found that physical activity partially mediated the relationship between greenness exposure and CVD [43]. This observation may be explained by the fact that green environments offer an intrinsic motivation for individuals to engage in outdoor activities, fostering increased opportunities for walking and active transportation [44]. People living in greener areas may benefit from reduced pollution, less noise, lower stress levels, and enhanced psychological restoration, all of which can improve health outcomes [44, 45]. Additionally, growing evidence shows that higher greenness is associated with lower BMI, a known risk factor for CVD [46, 47]. It is plausible that greenness indirectly affects cardiovascular health by influencing BMI. Although some studies have shown that BMI mediates the relationship between residential greenness and hypertension or ASCVD [11, 37], the exact mechanisms remain unclear. Some research indicates that green environments may provide convenient spaces for physical activity, helping to reduce obesity and lower blood pressure [48]. Nevertheless, comparing the proportions mediated across various studies poses a challenge owing to variations in exposure assessment, the delineation of buffer zones around individual's residential address, and selection of analytical methods.

Additionally, we failed to find the mediation effect of $PM_{2.5}$ on the association between greenness and the risk of CVD incidence in this current study, which was in accordance with prior research [19]. On the contrary, another study, enrolling 121,701 female registered nurses aged 30 to 55 years, has found support for the conclusion that air pollution might act as a mediator in the association between greenness and mortality [49]. This inconsistency could arise from variations in population Page 10 of 12

selection and the metrics used to assess air pollution. Furthermore, the diversity, quality, and accessibility of greenspaces within residential neighborhoods may exert a substantial influence on the estimation of observed effects [14], thus warranting further investigation as potential factors contributing to the existing inconsistencies in epidemiological findings. We also found that the association of greenness with the risk of CVD was partially mediated by diabetes mellitus and HDL-C. To our best knowledge, only a few studies have indicated that cardiometabolic risk factors partially mediated the association between greenness and CVD risk [20, 21]. For example, partially consistent with our findings, the 33 Communities Chinese Health Study suggested that hypertension, type 2 diabetes mellitus, obesity and hypercholesterolemia mediated the association of residential greenness (NDVI) with CVD prevalence of 4.5%, 4.1%, 3.1% and 12.7%, respectively [20]. Another study revealed that the association of NDVI levels with myocardial infarction was partially mediated by dyslipidemia and type 2 diabetes mellitus but not hypertension [21]. Further studies are warranted to bridge the aforementioned gap and broaden our understanding of the mediation effect of modifiable risk factors on the association of residential green space with cardiovascular health.

This present study has several strengths. The findings of this current study are noteworthy because this study was based on a large-scale, representative longitudinal survey with standardized procedures and strict quality control, which is essential for accurate estimation of the association between greenness and CVD and mortality. Incorporating a large range of NDVI levels at individual-level enables us to estimate the associations with increased precision. Moreover, mediation analysis with proportional hazard models was applied based on the counterfactual approaches [31], which could provide a mathematical decomposition of the total association into direct and indirect associations with clear explanations. However, we acknowledge some specific limitations. First, although the greenness levels were obtained based on high resolution satellite images, the information on the types, quality and access of greenspace in residential neighbourhoods, and changes in residential greenness during the follow-up period were not distinguished in this present study, which may lead to misclassification of greenness exposure. Previous study also showed that higher access to greenspace and pocket park were associated with decreased all-cause mortality risk [14]. Second, information on the duration of greenness exposure, and noise were not available, which could confound the association of greenness with CVD events and all-cause mortality and have potential influence on the mediation analyses. Third, the information on subjects'residential mobility or length of time spent at reported baseline addresses was unavailable, which might be an important determinant of the association. Fourth, the relatively short follow-up duration limited the number of CVDspecific mortality observed, so this study did not examine the association between residential greenness and CVD-specific mortality. We will further investigate this relationship in future studies with extended follow-up durations. Finally, we didn't take into account seasonal variability in greenspace. We are currently designing a further analysis examining seasonal effect modification using monthly NDVI data. These findings will be reported in future work investigating temporal variations in greenness-health relationships.

Conclusion

In this prospective cohort study of adults in China, we observed that higher residential greenness level was independently associated with a reduced risk of CVD. This association appears to be partially mediated by some modifiable risk factors, such as diabetes mellitus, BMI continuous, HDL-C, and physical activity. These findings highlight the importance of interventions promoting green environments in the primary prevention, providing valuable insights for policymakers regarding the potential role of green spaces in mitigating cardiovascular health risks amidst challenges posed by climate change.

Abbreviations

CVD	Cardiovascular disease
NDVI	Normalized Difference Vegetation Index
CI	Confidential interval
HR	Hazard ratio
CHS	China Hypertension Survey
SRS	Simple random sampling
PM	Particulate matter
SBP	Systolic blood pressure
DBP	Diastolic blood pressure
FPG	Fasting plasma glucose
BMI	Body mass index
TC	Total cholesterol
TG	Triglyceride
LDL-C	Low-density lipoprotein cholesterol
HDL-C	High-density lipoprotein cholesterol
METs	Metabolic equivalent tasks
GDP	Gross Domestic Product
CHD	Coronary heart disease
ICD-10	International Classification of Diseases 10th Revision
SD	Standard deviation
NDE	Natural direct effect
NIE	Natural indirect effect
ASCVD	Atherosclerotic Cardiovascular Disease

Supplementary Information

The online version contains supplementary material available at https://doi. org/10.1186/s12889-025-22899-7.

Supplementary Material 1.

Supplementary Material 2.

Authors' contributions

Xue Cao, Nuerguli Tuerdi: Conceptualization, Methodology, Software, Formal analysis, Visualization, Writing—Original Draft. Haosu Tang, and Yujie Zhang: Methodology, Visualization, and Investigation. Xin Wang, Congyi Zheng, and Gang Huang: Validation, Investigation, and Writing—Review & Editing. Yixin Tian, Xue Yu, Xuyan Pei: Investigation. Zengwu Wang: Conceptualization, Resources, Writing—Review & Editing, and Project administration.

Funding

This work was supported by Project Entrusted by the National Health Commission of the People's Republic of China (NHC2020-609), the National High Level Hospital Clinical Research Funding (2023-GSP-RC-18), the China National Science & Technology Pillar Program (Grant No. 2011BAI11B01) and the CAMS Innovation Fund for Medical Sciences (Grant No. 2017-I2M-1–004). The funder had no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Data availability

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

This study was approved by the ethics committee of Fuwai Hospital, Beijing, China, and all participants provided written informed consent.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹ Division of Prevention and Community Health, National Center for Cardiovascular Disease, National Clinical Research Center of Cardiovascular Disease, State Key Laboratory of Cardiovascular Disease, Fuwai Hospital, Peking Union Medical College & Chinese Academy of Medical Sciences, Beijing 102308, China. ² State Key Laboratory of Numerical Modeling for Atmospheric Sciences and Geophysical Fluid Dynamics, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China. ³University of Chinese Academy of Sciences, Beijing 100049, China. ⁴Division of Prevention and Community Health, National Center for Cardiovascular Disease, National Clinical Research Center of Cardiovascular Disease, State Key Laboratory of Cardiovascular Disease, Fuwai Hospital, Peking Union Medical College & Chinese Academy of Medical Sciences, No. 15 (Lin), Fengcunxili, Mentougou District, Beijing 102308, China.

Received: 19 September 2024 Accepted: 23 April 2025 Published online: 03 May 2025

References

- Roth GA, Mensah GA, Johnson CO, et al. Global Burden of Cardiovascular Diseases and Risk Factors, 1990–2019: Update From the GBD 2019 Study. J Am Coll Cardiol. 2020;76(25):2982–3021.
- Wang W, Liu Y, Liu J, et al. Mortality and years of life lost of cardiovascular diseases in China, 2005–2020: Empirical evidence from national mortality surveillance system. Int J Cardiol. 2021;340:105–12.
- Moran A, Gu D, Zhao D, Coxson P, et al. Future cardiovascular disease in china: markov model and risk factor scenario projections from the coronary heart disease policy model-china. Circ Cardiovasc Qual Outcomes. 2010;3(3):243–52.
- Yusuf S, Joseph P, Rangarajan S, et al. Modifiable risk factors, cardiovascular disease, and mortality in 155 722 individuals from 21 high-income, middle-income, and low-income countries (PURE): a prospective cohort study. Lancet. 2020;395(10226):795–808.
- Sun Y, Mu J, Wang DW, et al. A village doctor-led multifaceted intervention for blood pressure control in rural China: an open, cluster randomised trial. Lancet. 2022;399(10339):1964–75.

- Zhou H, Wang X, Yang Y, et al. Effect of a Multicomponent Intervention Delivered on a Web-Based Platform on Hypertension Control: A Cluster Randomized Clinical Trial. JAMA Netw Open. 2022;5(12):e2245439.
- Keith RJ, Hart JL, Bhatnagar A. Greenspaces And Cardiovascular Health. Circ Res. 2024;134(9):1179–96.
- Ji JS, Zhu A, Lv Y, Shi X. Interaction between residential greenness and air pollution mortality: analysis of the Chinese Longitudinal Healthy Longevity Survey. Lancet Planet Health. 2020;4(3):e107–15.
- Scheer C, Plans-Beriso E, Pastor-Barriuso R, et al. Exposure to green spaces, cardiovascular risk biomarkers and incident cardiovascular disease in older adults: The Seniors-Enrica II cohort. Environ Int. 2024;185:108570.
- Wang X, Zhang Y, Tarik B, et al. The effect of residential greenness on cardiovascular mortality from a large cohort in South China: An in-depth analysis of effect modification by multiple demographic and lifestyle characteristics. Environ Int. 2024;190:108894.
- 11. Dong X, Tu R, Zhang L, et al. Residential greenness and atherosclerotic cardiovascular disease risk in a rural Chinese adult population. Ecotoxicol Environ Saf. 2021;222:112458.
- 12. Liu T, Cai B, Peng W, et al. Association of neighborhood greenness exposure with cardiovascular diseases and biomarkers. Int J Hyg Environ Health. 2021;234:113738.
- Hu HB, Hou ZH, Huang CH, LaMonte MJ, Wang M, Lu B. Associations of exposure to residential green space and neighborhood walkability with coronary atherosclerosis in Chinese adults. Environ Pollut. 2022;292(Pt A):118347.
- Hajna S, Nafilyan V, Cummins S. Associations between residential greenspace exposure and mortality in 4 645 581 adults living in London, UK: a longitudinal study. Lancet Planet Health. 2023;7(6):e459–68.
- Chien JW, Wu C, Chan CC. The association of hypertension and prehypertension with greenness and PM(2.5) in urban environment. Sci Total Environ. 2022;821:153526.
- Paoin K, Pharino C, Phosri A, et al. Association between greenness and cardiovascular risk factors: Results from a large cohort study in Thailand. Environ Res. 2023;220:115215.
- Chen H, Burnett RT, Bai L, et al. Residential Greenness and Cardiovascular Disease Incidence, Readmission, and Mortality. Environ Health Perspect. 2020;128(8):87005.
- Seo S, Choi S, Kim K, Kim SM, Park SM. Association between urban green space and the risk of cardiovascular disease: A longitudinal study in seven Korean metropolitan areas. Environ Int. 2019;125:51–7.
- Yu W, Liu Z, La Y, et al. Associations between residential greenness and the predicted 10-year risk for atherosclerosis cardiovascular disease among Chinese adults. Sci Total Environ. 2023;868:1879–2026.
- 20. Yang BY, Hu LW, Jalaludin B, et al. Association Between Residential Greenness, Cardiometabolic Disorders, and Cardiovascular Disease Among Adults in China. JAMA Netw Open. 2020;3(9):e2017507.
- Yitshak-Sade M, Kloog I, Novack V. Do air pollution and neighborhood greenness exposures improve the predicted cardiovascular risk? Environ Int. 2017;107:147–53.
- Wang K, Lombard J, Rundek T, et al. Relationship of Neighborhood Greenness to Heart Disease in 249 405 US Medicare Beneficiaries. J Am Heart Assoc. 2019;8(6):e010258.
- Wang Z, Chen Z, Zhang L, et al. Status of Hypertension in China: Results From the China Hypertension Survey, 2012–2015. Circulation. 2018;137(22):2344–56.
- Hao G, Wang X, Chen Z, et al. Prevalence of heart failure and left ventricular dysfunction in China: the China Hypertension Survey, 2012–2015. Eur J Heart Fail. 2019;21(11):1329–37.
- Wang Z, Zhang L, Chen Z, et al. Survey on prevalence of hypertension in China: background, aim, method and design. Int J Cardiol. 2014;174(3):721–3.
- Dierckx W, Sterckx S, Benhadj I, et al. PROBA-V mission for global vegetation monitoring: standard products and image quality. Int J Remote Sens. 2014;35(7):2589–614.
- Zhang H, Liu L, Zeng Y, Liu M, Bi J, Ji JS. Effect of heatwaves and greenness on mortality among Chinese older adults. Environ Pollut. 2021;290:118009.
- Chinese Diabetes Society. Guidelines for the prevention and control of type 2 diabetes in China(2017 Edition) [in Chinese]; 2018.

- Joint Committee on the Chinese Guidelines for Lipid Management. Chinese Guidelines for Lipid Management (2023). Chin Circ J. 2023;38(3):237–71.
- Kong L, Tang X, Zhu J, et al. A 6-year-long (2013–2018) high-resolution air quality reanalysis dataset in China based on the assimilation of surface observations from CNEMC. Earth System Science Data. 2021;13:529–70.
- VanderWeele TJ. Causal mediation analysis with survival data. Epidemiology. 2011;22(4):582–5.
- Valeri L, Vanderweele TJ. Mediation analysis allowing for exposuremediator interactions and causal interpretation: theoretical assumptions and implementation with SAS and SPSS macros. Psychol Methods. 2013;18(2):137–50.
- Rajagopalan S, Vergara-Martel A, Zhong J, et al. The Urban Environment and Cardiometabolic Health. Circulation. 2024;149(16):1298–314.
- Mannucci PM. Air pollution, cardiovascular disease, and urban greening: an ecological blueprint. Eur J Prev Cardiol. 2023;30(15):1608–11.
- Nieuwenhuijsen MJ. Influence of urban and transport planning and the city environment on cardiovascular disease. Nat Rev Cardiol. 2018;15(7):432–8.
- Liu C, Zhang B, Liu C, et al. Association of ambient ozone exposure and greenness exposure with hemorrhagic stroke mortality at different times: A cohort study in Shandong Province. China Ecotoxicol Environ Saf. 2024;278:116356.
- Huang B, Xiao T, Grekousis G, et al. Greenness-air pollution-physical activity-hypertension association among middle-aged and older adults: Evidence from urban and rural China. Environ Res. 2021;195:110836.
- Liu C, Yu Y, Liu C, et al. Effect of neighbourhood greenness on the association between air pollution and risk of stroke first onset: A casecrossover study in shandong province. China Int J Hyg Environ Health. 2023;254:114262.
- Klompmaker JO, Laden F, Browning M, et al. Associations of parks, greenness, and blue space with cardiovascular and respiratory disease hospitalization in the US Medicare cohort. Environ Pollut. 2022;312:120046.
- de Bont J, Pickford R, Åström C, et al. Mixtures of long-term exposure to ambient air pollution, built environment and temperature and stroke incidence across Europe. Environ Int. 2023;179:108136.
- 41. Katzmarzyk PT, Friedenreich C, Shiroma EJ, Lee IM. Physical inactivity and non-communicable disease burden in low-income, middle-income and high-income countries. Br J Sports Med. 2022;56(2):101–6.
- 42. Markevych I, Schoierer J, Hartig T, et al. Exploring pathways linking greenspace to health: Theoretical and methodological guidance. Environ Res. 2017;158:301–17.
- 43. Jia X, Yu Y, Xia W, et al. Cardiovascular diseases in middle aged and older adults in China: the joint effects and mediation of different types of physical exercise and neighborhood greenness and walkability. Environ Res. 2018;167:175–83.
- Marquet O, Floyd MF, James P, et al. Associations Between Worksite Walkability, Greenness, and Physical Activity Around Work. Environ Behav. 2020;52(2):139–63.
- James P, Kioumourtzoglou MA, Hart JE, Banay RF, Kloog I, Laden F. Interrelationships Between Walkability, Air Pollution, Greenness, and Body Mass Index. Epidemiology. 2017;28(6):780–8.
- Huang WZ, Yang BY, Yu HY, et al. Association between community greenness and obesity in urban-dwelling Chinese adults. Sci Total Environ. 2020;702:135040.
- Ortega FB, Lavie CJ, Blair SN. Obesity and Cardiovascular Disease. Circ Res. 2016;118(11):1752–70.
- Bao WW, Jiang N, Zhao Y, et al. Urban greenspaces and child blood pressure in China: Evidence from a large population-based cohort study. Environ Res. 2024;244:117943.
- James P, Hart JE, Banay RF, Laden F. Exposure to Greenness and Mortality in a Nationwide Prospective Cohort Study of Women. Environ Health Perspect. 2016;124(9):1344–52.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.