



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/226780/>

Version: Accepted Version

Article:

Birmpas, K., O'Reilly, D. and Delis, I. (2025) Integrative sensorimotor coding and the neurophysiology of intent: Comment on "Kinematic coding: Measuring information in naturalistic behaviour" by Becchio, Pullar, Scaliti, and Panzeri. *Physics of Life Reviews*, 53. pp. 307-309. ISSN: 1571-0645

<https://doi.org/10.1016/j.plrev.2025.04.007>

This is an author produced version of an article published in *Physics of Life Reviews*, made available under the terms of the Creative Commons Attribution License (CC-BY), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Integrative sensorimotor coding and the neurophysiology of intent: Comment on “Kinematic coding: Measuring information in naturalistic behaviour” by Becchio, Pullar, Scaliti, and Panzeri

Kyriakos Birmpas^{a*}, David O'Reilly^{a*}, Ioannis Delis^a

^aSchool of Biomedical Sciences, University of Leeds, LS2 9JT, United Kingdom

*Primary authorship

Corresponding author: Ioannis Delis, School of Biomedical Sciences, University of Leeds, LS2 9JT, Leeds, United Kingdom. Email: i.delis@leeds.ac.uk

Becchio et al. introduce a compelling framework conceptualizing movement patterns as cognitive information carriers (*‘kinematic codes’*) read-out by observers as a form of non-verbal communication. This kinematic coding approach has broad implications for cognitive and behavioural neuroscience, providing a computational link between action and perception in social settings. Specifically, it carves out a path for principled investigation into agent-environment interactions, which we aim to employ to link together the neural mechanism of action, perception and decision-making [1,2]. Here, we outline this research endeavor aiming to characterise the role of perceptual and motor neural processes in sensorimotor decision-making and, by extension, in communicating cognitive intention to others.

Motor patterns, whether they emerge from muscle activations, motor unit spike trains, brain oscillations or kinematic trajectories, are conventionally seen to reflect decision-making outcomes. Recent work promotes a more integrated viewpoint where perceptual and decision-formation processes operate along an action continuum [3]. From this continuous and parallel perspective, humans dynamically interact with their environment, sampling sensory information, forming perceptual decisions and executing actions. Hence, motor patterns encode information about a variety of cognitive processes. Supporting this claim, in an active-sensing experiment where participants judged object heaviness, perceptual variance was accounted for by lifting arm muscle activation patterns [4,5]. Crucially, diverse types of functional muscle interactions were found to encode distinct forms of decision-related information in this task [6]. Similarly, during object-wielding, body-wide postural fluctuations were shown to correlate with perceptual judgments [7], and during multisensory decision-making, active exploration of visuo-haptic information was shown to enhance perceptual accuracy [1].

However, to mechanistically understand the structure-function relationships of motor patterns in naturalistic behaviors, embedding the agent in their environment is a necessary additional research step. Becchio et al. propose this conceptual shift, moving from interacting neural subsystems encoding and decoding decisions to the behavioural encoding of decisions that are read-out by an observer. This reframing is directly pertinent to understanding neural mechanisms geared towards communication. For instance, neurophysiological evidence suggests speech processing entails neural encoding of articulatory kinematics (i.e. tongue and lip movements), even when these are not visually available [8]. Hence, representing others' bodily movements is central to effective information transmission. Importantly, other motor behaviours not overtly intended for communication may also be disambiguated. For instance, pixel covariations in video recordings of psychiatric patients' bodily movements in dyadic interaction were associated with treatment outcomes and affective states [9,10]. This nonverbal synchrony exemplifies the subtle behavioural manifestations of cognitive states that may not be consciously visible to observers but may still impact the quality of their social interactions. We suggest that the kinematic coding framework represents a principled approach with enhanced specificity to address meaningful questions across these research streams.




Figure 1: The active-sensing experimental paradigm [4]. Including an observer and incorporating visual occlusion combined with observer feedback enables kinematic coding concepts to be readily accessed.

In future work, we aim to capitalize on our newly developed computational methodologies combined with neurophysiological measurements to characterize the neural mechanisms underlying this new “*decision-during-action*” view [3,11,12]. By mapping cortical, spinal and corticospinal neural interactions during active-sensing tasks, we wish to offer an in-depth characterization of the neural codes of sensorimotor decision-making and learning. As Becchio et al. suggest, including observers in the active-sensing paradigm with visual occlusions or as an active part of decision outcomes (e.g. feedback) will uncover what information is available (or not) to others and how sensorimotor decisions are shaped in social settings (Fig.1), ultimately bringing to bear a more comprehensive understanding of agent-environment interactions.

[1] Delis I, Ince RA, Sajda P, Wang Q. Neural encoding of active multi-sensing enhances perceptual decision-making via a synergistic cross-modal interaction. *Journal of Neuroscience*. 2022 Mar 16;42(11):2344-55.

[2] Delis I, Dmochowski JP, Sajda P, Wang Q. Correlation of neural activity with behavioral kinematics reveals distinct sensory encoding and evidence accumulation processes during active tactile sensing. *NeuroImage*. 2018 Jul 15;175:12-21.

[3] Balsdon T, Verdonck S, Loossens T, Philiastides MG. Secondary motor integration as a final arbiter in sensorimotor decision-making. *PLoS Biology*. 2023 Jul 17;21(7):e3002200.

[4] Toma S, Simonelli F, Delis I, Santello M. A novel experimental and analytical framework for investigating the interplay among sensory encoding, motor control and perceptual decision formation, under review.

- [5] Toma S, Santello M. Motor modules account for active perception of force. *Scientific reports*. 2019 Jun 20;9(1):8983.
- [6] Birmpas K, Balntoukas K, O'Reilly D, Toma S, Simonelli F, Santello M, & Delis I. Associating motor signals with decision-making dynamics during active perception. *UK Sensorimotor Conference 2024*.
- [7] Mangalam M, Chen R, McHugh TR, Singh T, Kelty-Stephen DG. Bodywide fluctuations support manual exploration: Fractal fluctuations in posture predict perception of heaviness and length via effortful touch by the hand. *Human Movement Science*. 2020 Feb 1;69:102543.
- [8] Pastore A, Tomassini A, Delis I, Dolfini E, Fadiga L, D'Ausilio A. Speech listening entails neural encoding of invisible articulatory features. *Neuroimage*. 2022 Dec 1;264:119724.
- [9] Ramseyer F, Tschacher W. Nonverbal synchrony of head-and body-movement in psychotherapy: different signals have different associations with outcome. *Frontiers in psychology*. 2014 Sep 5;5:979.
- [10] Tschacher W, Rees GM, Ramseyer F. Nonverbal synchrony and affect in dyadic interactions. *Frontiers in psychology*. 2014 Nov 24;5:1323.
- [11] O'Reilly D, Delis I. Dissecting muscle synergies in the task space. *Elife*. 2024 Feb 26;12:RP87651.
- [12] O'Reilly D, Shaw W, Hilt P, de Castro Aguiar R, Astill SL, Delis I. Quantifying the diverse contributions of hierarchical muscle interactions to motor function. *iScience*. 2025 Jan 17;28(1).