



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/226968/>

Version: Accepted Version

---

**Article:**

García, Olga P., Zavala, Gerardo A., Campos-Ponce, Maiza et al. (2025) Intestinal parasitic infections are related to micronutrient status and body composition in Mexican school-age children: results from a cross-sectional study. *Journal of parasitic diseases*. ISSN: 0971-7196

<https://doi.org/10.1007/s12639-025-01814-2>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

**Intestinal parasitic infections are related to micronutrient status and body composition in Mexican school-age children: results from a cross-sectional study**

Olga P. García<sup>a,\*</sup>, Gerardo A. Zavala<sup>b,1</sup>, Maiza Campos-Ponce<sup>b</sup>, Colleen M. Doak<sup>b,2</sup>, Alex Brito<sup>c</sup>, Katja Polman<sup>d</sup>, Lindsay H. Allen<sup>e</sup>, Jorge L. Rosado<sup>a</sup>

<sup>a</sup> School of Natural Sciences, Universidad Autónoma de Querétaro, Av. de la Ciencias S/N, Querétaro, Querétaro, CP 76230, Mexico;

<sup>b</sup> Faculty of Earth and Life Sciences, VU Amsterdam University, De Boelelaan 1085, 1081 HV, Amsterdam, The Netherlands;

<sup>c</sup> Laboratory of Pharmacokinetics and Metabolomic Analysis. Department of Translational Medicine and Biotechnology. I.M. Sechenov First Moscow State Medical University, 8-2 Trubetskaya str. 119991, Moscow, Russian Federation;

<sup>d</sup> Department of Biomedical Sciences, Institute of Tropical Medicine Antwerp, Nationalestraat 155 – 2000, Antwerp, Belgium;

<sup>e</sup> USDA, ARS, Western Human Nutrition Research Center, 430 West Health Sciences Drive, Davis CA 95616, USA

**\*Corresponding author:** Gerardo A Zavala and Olga P. García

Tel:

Email [g.zavala@york.ac.uk](mailto:g.zavala@york.ac.uk)

e-mail: [olga.garcia@uaq.mx](mailto:olga.garcia@uaq.mx)

<sup>1</sup> Present address: Department of Health Sciences, University of York, Heslington, York, YO10 5DD, United Kingdom

<sup>2</sup> Present address: Public Health, St Ambrose University, 518 Locust St, Davenport, IA 52803, USA  
The first two authors should be regarded as joint First Authors.

## **Abstract**

**Background:** Intestinal parasitic infections remain a significant public health issue, particularly in low-resource settings. They have been linked to poor micronutrient status and body composition, which are critical determinants of child health and development. The aim of this cross-sectional study was to determine the relationship between intestinal parasitic infections and micronutrient status, and whether these differ according to the body composition .

**Methods:** Serum concentrations of zinc, iron, ferritin, vitamins A, E, C, D, folate, B12 and CRP, were determined in 269 school-aged children from rural Mexico. Infection with soil transmitted helminths (STHs) and intestinal protozoa was screened in a fecal sample. Anthropometric and body composition measurements were taken.

**Results:** Lower ferritin, zinc and vitamin C concentrations were found in children infected with any STHs or *A. lumbricoides* compared to parasite-free children ( $p < 0.05$ ). Children infected with any intestinal protozoa, *Endolimax nana* or *Entamoeba coli* had higher concentrations of ferritin and B12 than parasite-free children ( $p < 0.05$ ). Vitamin E:lipid concentration was higher in children infected with any intestinal protozoa and *E. nana*. Among the children with high body fat percentage, those infected with STH had lower zinc, and those infected with intestinal protozoa had lower vitamin A than parasite-free children ( $p < 0.05$ ).

**Conclusion:** STH infection was associated with lower concentrations of ferritin, zinc and vitamin C, whereas intestinal protozoa infection with higher concentrations of ferritin, vitamin E:lipids, and B12. These associations differed according to body fat percentage.

**Key words:** Body Composition, Helminths, Intestinal Parasites, Micronutrients, Protozoa.

## **Introduction**

Globally, intestinal parasitic infections represent a major health burden, contributing to malnutrition and impaired growth in children, particularly in low- and middle-income countries. Intestinal parasite (helminths and protozoa) infections contribute one fourth of the infectious diseases in humans, and they are a recognized worldwide public health problem. (Alum, Rubino, & Ijaz, 2010). Socioeconomic status, health conditions, education, the presence of domestic animals, poor water supply and food hygiene are factors that affect the risk of having intestinal parasites (Black et al., 2013; García, Zavala, Campos-Ponce, & Polman, 2023). Intestinal parasite infections have been associated with malnutrition, and inflammation (Gutiérrez-Jiménez, Luna-Cazáres, & Vidal, 2017; Zavala et al., 2018). Additionally, intestinal parasite infections (5.7%), obesity (3.9 %), and overall micronutrient deficiencies (6.1%) account for more than 15% of the global disability-adjusted-life-years (Black, 2003; Hotez et al., 2014; Ng et al., 2014).

Intestinal parasitic infections affect the nutritional status of the host through several mechanisms, including reducing food intake (anorexia) and nutrient absorption, and increasing the loss of nutrients, leading to malnutrition and higher risk of micronutrient deficiencies (García, Zavala, Campos-Ponce, et al., 2023; Katona & Katona-Apte, 2008). Multiple micronutrient deficiencies negatively impact both disease progression and infection (Bhaskaram, 2002; Katona & Katona-Apte, 2008), and infection in turn may exacerbate malnutrition (Stephenson, Latham, & Ottesen, 2000). Micronutrient deficiencies and intestinal parasitic infections in children both affect growth and cognitive development (Schaible & Stefan, 2007). Most studies that have evaluated the association between poor micronutrient status and intestinal parasitic infection in children have focused on zinc, iron and vitamin A status (Brecht de Gier et al., 2014). The relationship between intestinal parasitic infections and other micronutrients that are essential for an adequate immune response (i.e. vitamins C, D or E) has not been extensively studied (Maggini et al., 2007). Moreover, the available evidence is centered on the relationships between intestinal parasites and micronutrients in undernourished populations; but not in children affected by a high prevalence of overweight and obesity. This is particularly important in transitional countries such as Mexico, where childhood obesity is the current main nutritional problem and where co-existing parasitic infections and micronutrient deficiencies are both public health problems.

The combined prevalence of overweight and obesity in Mexican school-age children is among the highest in the world (~34%) (Cuevas-Nasu et al., 2017). In addition, the prevalence of overall micronutrient deficiencies is high, especially among children (Rivera Dommarco, 2012). For the micronutrients on which there is national data, the most prevalent

deficiencies in school-age Mexican children (5 to 11 years) are iron (9.3%; serum ferritin <15 µg/L) and zinc (23.6%; <65 µg/dL) deficiencies (Shamah-Levy et al., 2012; Villalpando et al., 2015). Also, intestinal parasitic infections in children have been reported to range from 30% to 70%, being highest in the southern states of the country (Morales-Espinoza et al., 2003; Zavala et al., 2020). Infection with intestinal parasites, in particular *G. lamblia* and STH, is highly associated with dual burden households (co-existing overweight/obesity and stunting) (Campos Ponce et al., 2012). We recently found that Mexican children with moderate to heavy infection with *E. coli* had higher body fat and higher reported food intake compared to non-infected children (Zavala et al., 2016).


The aim of this study was to determine the relationships between micronutrient status and intestinal parasitic infections, and whether these differ across the range of body composition in children.

## **Materials and methods**

### **Study population and design**

A cross-sectional study was conducted in 269 school-aged children (6-10 y) recruited from two rural communities (Santa Cruz and Santa María Begoña), State of Querétaro, Mexico. All children from 2 local schools were invited to participate in the study. Parents or caretakers received oral and written information about their participation and the children's participation in the study. Written signed informed consent was obtained before entry into this research. Children also provided their assent to participate. The study was conducted according to the guidelines of the Declaration of Helsinki, and the study protocol was approved by the Bioethics Committee of the School of Natural Sciences at the Universidad Autónoma de Querétaro (UAQ) (protocol # FNN-2014-02).

Children who received deworming treatment in the last 4 months or had a physical or mental disability were excluded. Children that met the age criteria, that lived in one of the two communities and whose parents signed consent forms were included in the study. Parents or caretakers were asked to attend their local health clinic to answer a medical history and a socioeconomic status questionnaire previously validated. The complete research process is illustrated in Figure 1.



**Figure 1.** Flow chart of the research procedures

### **Blood sample collection**

Children’s parents/caretakers were instructed not to feed the children after dinner (8:00 PM) one day before the blood sample collection to ensure a 12 h fasting before the blood sample was collected by venipuncture in the early morning. The samples were stored in a cooler and transported the same morning to the Human Nutrition Laboratory, at UAQ. Plasma and serum were separated by centrifugation at 1800–2000 rpm for 15 min (Beckman Allegra 21R, Palo Alto, CA, USA), and aliquots stored at  $-70^{\circ}\text{C}$  for later analysis. All the biochemical analyses were performed in duplicate by trained personnel at the Human Nutrition Laboratory, UAQ, except vitamin B12 and folate that were measured at the measured at the U.S. Department of Agriculture (USDA), Agricultural Research Service (ARS) Western Human Nutrition Research Center in Davis, California.

### **Micronutrient status evaluation**

Vitamins A and E ( $\alpha$ -tocopherol) were measured simultaneously in serum using reverse phase high pressure liquid chromatography (HPLC) (Mod 2996, Waters Associates, Milford, MA, USA) using corresponding U.S. Pharmacopeia (USP) certified retinol and alpha-tocopherol standards. Vitamin A status was considered deficient with serum retinol concentrations  $<10\ \mu\text{g/dL}$  and low if  $<20\ \mu\text{g/dL}$ .(Gibson, 2005) Vitamin E deficiency was defined as serum alpha-tocopherol  $<3\ \mu\text{g/mL}$  and as low status at concentrations  $<5\ \mu\text{g/mL}$ .(Fares et al., 2011) Using the vitamin E:lipids ratio, vitamin E deficiency was classified as  $<0.8\ \text{mg/g}$ .(Drewel et al., 2006) Serum vitamin C was analyzed using reverse phase HPLC, with an ascorbic acid USP standard, and a cut point for deficiency of  $<2\ \mu\text{g/mL}$  and low status

as  $<4 \mu\text{g/mL}$ .(Gibson, 2005) Serum vitamin D was determined by a commercial 25(OH)-Vitamin D direct ELISA kit (Immundiagnostik AG, Bensheim, Germany) and a microplate spectrophotometer (Multiskan Ascent, Thermo Electron Corporation, MA, USA). Vitamin D deficiency was defined as  $<20 \text{ ng/mL}$  and insufficient status if  $<30 \text{ ng/mL}$ .(LeBlanc et al., 2014) Plasma vitamin B12 and folate concentrations were measured by enzyme-based immunoanalysis using a Cobas® e41 analyzer (Roche, Switzerland) and the reagents vitamin B12 (theoretical normality values 95%CI: 191-663 pg/mL) and folate III (4.6-18.7 ng/mL). Low serum folate was defined as  $<4 \text{ ng/mL}$  and high as  $>20 \text{ ng/mL}$  (the highest limit of detection of the instrument) and low and marginal B-12 concentrations were defined as  $<200 \text{ ng/mL}$  and between 200 and 300 ng/mL, respectively.(Brito et al., 2015)

Serum ferritin concentrations were determined by a commercial ELISA kit (ab108837, Abcam, UK) and a microplate spectrophotometer (Multiskan Ascent, Thermo Electron Corporation, MA, USA). Low iron stores were defined as serum ferritin  $<15 \mu\text{g/L}$ . Total serum iron concentrations were measured using a commercial kit (Iron Ferrozine, Elitech, Sées, France). Low iron status was defined as  $<60 \mu\text{g/dL}$  and iron deficiency as  $<45 \mu\text{g/dL}$  (Organization, 2011). Zinc concentrations were measured in serum by atomic absorption spectrometry (AAAnalyst 7000, Perkin Elmer Instruments, Norwalk, CT, USA), using its corresponding standard (Perkin Elmer) and deficiency was defined as serum zinc  $<65 \mu\text{g/dL}$  (Hotz, Peerson, & Brown, 2003).

High sensitive C-reactive protein (hsCRP) was measured in plasma using commercial ELISA kits (High Sensitivity C-Reactive Protein ELISA Kit, Bioquant) and analyzed in a Multiskan Ascent microplate photometer (Thermo Electron Corporation, Ma, EUA).

### **Anthropometry and body composition evaluation**

Children and one parent or caretaker were transported from their local communities to the Nutrition Clinic at the UAQ. Weight and height were measured by trained personnel following the World Health Organization (WHO) procedures. Weight was measured using a calibrated digital scale (SECA Mod. 813, Hamburg, Germany) and height with a stadiometer (SECA Mod 206, Hamburg, Germany). Underweight was defined as BMI two standard deviations (SD) below, and overweight and obesity as one and two SDs above, the WHO reference median of the BMI-for-age z-scores for children aged 5 to 19 years, respectively. Stunting was defined as two SDs below the WHO reference median for height-for-age (Onis, 2006).

Whole body composition was measured to determine body fat percentage by a certified technician using dual-energy X-ray absorptiometry (DXA) (Hologic Mod Explorer, Bedford, MA, USA). Children were asked to remove any

jewelry or anything metal, and once in the machine, they were asked to remain still for the duration of the scanning. Total body fat percent was obtained from the DXA absorptiometry values; high body fat for girls was considered to be above 30%, and for boys, above 25% (Ellis, Abrams, & Wong, 1997).

### **Evaluation of intestinal parasitic infections**

A single stool sample was collected from each participant and analyzed on the day of collection. A direct copro-parasitological test consisting of a wet mount with iodine staining of slides was performed to determine the presence of protozoan intestinal parasites, as described by WHO (WHO, 1991). These protozoa included *Entamoeba coli* (*E.coli*), *Endolimax nana* (*E. nana*), *Balantidium coli* (*B. coli*), *Entamoeba histolytica*, *Iodamoeba Bütschlii* and *Giardia lamblia*. Two Kato-Katz smears (2 x 41.7g) were made according to standard procedures to determine the presence and number of eggs of soil transmitted helminths (STHs).(Willcox & Coura, 1991) These parasites included *Ascaris lumbricoides* (*A. lumbricoides*), *Trichuris trichiura* (*T. trichiura*) and hookworm. The intensity of infection was characterized following WHO recommendations. Light- and moderate-intensity infection respectively were defined as 1-4999 and >5000 epg feces for *A. lumbricoides*; 1-999 and  $\geq 1000$  epg for *T. trichiura*; and; 1-1999 and  $\geq 2000$  epg for hookworm. Children with no intestinal protozoa or STH eggs were categorized as parasite-free. Those infected with more than one species of intestinal parasite were combined and categorized as having “multiple infections”.

### **Statistical Analysis**

The Shapiro-Wilk test was used to check normality of the distributions of the variables. Chi- squared tests were performed to compare the proportion of children with parasitic infections versus parasite-free children and to compare children with and without micronutrient deficiencies. After checking linearity and normal distribution of the dependent variables, one way, multivariate analyses of co-variance (MANCOVA) were used to compare the concentrations of micronutrients in children infected with any STH, any intestinal protozoa or specific parasites with prevalence above 10% (i.e. *A. lumbricoides*, *E. coli*, *B. coli* and *E. nana*) versus parasite-free children and CRP concentration. Interactions were also assessed through MANCOVA defining the absence/presence of infection as factor (independent variable); micronutrient concentrations as the response (dependent variable) and considering the category of normal/high body fat percent as covariate. Both the MANCOVAs to compare groups and to assess interactions were adjusted by age (months), gender (male/female), mother’s education level (years of education) and CRP concentrations. A *p* value <0.05 was considered statistically significant. We analyzed the interactions between body

fat and each micronutrient, and conducted post hoc analysis to further investigate the interaction for those micronutrients with a significant interaction term. Serum folate concentrations were not incorporated in the regression models because most of the values were above the limit of detection of the assay, affecting the distribution of the variable. All statistical analyses were performed using SPSS 21.0 (SPSS, /Chicago IL) and the figure plotted using Graph Pad Prism 8 (Graph Pad Software Inc, San Diego, CA).

## Results

A total of 269 children ( $8.0 \pm 1.6$  y, 47% girls) participated in the study (**Table 1**). The most common micronutrient deficiencies were vitamin D (19.3%) and zinc (15.2%). The prevalences of vitamin B-12 and vitamin E deficiencies were very low (1.1% and 0.4%, respectively). No child had folate deficiency and most had high folate (84.8%). No deficiencies were observed for vitamin A, vitamin C or iron. A total of 18.4% of the children were overweight, 9.6% obese, 51% had high body fat, 2% were underweight and 4% were stunted. C-reactive protein concentration was directly related to iron ( $\beta=-0.13$ , 95% C.I.=-0.20 to -0.06  $p < 0.001$ ) and zinc ( $\beta=-2.23$ , 95% C.I.=0.14 to 4.32  $p=0.036$ ) and negatively to ferritin ( $\beta= 0.01$ , 95% C.I.=0.01 to 0.02  $p < 0.001$ ).

**Table 1.** Main characteristics of the population (n=269)

Variable	Mean (SD) <sup>a</sup>	Proportion children with micronutrient deficiency (n)
Age (years)	8.0 (1.6)	NA
<b>Anthropometry and body composition</b>		
Weight (kg)	27.6 (8.2)	NA
Height (cm)	126.1 (9.9)	NA
Height-for-age (Z-Score)	-0.6 (1.0)	NA
BMI-for-age (Z-Score)	0.3 (1.3)	NA
Total body fat (%)	4.0 (2.7)	NA

### Biochemical measurements

Ferritin (µg/L)	99.0 (51.3)	NA
Iron (µg/dL)	92.1 (24.5)	0% (0)
Zinc ( µg/dL)	74.2 (11.2)	15.2% (41)
Retinol (µg/dL)	32.5 (6.7)	0% (0)
Vitamin D (ng/mL)	17.9 (3.1)	19.3% (52)
Vitamin E (µg/mL)	2.7 (1.0)	NA
Vitamin E lipids ratio (mg/g )	2.7 (1.0)	.04% (1)
Vitamin C (µg/mL)	7.6 (1.8)	0% (0)
Vitamin B-12 (pg/mL)	662.3 (258.1)	1.1% (3)
C-reactive protein (mg/L)	0.97 (1.6)	NA

---

<sup>a</sup> SD=Standard Deviation

Intestinal infections were common, with 60% of children being infected with at least one intestinal parasite and 46% with at least one protozoan. Among the protozoans, *E. coli* (19 %), *E. nana* (16%) and *B. coli* (12%) were the most common. The prevalence of any STH infection was 19%; *A. lumbricoides* was the most prevalent (16.4%). Multiple-intestinal parasitic infections affected 14.5%.

Children infected with any intestinal parasite exhibited higher serum concentrations of vitamin B12 and ferritin compared to parasite-free children (Table 2). Those infected with any STH or *A. lumbricoides* had lower concentrations of plasma ferritin, zinc, and serum vitamin C. Conversely, children infected with intestinal protozoa showed higher levels of serum ferritin, vitamin B12, and vitamin E:lipids compared to parasite-free peers. Specifically, *E. coli*-infected children had higher serum ferritin retinol and vitamin B12 concentrations, while those infected with *E. nana* had higher levels of serum ferritin, retinol vitamin E:lipids, and vitamin B12. Finally, children with multiple infections demonstrated lower serum ferritin but higher vitamin B12 concentrations compared to parasite-free children ( $p<0.05$ )."

1 **Table 2.** Serum or plasma micronutrient concentrations by parasite infection group (n=269)<sup>a</sup>


Micronutrient	Parasite-Free (107)	Infected <sup>b</sup> (162)	Soil						
			transmitte d helminths (52)	A. <i>lumbricoides</i> (44)	Intestinal Protozoa (123)	<i>E. coli</i> (52)	<i>B. coli</i> (33)	<i>E. nana</i> (44)	Multiple infection <sup>c</sup> (39)
Ferritin (µg/L)	98.88 ± 52.83	99.54 ± 50.67*	92.22 ± 45.32*	93.64 ± 48.28*	101.03 ± 51.20*	101.53 ± 54.66*	92.31 ± 35.96	108.56 ± 64.76*	93.91 ± 54.61*
Iron (µg/dL)	89.07 ± 28.10	94.67 ± 31.92	91.73 ± 29.12	93.70 ± 29.23	95.96 ± 32.26	98.71 ± 26.17	92.94 ± 37.65	100.58 ± 33.33	100.27 ± 27.56
Zinc (mg/L)	0.75 ± 0.09	0.74 ± 0.11	0.71 ± 0.12*	0.70 ± 0.12*	0.75 ± 0.10	0.77 ± 0.10	0.73 ± 0.13	0.75 ± 0.11	0.75 ± 0.11
Retinol (µg/dL)	32.59 ± 7.67	32.98 ± 6.01	32.72 ± 5.49	32.85 ± 5.56	32.91 ± 6.03	32.85 ± 5.89*	32.75 ± 6.55	33.43 ± 5.42*	32.46 ± 4.21
Vitamin D (ng/mL)	17.40 ± 3.32	18.25 ± 2.93	18.17 ± 2.75	18.22 ± 2.70	18.24 ± 3.03	18.38 ± 2.75	18.34 ± 3.70	17.52 ± 5.28	18.11 ± 3.84
Vitamin E (µg/mL)	5.35 ± 1.66	5.34 ± 1.66	5.11 ± 1.53	5.24 ± 1.53	5.46 ± 1.98	5.27 ± 1.59	5.36 ± 1.71	5.57 ± 1.77	5.65 ± 1.57
Vitamin E lipids (µg/mL)	2.67 ± 0.97	2.72 ± 1.03	2.64 ± 1.08	2.74 ± 1.11	2.74 ± 1.02*	2.66 ± 1.14	2.66 ± 0.97	2.83 ± 1.06*	2.69 ± 1.06
Vitamin C (mg/mL)	7.81 ± 1.91	7.57 ± 1.81	7.16 ± 1.92*	7.09 ± 1.92*	7.73 ± 1.74	7.72 ± 1.64	7.34 ± 2.26	7.81 ± 1.86	7.89 ± 2.01
Vitamin B-12 (pg/mL)	633.22 ± 204.92	678.13 ± 289.48*	646.39 ± 241.59	641.42 ± 237.54	699.75 ± 302.84*	719.88 ± 325.19*	612.62 ± 170.41	727.93 ± 335.64*	648.42 ± 260.66*

<sup>a</sup> MANCOVA, Mean ± Standard Deviation, adjusted by age, sex, mother's educational level and C-reactive protein concentration <sup>b</sup> Infected with any intestinal parasite <sup>c</sup>

Infection with 2 or more intestinal parasites; \* Significantly different from parasite-free with MANCOVA (p<0.05)

3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13

Of the micronutrients studied, a significant interaction with body fat and parasitic infection was observed only with zinc and vitamin A. Total body fat percentage interacted with STHs infection in relation to serum zinc concentrations and with intestinal protozoa in relation to serum retinol concentration ( $p$  value  $< 0.05$ ) (**Figure 2**). Among the children with high body fat percentage, those infected with STH had lower serum zinc concentrations than parasite-free children, while those infected with intestinal protozoa had lower serum retinol than parasite-free children ( $p < 0.05$ ). In addition, among the children with normal body fat percentage, those infected with STH had similar serum zinc concentrations compared to parasite-free children and those infected with intestinal protozoa had higher vitamin A concentration than parasite-free children ( $p < 0.05$ ).



14

**Fig. 2** Interaction between high/normal body fat with A) helminth infection and serum zinc; B) protozoan infection and serum retinol concentrations. Lines represent 95% confidence interval

15  
16  
17

### Discussion

18  
19

The present study aimed to evaluate the relationship between intestinal parasites and micronutrient status in a population of school-aged children in Mexico, where the prevalence of overweight and obesity is high, lower

20 concentrations of ferritin, zinc, and vitamin C were found in children infected with *A. lumbricoides* which may be  
21 explained by different physiological and immunological pathways. For instance, impaired micronutrient absorption  
22 has been related to reduced gastrointestinal function, to damage of the gut mucosa and to competition for available  
23 micronutrients when infected by STHs (Hesham, Edariah, & Norhayati, 2004). In addition, lower intake of  
24 micronutrients and food has been associated with STH infection, due to symptoms such as abdominal pain and loss  
25 of appetite (Crompton & Nesheim, 2002; Greichus & Greichus, 1980; Northrop et al., 1987; Stephenson, Latham, &  
26 Ottesen, 2000; Zavala et al., 2017). However, the possibility that children with lower ferritin, zinc and vitamin C  
27 concentrations are more likely to be infected should also be considered. Iron, zinc and vitamin C are immuno-  
28 modulators and play a role in the immune system responses that occur at mucosal membranes (Liu et al., 2016;  
29 Maggini, Wenzlaff, & Hornig, 2010). Additionally, vitamin C is a exogenous antioxidant, necessary for collagen  
30 synthesis required for the adequate function of the epithelial barrier (Liu et al., 2016; Maggini, Wenzlaff, & Hornig,  
31 2010). The association found between STH infection and lower plasma zinc and ferritin concentrations is in line with  
32 several studies in different age-groups and populations that have found similar associations (Adebara, Ernest, &  
33 Ojuawo, 2011; G. O. Arinola et al., 2015; de Gier et al., 2015). In contrast, there are no available studies evaluating  
34 the relationship between vitamin C and STHs. No association between any of the STH infections and serum vitamin  
35 E was found. According to a systematic review, few available studies have addressed the relationship between vitamin  
36 E and STH infection.(B. de Gier et al., 2014).

37 The higher concentrations of circulating ferritin, vitamin B-12 and vitamin E:lipids found in children infected with  
38 intestinal protozoa may be explained by possible a higher intake of animal source foods. In a previous study from the  
39 same cohort, we observed that children infected with *E.coli* have higher intakes of meat, poultry, fish and eggs which  
40 are the main sources of vitamin B-12 and bioavailable iron in the diet (Zavala et al., 2017). To the best of our  
41 knowledge, only one study has evaluated the association between intestinal protozoa and micronutrient concentrations.  
42 In that study no association was observed between *E. coli* infection and serum iron, retinol, zinc or vitamin B-12  
43 concentrations in the children (Boeke et al., 2010). Since it has been shown that *E. coli* infection is associated with  
44 higher food intake, discrepancy among studies may be attributed to differences in food group intake (i.e. seafood,  
45 meat, egg and legumes), food quality, availability and affordability. Is important to highlight that despite the retinol  
46 associations with *E. coli* and *E. nana* being significant they may not be biologically or clinically relevant..

47 Given the distinct pathogenic profiles of the parasites studied, our findings require careful interpretation. *A.*  
48 *lumbricoides* and *B. coli*, which are recognized as pathogenic, were associated with lower serum concentrations of  
49 ferritin, zinc, and vitamin C, supporting their known impact on nutrient absorption and host health. In contrast, *E. coli*  
50 and *E. nana*, generally considered non-pathogenic, were associated with higher serum levels of vitamin B12 and  
51 ferritin. This may reflect differences in dietary intake or host responses among infected children, as previously reported  
52 in studies linking *E. coli* infection to higher consumption of animal-source foods (Zavala et al., 2017). These findings  
53 highlight the importance of differentiating between pathogenic and non-pathogenic parasites when interpreting the  
54 health and nutritional impacts of intestinal infections. Further research is warranted to explore whether these  
55 associations are mediated by dietary, immunological, or microbiota-related mechanisms.

56 Plasma concentrations of zinc were sensitive to STH infections in children with high body fat. The lower zinc  
57 concentrations in the STH infected children that had high body fat content might be caused by zinc malabsorption  
58 associated with both obesity and STH infection (Ganiyu Olatunbosun Arinola et al., 2015). In addition, as zinc is  
59 necessary for immune function, children with lower zinc concentrations are more likely to become infected with  
60 intestinal parasites (Brown et al., 2004; Maares & Haase, 2016). Thus, obesity together with low zinc concentrations  
61 may have a larger weakening effect on the immune response and defense against STHs.

62 The interaction between serum retinol and intestinal protozoa infection differed by body fat content. In the children  
63 with high body fat, the lower serum retinol concentrations in the intestinal protozoa infected children might be  
64 attributed to an inflammation process, specifically a T-helper type 1 (Th1) pro-inflammatory response, associated to  
65 both infection and obesity (Kirkpatrick et al., 2002). A Th1 response has been associated with lower serum retinol  
66 concentrations (García, 2012; Thurnham et al., 2003; Zavala et al., 2013). In contrast, in the children with normal  
67 body fat, the higher serum retinol concentrations observed in the protozoa infected children might be attributed to a  
68 higher food intake (Zavala et al., 2017).

69 As far as we know, this is the first study assessing the relationship between micronutrients and intestinal parasites  
70 while taking into consideration the interaction with body composition. These findings and possible underlying  
71 mechanisms that may lead to the observed differences, especially those between serum zinc and serum retinol  
72 concentrations, parasitic infections and body fat need to be further explored. Some methodological limitations of our  
73 study need to be addressed. Due to the cross-sectional study design, we were unable to determine the causal  
74 relationship between intestinal parasitic infections, micronutrient concentrations and body composition. In this

75 population the most prevalent STH was *A. lumbricoides* with 44 cases, followed by hookworm with only 6 cases, thus  
76 the association between STH and micronutrients in this study are mostly driven by *A. lumbricoides* infection, which  
77 should be taken into consideration while interpreting the results. Parasitic infection prevalence was determined by  
78 parasitological examination of one stool sample per individual which may underestimate the number of infected  
79 children (Staat et al., 2011). However, the prevalence of the studied intestinal parasites was high and similar to that  
80 reported previously in rural Mexico (Quihui-Cota & Morales-Figueroa, 2012). Our analyses and interpretations were  
81 limited for several micronutrients because of high status (i.e. folate, vitamin E) or because of absence of deficiency  
82 (i.e. vitamin B-12), affecting our ability to detect associations or interactions. Also, we did not measure all the  
83 biochemical markers available to determine status, but this study covered a large number of vitamins and minerals as  
84 it was intended to contribute to the knowledge of overall micronutrient deficiencies.

85 It is important to consider that Mexico has moved from a situation where a high prevalence of micronutrient  
86 deficiencies was a major concern to one where status of many has improved. For example, iron deficiency (serum  
87 ferritin <15 µg/L) has gone from 13% to 9.3% in school-aged children in a period of 6 years.<sup>23,24</sup> The prevalence of  
88 marginal and deficient vitamin B12 status also has been shown to be much lower in recent published data than two  
89 decades ago (Anaya-Loyola et al., 2019). Improvement has been achieved by the success of national programs such  
90 as Prospera, and voluntary fortification of foods (such as ready-to-eat cereals, dairy products among others) and flour  
91 fortification with iron, zinc and folic acid. Although there have been improvements in the micronutrient status of the  
92 Mexican population, overweight and obesity have increased, but we also found a high proportion of children with  
93 elevated folate status. None of them presented folate values <4.41 ng/mL; 99.7% of them had values above 13.2  
94 ng/mL, being 84.8% above the upper limit of the standard curve for the assay (>19.9 ng/mL). Our finding on excessive  
95 folate represents an alarm for public health in Mexico and this issue should be addressed in the near future. In the  
96 same sense, the prevalence of parasitic infections remains high. While some associations with micronutrient status  
97 were observed, the low prevalence of deficiency of some micronutrients and the situation of folate repletion likely  
98 made it difficult to detect more associations of the parasites on status that might occur in more micronutrient deficient  
99 populations.

## 100 **Conclusion**

101 In conclusion, STH and intestinal protozoa have different and opposite associations with micronutrient concentrations.  
102 STH infection was associated with lower concentrations of zinc and vitamin C and intestinal protozoa infection with

103 higher concentrations of iron and vitamin B12. The associations between zinc and vitamin A status with intestinal  
104 parasites differed by body fat percentage. Interestingly, non-pathogenic parasites like *E. coli* and *E. nana* were linked  
105 to higher serum levels of vitamin B12 and ferritin. While these parasites are not considered harmful, they may still  
106 influence gut microbiome composition, appetite, and dietary intake, potentially altering nutrient absorption or  
107 metabolism. Further studies with better epidemiological design are needed to explore the relationship between multiple  
108 micronutrients and parasites in populations with co-existing high prevalence of micronutrient deficiencies and  
109 overweight/obesity.

## 110 **Statements and Declarations**

111 **Funding:** The present study was partially funded by Consejo Nacional de Ciencia y Tecnología (CONACYT),  
112 Mexico.

113 **Competing interest:** None

114 **Authors contributions:** OPG, GAZ, MCP and JLR designed and conceived the study. MC and AB conducted  
115 laboratory analyses at UAQ in Querétaro, Mexico, and at the WHNRC in Davis, USA, respectively. OPG and GAZ  
116 performed statistical analyses. OPG, GAZ, MCP, CMD, AB, KP, JLR and LHA interpreted data and provided  
117 intellectual input to the drafts of the manuscript. All authors were involved in writing the paper. All authors read and  
118 approved the final version of the manuscript. OPG has final responsibility for all parts of the manuscript.

119 **Data availability:** The datasets generated during and/or analysed during the current study are available from the  
120 corresponding author on reasonable request.

121 **Ethical approval:** The study was conducted according to the guidelines of the Declaration of Helsinki, and the study  
122 protocol was approved by the Bioethics Committee of the School of Natural Sciences at the Universidad Autónoma  
123 de Querétaro (UAQ) (protocol # FNN-2014-02). The authors assert that all procedures contributing to this work  
124 comply with the ethical standards of the relevant national and institutional committees on human experimentation and  
125 with the Helsinki Declaration of 1975, as revised in 2008.

126 **Consent to participate:** Informed consent was obtained from all parents of children that participants in the study.

127 **Acknowledgements:** We appreciate the general input and proofreading provided by Professor Michael La Frano from  
128 California Polytechnic State University. We thank the children and their parents for their consent to participate in our  
129 study. Also, we thank Martha Soto, Karen Torres, Andrea Hernández, Paola Flores, Shadia Elian, Lorena Oropeza,

130 Sara de los Cobos, Guadalupe Martínez and Ana Pérez from the Human Nutrition Department, School of Natural  
131 Sciences, UAQ, for their assistance in the collection and analysis of the samples.

132

133 **References**

- 134 Adebara, O. V., Ernest, S. K., & Ojuawo, I. A. (2011). Association between intestinal helminthiasis and serum  
 135 ferritin levels among school children. *Open Journal of Pediatrics, 1*(02), 12.
- 136 Alum, A., Rubino, J. R., & Ijaz, M. K. (2010). The global war against intestinal parasites—should we use a holistic  
 137 approach? *International Journal of Infectious Diseases, 14*(9), e732-e738.
- 138 Anaya-Loyola, M. A., Brito, A., Villalpando, S., & Allen, L. H. (2019). Prevalence of low serum vitamin B12 in  
 139 Mexican children and women: results from the first National Nutrition Survey (1999) as a basis for  
 140 interventions and progress. *International Journal for Vitamin and Nutrition Research.*
- 141 Arinola, G. O., Morenikeji, O. A., Akinwande, K. S., Alade, A. O., Olateru-Olagbegi, O., Alabi, P. E., & Rahamon,  
 142 S. K. (2015). Serum micronutrients in helminth-infected pregnant women and children: suggestions for  
 143 differential supplementation during anti-helminthic treatment. *Annals of global health, 81*(5), 705-710.
- 144 Arinola, G. O., Morenikeji, O. A., Akinwande, K. S., Alade, A. O., Olateru-Olagbegi, O., Alabi, P. E., & Rahamon,  
 145 S. K. (2015). Serum Micronutrients in Helminth-infected Pregnant Women and Children: Suggestions for  
 146 Differential Supplementation During Anti-helminthic Treatment. *Ann Glob Health, 81*(5), 705-710.  
 147 <https://doi.org/10.1016/j.aogh.2015.10.001>
- 148 Bhaskaram, P. (2002). Micronutrient malnutrition, infection, and immunity: an overview. *Nutrition Reviews, 60*(s5).
- 149 Black, R. (2003). Micronutrient deficiency: an underlying cause of morbidity and mortality. *Bulletin of the World  
 150 Health Organization, 81*(2), 79-79.
- 151 Black, R. E., Victora, C. G., Walker, S. P., Bhutta, Z. A., Christian, P., De Onis, M., Ezzati, M., Grantham-  
 152 McGregor, S., Katz, J., & Martorell, R. (2013). Maternal and child undernutrition and overweight in low-  
 153 income and middle-income countries. *The lancet, 382*(9890), 427-451.
- 154 Boeke, C. E., Mora-Plazas, M., Forero, Y., & Villamor, E. (2010). Intestinal protozoan infections in relation to  
 155 nutritional status and gastrointestinal morbidity in Colombian school children. *Journal of tropical  
 156 pediatrics, 56*(5), 299-306.
- 157 Brito, A., Mujica-Coopman, M. F., Olivares, M., Lopez de Romana, D., Cori, H., & Allen, L. H. (2015). Folate and  
 158 vitamin B12 status in Latin America and the Caribbean: an update. *Food and nutrition bulletin,  
 159 36*(2\_suppl), S109-S118.
- 160 Brown, K. H., Rivera, J. A., Bhutta, Z., Gibson, R. S., King, J. C., Lonnerdal, B., Ruel, M. T., Sandtrom, B.,  
 161 Wasantwisut, E., & Hotz, C. (2004). International Zinc Nutrition Consultative Group (IZiNCG) technical  
 162 document #1. Assessment of the risk of zinc deficiency in populations and options for its control. *Food  
 163 Nutr Bull, 25*(1 Suppl 2), S99-203.
- 164 Campos Ponce, M., Incani, R. N., Pinelli, E., ten Kulve, N., Ramak, R., Polman, K., & Doak, C. M. (2012). Are  
 165 intestinal parasites fuelling the rise in dual burden households in Venezuela? *Transactions of The Royal  
 166 Society of Tropical Medicine and Hygiene, 107*(2), 119-123.
- 167 Crompton, D. W., & Nesheim, M. C. (2002). Nutritional impact of intestinal helminthiasis during the human life  
 168 cycle. *Annu Rev Nutr, 22*, 35-59. <https://doi.org/10.1146/annurev.nutr.22.120501.134539>
- 169 Cuevas-Nasu, L., Levy, T. S., Arcos, M. A. A., Gomez, I. M., & Dommarco, J. R. (2017). Overweight and obesity  
 170 in Mexican children and adolescents: Data from National Nutrition and Health Surveys in 2016. *The  
 171 FASEB Journal, 31*(1 Supplement), 640.635-640.635.
- 172 de Gier, B., Campos Ponce, M., van de Bor, M., Doak, C. M., & Polman, K. (2014). Helminth infections and  
 173 micronutrients in school-age children: a systematic review and meta-analysis. *Am J Clin Nutr, 99*(6), 1499-  
 174 1509. <https://doi.org/10.3945/ajcn.113.069955>
- 175 de Gier, B., Mpabanzi, L., Vereecken, K., van der Werff, S. D., D'Haese, P. C., Fiorentino, M., Khov, K., Perignon,  
 176 M., Chamnan, C., Berger, J., Parker, M. E., Diaz, R. J., Nunez, F. A., Rivero, L. R., Gorbea, M. B., Doak,  
 177 C. M., Ponce, M. C., Wieringa, F. T., & Polman, K. (2015). Height, zinc and soil-transmitted helminth  
 178 infections in schoolchildren: a study in Cuba and Cambodia. *Nutrients, 7*(4), 3000-3010.  
 179 <https://doi.org/10.3390/nu7043000>
- 180 de Gier, B., Ponce, M. C., van de Bor, M., Doak, C. M., & Polman, K. (2014). Helminth infections and  
 181 micronutrients in school-age children: a systematic review and meta-analysis. *The American Journal of  
 182 Clinical Nutrition, 99*(6), 1499-1509.
- 183 Drewel, B. T., Giraud, D. W., Davy, S. R., & Driskell, J. A. (2006). Less than adequate vitamin E status observed in  
 184 a group of preschool boys and girls living in the United States. *J Nutr Biochem, 17*(2), 132-138.  
 185 <https://doi.org/10.1016/j.jnutbio.2005.06.003>
- 186 Ellis, K. J., Abrams, S. A., & Wong, W. W. (1997). Body composition of a young, multiethnic female population.  
 187 *The American journal of clinical nutrition, 65*(3), 724-731.

188 Fares, S., Chahed, M. K., Feki, M., Beji, C., Traissac, P., El Ati, J., & Kaabachi, N. (2011). Status of vitamins A and  
189 E in schoolchildren in the centre west of Tunisia: a population-based study. *Public health nutrition*, 14(02),  
190 255-260.

191 García, O. P. (2012). Effect of vitamin A deficiency on the immune response in obesity. *Proceedings of the*  
192 *Nutrition Society*, 71(2), 290-297.

193 García, O. P., Zavala, G. A., Campos-Ponce, M., Doak, C. M., Polman, K., & Rosado, J. L. (2023). Parasitic  
194 Infection, Obesity, and Micronutrient Deficiencies in School-Aged Children in Mexico. In *Human Growth*  
195 *and Nutrition in Latin American and Caribbean Countries* (pp. 387-396). Springer.

196 García, O. P., Zavala, G. A., Campos-Ponce, M., & Polman, K. (2023). Nutrition and parasitism. In *Encyclopedia of*  
197 *Human Nutrition: Volume 3* (pp. 541-548). Elsevier.

198 Gibson, R. S. (2005). *Principles of nutritional assessment*. Oxford university press, USA.

199 Greichus, A., & Greichus, Y. A. (1980). Identification and quantification of some elements in the hog roundworm,  
200 *Ascaris lumbricoides suum*, and certain tissues of its host. *Int J Parasitol*, 10(2), 89-91.

201 Gutiérrez-Jiménez, J., Luna-Cazáres, L. M., & Vidal, J. E. (2017). Malnutrition and Intestinal Parasites: Mexico  
202 Perspectives. *Handbook of Famine, Starvation, and Nutrient Deprivation. From biology to policy*. Cham:  
203 Springer International Publishing, 1-18.

204 Hesham, M., Edariah, A., & Norhayati, M. (2004). Intestinal parasitic infections and micronutrient deficiency: a  
205 review. *Med J Malaysia*, 59(2), 284-293.

206 Hotez, P. J., Alvarado, M., Basáñez, M.-G., Bolliger, I., Bourne, R., Boussinesq, M., Brooker, S. J., Brown, A. S.,  
207 Buckle, G., & Budke, C. M. (2014). The global burden of disease study 2010: interpretation and  
208 implications for the neglected tropical diseases. *PLoS Negl Trop Dis*, 8(7), e2865.

209 Hotz, C., Peerson, J. M., & Brown, K. H. (2003). Suggested lower cutoffs of serum zinc concentrations for assessing  
210 zinc status: reanalysis of the second National Health and Nutrition Examination Survey data (1976-1980).  
211 *Am J Clin Nutr*, 78(4), 756-764.

212 Katona, P., & Katona-Apte, J. (2008). The interaction between nutrition and infection. *Clinical Infectious Diseases*,  
213 46(10), 1582-1588.

214 Kirkpatrick, B. D., Daniels, M. M., Jean, S. S., Pape, J. W., Karp, C., Littenberg, B., Fitzgerald, D. W., Lederman,  
215 H. M., Nataro, J. P., & Sears, C. L. (2002). Cryptosporidiosis stimulates an inflammatory intestinal  
216 response in malnourished Haitian children. *The Journal of infectious diseases*, 186(1), 94-101.

217 LeBlanc, E., Chou, R., Zakher, B., Daeges, M., & Pappas, M. (2014). U.S. Preventive Services Task Force Evidence  
218 Syntheses, formerly Systematic Evidence Reviews. In *Screening for Vitamin D Deficiency: Systematic*  
219 *Review for the U.S. Preventive Services Task Force Recommendation*. Agency for Healthcare Research and  
220 Quality (US).

221 Liu, F., Cottrell, J. J., Furness, J. B., Rivera, L. R., Kelly, F. W., Wijesiriwardana, U., Pustovit, R. V., Fothergill, L.  
222 J., Bravo, D. M., Celi, P., Leury, B. J., Gabler, N. K., & Dunshea, F. R. (2016). Selenium and Vitamin E  
223 together improve intestinal epithelial barrier function and alleviate oxidative stress in heat stressed pigs.  
224 *Exp Physiol*. <https://doi.org/10.1113/ep085746>

225 Maares, M., & Haase, H. (2016). Zinc and immunity: An essential interrelation. *Archives of biochemistry and*  
226 *biophysics*, 611, 58-65.

227 Maggini, S., Wenzlaff, S., & Hornig, D. (2010). Essential role of vitamin C and zinc in child immunity and health. *J*  
228 *Int Med Res*, 38(2), 386-414.

229 Maggini, S., Wintergerst, E. S., Beveridge, S., & Hornig, D. H. (2007). Selected vitamins and trace elements support  
230 immune function by strengthening epithelial barriers and cellular and humoral immune responses. *British*  
231 *Journal of Nutrition*, 98(S1), S29-S35.

232 Morales-Espinoza, E. M., Sánchez-Pérez, H. J., García-Gil, M. d. M., Vargas-Morales, G., Méndez-Sánchez, J. D.,  
233 & Pérez-Ramírez, M. (2003). Intestinal parasites in children, in highly deprived areas in the border region  
234 of Chiapas, Mexico. *Salud Publica Mex*, 45(5), 379-388.

235 Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., Margono, C., Mullany, E. C., Biryukov, S., Abbafati,  
236 C., & Abera, S. F. (2014). Global, regional, and national prevalence of overweight and obesity in children  
237 and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The*  
238 *Lancet*, 384(9945), 766-781.

239 Northrop, C., Lunn, P., Wainwright, M., & Evans, J. (1987). Plasma albumin concentrations and intestinal  
240 permeability in Bangladeshi children infected with *Ascaris lumbricoides*. *Trans R Soc Trop Med Hyg*,  
241 81(5), 811-815.

242 Onis, M. (2006). WHO Child Growth Standards based on length/height, weight and age. *Acta paediatrica*,  
243 95(S450), 76-85.

244 Organization, W. H. (2011). Assessing the iron status of populations: including literature reviews: report of a Joint  
245 World Health Organization/Centers for Disease Control and Prevention Technical Consultation on the  
246 Assessment of Iron Status at the Population Level Geneva, Switzerland, 6–8 April 2004.–. In.  
247 Quihui-Cota, L., & Morales-Figueroa, G. G. (2012). Persistence of intestinal parasitic infections during the national  
248 de-worming campaign in schoolchildren of northwestern Mexico: a cross-sectional study. *Ann*  
249 *Gastroenterol*, 25(1), 57-60.

250 Rivera Dommarco, J. Á. (2012). Deficiencias de micronutrientos en México: un problema invisible de salud  
251 pública. *Salud Publica Mex*, 54(2), 101-102.

252 Schaible, U. E., & Stefan, H. (2007). Malnutrition and infection: complex mechanisms and global impacts. *PLoS*  
253 *medicine*, 4(5), e115.

254 Shamah-Levy, T., Villalpando, S., Jáuregui, A., & Rivera, J. A. (2012). Overview of the nutritional status of  
255 selected micronutrients in Mexican children in 2006. *Salud Publica Mex*, 54(2), 146-151.

256 Staat, M. A., Rice, M., Donauer, S., Mukkada, S., Holloway, M., Cassedy, A., Kelley, J., & Salisbury, S. (2011).  
257 Intestinal parasite screening in internationally adopted children: importance of multiple stool specimens.  
258 *Pediatrics*, 128(3), e613-622. <https://doi.org/10.1542/peds.2010-3032>

259 Stephenson, L. S., Latham, M. C., & Ottesen, E. (2000). Malnutrition and parasitic helminth infections.  
260 *Parasitology*, 121(S1), S23-S38.

261 Thurnham, D. I., McCabe, G., Northrop-Clewes, C., & Nestel, P. (2003). Effects of subclinical infection on plasma  
262 retinol concentrations and assessment of prevalence of vitamin A deficiency: meta-analysis. *The Lancet*,  
263 362(9401), 2052-2058.

264 Villalpando, S., Cruz, V. d. l., Shamah-Levy, T., Rebollar, R., & Contreras-Manzano, A. (2015). Nutritional status  
265 of iron, vitamin B12, folate, retinol and anemia in children 1 to 11 years old: Results of the Ensanut 2012.  
266 *salud pública de méxico*, 57(5), 372-384.

267 WHO. (1991). *Basic laboratory methods in medical parasitology*.

268 Willcox, H. P., & Coura, J. R. (1991). The efficiency of Lutz, Kato-Katz and Baermann-Moraes (adapted)  
269 techniques association to the diagnosis of intestinal helminths. *Mem Inst Oswaldo Cruz*, 86(4), 457-460.

270 Zavala, G., García, O., Camacho, M., Ronquillo, D., Campos-Ponce, M., Doak, C., Polman, K., & Rosado, J.  
271 (2018). Intestinal parasites: Associations with intestinal and systemic inflammation. *Parasite Immunology*,  
272 40(4), e12518.

273 Zavala, G., García, O., Campos-Ponce, M., Ronquillo, D., Caamaño, M., Doak, C., & Rosado, J. (2016). Children  
274 with moderate-high infection with *Entamoeba coli* have higher percentage of body and abdominal fat than  
275 non-infected children. *Pediatric obesity*, 11(6), 443-449.

276 Zavala, G., Long, K. Z., García, O. P., del Carmen Caamaño, M., Aguilar, T., Salgado, L. M., & Rosado, J. L.  
277 (2013). Specific micronutrient concentrations are associated with inflammatory cytokines in a rural  
278 population of Mexican women with a high prevalence of obesity. *British Journal of Nutrition*, 109(4), 686-  
279 694.

280 Zavala, G. A., Rosado, J. L., Doak, C. M., del Carmen Caamaño, M., Campos-Ponce, M., Ronquillo, D., Polman,  
281 K., & García, O. P. (2017). Energy and food intake are associated specific intestinal parasitic infections in  
282 children of rural Mexico. *Parasitol Int*.

283 Zavala, G. A., van Dulm, E., Doak, C. M., García, O. P., Polman, K., & Campos-Ponce, M. (2020). Ascariasis,  
284 Amebiasis and Giardiasis in Mexican children: distribution and geographical, environmental and  
285 socioeconomic risk factors. *Journal of parasitic diseases*, 44, 829-836.

286

