



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/227040/>

Version: Accepted Version

Article:

Mason-Jones, Amanda Jayne, Bottomley-Wise, Rachel Margaret, Okanlawon, Adenike A et al. (2025) Accessible, acceptable and equitable: a range of contraceptive methods are still needed. *BMJ Sexual & Reproductive Health*. ISSN: 2515-1991

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Correspondence in response to:

<https://srh.bmj.com/content/early/2025/01/01/bmj-srh-2024-202573.long>

Title: Accessible, acceptable and equitable: a range of contraceptive methods are still needed

Amanda J. Mason-Jones, Department of Health Sciences, University of York, YO10 5DD. ORCID <https://orcid.org/0000-0002-4292-3183>

Rachel Margaret Bottomley-Wise. York Trials Unit, Department of Health Sciences, University of York, York, YO10 5DD, United Kingdom. ORCID <https://orcid.org/0000-0001-6972-0100>

Adenike A. Okanlawon, York Trials Unit, Department of Health Sciences, University of York, York, YO10 5DD. ORCID <https://orcid.org/0009-0002-5839-3292>

Amie Woodward, Institute for Health and Care Improvement, York St John University, Lord Mayor's Walk, YO31 7EX. ORCID is <https://orcid.org/0000-0002-9579-4012>.

Rosemary Greenwood, Senior Statistician, York Trials Unit, Department of Health Sciences, University of York, YO10 5DD

McNee and colleagues explore the change in contraceptive methods among those seeking abortion in 2018 and 2023 using cross-sectional self-reported data from the British Pregnancy Advisory Service (BPAS). The BPAS is highly valued and provides much needed accessible reproductive health services and the paper raises important issues.

The analysis highlights the more commonplace use of medical terminations earlier in pregnancy. They also report that the use of long-acting reversible contraception fell from 3% to 0.6%, and hormonal contraception methods from 18.8% to 11.3% in their population. However, the analysis does not necessarily support their conclusion that use of 'fertility awareness-based methods' (FABM) are associated with attending BPAS for abortion. The data suggest that the decline between the chosen years in the use of hormonal contraceptives and the increase in FABM is a trend but it does not imply a causal link with more people seeking abortions.

The authors seem to selectively focus on FABM. For example, in "What this Study Adds" the focus is on the shift from 'reliable hormonal methods' to 'less reliable FABM' with no mention of the greater increase in women not using contraception. Meanwhile, 'other method' use decreased from 22% to 16%. This decrease potentially goes against the argument that people are shifting from hormonal to non-hormonal based methods. The authors do not consider that with FABM, people often use a mixed method approach e.g., a condom on fertile days, that greater awareness through FABM might be driving the earlier less invasive forms of termination or that the older age of those seeking terminations might suggest success in avoiding pregnancy in the younger age groups.

The analysis which includes limited detail, shows that there is simply an association between the year and method of contraception recorded. Although they acknowledge

that the transition to different data collection methods could result in bias, it would have been ideal to account for such confounding factors through multivariate or sensitivity analyses. ¹ Additionally, the category 'None', includes 'no method reported' or 'unsure' which is not the same as 'none'. This is likely to have biased their analysis.

The 'Women's Health Strategy for England', has highlighted contraception as a crucial component. ² The reality for many is that this has not necessarily translated into accessible equitable preventative healthcare. Despite women needing and wanting contraception many women do not have access to modern, or indeed any contraception methods or the ability to control their fertility. ³ This is particularly pronounced for some populations including young women, ⁴ minoritised communities,⁵ or those who are incarcerated, who encounter additional challenges restricting access. ⁶

'Hormonal hesitancy' concerns are very real, ⁵ and as evidence of other health risks, related to hormonal contraception, such as breast cancer, are published ⁷ simply attributing this to the influence of social media is simplistic; Users also draw on embodied knowledge and personal experience to make decisions. ^{8,9} We concur that although developments in reproductive health technology, such as FABM have taken place, we have not reached the 'techno-utopia' ¹⁰ we might dream of. This technology often over-promises and underdelivers.

The paper fails to provide sufficient information to determine the multifactorial drivers. A missed opportunity. To address these issues, a concerted effort is needed to conduct and publish responsible and robust research that can help us to understand contraceptive decision making and barriers to access and use. The good news story here may be about the increase in the proportion of medically managed terminations rather than mistrusting women's ability to manage technology.

- 1 Pérez-Guerrero EE, Guillén-Medina MR, Márquez-Sandoval F, Vera-Cruz JM, Gallegos-Arreola MP, Rico-Méndez MA, Aguilar-Velázquez JA, Gutiérrez-Hurtado IA. Methodological and statistical considerations for cross-sectional, case-control, and cohort studies. *Journal of Clinical Medicine* 2024; 13: 4005
- 2 Department of Health and Social Care. Women's Health Strategy for England volume 2023, 2022
- 3 Paltrow LM, Harris LH, Marshall MF. Beyond abortion: The consequences of overturning Roe. *The American Journal of Bioethics* 2022; 22: 3-15
- 4 Turnbull G, Scott RH, Mann S, Wellings K. Accessing emergency contraception pills from pharmacies: the experience of young women in London. *BMJ Sexual & Reproductive Health* 2021; 47: 27-31
- 5 Cory RJ, Mawson R, Linton E, Al-Bazz DY, Fryer K, Ma R, Mitchell CA. Influences on ethnic minority women's experiences and access to contraception in the UK: a systematic qualitative evidence synthesis. *BMJ Sexual & Reproductive Health* 2025; 51: 64-73

- 6 Megha Ramaswamy H-FC, Karen L. Cropsey, Jennifer G. Clarke, and Patricia J. Kelly. Highly Effective Birth Control Use Before and After Women's Incarceration. *Journal of Women's Health* 2015; 24: 530-539
- 7 Fitzpatrick D, Pirie K, Reeves G, Green J, Beral V. Combined and progestagen-only hormonal contraceptives and breast cancer risk: A UK nested case–control study and meta-analysis. *Plos Medicine* 2023; 20: e1004188
- 8 Littlejohn KE. Hormonal contraceptive use and discontinuation because of dissatisfaction: differences by race and education. *Demography* 2012; 49: 1433-1452
- 9 Lowe P. Embodied expertise: women's perceptions of the contraception consultation. *Health*: 2005; 9: 361-378
- 10 Willems W, Heltzel A, Nabuurs J, Broerse J, Kupper F. Welcome to the fertility clinic of the future! Using speculative design to explore the moral landscape of reproductive technologies. *Humanities & Social Sciences Communications* 2023; 10