

This is a repository copy of *Exploring perspectives regarding death cafés for people experiencing homelessness:a qualitative study*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/227533/>

Version: Published Version

---

**Article:**

Crooks, Jodie, Davis, Gareth, Pattinson, Mandy et al. (4 more authors) (2025) Exploring perspectives regarding death cafés for people experiencing homelessness:a qualitative study. BMJ Open. e096990. ISSN 2044-6055

<https://doi.org/10.1136/bmjopen-2024-096990>

---


**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here:  
<https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# BMJ Open Exploring perspectives regarding death cafés for people experiencing homelessness: a qualitative study

Jodie Crooks <sup>1</sup>, Gareth Davis,<sup>2</sup> Mandy Pattinson,<sup>2</sup> Emma Casey,<sup>3</sup> Caroline Shulman,<sup>2,4</sup> Kate Flemming,<sup>5</sup> Briony Hudson<sup>1,2,4</sup>

**To cite:** Crooks J, Davis G, Pattinson M, *et al.* Exploring perspectives regarding death cafés for people experiencing homelessness: a qualitative study. *BMJ Open* 2025;**15**:e096990. doi:10.1136/bmjopen-2024-096990

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-096990>).

Received 22 November 2024  
Accepted 28 May 2025

## ABSTRACT

**Background** People experiencing homelessness (PEH) often die at a younger age than the general population. Advanced ill-health often occurs late in its trajectory (if at all), leaving many PEH to die without adequate support from hospital-based, hospice-based or community-based palliative care services. Despite the high rate of bereavement and exposure to death among PEH, there are rarely opportunities for them to reflect on their experiences, thoughts and preferences around death or receive bereavement support. Death cafés are a global social franchise, providing a space for people to participate in an open group discussion about death, dying and bereavement. They are free to attend and unstructured but facilitated.

**Objectives** To explore the perspectives of PEH and the professionals that support them with regards to the potential acceptability, benefits and challenges of death cafés.

**Methods** A qualitative study comprising of interviews with nine homelessness and/or palliative care professionals with experience of running death cafés for, or supporting PEH in other ways, and two focus groups with nine people with lived experience of homelessness. The research was co-produced with a lived experience co-researcher forming a core part of the research team. Reflexive thematic analysis was utilised to develop themes.

**Results** Five themes were developed to highlight the perspectives and experiences of death cafés for PEH: choice; expertise in facilitation: recognising risk and the unstructured nature; promoting safety against potential risk; potential benefits and potential practical considerations. Themes highlight the need for a trauma-informed approach. Death cafés could be impactful, where done well, but they are not without risk.

**Conclusions** Death cafés for PEH pose risks that require comprehensive consideration and trauma-informed, expert facilitation. However, this research highlighted that a low-pressure space to explore thoughts around death, dying and bereavement is currently an unmet need within this population.

## BACKGROUND

### Homelessness, death & bereavement

People experiencing homelessness (PEH) often die prematurely at a younger age than the general population.<sup>1 2</sup> People with long histories of homelessness may also experience accelerated ageing and associated life-limiting, long-term conditions (including

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This research was co-produced by academic researchers and people with lived experience of homelessness.
- ⇒ Relationships developed through the coproduction process enabled fast rapport with participants throughout data collection and adds nuance to the analysis.
- ⇒ As recruitment occurred opportunistically, there is a risk of sampling bias including self-selection bias and/or bias introduced through snowball methods.
- ⇒ Further research may benefit from a larger, more diverse sample of people with lived experience of homelessness, as evidence demonstrates the complexity and heterogeneity of experiences.

frailty) as early as in their 40s.<sup>3</sup> International evidence consistently demonstrates the high prevalence of multimorbidity among people with experience of homelessness.<sup>4-7</sup> For many PEH, healthcare access and outcomes are influenced by intersecting factors related to trauma, physical health, mental health and exclusion.<sup>8</sup>

Many PEH are not recognised as living with advanced ill-health until late in its trajectory, if at all, for reasons including barriers to health service access and unpredictable illness trajectories.<sup>9 10</sup> In the UK in 2023, 1474 people died while being homeless; the equivalent of one person every 6 hours.<sup>11</sup> Importantly, one study looking at 3882 PEH in England found that one in three deaths among PEH were due to causes amenable to being prevented or managed by timely and effective healthcare.<sup>12</sup>

Many PEH with advanced ill-health die without adequate support from relevant health and social care services (including palliative care), leading to poor experiences and undignified deaths.<sup>10 13</sup> In addition, the majority of people with multiple-exclusion homelessness have experienced significant trauma in their lives, often starting in



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

<sup>1</sup>Marie Curie, London, UK

<sup>2</sup>Pathway, London, UK

<sup>3</sup>Groundswell, London, UK

<sup>4</sup>Marie Curie Palliative Care Research Department, University College, London, UK

<sup>5</sup>University of York, York, UK

## Correspondence to

Dr Briony Hudson;  
[briony.hudson@mariecurie.org.uk](mailto:briony.hudson@mariecurie.org.uk)

childhood: one study reported up to 94% of people who have experienced homelessness also identified as having at least one traumatic experience.<sup>14</sup> Furthermore, PEH have often experienced bereavements and loss among their peers: in the UK, the estimated number of deaths among PEH has increased by 54% since records began in 2013, representing an extra 259 deaths.<sup>15</sup>

Despite the high rate of bereavement and exposure to death,<sup>16</sup> a high palliative care need and limited access to health and social care services among this group, there are infrequent opportunities for PEH to explore their own thoughts and preferences around death and the end of their lives or receive bereavement support.

### Death Cafés

Death cafés have become a social franchise, expanding quickly across Europe, North America and Australasia; to date, over 16000 cafés have been held.<sup>17</sup> Death cafés are unstructured meetings which aim to provide a space for people to participate in an open group discussion of death alongside refreshments.<sup>17</sup> It is a space ‘where people talk about whatever is on their minds related to death, dying, and bereavement’ with the aim of helping ‘people to make the most of their (finite) lives’<sup>17 18</sup> (page 153). There is no specified agenda to the meetings, with no intention of directing people to any course of (medical) action. Importantly, death cafés are intentionally not grief support groups, educational sessions or a source for information on end-of-life topics.<sup>18</sup>

Very few death cafes, thus far, have been held specifically for PEH, despite the outlined increased exposure to death and bereavement. This could be for various reasons, including: low engagement with services and competing priorities day-to-day, such as sourcing food or shelter, which are likely to take priority over discussions of death and bereavement.<sup>19</sup> In addition, professionals from health and homelessness services and PEH have reported finding death difficult to discuss with staff, fearing that raising issues related to palliative care may be detrimental to service users’ well-being and their relationships with them.<sup>20</sup> Furthermore, hostels and other frontline services that support PEH are often recovery-focused in nature, aiming to support people ‘out of’ homelessness. This forms a juxtaposition between the nature of many homelessness services against talking about death and dying, which has been cited as a barrier for PEH to access palliative care services.<sup>20</sup>

Homelessness is associated with loss in a variety of ways.<sup>21</sup> Loss may be related to stable housing, loss of identity due to changed circumstances or the loss of family, friends or peers. Bereavement has been identified as a common route to homelessness.<sup>22</sup>

Given the likely high exposure to death and dying among PEH, exploration and consideration need to be given to ways in which conversations around these issues can be facilitated for people that wish to engage with them. As formal bereavement services may be less accessible to this population, consideration needs to be given

to community-based initiatives, such as death cafes, that can provide space for people to discuss issues around dying and loss. However, in light of the high prevalence of ill health, trauma, exposure to loss and the challenges to accessing health and social care support, it is crucial that we understand the views of PEH and those that support them with regards to death cafés. This paper explores whether death cafés could be helpful for PEH, and what adaptations may be needed to traditional death café approaches to tailor them for this group.

### Aim

This research aimed to explore the perspectives of PEH and the professionals that support them with regards to the potential acceptability, benefits and challenges of death cafes.

The core research question was: what are the views and experiences of people with lived experience of homelessness, health and social care professionals, including health and homelessness professionals, around death cafés for PEH?

## METHODOLOGY

### Design

A coproduced, qualitative approach was taken to explore the perspectives of people with lived experience and the professionals that support them. An exploratory stance was taken to begin to tentatively investigate an area yet to be addressed by research.<sup>23</sup> This approach is defined by Stebbins<sup>24</sup> as ‘a broad-ranging, purposive undertaking designed to maximise the discovery of generalisations leading to description and understanding of an area of social or psychological life’. Reporting of this research is guided by the Reflexive Thematic Analysis Reporting Guidelines<sup>25</sup>. Ethical approval was sought and approved by University College London (ID: 6202/011).

### Population

Participants were people with lived experience of homelessness and associated professionals who worked within the sectors of homelessness and/or palliative care. Homelessness was self-defined by participants and included street homelessness (ie, people who sleep on the streets, without shelter), hostel or other temporary accommodation and sofa-surfing.<sup>26</sup> Some professionals had been involved in death cafés for PEH previously. However, given the extremely small number of previous death cafés for this group, professionals were eligible to participate if they had experience of working closely with PEH. Participant demographics can be seen in [table 1](#). The relatively small number of participants reflects the infancy of considering death cafés for PEH, while representing a diverse range of professional and lived experience perspectives. The study took place remotely, within participants located across England, UK.

### Recruitment

An opportunistic approach was taken to sampling. Project details were shared within existing networks attended by

**Table 1** Participant demographics

Demographic	Categories	Number
Professionals		
Gender	Female	9
	Male	0
Employer organisation type	Charity	3
	Hospice	2
	Housing association / hostel	4
Role	Palliative care professional	3
	Clinical psychologist	2
	Counsellor	1
	Social worker	1
	Hostel staff	2
	Registered nurse	1
People with lived experience homelessness		
Gender	Female	6
	Male	3

professionals who may be eligible and interested in the study. These included the National Homelessness and Palliative Care Network and informal working groups.

Groundswell, a UK leading peer advocacy organisation, led the recruitment of people with lived experience of homelessness. Groundswell works collaboratively over the long term with a large group of experts by experience, supporting them to get involved in research projects. A peer coordinator employed by Groundswell shared project details with the group, both via email and in-person at regular meetings.

Interested and eligible participants were provided with a participant information sheet and given a minimum of 24 hours to consider participation. People with lived experience of homelessness also discussed the information sheet with the peer coordinator at Groundswell. All participants were given the opportunity to ask questions. Informed consent was gathered and recorded via RedCap online software.

### Data collection

Data were collected between March and May 2024. Online, semistructured interviews were conducted with professionals via Microsoft Teams. Data were collected remotely to enable recruitment of professionals situated across England, UK within the budgetary scope of the research. Interviews were co-delivered by the principal researcher (JC) and the lived-experience co-researcher (GD). A semi-structured approach was selected to 'permit interviews to be focused while still giving the investigator the autonomy to explore pertinent ideas that may come up in the course of the interview'.<sup>27</sup> The interview schedule was developed based on discussions within a previous research project by the authors.<sup>26</sup> Professionals were asked about their experiences of death cafés (including venue, attendees and

content), which professionals were involved in preparation and facilitation and general reflections on the death café. Where professionals hadn't been involved in a death café previously, they were asked about their views of death cafés, covering similar topics. Interviews lasted between 30 and 60 min.

People with lived experience of homelessness attended one of two online focus groups. These lasted for 90 min and were organised and facilitated by Groundswell. Discussions were semistructured using a predefined list of prompts based on the interview schedule and covered topics such as reviewing example advertisements for death cafés, perceived benefits and concerns surrounding death cafés for PEH, and practicalities such as venue, professionals and safety. Interviews and focus groups were audio recorded and transcribed. Interview and focus group schedules can be found in online supplemental material 1.

### Data analysis

Given the exploratory nature of the research generated from a relatively small number of participants' individual perspectives, inductive reflexive thematic analysis was chosen to emphasise the inescapable subjectivity of data interpretation.<sup>28</sup>

The six recursive phases of reflexive thematic analysis guided the inductive interpretation of data. Initially, immediately following each interview co-conducted by JC and GD, they reflected on the interview content and produced short, written reflective notes on pertinent or interesting aspects. To begin analysis, JC familiarised herself with all transcripts through reading and re-reading. Then, JC manually coded transcripts in NVivo qualitative analysis software, detailing inductive descriptive codes by marking similar phrases or words from the interviews and focus groups. GD also engaged in this process for 5/11 transcripts. Reflexive thematic analysis was a useful approach for enabling GD to consider the data in relation to their own lived experiences. Codes from each researcher were reviewed and combined where appropriate. JC, GD and BH collaboratively generated key themes, before refining, defining and naming themes.<sup>28</sup>

### Patient and public involvement

Involvement of people with lived experience of homelessness was central to this research project. A lived experience co-researcher was employed on a part-time (0.1 FTE), fixed-term basis. They were recruited through a partner organisation, Pathway, a homeless and inclusion health charity who advocate for and work toward equitable healthcare for PEH. One aspect of Pathway's work is to support people with lived experience of homelessness to be involved in research through working in partnership with organisations to carry out in-depth research and increase collective understanding of the links between homelessness, exclusion and health.<sup>29</sup> The lived experience co-researcher was a core member of the research team involved in conceptualising the study and its



methodology, developing the interview and focus group guides, co-interviewing during the data collection phase, independently coding transcripts and supporting analysis and inputting to this manuscript. To plan and carry out this involvement, we used our previously published TIFFIN recommendations that aim to promote the safe involvement of people with lived experience of homelessness in palliative care research.<sup>26</sup>

Additionally, four further people with lived experience of homelessness supported the analysis. Recruited through the working partnership with Pathway, Contributors with lived experience of homelessness gave feedback on initial themes and supported refining of the themes through an additional online focus group. All lived experience contributors were reimbursed in line with the National Institute for Health and Care Research guidelines for involving people in research.<sup>30</sup>

## RESULTS

Nine interviews and two focus groups with a total of 18 participants (nine professionals, nine people with lived experience of homelessness) were carried out. Professionals were from a diverse range of backgrounds including hospice staff, clinicians, mental health professionals and staff from homelessness services. Five out of the nine professionals had experience of facilitating, observing or attending a death café for PEH. Five themes were generated from the combined data: choice, expertise in facilitation, promoting safety against potential risk, potential benefits and potential practical considerations. These themes represent the participants' experiences of each stage of a death café, including preparation, delivery/attendance and aftercare. They highlight both participants' interests and concerns regarding death cafés for this population.

### Choice

Choice was seen as integral to the concept of death cafés for PEH. There was unanimous agreement from participants that attendance cannot be compulsory due to the potential risks associated with forcing people to face the emotionality of this topic, particularly where they are likely to have experienced previous trauma. To ensure autonomy in decision making is upheld, clear information should be provided on what a death café is and isn't as people cannot make an informed choice about attendance without adequate information.

As with any population, not everyone experiencing homelessness will want to attend a death café. Participants discussed how many PEH often face severe exclusion and barriers to accessing services and thus may not wish to engage with a death café. Furthermore, professionals felt that previous negative experiences of accessing services could negatively influence engagement with or expectations of a death café if it is perceived to be another service that is delivered to them and not for them.

The crew that I'm working with at the minute are the severe and multiple disadvantaged. So my cohort are the people who do not engage. They don't engage with their GP. They don't engage with secondary care...they will struggle with anything like this. (Registered Nurse, hostel)

Another reason highlighted by professionals for not wishing to engage could be people's perceptions that it was unnecessary because they were 'not dying yet'. However, PWLE homelessness expressed how they were often aware of their heightened risk of death, making attending a death café "too morbid" (PWLE).

I wouldn't have bothered going. It's more likely for me to die when I'm on the streets, and I knew that, so I wouldn't have wanted to go into somewhere and speak about death when I was homeless. (PWLE)

For PEH, their willingness to attend may depend on what is happening in their life at that time. Both professionals and PWLE suggested that death cafés "might attract the people who are further along their recovery journey [out of homelessness and/or addiction] ... who are ready for something like this" (Palliative Care professional).

### Expertise in facilitation: recognising risk and the unstructured nature

Participants emphasised the importance of a confident, skilled facilitator to lead and manage a death café. It was recognised that more than one facilitator may be required, bringing expertise from professionals from different backgrounds together, for example, professionals across the fields of palliative care, homelessness services or mental health working in partnership. Professionals acknowledged that their experience in one area did not automatically mean they were a suitable or appropriate facilitator: "just because a couple of us here have got good knowledge of death, or grief or palliative care doesn't necessarily mean that we're absolutely ace at facilitating groups like this." (Palliative Care Professional).

Importantly, participants highlighted that facilitation needs to be trauma informed. It is important that professionals within a death café can recognise the potential for conversations to trigger an emotional response or lead to behaviours that may be harmful. That is, attendees can't be expected to "leave their complex trauma and chaotic lifestyles at the door" (Social Worker). Therefore, due to the typically unstructured nature of death cafés, confident facilitation is key to "steer conversations if they look like they're becoming too difficult or if there are things which you can hear are likely to trigger other people." (Clinician and researcher).

This skilled facilitation was also thought to be integral to the well-being support provided both during and after death cafés. Traditionally, death cafés don't offer well-being follow-ups. However, all participants agreed that the availability of post-session support was important due to the potential for emotional or sensitive discussions during death cafés, coupled with the limited support

networks or previous negative experiences that attendees may have. There was a perceived risk that following a death café, people may return to the streets or temporary accommodation while experiencing distress. Professionals experienced in facilitating death cafés within a hostel environment described methods for providing support, such as debriefs, informal check-ins or staying behind following a session. This also highlighted the need for building trust and rapport between facilitators and attendees to improve support offered both during and after death cafés.

If you are facilitating these sorts of conversations we always hang around for about 20 minutes after just with a cup of tea. It's just something that somebody needs to chat around in private or they've been distressed by the conversations that have taken place (Bereavement Counsellor).

Signposting (ie, helping people or directing them to access appropriate support if needed) people to where they may access support was debated by participants. PWLE homelessness often felt they were 'over-signposted', leading to them feeling dismissed. However, professionals highlighted the necessity of helping people to access further support and suggested non-tokenistic signposting such as referrals made at the time, if possible, or continued support in accessing services, rather than just providing information that could be beneficial.

### Promoting safety against potential risk

Participants discussed the need to manage the possibility that attendees may have active addiction issues. First, they outlined the need to consider how to manage people at varying stages of recovery. People with lived experience expressed that attending a session during a period of sobriety with a peer who was actively using substances could be triggering. Furthermore, participants recognised the need to manage intoxication at sessions. The general consensus was that banning substance use is likely to be ineffective, as it would lead people who were dependent on alcohol to simply not attend. Instead, both professionals and PWLE homelessness participants suggested that facilitators would need to be confident making behaviour-based decisions on attendance on a case-by-case basis.

If somebody's in addiction, there's no point saying to them, you can't be on drink, or you can't be high on drugs. Because that's their life. That's, that's how they function in, in their daily lives. So, having a limit- if you're overly intoxicated, or you're overly aggressive while you're that intoxicated (PWLE)

Participants also discussed the potential risk for retraumatisation during a death café. The potential for discussions around death to resurface emotions and trigger distress was recognised.

The process of looking after themselves or exploring their own needs or exploring things that has the propensity to bring up really difficult emotions, then they might not have the skills or resources to be able to sit with and manage and might escalate their drug use or lead to self-harm. (Clinical Psychologist)

Participants outlined how this could be unexpected, even for people who felt able to attend. Given the unstructured nature of a death café, it is not possible to control the direction of the conversation or what topics people may be exposed to: *"they may be triggered by just being present in this room where people might be opening up, it maybe what other people might say might trigger them"* (Clinical Psychologist)

### Potential benefits of death cafés for PEH

Despite the potential for risk, participants agreed that, if done safely, death cafés could provide a space for PEH to talk about death and dying. An absence of other spaces where they felt welcome and free to talk about death and dying without judgement was noted.

Even though he was only in his 30s, he had a big fear of what was going to happen to him when he died, and there was nowhere he had ever talked about that. But he was able to do so in that group. He felt safe to raise it. (Hostel staff)

An example of a successful, hostel-based death café that had been facilitated by a psychologist working within a community palliative care team outlined that the event allowed residents to discuss things they hadn't previously: *"Some of the discussions that came up were areas that absolutely were concerning the residents, but they never felt that they were things they could raise with their support worker."* (Hostel staff)

Both professionals and people with lived experience felt that through holding a death café, a supportive space was created in which existing power dynamics were reduced: *"there was no staff vs residents, everyone was able to share experiences and share on a very level playing field really."* (Palliative care clinician). Furthermore, PEH expressed that they *"didn't know (hostel staff) cared that much"* (PWLE) about them, with regards to holding the event and sharing their own thoughts and feelings.

Participants highlighted how a death café could provide an opportunity for attendees to begin to consider their own preferences for end of life. Although attendees at a death café traditionally will not complete any formal advance care planning, it could be a catalyst for initial exploration of preferences. For some people, in time, this could be a precursor to an openness to having more detailed discussions about making their preferences for care and treatment known. Giving people space to discuss their thoughts may also support them to feel valued, seen and heard:

Some people might not be able to think about it, but for some people, having a space to at least talk about how they want to live and how they want to be and

what they want to do in the rest of their life. (Clinical Psychologist)

### Potential practical considerations

Initial considerations regarding practicalities for holding a death café were discussed. Given the exploratory nature of this research, these should not be considered as recommendations for developing a death café, but points for consideration.

Both professionals and PWLE homelessness expressed a preference for fewer attendees per session, allowing space for each attendee to talk without the environment feeling overwhelming: *"It shouldn't be crowded because if it is crowded there'll be so many questions and opinions and you wouldn't, there'd be panic."* (PWLE).

People with lived experience highlighted they would be put off by negative, dark imagery in advertisements for the event. Instead, they outlined a preference for colourful advertisements (such as *"a sunset, a tree of life...a bit light and bright with a bit of hope to it"* (PWLE)) that importantly, contained clear information on key elements of the session.

To create a safe space, participants discussed the need for ground rules, termed as a *"group agreement"* (PWLE). Participants suggested this could cover topics such as confidentiality, substance use and being respectful, with the aim *"to try and help people understand that this is part of keeping the safe space. If we didn't have this rule, then it could lead to the space and individuals feeling more unsafe."* (Clinical Psychologist).

When considering venues, some participants felt it would be beneficial to hold the event somewhere known to potential attendees: *"I would always think that a Death Cafe should be undertaken where somebody is comfortable and where a locality that they are familiar with but also their needs. So, for example, if it's for people living in a hostel, I would think you go to a hostel."* (Palliative care clinician). However, some participants had concerns that this may *"force other people who don't want to necessarily participate to feel that they're being forced to participate."* (Hostel staff).

Participants debated their preferred name for these events. Some felt that 'death café' was too negative, intimidating and off-putting.

I think because we, as a society, don't talk about death, so death's quite taboo. So even the word 'death' would probably be quite off-putting to some people. (Registered nurse)

Going into something that sort of says death café, forgive the pun, is like adding another nail to the coffin, isn't it? (PWLE)

However, others expressed the need for clear, transparent naming to ensure participants understood the brief of the event and were able to give informed consent to attend. *"If you're going to call it something different, you would still need to help people to be aware that it is going to be*

*talking about death. Otherwise, they might get a bit of a shock when they turn up"* (Palliative care professional).

### DISCUSSION

This qualitative research used interviews with professionals and focus groups with people with lived experience of homelessness to explore perspectives on death cafés for PEH. Five themes were generated: choice; expertise in facilitation; recognising risk and the unstructured nature; promoting safety against potential risk; potential benefits and potential practical considerations.

#### Trauma-informed care and well-being support

A key finding is the need for death cafés to be trauma informed. This is supported by a plethora of existing evidence connecting trauma and homelessness,<sup>31</sup> which consistently recommends that 'ervice providers must consider trauma when designing and offering services to persons with a history of homelessness',<sup>32</sup>

Pertinent to death cafés, PEH have often witnessed a high number of deaths, both prior to and during periods of homelessness.<sup>16</sup> Death experienced on the streets can be traumatic<sup>16 33 34</sup> and are often sudden or unexpected.<sup>35</sup> Furthermore, the death of loved ones is often described by PEH as one of the negative 'turning points' in their lives, acting as a significant risk factor for their homelessness.<sup>21 22 36 37</sup> This potential history of traumatic experiences surrounding death, dying and bereavement means that extra considerations of psychological safety are required in the planning, conduct and follow-up from a death café.

Additionally, experiences of trauma can heighten the importance of adequate emotional support in services or spaces such as death cafés. Repeated or multiple traumatic experiences can 'overwhelm a person's available internal and/or external resources', leading to ineffective coping and reduced capacity for emotional regulation.<sup>38</sup> Clinical psychologists (among other professionals) in our research suggested techniques for well-being support during sessions (including regular breaks, non-tokenistic signposting and referral at time where possible) and after sessions (such as debriefs, staying in place for one-to-one conversations and checking in with attendees where they are known).

#### Facilitation

This research highlights that expert facilitation of death cafés for PEH is essential. This is supported by previous research from death cafés aimed at the general population. Facilitation can promote meaningful conversations and guide attendees through an event. A death café facilitator needs to be able to: 'manage speaking turns and pay close attention to conversational flow, group climate, and expressions of emotion'.<sup>39</sup>

Traditionally, death cafés propose that facilitators 'leave their professional identity at the door', aiming to enable death cafés to be egalitarian spaces. However, based



on the findings of this exploratory research, it may be important that facilitators for death cafés for PEH have adequate professional experience that they use in the events to support attendees.

Participants felt that no one group of professionals could adequately run a death café for this population and that a multidisciplinary approach was seen as useful. Previous research has<sup>40</sup> partnered palliative care professionals with hostels aiming to build relationships and introduce the benefits of a palliative approach to support for people with advanced ill health. This found benefits such as better working partnerships, hostel staff feeling empowered to discuss death and dying, and integration of the ethos of each place. A death café was held as part of this work, which was found to address some anxiety within hostel staff when discussing death with residents. This supports the findings of this work which suggest co-facilitation across the fields of homelessness and palliative care can be beneficial.

## Limitations

As recruitment occurred opportunistically through advertising at networks and allowing people to put themselves forward, there is a risk of sampling bias including self-selection bias. Further bias may have been introduced by the snowball nature of sampling.

Furthermore, although participants bought a range of experiences, including lived experience of homelessness, further research may be required with a larger sample size. This would enable the participant sample to represent a more diverse sample of people with lived experience of homelessness, or involvement in death cafés, as evidence demonstrates the complexity and heterogeneity of experiences.

## Recommendations

To the best of the authors' knowledge, this is the first research to qualitatively explore the idea of death cafés for PEH. These initial explorations have raised further points for consideration. For example, while recognising the potential benefits of death cafés for PEH, it is important to fully explore the potential risks and how these can be managed. This may include an in-depth exploration of the facilitation of death cafés for PEH, taking into account issues raised in this research such as previous experiences of attendees, risk moderation and the importance of partnership working, with the potential to develop training for facilitators in this space.

## CONCLUSIONS

As death cafés continue to grow, exploration of their applicability to varying populations facing multi-faceted social exclusion is key. Death cafés for PEH may pose risks that require comprehensive consideration and trauma-informed, expert facilitation and support. However, this research highlighted that a low-pressure space to explore thoughts around death, dying and bereavement

is currently an unmet need within this population, which warrants further exploration.

**Contributors** JC, BH and GD conceptualized the study. JC and GD carried out interview data collection. EC facilitated focus groups. JC and GD led on data analysis. BH supported data analysis. JC led on writing the manuscript. BH, GD, MP, EC, CS, KF (all authors) reviewed the final manuscript. JC is guarantor.

**Funding** This work was supported by Marie Curie grant number MCSGS-23-101. The funder didn't influence the results/outcomes of the study despite author affiliations with the funder.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants and was approved by Ethical approval was sought and obtained by University College London's ethics committee (ID: 6202/011). Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID ID

Jodie Crooks <http://orcid.org/0000-0003-2574-4420>

## REFERENCES

- 1 Fine DR, Dickins KA, Adams LD, *et al*. Mortality by Age, Gender, and Race and Ethnicity in People Experiencing Homelessness in Boston, Massachusetts. *JAMA Netw Open* 2023;6:e2331004.
- 2 Mostowska M. Researching Mortality and Deaths of People Experiencing Homelessness. *Eur J Homelessness* 2023;17:1.
- 3 Mantell R, Hwang YIJ, Radford K, *et al*. Accelerated aging in people experiencing homelessness: A rapid review of frailty prevalence and determinants. *Front Public Health* 2023;11:1086215.
- 4 Keogh C, O'Brien KK, Hoban A, *et al*. Health and use of health services of people who are homeless and at risk of homelessness who receive free primary health care in Dublin. *BMC Health Serv Res* 2015;15:58.
- 5 Rogans-Watson R, Shulman C, Lewer D, *et al*. Premature frailty, geriatric conditions and multimorbidity among people experiencing homelessness: a cross-sectional observational study in a London hostel. *HCS* 2020;23:77–91.
- 6 Vallesi S, Tuson M, Davies A, *et al*. Multimorbidity among People Experiencing Homelessness-Insights from Primary Care Data. *Int J Environ Res Public Health* 2021;18:6498.
- 7 Harrison C, Henderson J, Miller G, *et al*. The prevalence of diagnosed chronic conditions and multimorbidity in Australia: A method for estimating population prevalence from general practice patient encounter data. *PLoS One* 2017;12:e0172935.
- 8 Liu M, Hwang SW. Health care for homeless people. *Nat Rev Dis Primers* 2021;7:5.
- 9 de Veer AJE, Stringer B, van Meijel B, *et al*. Access to palliative care for homeless people: complex lives, complex care. *BMC Palliat Care* 2018;17:119:119.



- 10 Shulman C, Hudson BF, Low J, *et al.* End-of-life care for homeless people: A qualitative analysis exploring the challenges to access and provision of palliative care. *Palliat Med* 2018;32:36–45.
- 11 Homelessness Mo. Dying Homeless Project. 2024.
- 12 Aldridge RW, Menezes D, Lewer D, *et al.* Causes of death among homeless people: a population-based cross-sectional study of linked hospitalisation and mortality data in England. *Wellcome Open Res* 2019;4:49.
- 13 Baggett TP, O'Connell JJ, Singer DE, *et al.* The unmet health care needs of homeless adults: a national study. *Am J Public Health* 2010;100:1326–33.
- 14 Irving AH. The Prevalence of Trauma among People who have Experienced Homelessness in England. A Report for Oasis Community Housing, 2022. Available: <https://www.oasiscommunityhousing.org/trauma-homelessness>
- 15 Link H. Rise in deaths of people experiencing homelessness is 'shameful', 2022. Available: <https://homeless.org.uk/news/rise-in-deaths-of-people-experiencing-homelessness-is-shameful>
- 16 Song J, Ratner ER, Bartels DM, *et al.* Experiences with and attitudes toward death and dying among homeless persons. *J Gen Intern Med* 2007;22:427–34.
- 17 Café D. Holding your own death café, 2023. Available: <https://deathcafe.com/how/>
- 18 Miles L, Corr CA. Death cafe: What is it and what we can learn from it. *OMEGA-J Death Dying* 2017;75:151–65.
- 19 Stajduhar KI, Mollison A, Giesbrecht M, *et al.* "Just too busy living in the moment and surviving": barriers to accessing health care for structurally vulnerable populations at end-of-life. *BMC Palliat Care* 2019;18:11.
- 20 Hudson BF, Shulman C, Low J, *et al.* Challenges to discussing palliative care with people experiencing homelessness: a qualitative study. *BMJ Open* 2017;7:e017502.
- 21 Monk J, Black J, Carter RZ, *et al.* Bereavement in the context of homelessness: A rapid review. *Death Stud* 2024;48:561–70.
- 22 Hansford L, Thomas F, Wyatt K. How does housing affect end-of-life care and bereavement in low-income communities? A qualitative study of the experiences of bereaved individuals and service providers in the United Kingdom. *Palliat Care Soc Pract* 2022;16:26323524221110248.
- 23 Elman C, Gerring J, Mahoney J. *The production of knowledge: enhancing progress in social science*. Cambridge University Press, 2020.
- 24 Stebbins RA. *Exploratory research in the social sciences*. Sage, 2001.
- 25 Braun V, Clarke V. Supporting best practice in reflexive thematic analysis reporting in *Palliative Medicine*: A review of published research and introduction to the *Reflexive Thematic Analysis Reporting Guidelines* (RTARG). *Palliat Med* 2024;38:608–16.
- 26 Crooks J, Flemming K, Shulman C, *et al.* Development of the TIFFIN recommendations for co-producing palliative and end-of-life care research with individuals with lived experience of homelessness: A qualitative study. *Palliat Med* 2024;38:746–54.
- 27 Adeoye-Olatunde OA, Olenik NL. Research and scholarly methods: Semi-structured interviews. *J Am Coll Clin Pharm* 2021;4:1358–67.
- 28 Braun V, Clarke V. Can I use TA? Should I use TA? Should I *not* use TA? Comparing reflexive thematic analysis and other pattern-based qualitative analytic approaches. *Couns and Psychother Res* 2021;21:37–47.
- 29 Pathway. Research, Policy and Practice, 2025. Available: <https://www.pathway.org.uk/research>
- 30 NIHR. Payment guidance for researchers and professionals, 2022. Available: <https://www.nihr.ac.uk/documents/payment-guidance-for-researchers-and-professionals/27392>
- 31 Pope ND, Buchino S, Ascienzo S. "Just like Jail": Trauma Experiences of Older Homeless Men. *J Gerontol Soc Work* 2020;63:143–61.
- 32 Wiewel B, Hernandez L. Traumatic Stress and Homelessness: A Review of the Literature for Practitioners. *Clin Soc Work J* 2022;50:218–30.
- 33 Mackelprang JL, Clifasefi SL, Grazioli RS, *et al.* Content Analysis of Health Concerns among Housing First Residents with a History of Alcohol Use Disorder. *J Health Care Poor Underserved* 2021;32:463–86.
- 34 Perry R, Adams EA, Harland J, *et al.* Exploring high mortality rates among people with multiple and complex needs: a qualitative study using peer research methods. *BMJ Open* 2021;11:e044634.
- 35 Lakeman R. How homeless sector workers deal with the death of service users: a grounded theory study. *Death Stud* 2011;35:925–48.
- 36 Burns VF, Sussman T, Bourgeois-Guérin V. Later-Life Homelessness as Disenfranchised Grief. *Can J Aging* 2018;37:171–84.
- 37 Mayock P, Parker S, Murphy A. Family 'turning point' experiences and the process of youth becoming homeless. *Child Fam Soc Work* 2021;26:415–24.
- 38 Macia KS, Moschetto JM, Wickham RE, *et al.* Cumulative Trauma Exposure and Chronic Homelessness Among Veterans: The Roles of Responses to Intrusions and Emotion Regulation. *J Trauma Stress* 2020;33:1017–28.
- 39 Cherry J, Black LW. "Talking About Funerals Won't Make You Dead": Dialogic Tensions in Death Café Design and Facilitation. *Omega (Westport)* 2024.:302228241272716.
- 40 Armstrong M, Shulman C, Hudson B, *et al.* The benefits and challenges of embedding specialist palliative care teams within homeless hostels to enhance support and learning: Perspectives from palliative care teams and hostel staff. *Palliat Med* 2021;35:1202–14.