



Deposited via The University of York.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/228564/>

Version: Published Version

Article:

Petersen Williams, Petal (2025) Associations of maternal alcohol and non-prescribed substance use with early child growth. South African Journal of Psychiatry. ISSN: 1608-9685

<https://doi.org/10.4102/sajpsychoiatry.v31i0.2486>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Associations of maternal alcohol and non-prescribed substance use with early child growth



Authors:

Yukiko Washio^{1,2} 
 Zugui Zhang³ 
 Mona Lisa Baishya⁴ 
 Marilyn T. Lake⁵ 
 Bronwyn Myers⁶ 
 Nadia Hoffman⁵ 
 Elizabeth Goddard⁷ 
 Heather J. Zar⁷ 
 Dan J. Stein⁵ 
 Petal Petersen Williams^{8,9} 

Affiliations:

¹Women's Global Health Imperative, RTI International, Media, Pennsylvania, United States

²Department of Obstetrics, Gynecology and Reproductive Sciences, Lewis Katz School of Medicine, Temple University, Philadelphia, Pennsylvania, United States

³REACH, Christiana Care, Newark, Delaware, United States

⁴College of Public Health, Temple University, Philadelphia, Pennsylvania, United States

⁵SAMRC Unit on Risk and Resilience in Mental Disorders, Department of Psychiatry and Neuroscience Institute, University of Cape Town, Cape Town, South Africa

⁶enAble Institute, Curtin University, Bentley, Australia

⁷SAMRC Unit on Child and Adolescent Health, Department of Pediatrics, University of Cape Town, Cape Town, South Africa

⁸Mental Health, Alcohol, Substance Use and Tobacco Research Unit, South African Medical Research Council, Cape Town, South Africa

Read online:



Scan this QR code with your smart phone or mobile device to read online.

Background: Perinatal alcohol and non-prescribed substance use may be detrimental to foetal and infant growth.

Aim: This observational study investigated how combined and continued alcohol and non-prescribed substance use throughout antenatal and 1-year postnatal periods were associated with adverse child length and weight outcomes up to 24 months.

Setting: Data from participants ($n = 1098$) with information on alcohol and non-prescribed substance use and infant and child outcomes, were drawn from a prospective birth cohort in the Drakenstein Child Health Study (DCHS), conducted in the Western Cape province of South Africa.

Methods: Generalised estimating equations were conducted on standardised child length and weight outcomes at 12, 18 and 24 months.

Results: Non-prescribed substances consisted mostly of tobacco use (77%). Child length and weight were significantly lower in those exposed to the combined use of alcohol and substances compared to no-use and all other use groups ($p < 0.001$), as confirmed by multivariable analyses. Child length and weight were also significantly lower in those exposed to alcohol and/or substance use throughout the antenatal and 1-year postnatal periods, as confirmed by multivariable analyses.

Conclusion: Interventions to address the potential long-term adverse effects of combined alcohol and substance use particularly tobacco use, as well as continuous use throughout antenatal and early postnatal periods on subsequent child growth, are needed.

Contribution: This study has contributed to the field by showing that combined and continued use of alcohol and other substances during pregnancy and postpartum is associated with impaired early child growth.

Keywords: pregnancy; early postpartum; infant child length; infant child weight; combined alcohol and substance use.

Introduction

Antenatal alcohol use is associated with adverse pregnancy, birth and child outcomes, including miscarriage, stillbirth, preterm birth, low birth weight, sudden infant death syndrome and development of foetal alcohol spectrum disorders (FASD).^{1,2} South Africa has the highest FASD rate in the world, followed by Eastern European regions, the United Kingdom and New Zealand.^{3,4} Notably, the Western Cape region of South Africa has detected 20% of biochemically verified antenatal alcohol use,⁵ comparable to the rate of 26% self-reported antenatal alcohol use risk among those receiving services from the Special Supplemental Nutrition Programme for Women, Infants and Children (WIC) in the United States of America (USA).⁶

Combined use of alcohol and other substances, including tobacco, cannabis and methamphetamine, is associated with adverse birth outcomes, including preterm birth, low birth and infant weight and smaller head circumference, compared to antenatal alcohol exposure alone.^{7,8,9} Alcohol use during lactation is associated with antenatal alcohol use and has been shown to adversely impact infant development, including lower verbal IQ scores and reduced infant weight, independently of antenatal alcohol use.¹⁰ In South Africa, antenatal alcohol and substance use is associated with

⁹Institute for Life Course Health Research, Department of Global Health, Stellenbosch University, Cape Town, South Africa

Corresponding author: Yukiko Washio, ywashio@rti.org

Dates: Received: 15 Feb. 2025 | Accepted: 14 Apr. 2025 | Published: 30 June 2025

How to cite this article: Washio Y, Zhang Z, Baishya ML, et al. Associations of maternal alcohol and non-prescribed substance use with early child growth. *S Afr J Psychiatr.* 2025;31(0), a2486. <https://doi.org/10.4102/sajpsychoiatry.v31i0.2486>

Copyright: © 2025. The Authors. Licensee: AOSIS. This work is licensed under the Creative Commons Attribution License.

intimate partner violence, lower socioeconomic status, lack of social support, trauma exposure and depression,^{11,12,13} which in turn may also impact postpartum parenting, thus also impacting infant and child growth.^{14,15}

It is not clear whether the combined use of alcohol with any type of non-prescribed substance(s) as well as the continued use of alcohol and/or substances throughout antenatal and postnatal periods is associated with adverse infant and child growth. The aim of this observational study was to investigate the association of perinatal alcohol and non-prescribed substance use with child height and weight through the first 2 years of life. Substances in the analysis of this observational study were non-prescribed substances with a majority being tobacco but also including methamphetamine, cocaine, MDMA, cannabis, opiates and mandrax.

Research methods and design

Study design and participants

Data were derived from a prospective birth cohort study of mother-child pairs enrolled in the Drakenstein Child Health Study (DCHS), conducted in a peri-urban area, in the Western Cape province, South Africa. The dataset includes antenatal and postnatal use of alcohol and substance use and child growth standardised outcomes.^{16,17} The DCHS included women during 20–28 weeks of gestation, who were enrolled between March 2012 and March 2015, and subsequently followed from birth as mother-child dyads. Pregnant women were recruited from public primary health care clinics, and all births occurred at a single public hospital serving the area. Pregnant women were eligible to participate in the parent study if they were 18 years and older, accessing antenatal care between 20 and 28 gestational weeks, and intended to remain in the area for at least a year.

Enrolled participants in the study completed interviewer administered self-report questionnaires in their preferred language of English, Afrikaans or isiXhosa (the three local languages of the Western Cape) assessing maternal psychosocial risk factors at antenatal baseline and postnatal time points and child length and weight at 12, 18 and 24 months.¹⁶ All child assessments were carried out in a private room in one of the study offices; where possible, interviews with the maternal participants were carried out in a separate room or private area. Participants were contacted telephonically or by home visit to schedule study visits.

Explanatory variables and outcomes

For antenatal (i.e. second trimester), 6-month postpartum and 12-month postpartum alcohol and substance use, ASSIST¹⁸ questions on use in the past 3 months were used, and a dichotomous variable of use versus non-use at any frequency was created. Urinalysis was conducted for tobacco, methamphetamine, cocaine, MDMA, cannabis, opiates and mandrax at the antenatal baseline visit. For antenatal non-prescribed substance use, a combination of urinalysis results and self-reported use in the past 3 months was used to

determine antenatal use, as represented using a dichotomous variable. The variable was categorised as negative only when both urinalysis and self-report indicated negative substance use; otherwise, the variable was categorised as positive of use if either urinalysis or self-report was positive of use. For postnatal use, either 6-month or 12-month positive use was counted as postnatal use at some point during the 1-year postnatal period. Lifetime alcohol and substance use was also assessed to confirm no lifetime use for the no-use groups.

Participants were grouped into four groups based on the type of alcohol and substance use: (1a) no use, (1b) alcohol use only, (1c) substance use only or (1d) alcohol and substance use. Participants were also grouped into four groups based on the timing of alcohol and/or substance use: (2a) no use, (2b) antenatal use only, (2c) postnatal use only or (2d) antenatal and postnatal use. The second grouping was not specific to the type of alcohol or substance use, and the use of any alcohol or substance use was included.

Outcomes included in this analysis were standardised child length and weight scores at 12, 18 and 24 months as continuous variables.^{19,20} Infants are defined as individuals less than 1-year-old and children as 1 year or older.²¹ The infant and child's length was measured in centimetres to the nearest completed 0.5 cm, using a Seca length-measuring mat (Seca, Hamburg, Germany) for recumbent length up to 18 months and a wall-mounted stadiometer for standing length at 24 months. Infant and child weight was measured in kilograms to the nearest 10 g, in light or no clothing, using the Tanita digital platform scale (TAN1584; IL, USA). Child weight and length measurements were converted to standardised *z*-scores, adjusted for infant age, sex and prematurity. The Fenton growth chart was applied to anthropometry measurements from premature infants (<37 weeks gestation) at birth and up to 50 weeks postmenstrual age. The anthro software of the World Health Organization was used to adjust for age and sex in measurements taken from full-term infants at birth up to 2 years, as well as age-corrected measurements taken from premature children captured between 50 weeks postmenstrual age and 2 years.²² Stunting was defined as the children with length/height-for-age *Z*-score of <-2SD with severe stunting defined as the score of <-3SD.²³

Sociodemographic variables comprised maternal age, maternal education, marital status, household income, parity, HIV status, maternal height and weight, maternal trauma exposure, maternal intimate partner emotional, physical and sexual violence exposure,²⁴ duration of exclusive and any breastfeeding, as well as maternal antenatal depression scores.²⁵ The variable of maternal education had two categories of those with less than secondary education level, versus those with at least secondary level education. Household income levels were dichotomised into equal to or less than R5000 and greater than R5000 per month. Marital status was dichotomised into being either single or married/cohabiting. Intimate partner violence exposure variables capturing emotional, physical or sexual abuse in the past 12 months were dichotomized into above and below threshold levels.^{26,27}

Statistical analysis

Descriptive statistics were used to describe the data. Data were summarised using the mean (standard deviations) for continuous variables and using frequency/percentages for categorical variables. A contingency table analysis (Chi-square test) was conducted for comparison of categorical variables, and a non-parametric Kruskal–Wallis test was used to compare continuous variables by each level within each grouping. Statistical significance was set at p -value < 0.05 .

The generalised estimating equations (GEE) approach was used to model longitudinal outcome measurements of standardised child weight and length at 12, 18 and 24 months against antenatal and postnatal alcohol and/or substance use in the first year of life. Adjusted covariates from participant characteristics included sociodemographic variables as well as those that were statistically significant by univariate comparisons. We included standardised birth height for child height measures and standardised birth weight for child weight measures. Additionally, unadjusted GEE analyses were conducted without covariates, except for grouping and time variables. All analyses were performed using SAS version 9.4 (SAS Institute, Cary NC).

Ethical considerations

Ethical clearance to conduct this study was obtained from the University of Cape Town Faculty of Health Sciences Human Research Ethics Committee (No. 401/2009), Stellenbosch University (No. N12/02/0002[A1]) and the Western Cape Provincial Health Research Committee (No. 2011RP45). The study was conducted in accordance with the Helsinki Declaration as revised in 2013. Written informed consent was obtained from the participants of the parent study.

Results

Participant characteristics

Of 1225 participants enrolled in the DCHS at the second trimester of pregnancy, there were 1143 live births. The current analyses included an analytic sample size of 1098 mother-child dyads, where 45 were excluded due to missing data on alcohol and substance use or infant and child outcomes. When looking at (1) substance type groupings: defined into (1a) no use, (1b) alcohol use only, (1c) substance use only or (1d) alcohol and substance use, 590 of 1098 (54%) participants reported no lifetime alcohol or substance use, 158 (14%) reported alcohol use only during pregnancy and/or postpartum, 88 (8%) reported at least one of the substances but no alcohol use during pregnancy and/or postpartum and 262 (24%) reported both alcohol and substance use during pregnancy and/or postpartum. A majority of substances used antenatally comprised of tobacco use (77%), with smaller proportions testing positive for cannabis and methamphetamine use (3%), mandrax use (2%), opioid, MDMA and cocaine use ($< 1\%$). With respect to (2) timing of alcohol and/or substance use, as defined by (2a) no use, (2b) antenatal use only, (2c) postnatal use only or (2d)

antenatal and postnatal use, 590 of 1098 (54%) participants reported no use, 91 (8%) reported antenatal use only, 176 (16%) reported postnatal use only and 241 (22%) reported use throughout antenatal and postnatal periods.

Participant characteristics by group comparison are shown in Table 1 and Table 2. The total cohort included in the analyses ($N = 1098$) had children with an average length of 50 centimetres and three kilograms at birth. In Table 1, with grouping by substance use type, substance use only and alcohol and substance use were significantly associated with reduced height measures compared to no use and alcohol use only ($p < 0.001$). Child weight measures showed progressively reduced weight measures in the order of no use, alcohol use only, substance use only and alcohol and substance use with statistically significant differences ($p < 0.001$). In Table 2, with grouping by timing of substance use, all child measures showed statistical differences by group, with the most compromised growth measures associated with the antenatal and postnatal use group compared to no-use, antenatal-only and postnatal-only groups ($p < 0.001$). All groups showed progressively reduced length measures from 12 to 24 months, while all groups showed reduced weight measures at 12 and 24 months compared to 18 months (Table 1 and Table 2).

In the cohort included in the analyses ($N = 1098$), participants were 27 years old on average with more than half having lower than secondary education, 87% earning R5000 per month or less as the household income, 41% married or cohabiting and 22% HIV seropositive. Approximately 25% were exposed to a traumatic event with 27% suffering from emotional abuse (Table 1 and Table 2). In the first grouping by substance use type, the no-use group was older than the other use groups and had higher levels of education (Table 1). A much higher rate of HIV seropositivity was found in the no-use group compared to the other use groups. Depression scores were similar across all groups. Progressively higher rates of past year emotional, physical and sexual abuse were observed in the order of no use, alcohol use only, substance use only and alcohol and substance use. In the second grouping by the timing of substance use, the no-use group was older and had a higher rate of HIV seropositivity than the other use groups (Table 2). Depression scores were similar across all groups. Higher rates of emotional, physical and sexual abuse were observed among antenatal-only and antenatal and postnatal use groups compared to no-use and postnatal-only groups.

Child length and weight associated with substance use type

Table 3 shows the multivariable analyses by alcohol and substance use type with adjusted estimates on the standardised child length and weight as repeated measures at 12, 18 and 24 months. Alcohol use only was not significantly associated with shorter child length but there was a statistically significant reduction in child weight compared to the no-use group ($p = 0.516$ and $p = 0.019$). Substance use only was not significantly associated with shorter child length

TABLE 1: Participant characteristics by substance use type (*N* = 1098).

Variable	Total (n)	%	Mean	s.d.	No use (n = 590)	%	Mean	s.d.	Alcohol use only (n = 158)	%	Mean	s.d.	Substance use only (n = 88)	%	Mean	s.d.	Alcohol and substance use (n = 262)	%	Mean	s.d.	<i>p</i>
Maternal age at enrollment	-	-	26.63	5.72	-	-	27.28	5.75	-	-	26.45	5.99	-	-	25.54	5.54	-	-	25.62	5.35	< 0.001
Lower than secondary education	676	61.58	-	-	342	57.97	-	-	95	60.13	-	-	64	72.73	-	-	175	66.79	-	-	0.012
Household income more or less R5000	951	86.61	-	-	511	86.61	-	-	132	83.54	-	-	79	89.77	-	-	229	87.40	-	-	ns
Married or cohabiting	453	41.29	-	-	259	43.97	-	-	44	27.85	-	-	41	46.59	-	-	109	41.60	-	-	0.002
Antenatal maternal height	-	-	159.41	6.79	-	-	160.07	6.87	-	-	158.75	6.80	-	-	157.95	5.35	-	-	158.85	6.92	0.005
Parity	-	-	1.04	1.05	-	-	1.06	1.01	-	-	1.00	1.10	-	-	1.13	1.22	-	-	0.97	1.07	ns
HIV-positive status	242	22.04	-	-	183	31.02	-	-	29	18.35	-	-	8	9.09	-	-	22	8.40	-	-	< 0.001
Depression scale score antenatal	-	-	9.45	5.28	-	-	9.52	4.70	-	-	9.19	5.29	-	-	8.49	5.59	-	-	9.79	6.24	ns
Exposed to traumatic events	255	25.63	-	-	152	29.23	-	-	33	22.15	-	-	18	21.69	-	-	52	21.40	-	-	ns
Past 12 months emotional abuse	265	26.69	-	-	84	16.25	-	-	48	32.00	-	-	28	33.73	-	-	105	43.21	-	-	< 0.001
Past 12 months physical abuse	217	21.85	-	-	83	16.05	-	-	33	22.00	-	-	22	26.51	-	-	79	32.51	-	-	< 0.001
Past 12 months sexual abuse	68	6.85	-	-	19	3.68	-	-	12	8.00	-	-	9	10.84	-	-	28	11.52	-	-	< 0.001
Infant length at birth (cm)	-	-	49.64	3.73	-	-	50.01	3.68	-	-	49.98	3.52	-	-	49.30	3.48	-	-	48.76	3.91	< 0.001
Standardised infant length at 12 months	-	-	-0.63	1.26	-	-	-0.38	1.16	-	-	-0.69	1.34	-	-	-0.96	1.35	-	-	-0.92	1.28	< 0.001
Standardised child length at 18 months	-	-	-0.88	1.45	-	-	-0.73	1.34	-	-	-0.95	1.54	-	-	-0.96	1.55	-	-	-1.12	1.52	< 0.001
Standardised child length at 24 months	-	-	-1.06	1.20	-	-	-0.88	1.15	-	-	-1.04	1.33	-	-	-1.41	1.31	-	-	-1.29	1.12	< 0.001
Infant weight at birth (kg)	-	-	3.02	0.60	-	-	3.08	0.59	-	-	3.09	0.55	-	-	2.97	0.60	-	-	2.87	0.63	< 0.001
Standardised infant weight at 12 months	-	-	-0.02	1.29	-	-	0.28	1.28	-	-	-0.06	1.30	-	-	-0.32	1.15	-	-	-0.46	1.21	< 0.001
Standardised child weight at 18 months	-	-	0.11	1.32	-	-	0.40	1.30	-	-	-0.03	1.33	-	-	-0.13	1.37	-	-	-0.29	1.22	< 0.001
Standardised child weight at 24 months	-	-	-0.29	1.23	-	-	-0.00	1.21	-	-	-0.37	1.23	-	-	-0.50	1.20	-	-	-0.68	1.15	< 0.001
Number of months exclusive breastfeeding	-	-	2.09	1.96	-	-	2.01	2.01	-	-	1.96	1.94	-	-	2.11	1.94	-	-	2.32	1.87	0.008
Number of months any breastfeeding	-	-	9.91	9.50	-	-	7.80	8.31	-	-	9.35	9.65	-	-	12.46	10.58	-	-	14.02	9.98	< 0.001

HIV, human immunodeficiency virus; s.d., standard deviation; ns, not significant.

TABLE 2: Participant characteristics by when alcohol and substance were used (N = 1098).

Variable	Total (n)	%	Mean	s.d.	No use (n = 590)	%	Mean	s.d.	Antenatal only (n = 91)	%	Mean	s.d.	Postnatal only (n = 176)	%	Mean	s.d.	Antenatal and postnatal (n = 241)	%	Mean	s.d.	p
Maternal age at enrollment	-	-	26.63	5.72	-	-	27.28	5.75	-	-	25.86	5.20	26.03	-	-	6.02	-	-	25.74	5.42	< 0.001
Lower than secondary education	676	61.58	-	-	342	57.97	-	-	64	70.33	-	-	99	56.25	-	-	171	70.95	-	-	< 0.001
Household income more or less R5000	951	86.61	-	-	511	86.61	-	-	82	90.11	-	-	145	82.39	-	-	213	88.38	-	-	ns
Married or cohabiting	453	41.29	-	-	259	43.97	-	-	40	43.96	-	-	53	30.11	-	-	101	41.91	-	-	0.011
Antenatal maternal height	-	-	159.41	6.79	-	-	160.07	6.87	-	-	158.97	5.12	-	-	158.60	6.98	-	-	158.59	6.89	0.006
Parity	-	-	1.04	1.05	-	-	1.06	1.01	-	-	1.07	1.11	-	-	0.94	1.13	-	-	1.03	1.09	ns
HIV-positive status	242	22.04	-	-	183	31.02	-	-	18	19.78	-	-	23	13.07	-	-	18	-	7.47	-	< 0.001
Depression scale score antenatal	-	-	9.45	5.28	-	-	9.52	4.70	-	-	9.56	5.88	-	-	8.65	5.26	-	-	9.74	6.15	ns
Exposed to traumatic events	255	25.63	-	-	152	-	-	29.23	22	24.18	-	-	29	20.28	-	-	52	21.58	-	-	0.049
Past 12 months emotional abuse	265	26.69	-	-	84	-	-	16.25	39	42.86	-	-	40	27.78	-	-	102	42.32	-	-	< 0.001
Past 12 months physical abuse	217	21.85	-	-	83	-	-	16.05	30	32.97	-	-	27	18.75	-	-	77	31.95	-	-	< 0.001
Past 12 months sexual abuse	68	6.85	-	-	19	-	-	3.68	12	13.19	-	-	9	6.25	-	-	28	11.62	-	-	< 0.001
Infant length at birth (cm)	-	-	49.64	3.73	-	-	50.01	3.68	-	-	48.79	3.48	-	-	50.20	3.45	-	-	48.70	3.93	< 0.001
Standardised infant length at 12 months	-	-	-0.63	1.26	-	-	-0.38	1.16	-	-	-0.71	1.46	-	-	-0.69	1.26	-	-	-1.00	1.30	< 0.001
Standardised child length at 18 months	-	-	-0.88	1.45	-	-	-0.73	1.34	-	-	-0.89	1.50	-	-	-0.90	1.53	-	-	-1.16	1.54	< 0.001
Standardised child length at 24 months	-	-	-1.06	1.20	-	-	-0.88	1.15	-	-	-1.25	1.42	-	-	-1.03	1.26	-	-	-1.35	1.14	< 0.001
Infant weight at birth (kg)	-	-	3.02	0.60	-	-	3.08	0.59	-	-	2.95	0.59	-	-	3.12	0.58	-	-	2.84	0.61	< 0.001
Standardised infant weight at 12 months	-	-	-0.02	1.29	-	-	0.28	1.28	-	-	-0.16	1.06	-	-	-0.08	1.26	-	-	-0.50	1.24	< 0.001
Standardised child weight at 18 months	-	-	0.11	1.32	-	-	0.40	1.30	-	-	-0.13	1.26	-	-	0.03	1.30	-	-	-0.33	1.25	< 0.001
Standardised child weight at 24 months	-	-	-0.29	1.23	-	-	-0.00	1.21	-	-	-0.40	1.16	-	-	-0.33	1.22	-	-	-0.75	1.15	< 0.001
Number of months exclusive breastfeeding	-	-	2.09	1.96	-	-	2.01	2.01	-	-	1.59	1.72	-	-	2.21	1.92	-	-	2.33	1.93	0.002
Number of months any breastfeeding	-	-	9.91	9.50	-	-	7.80	8.31	-	-	7.90	9.51	-	-	10.49	9.51	-	-	14.99	10.09	< 0.001

HIV, human immunodeficiency virus; s.d., standard deviation; ns, not significant.

but there was a statistically significant reduction in child weight compared to the no-use group ($p = 0.094$ and $p = 0.022$). Alcohol and substance combined use was significantly associated with reduced child length and weight compared to the no-use group ($p = 0.002$ and $p < 0.001$). Findings from unadjusted GEE analyses mostly confirmed the findings of the multivariable analyses (see Supplementary Table 1).

Child length and weight associated with use timing

Table 4 shows the multivariable analyses by when use occurred with adjusted estimates on the standardised child length and weight as repeated measures at 12, 18 and 24 months. Antenatal use only and postnatal use only were not significantly associated with reduced child length compared to the no-use group ($p = 0.374$ and $p = 0.219$); however, use throughout antenatal and postnatal periods

showed a significant association with reduced child length compared to the no-use group ($p = 0.004$). Reduced child weight, on the other hand, was significantly associated with antenatal use only, postnatal use only and use throughout antenatal and postnatal periods compared to no use ($p = 0.019$, $p = 0.009$ and $p < 0.001$). Findings from unadjusted GEE analyses confirmed the findings of the multivariable analyses (see Supplementary Table 2).

Child length and weight associated with other covariates

Statistically significant covariates besides the grouping variables on the standardised child length and weight included months (i.e. 12, 18 and 24), maternal education and height, birth length or weight and any breastfeeding duration (Table 3 and Table 4). Maternal education and height, as well as birth length or weight, were positively associated with child length and weight, indicating that

TABLE 3: Child length and weight associated with substance use type ($N = 1098$).

Variables	Estimate	s.e.	95% CI	P
Child length				
Intercept	-10.91	0.92	-12.71, -9.11	< 0.01
Substance use type				
Alcohol use only	-0.07	0.11	-0.28, 0.14	ns
Substance use only	-0.25	0.15	-0.54, 0.04	ns
Alcohol and any other substance use	-0.28	0.09	-0.45, -0.10	< 0.01
Months (12, 18 and 24)	-0.04	< 0.01	-0.04, -0.03	< 0.01
Maternal age				
Maternal education less than secondary	-0.31	0.07	-0.45, -0.17	< 0.01
Marital status not married or cohabiting	-0.06	0.08	-0.21, 0.10	ns
Antenatal maternal height	0.04	0.01	0.03, 0.05	< 0.01
Maternal HIV-positive	-0.23	0.10	-0.43, -0.03	0.02
Trauma exposure				
Past 12 months emotional abuse exposure	0.09	0.10	-0.11, 0.28	ns
Past 12 months physical abuse exposure	< 0.01	0.10	-0.20, 0.20	ns
Past 12 months sexual abuse exposure	-0.01	0.16	-0.32, 0.31	ns
Infant length at birth (cm)	0.09	0.01	0.07, 0.11	< 0.01
Number of months exclusive breastfeeding	-0.03	0.02	-0.06, 0.01	ns
Number of months any breastfeeding	-0.01	0.01	-0.02, < 0.01	0.01
Child weight				
Intercept	-5.67	0.90	-7.43, -3.90	< 0.01
Substance use type				
Alcohol use only	-0.24	0.10	-0.44, -0.04	0.02
Substance use only	-0.32	0.14	-0.59, -0.05	0.02
Alcohol and any other substance use	-0.43	0.09	-0.61, -0.24	< 0.01
Months (12, 18 and 24)	-0.02	< 0.01	-0.02, -0.01	< 0.01
Maternal age				
Maternal education less than secondary	-0.33	0.08	-0.48, -0.18	< 0.01
Marital status not married or cohabiting	0.02	0.08	-0.14, 0.18	ns
Antenatal maternal height	0.03	0.01	0.02, 0.04	< 0.01
Maternal HIV-positive	-0.18	0.11	-0.39, 0.03	ns
Trauma exposure				
Past 12 months emotional abuse exposure	0.05	0.08	-0.12, 0.22	ns
Past 12 months physical abuse exposure	0.13	0.10	-0.08, 0.34	ns
Past 12 months sexual abuse exposure	< 0.01	0.11	-0.21, 0.21	ns
Past 12 months sexual abuse exposure	-0.22	0.15	-0.52, 0.09	ns
Infant weight at birth (kg)	0.72	0.07	0.58, 0.86	< 0.01
Number of months exclusive breastfeeding	-0.02	0.02	-0.06, 0.01	ns
Number of months any breastfeeding	-0.02	< 0.01	-0.03, -0.02	< 0.01

HIV, human immunodeficiency virus; CI, confidence interval; s.e., standard error; ns, not significant.

TABLE 4: Child length and weight associated with period of use (*N* = 1098).

Variables	Estimate	s.e.	95% CI	<i>P</i>
Child length				
Intercept	-10.88	0.92	-12.67, -9.08	< 0.01
Substance use timing				
Antenatal use only	-0.14	0.16	-0.45, 0.17	ns
Postpartum use only	-0.13	0.11	-0.34, 0.08	ns
Use throughout pregnancy and postpartum periods	-0.26	0.09	-0.44, -0.09	< 0.01
Months (12, 18 and 24 months)	-0.04	< 0.01	-0.04, -0.03	< 0.01
Maternal age				
Maternal education less than secondary	-0.31	0.07	-0.45, -0.16	< 0.01
Marital status not married or cohabiting	-0.05	0.08	-0.20, 0.11	ns
Antenatal maternal height	0.04	0.01	0.03, 0.05	< 0.01
Maternal HIV-positive	-0.23	0.10	-0.43, -0.03	0.03
Trauma exposure	0.11	0.08	-0.05, 0.26	ns
Past 12 months emotional abuse exposure	0.09	0.10	-0.11, 0.28	ns
Past 12 months physical abuse exposure	< 0.01	0.10	-0.20, 0.20	ns
Past 12 months sexual abuse exposure	< 0.01	0.16	-0.32, 0.31	ns
Infant length at birth (cm)	0.09	0.01	0.07, 0.11	< 0.01
Number of months exclusive breastfeeding	-0.03	0.02	-0.07, 0.01	ns
Number of months any breastfeeding	-0.01	0.01	-0.02, < 0.01	0.01
Child weight				
Intercept	-5.63	0.90	-7.40, -3.86	< 0.01
Substance use timing				
Antenatal use only	-0.34	0.15	-0.63, -0.06	0.02
Postpartum use only	-0.27	0.10	-0.47, -0.07	0.01
Use throughout pregnancy and postpartum periods	-0.40	0.09	-0.58, -0.21	< 0.01
Months (12, 18 and 24 months)	-0.02	< 0.01	-0.02, -0.01	< 0.01
Maternal age				
Maternal education less than secondary	-0.32	0.08	-0.47, -0.17	< 0.01
Marital status not married or cohabiting	0.03	0.08	-0.13, 0.18	ns
Antenatal maternal height	0.03	0.01	0.02, 0.04	< 0.01
Maternal HIV positive	-0.17	0.11	-0.38, 0.03	ns
Trauma exposure	0.05	0.08	-0.11, 0.22	ns
Past 12 months emotional abuse exposure	0.13	0.11	-0.08, 0.34	ns
Past 12 months physical abuse exposure	< 0.01	0.11	-0.21, 0.20	ns
Past 12 months sexual abuse exposure	-0.21	0.15	-0.51, 0.09	ns
Infant weight at birth (kg)	0.72	0.07	0.58, 0.86	< 0.01
Number of months exclusive breastfeeding	-0.02	0.02	-0.06, 0.01	ns
Number of months any breastfeeding	-0.02	< 0.01	-0.03, -0.02	< 0.01

HIV, human immunodeficiency virus; CI, confidence interval; s.e., standard error; ns, not significant.

higher maternal height and education and greater birth length or weight were associated with increased child length and weight. Longer duration of any breastfeeding was inversely associated with child length and weight. Maternal trauma exposure and past 12 month abuse exposure variables did not show any significant associations with child length and weight.

Discussion

Key findings

The current study was a secondary analysis of data based on an ongoing prospective study of mother-child pairs enrolled in the DCHS conducted in Western Cape, South Africa. Reduced standardised child length and weight outcomes at 12, 18 and 24 months were significantly associated with combined alcohol and non-prescribed substance use and continued use throughout antenatal and 1-year postnatal periods compared to no use. Alcohol

use-only, substance use-only, antenatal use-only and postnatal use-only groups were significantly associated with reduced infant and child weight compared to no use, however, not with infant and child length. Multivariable models showed that time (i.e. 12, 18 and 24 months), maternal height and education, birth length or weight, and breastfeeding duration were significantly associated with child outcomes.

Discussion of key findings

None of the average standardised stunting scores across the groups was within the range of severe stunting. The current report provided evidence that both combined use of alcohol and non-prescribed substances and continued use throughout antenatal and 1-year postnatal periods were significantly associated with reduced child length and weight outcomes up to 24 months, compared to no use. Notably, child weight was significantly less in the

alcohol use-only group and in the antenatal-only and postnatal-only groups, compared to the no-use groups. A majority of non-prescribed substances used comprised tobacco, which adversely impacts infant birth weight²⁸; thus, the weight outcomes reflected the high prevalence of tobacco use in the samples. Combined use of alcohol and tobacco worsens birth outcomes, including infant weight compared to antenatal alcohol exposure alone.⁹ The current study expands the previous evidence with findings that combined use showed more compromised growth up to 24 months of life compared to alcohol use alone.

The current report showed that a longer duration of breastfeeding was independently significantly associated with decreased child length and weight outcomes, and having at least a secondary level of education was associated with positive child length and weight outcomes. These findings were consistent with the existing body of literature on food insecurity's impact on breastfeeding in underserved communities²⁹ and social determinants of health on child growth.^{30,31,32}

Although associations of maternal trauma and past 12-month interpersonal violence with child length and weight were not significant, addressing maternal substance and alcohol use as well as relevant contextual issues such as violence and trauma exposure as a syndemic is important.

Strengths and limitations

The current report contributes to the field of maternal alcohol and non-prescribed substance use and infant and child outcomes, showing that combined use of alcohol and non-prescribed substances was associated with the most compromised child length and weight outcomes up to 24 months. The current report also added to the body of the evidence on perinatal alcohol and/or non-prescribed substance use with the evidence that use throughout antenatal and 1-year postnatal periods was also associated with the most compromised child length and weight outcomes up to 24 months.

Several limitations of the current report deserve emphasis. Firstly, some of the responses to the alcohol and substance use questions were missing, which lowered the sample size. Secondly, alcohol use was addressed only by self-report without biochemical verification although ASSIST questions on the past 3-month use are well validated across multiple populations and settings.³³ Substance use during postpartum was also only by self-report without biochemical verification. Thirdly, multiple substance use, without a sufficient number of participants using each, could not be dismantled to investigate the independent adverse effects of each substance use on child growth. Fourthly, dietary diversity was not investigated in this study cohort. Fifthly, this was an observational study and could not investigate a causal relationship between variables and child growth beyond associations.

Conclusion

The current study demonstrated the adverse associations of maternal alcohol and non-prescribed substance use with early child growth. Most of the non-prescribed substance comprised tobacco use, and combined adverse effects might have resulted from alcohol and tobacco co-use. Much emphasis on addressing combined and continued use of alcohol and substances during pregnancy and early postpartum needs to be made in terms of the potential long-term synergetic adverse effects on subsequent child growth. Proper education, guidance and intervention to reduce antenatal and postnatal combined use of alcohol and substances are urgently needed for optimal child length and weight outcomes. Other associated mental health and violence-related issues should also be addressed as contextual issues.

Acknowledgements

Competing interests

D.J.S. has received consultancy honoraria from Discovery Vitality, Kanna, L'Oreal, Lundbeck, Orion, Servier, Seaport Therapeutics, Takeda, and Wellcome. The author, D.J.S., serves as an editorial board member of this journal. D.J.S. has no other competing interests to declare.

Authors' contributions

Y.W. conceptualised the study and analyses, supervised the analytical process, developed the initial draft of the article and finalised the article. Z.Z. supervised the entire data analytic process and conducted the confirmatory analyses. M.L.B. conducted analyses under Z.Z.'s supervision. M.T.L. assisted with data management and the analytical process and reviewed the article draft. B.M. contributed to the conceptualisation of the study, facilitated team communication and reviewed the article. N.H. supported team communication and reviewed the article draft. E.G., H.J.Z. and D.J.S. assisted in the analytical process and reviewed the article draft. P.P.W. helped conceptualise the study and analyses, facilitated team communication and reviewed the article draft. H.J.Z. is the principal investigator of the DCHS and D.J.S. is the psychosocial principal investigator of the DCHS.

Funding information

The core study has been funded by the Gates Foundation (INV. 006732). Additional support for H.J.Z. and D.J.S. was provided by the Medical Research Council of South Africa.

Data availability

De-identified data will be made available from the principal investigator of the DCHS, author H.J.Z., upon reasonable request. Requests for collaborations to undertake data analysis are welcome. More information can be found on the website at <http://www.paediatrics.uct.ac.za/scah/dchls>.

Disclaimer

The views and opinions expressed in this article are those of the authors and are the product of professional research. It does not necessarily reflect the official policy or position of any affiliated institution, funder, agency or that of the publisher. The authors are responsible for this article's results, findings and content.

References

- Bailey BA, Sokol RJ. Prenatal alcohol exposure and miscarriage, stillbirth, preterm delivery, and sudden infant death syndrome. *Alcohol Res Health*. 2011;34(1):86–91.
- CDC. About Fetal Alcohol Spectrum Disorders (FASDs) [homepage on the Internet]. 2024 [cited 2024 Oct 24]. Available from: <https://www.cdc.gov/fasd/about/index.html>
- Lange S, Probst C, Gmel G, Rehm J, Burd L, Popova S. Global prevalence of fetal alcohol spectrum disorder among children and youth. *JAMA Pediatr*. 2017;171(10):948. <https://doi.org/10.1001/jamapediatrics.2017.1919>
- Olivier L, Curfs L, Viljoen D. Fetal alcohol spectrum disorders: Prevalence rates in South Africa. *S Afr Med J*. 2016;106(6 Suppl 1):S103–S106. <https://doi.org/10.7196/SAMJ.2016.v106i6.11009>
- Petersen-Williams P, Jordaan E, Mathews C, Lombard C, Parry CDH. Alcohol and other drug use during pregnancy among women attending midwife obstetric units in the Cape Metropole, South Africa. *Adv Prev Med*. 2014;2014:871427. <https://doi.org/10.1155/2014/871427>
- Washio Y, Mericle AA, Cassey H, Daubert AM, Kirby KC. Characteristics of low-income racial/ethnic minority pregnant women screening positive for alcohol risk. *J Immigr Minor Health*. 2016;18(4):850–855. <https://doi.org/10.1007/s10903-015-0238-5>
- Fish EW, Murdaugh LB, Zhang C, et al. Cannabinoids exacerbate alcohol teratogenesis by a CB1-Hedgehog Interaction. *Sci Rep*. 2019;9(1):16057. <https://doi.org/10.1038/s41598-019-52336-w>
- Brink LT, Springer PE, Nel DG, Potter MD, Odendaal HJ. The tragedy of smoking, alcohol, and multiple substance use during pregnancy. *S Afr Med J*. 2022;112(8):526–538. <https://doi.org/10.7196/SAMJ.2022.v112i8.16480>
- Odendaal H, Wright C, Schubert P, et al. Associations of maternal smoking and drinking with fetal growth and placental abruption. *Eur J Obstet Gynecol Reprod Biol*. 2020;253:95–102. <https://doi.org/10.1016/j.ejogrb.2020.07.018>
- May PA, Hasken JM, Blankenship J, et al. Breastfeeding and maternal alcohol use: Prevalence and effects on child outcomes and fetal alcohol spectrum disorders. *Reprod Toxicol*. 2016;63:13–21. <https://doi.org/10.1016/j.reprotox.2016.05.002>
- Skagerstrom J, Chang G, Nilsen P. Predictors of drinking during pregnancy: A systematic review. *J Womens Health (Larchmt)*. 2011;20(6):901–913. <https://doi.org/10.1089/jwh.2010.2216>
- Davis EC, Rotheram-Borus MJ, Weichle TW, Rezai R, Tomlinson M. Patterns of alcohol abuse, depression, and intimate partner violence among township mothers in south africa over 5 years. *AIDS Behav*. 2017;21(S2):174–182. <https://doi.org/10.1007/s10461-017-1927-y>
- Myers B, Koen N, Donald KA, et al. Effect of hazardous alcohol use during pregnancy on growth outcomes at birth: Findings from a South African Cohort Study. *Alcohol Clin Exp Res*. 2018;42(2):369–377. <https://doi.org/10.1111/acer.12356>
- Farewell CV, Donohoe R, Thayer Z, et al. Maternal depression trajectories and child BMI in a multi-ethnic sample: A latent growth modeling analysis. *BMC Pregnancy Childbirth*. 2021;21(1):827. <https://doi.org/10.1186/s12884-021-04308-0>
- Choi KW, Denckla CA, Hoffman N, et al. Influence of maternal childhood trauma on perinatal depression, observed mother-infant interactions, and child growth. *Matern Child Health J*. 2022;26(8):1649–1656. <https://doi.org/10.1007/s10995-022-03417-2>
- Stein D, Koen N, Donald K, et al. Investigating the psychosocial determinants of child health in Africa: The Drakenstein Child Health Study. *J Neurosci Methods*. 2015;252:27–35. <https://doi.org/10.1016/j.jneumeth.2015.03.016>
- Zar HJ, Barnett W, Myer L, Stein DJ, Nicol MP. Audit, research and guideline update: Investigating the early-life determinants of illness in Africa: The Drakenstein Child Health Study. *Thorax*. 2015;70(6):592. <https://doi.org/10.1136/thoraxjnl-2014-206242>
- Ali R, Awwad E, Babor TF, et al. The Alcohol, Smoking and Substance Involvement Screening Test (ASSIST): Development, reliability and feasibility. *Addiction*. 2002;97(9):1183–1194. <https://doi.org/10.1046/j.1360-0443.2002.00185.x>
- Fenton TR, Kim JH. A systematic review and meta-analysis to revise the Fenton growth chart for preterm infants. *BMC Pediatr*. 2013;13(1):1–13. <https://doi.org/10.1186/1471-2431-13-59>
- WHO. WHO child growth standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development [homepage on the Internet]. 2006 [cited 2023 Oct 16]. Available from: <https://iris.who.int/handle/10665/43413>
- CDC. Positive parenting tips: Infants (0–1 years) [homepage on the Internet]. Child Development. CDC; 2014 [cited 2024 Jun 11]. Available from: https://www.cdc.gov/child-development/positive-parenting-tips/infants.html?CDC_ARef_Val=https://www.cdc.gov/ncbddd/childdevelopment/positiveparenting/infants.html
- Wedderburn CJ, Bondar J, Lake MT, et al. Risk and rates of hospitalisation in young children: A prospective study of a South African birth cohort. *PLoS Glob Public Health*. 2024;4(1):e0002754 <https://doi.org/10.1371/journal.pgph.0002754>
- Sinha R, Dua R, Bijalwan V, Rohatgi S, Kumar P. Determinants of stunting, wasting, and underweight in five high-burden pockets of four Indian States. *Indian J Community Med*. 2018;43(4):279. https://doi.org/10.4103/ijcm.UJCM_151_18
- Jewkes R. Intimate partner violence: Causes and prevention. *Lancet*. 2002;359(9315):1423–1429. [https://doi.org/10.1016/S0140-6736\(02\)08357-5](https://doi.org/10.1016/S0140-6736(02)08357-5)
- Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. *Br J Psychiatry*. 1987;150:782–786. <https://doi.org/10.1192/bjp.150.6.782>
- Foa EB, Riggs DS, Dancu CV, Rothbaum BO. Reliability and validity of a brief instrument for assessing post-traumatic stress disorder. *J Trauma Stress*. 1993;6(4):459–473. <https://doi.org/10.1007/BF00974317>
- Foa EB, McLean CP, Zang Y, et al. Psychometric properties of the posttraumatic stress disorder symptom scale interview for DSM-5 (PSSI-5). *Psychol Assess*. 2016;28(10):1159–1165. <https://doi.org/10.1037/pas0000259>
- Cnattingius S. The epidemiology of smoking during pregnancy: Smoking prevalence, maternal characteristics, and pregnancy outcomes. *Nicotine Tob Res*. 2004;6:125–140. <https://doi.org/10.1080/14622200410001669187>
- Budree S, Goddard E, Brittain K, Cader S, Myer L, Zar HJ. Infant feeding practices in a South African birth cohort – A longitudinal study. *Matern Child Nutr*. 2017;13(3):1–9. <https://doi.org/10.1111/mcn.12371>
- CDC. Frequently asked questions. Social determinants of health [homepage on the Internet]. NCHHSTP. CDC; 2019 [cited 2022 Jan 10]. Available from: <https://www.cdc.gov/nchhstp/socialdeterminants/faq.html>
- Higgins ST, Chilcoat HD. Women and smoking: An interdisciplinary examination of socioeconomic influences. *Drug Alcohol Depend*. 2009;104 Suppl:S1–S5. <https://doi.org/10.1016/j.drugalcdep.2009.06.006>
- Higgins ST, Heil SH, Badger GJ, Skelly JM, Solomon LJ, Bernstein IM. Educational disadvantage and cigarette smoking during pregnancy. *Drug Alcohol Depend*. 2009;104 Suppl:S100–S105. <https://doi.org/10.1016/j.drugalcdep.2009.03.013>
- Da Silva AC, Lucchese R, Vargas LS, Benício PR, Vera I. Application of the Alcohol, Smoking and Substance Involvement Screening Test (ASSIST) instrument: An integrative review. *Rev Gaucha Enferm*. 2016;37(1):e52918. <https://doi.org/10.1590/1983-1447.2016.01.52918>