

This is a repository copy of *The pan-tropical age distribution of regenerating tropical moist forest*.

White Rose Research Online URL for this paper: <u>https://eprints.whiterose.ac.uk/229160/</u>

Version: Published Version

#### Article:

Bousfield, C.G. orcid.org/0000-0003-3576-9779 and Edwards, D.P. orcid.org/0000-0001-8562-3853 (2025) The pan-tropical age distribution of regenerating tropical moist forest. Nature Ecology & Evolution, 9 (7). pp. 1205-1213. ISSN 2397-334X

https://doi.org/10.1038/s41559-025-02721-8

#### Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here: https://creativecommons.org/licenses/

#### Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



# The pan-tropical age distribution of regenerating tropical moist forest

Received: 27 November 2024

Accepted: 14 April 2025

Published online: 20 May 2025

Check for updates

Christopher G. Bousfield <sup>1,2,3</sup> & David P. Edwards <sup>1,2</sup>

Natural forest regeneration in the tropics is a key element of restoration pledges. Protecting older regenerating forests that already hold substantial carbon and biodiversity value, while promoting natural regeneration in young secondary forests in regions where forests are likely to persist long term, is vital for effective forest restoration. Key questions therefore include understanding the age distribution of naturally regenerating forests pan-tropically and which environmental or socio-economic conditions predict increased longevity in regenerating forests. Here, using a time series of forest cover data (1990-2023) to map the age of regenerating tropical moist forests, we identify 51 Mha of regenerating tropical moist forest, of which >50% is  $\le 5$  years old and under high deforestation pressure, whereas only 6% (3 Mha) is  $\ge 20$  years old and located predominantly in the tropical Americas. Location and forest characteristics in the surrounding landscape best predict the age of regenerating forests, with older forests located in areas with high forest integrity and extent, and low forest loss. Realizing the environmental and social values of naturally regenerating forests requires urgent financial, political and societal mechanisms to facilitate the long-term persistence of restoration.

Large-scale restoration of forest ecosystems has immense capacity to sequester globally important amounts of carbon to combat climate change<sup>1</sup>, create millions of hectares of habitat to safeguard biodiversity<sup>2</sup> and contribute towards multiple Sustainable Development Goals<sup>3</sup>. Many global initiatives promote the restoration of degraded ecosystems, including the 2021–2030 United Nations Decade on Ecosystem Restoration<sup>4</sup>, Target 2 of the Global Biodiversity Framework to restore 30% of all degraded ecosystems by 2030<sup>5</sup> and the Bonn Challenge to bring 350 million hectares (Mha) of degraded landscapes under restoration by 2030<sup>6</sup>. Such international commitments have created a critical opportunity to restore degraded ecosystems globally.

Forest restoration in the tropics is a key element of global restoration strategies, as it offers the largest benefits for both carbon and biodiversity at comparatively low economic cost<sup>2,7,8</sup>. Natural forest regeneration–naturally occurring regrowth on previously deforested and degraded land, sometimes assisted by direct planting of native seedlings–accounts for 34% (64 Mha) of all tropical restoration pledges<sup>3</sup>. The remainder consists of commercial tree plantations (for example, for timber) or agroforestry<sup>3</sup>, yet national-level pledges are often ambitious, untracked and can even exceed the total area available for restoration in many countries<sup>9</sup>.

Naturally regenerating forests store more carbon, support more biodiversity and provide greater ecosystem services (that is, water provision and soil erosion control) than plantations<sup>10</sup>. Recovery of forest biomass and structure in naturally regenerating tropical forests is rapid, up to 100 times faster than in slower-growing regions (for example, boreal forest)<sup>7</sup>, with carbon stocks capable of reaching 85% of old-growth values in just 20 years, and 90% of old-growth values in 66 years on average<sup>11</sup>. Biodiversity of most plant and animal taxa can also recover rapidly, with up to 80% of old-growth species richness levels recovered in 20 years, and full recovery after 50 years<sup>12-15</sup> (though full recovery of species composition can take centuries)<sup>12</sup>.

<sup>1</sup>Department of Plant Sciences and Centre for Global Wood Security, University of Cambridge, Cambridge, UK. <sup>2</sup>Conservation Research Institute, University of Cambridge, Cambridge, UK. <sup>3</sup>Ecology and Evolutionary Biology, School of Biosciences, University of Sheffield, Sheffield, UK. ©e-mail: cgb48@cam.ac.uk This rapid recovery of biodiversity, carbon and forest function makes natural regeneration of tropical forests one of the most effective and cost-effective forms of ecosystem restoration.

Natural forest regeneration is feasible across an estimated 215 Mha of deforested tropical land<sup>16</sup>, indicating widespread opportunities to restore natural forest ecosystems across the tropics. Restoration would be most effective if these initiatives ensure the longevity of the new forests they promote or target the protection and further recovery of existing older naturally regenerating forests that are in danger of being deforested<sup>17</sup>. But regenerating forests are frequently at risk of clearance. In the Amazon, for example, almost half of naturally regenerating forests are  $\leq 5$  years old<sup>17,18</sup> and subject to increased clearance<sup>19</sup>, and the probability of persistence is strongly linked to surrounding tree cover and proximity to existing forests in southern Costa Rica (CRI) and the Brazilian Atlantic Forest<sup>20,21</sup>. However, the age distribution of regenerating forests, and conditions that promote their persistence across the tropics remains unclear.

A key question is understanding the age and spatial distribution of regenerating tropical forests and the environmental and social conditions that support recovery. This would allow for targeted conservation of older regenerating forests and focused restoration efforts in areas that are more likely to support longer-term forest recovery. Here, we develop a 30 m-resolution mapped time series of the age of regenerating tropical moist forest (TMF) growing on previously deforested land using the latest version of a remotely sensed land-use change dataset covering the period 1990 and 2023<sup>22</sup>. We use this to answer the following questions: (1) where are the oldest regenerating TMFs, (2) which regions and nations are most important for harbouring older regenerating forests? and (3) what socio-environmental landscape characteristics best predict regenerating forest age?

#### Results

#### Age distribution of regenerating forests across the tropics

We used a time series (1990–2023) of the annual extent of TMF<sup>22</sup> (defined as a closed forest with >90% canopy cover located in the humid tropics, not including tropical dry forests) to map and age regenerating forests across the tropics at 30 m resolution. In 2023, there was 51 Mha of regenerating TMF, with 21.5 Mha found in the tropical Americas, followed by 19.4 Mha in Asia–Pacific and 9.8 Mha in Africa (Fig. 1). Just five countries–Brazil (BRA, 10.6 Mha), Indonesia (IDN, 7.2 Mha), Democratic Republic of Congo (COD, 3.3 Mha), Colombia (COL, 2.6 Mha) and Myanmar (MMR, 2.2 Mha)–account for over half (52%) of the world's regenerating TMFs. Globally, regenerating TMF has a mean age of 7.6 ± 5.9 years (median age of 5 years) but demonstrates regional variation, with mean ages of 8.4 ± 6.7, 6.0 ± 4.2 and 7.4 ± 5.3 years (median ages 6, 5 and 5 years) in the Americas, Africa and Asia–Pacific, respectively.

Only 6% (3 Mha) of regenerating TMF is  $\ge 20$  years old. Of this 'old' regenerating forest, 63% (1.9 Mha) is located in the American tropics, 31% (0.9 Mha) in the Asian-Pacific tropics and only 7% (0.2 Mha) in the African tropics (Fig. 1b). Hotspots of old regenerating TMF extent are located predominantly in the Amazon Basin, Brazilian Atlantic Forest, Central America and Congo Basin as well as Borneo, Indo-Burma and southern Papua New Guinea (PNG). Many of these areas overlap with global biodiversity and conservation hotspots<sup>23</sup>, areas with the highest carbon accumulation potential in regenerating forests<sup>7</sup>, and restoration opportunity hotspots<sup>24</sup>, highlighting the important role that older regenerating TMFs could play in global conservation and restoration efforts.

By contrast, 52% (26.3 Mha) of regenerating TMF is 5 years old or younger, of which 39% (10.1 Mha), 24% (6.3 Mha) and 38% (9.9 Mha) is located in the Americas, Africa and Asia–Pacific, respectively (Fig. 1c; see Supplementary Figs. 1 and 2 for national-level distributions). As seen in Brazil<sup>17,21</sup>, this young forest is probably at high risk of conversion. In Costa Rica, where the relative extent of old regenerating forest was comparatively large, regional studies still suggest that >50% of naturally regenerating forests are cleared within 20 years of establishment<sup>20</sup>. Similarly, in the Brazilian Atlantic, a hotspot of older regenerating forests, the turnover of 'ephemeral' regenerating forests is high, with a mean age of just 7.9 years<sup>21</sup>, while 70% of cleared regenerating forest in the Amazon are 5 years old or younger<sup>17</sup>.

#### Older regenerating forests are concentrated in the Americas

At a national scale, Brazil accounts for 37% (1.1 Mha) of old regenerating TMF ( $\geq$ 20 years), and combined with Indonesia (0.3 Mha) and Colombia (0.2 Mha), represents >50% of global old regenerating TMF (Fig. 2a). Of the 30 countries with the largest extent of old regenerating forest, 14 are tropical American, 10 Asian-Pacific and only 6 African (Fig. 2a). Tropical American countries also account for a higher share of pan-tropical older regenerating TMF area than would be expected based on their pan-tropical share of regenerating forest area of all ages (Fig. 2b), holding 63% of old regenerating TMF in just 42% of the total regenerating TMF area. Conversely, Africa and Asia-Pacific hold less old regenerating forest than would be expected given their share of pan-tropical regenerating forest extent (Africa is composed of 6% of old regenerating forest on 19% of regenerating forest; Asia-Pacific is composed of 31% of old regenerating forest on 38% of regenerating forest). The drivers behind these continental-scale differences are unclear, but could be due to greater agricultural intensification<sup>25</sup>, increased shares of populations living in urban areas and declines of rural populations<sup>26,27</sup> or forest transitions as a result of greater economic growth<sup>28</sup>. Similar deforestation trajectories across the regions since 1990 (Fig. 1c) suggest that the observed variance in regenerating forest age distributions is not simply a legacy of the timing of past deforestation events.

Of the 50 countries with regenerating TMF in the Americas, 21 have more old regenerating forest than would be expected based on their total regenerating forest extent, with Costa Rica, Honduras (HND) and Brazil supporting 3.7, 2.6 and 1.8 fold more old regenerating forest than expected, respectively (Fig. 2b). High extents of older regenerating TMFs in countries, such as Costa Rica, suggests national-level policies focusing on forest conservation (for example, the 1996 Forest Law) may impact the age distribution of regenerating forest extent<sup>29</sup>. By contrast, only 5 of 35 African countries have more old regenerating TMF than expected, with many West African countries (for example, Ghana and Sierra Leone) skewed towards younger forests and holding ten times less older-regenerating forest than expected, probably owing to the dominance of shifting cultivation in these countries and sub-Saharan Africa more broadly<sup>30</sup> (Fig. 2b). Shifting cultivation itself is an important land-use system that occurs over 280 Mha worldwide<sup>31</sup>, and produces a mosaic of habitat capable of sustaining high species richness and carbon stocks<sup>32</sup>, especially when compared with intensive agriculture<sup>33</sup>. Tropical Asia-Pacific is more mixed, with 8 of 23 countries having more old regenerating tropical forest than expected, including Bangladesh (BGD) and Papua New Guinea with 2.5 and 1.3 times more old regenerating forest, respectively, while other countries such as Myanmar and Vietnam (VNM) have only half the expected old forest (Fig. 2b). Myanmar, Laos (LAO) and Vietnam are the three countries with the greatest extent of shifting cultivation in tropical Asia-Pacific<sup>34</sup>, which could explain their younger regenerating forest age distributions, while Papua New Guinea has larger remaining areas of undisturbed primary forest and less historical forest loss compared with other regions, such as Borneo<sup>22</sup>, perhaps leading to greater prevalence of older regenerating forest.

#### Landscape forest characteristics best predict forest age

We used random forest models with 32 predictor variables (see Supplementary Table 1 for more detail), spanning factors related to forest, environment, location and human pressure, to determine which are most important for predicting regenerating TMF age at a 30 m resolution. Location variables, including longitude, latitude and country



**Fig. 1** | **Age distribution of regenerating TMF. a**, The mean age of regenerating TMF per 30 km grid cell. **b**, The hotspots of total area of older (≥20 years) regenerating TMF within 30-km cells. Cells in the 75th percentile for total area of old regenerating TMF are coloured in light green and cells in the 90th percentile are coloured in dark green. **c**, The area distribution (Mha) of regenerating TMFs

among age classes in the tropical Americas (left), Africa (middle) and Asia–Pacific (right). The dashed line represents the proportion of all deforestation (1990–2023) that occurred during the corresponding time period (for example, height of the line at 6–10 years represents the proportion of deforestation between 1990 and 2023 that occurred 6–10 years ago).

had high predictive power, indicating that older regenerating forests are spatially clustered across landscapes and regions, and that national-level policy and cultural practices can have strong impacts on the age distribution of regenerating forest<sup>29</sup> (Fig. 3). Additionally, elevation was the most important predictor in Africa (but less so in the Americas and Asia–Pacific), where forests tended to be increasingly old as altitude increased to -2,000 metres above sea level (Fig. 4), probably owing to the lower deforestation threat at higher elevations as land becomes increasingly marginal for agriculture<sup>35</sup>.

Forest landscape characteristics tended to be very important in predicting regenerating forest age, particularly in the Americas (Fig. 3; see Supplementary Fig. 3 for pan-tropical model). Landscapes with higher Forest Landscape Integrity Index (FLII) scores<sup>36</sup>, a larger extent of regenerating forests (especially in the Americas) and lower levels of forest loss tended to have older regenerating forests (Fig. 4; see Supplementary Fig. 4 for partial-effects of all variables). This supports regional studies in Costa Rica and Brazil, which found that nearby forest cover was one of the most important predictors of regenerating forests are important predictors of natural forest regeneration potential at the pan-tropical scale<sup>16</sup>. Older regenerating forests located in high forest density landscapes are also likely to be supporting elevated biodiversity given landscape forest extent is one of the most important predictors of biodiversity recovery in naturally regenerating forests<sup>37</sup>. Our results therefore indicate that restoration efforts focusing on natural regeneration should target areas with high surrounding forest landscape integrity to increase the chances of long-term restoration success.

Some environmental variables were also important in predicting the age of regenerating TMF, in particular temperature seasonality (Fig. 4), whereby predicted age increased as seasonality grew to intermediate levels before either plateauing (Americas) or dropping (Africa and Asia-Pacific) as seasonality became more extreme. Factors relating directly to forms of human pressure on the landscape (for example, deforestation drivers, agricultural suitability, cropland extent and protected area status) had little influence. This poor predictive power may be due to coarse-resolution mapping of some variables (for example, deforestation driver at 10 km<sup>2</sup>)<sup>30</sup> rendering them unable to capture the substantial age variation within landscapes or because other predictors without readily available pantropical data (for example, land tenure security and land-use history)<sup>38</sup> may have stronger influences over the longevity of regenerating forest. Furthermore, the high predictive power of variables, such as forest loss, regenerating forest extent and FLII, are indirectly reflective of human pressures. Natural regeneration potential is also highest in heavily forested landscapes<sup>16</sup>, and plays a vital role even where tree planting is used to initiate forest recovery<sup>39</sup>. Our results indicate that restoration efforts focusing on natural regeneration should target



**Fig. 2** | **National-level distribution of older regenerating TMF. a**, Total area (in Mha) of the ten countries with the largest extent of old ( $\geq$ 20 years) regenerating TMF by region (Americas, Africa and Asia–Pacific). **b**, The national-level ratio of the proportion of all pan-tropical old regenerating TMF compared to the proportion of all regenerating TMF of any age. A ratio of 1 means a country has the same share of all old regenerating forest as it does all regenerating forest of any age across the tropics (for example, 10% of all old regenerating forest and 10% of all regenerating forest). Blue colours represent ratios >1, whereby the country

has more old regenerating forest than would be expected based on its share of all regenerating forest, red colours represent ratios <1, whereby the country has less old regenerating forest than would be expected. MEX, México; BOL, Bolivia; PER, Peru; GTM, Guatemala; VEN, Venezuela; PAN, Panama; AGO, Angola; CAF, Central African Republic; CIV Côte d'Ivoire; ETH, Ethiopia; CMR, Cameroon; NGA, Nigeria; KEN, Kenya; MDG, Madagascar; COG, Republic of the Congo; MYS, Malaysia; THA, Thailand; IND, India; PHL, Philippines.

areas with high surrounding forest landscape integrity to increase the chances of long-term restoration success.

The greatest uncertainty in our study originates in the difficulty of accurately distinguishing naturally regenerating forest from tree plantations, which have expanded across the tropics at a similar rate to naturally regenerating forests<sup>40</sup>. To avoid confusion with oil palm, we used a pan-tropical dataset of smallholder and industrial oil palm plantations to mask out known oil palm plantations from our dataset<sup>41</sup>. We compared our data on regenerating TMFs with a recent global map of forest management<sup>42</sup> and found minimal overlap with areas confidently identified as tree plantations or agroforestry (Supplementary Note 1 and Supplementary Table 2). In addition, the short rotation times of tropical tree plantations (for example, <10 years in Eucalyptus and Acacia plantations across much of the tropics)43-45 means that areas of old regenerating forests with detectable tree cover for  $\geq$ 20 years are unlikely to have been misidentified tree plantations. We also compared our results with what is, to our knowledge, the only publicly available large dataset of regenerating forest age compiled for Brazil as part of the MapBiomas project<sup>46,47</sup> (Supplementary Note 2 and Supplementary Fig. 5). Despite substantially different definitions of regenerating forest between the two datasets, we found moderate agreement between the datasets (Supplementary Fig. 5) in both regenerating forest area ( $R^2 = 0.49$ ) and age ( $R^2 = 0.37$ ). Additional analyses assessing the age distribution of regenerating forests at regional scales also found similar age profiles for regenerating forests across the Brazilian Amazon<sup>48</sup> and Amazon Biome<sup>17</sup>.

#### Discussion

Our study identifies 3 Mha of old regenerating TMF ( $\geq$ 20 years) located predominantly across the tropical Americas, typically in regions with high forest integrity and low deforestation rates, that will already hold substantial biodiversity and carbon value<sup>11,12</sup> and should be priority targets for conservation and further restoration. However, we also find that the majority of naturally regenerating TMF is  $\leq$ 5 years old, and probably caught in continuous cycles of farm abandonment, regeneration, deforestation and agricultural production, leaving them unable to accrue substantial carbon stocks or biodiversity in the long term. Using spatially explicit models of the carbon accumulation potential of naturally regenerating forests<sup>7</sup>, we estimate that protecting all 26 Mha of young regenerating TMF for the next 25 years would result in a potential above-ground sequestration of ~3 gigatons of carbon (GtC) by 2050. Ensuring the protection of naturally regenerating forests from re-clearance through financial, political and societal mechanisms should thus be key a target for the global restoration agenda to ensure long-term provision of forest ecosystem services<sup>49</sup>.

Carbon-financing schemes through Reducing Emissions from Deforestation and Forest Degradation (REDD+), offer a route towards funding the protection of regenerating forests, potentially creating a financially viable alternative to re-clearance for agricultural production<sup>50</sup>. Such schemes can also be used to promote assisted natural regeneration (ANR), which could enhance the recovery of carbon and biodiversity relative to passive regeneration<sup>49</sup> in a cost-effective



**Fig. 3** | **Variable importance scores for predicting regenerating TMF age.** The variable importance scores from the random forest models for each region, with variables ordered by their total predictive power across all three regional models and coloured by type: environmental variables (purple), variables concerning the surrounding forest landscape (green), variables relating to levels of human

pressure (orange) and variables relating to the location of the regenerating forest (blue). Insets: random forest model error (RMSE) and variance explained ( $R^2$ ) for each region. GDP, gross domestic product; HDI, human development; SOC, soil organic carbon.



Fig. 4 | Partial effects of the six most important predictors of regenerating TMF age. The partial effect plots on regenerating forest age of the top six most important variables across the three models (excluding longitude, latitude and country) in order of importance (left to right). The regional-level partial effects are coloured by region: Americas (green), Africa (orange) and Asia–Pacific (purple). The distribution of data points for modelling partial effects in each region is shown at the bottom of each plot. Predicted age values represent predictions from the stratified sample of secondary forest points used to train the random forest models, and not the raw data, which was heavily skewed towards younger forests.

manner<sup>51,52</sup> that also generates employment opportunities across the cycle of activities (for example, seed collection, nursery, land preparation, direct carbon payments to landowners and land-holding communities), thus incentivising the longer-term protection of these forests. However, methodologies to certify natural forest regeneration are lacking. Requirements for ten consistent years of forested or unforested land cover before an area qualifies for REDD+ or afforestation, reforestation and revegetation, and the premise that regenerating forests provide no additionality because they are already growing (despite their elevated deforestation risk<sup>19</sup>) prevents most young regenerating forests from qualifying for any form of carbon-financing scheme<sup>53</sup>. Under current regulations, regenerating forests therefore have limited economic value and cannot counter the often-marginal opportunity cost of conversion to agricultural land<sup>21,50</sup>. Unlocking this funding gap and prescribing economic value to regenerating forests will be vital for preventing ongoing future losses of ecologically valuable forest systems.

Effective regulatory protection for regenerating forests and community engagement in restoration and protection is also key. The combination of strict legislation in Brazil's Forest Code and specific legal protections in the Brazilian Atlantic Forest that prevent clearance of regenerating forests >10 years old<sup>54</sup> may be contributing towards the widespread natural regeneration occurring there<sup>55</sup>. Expanding these protections across the tropics through coherent and consistent legal frameworks<sup>56,57</sup> could promote increased forest longevity. However, protections must be carefully designed to avoid perverse outcomes. Environmental service payments in Costa Rica economically favour the replacement of naturally regenerating forests with native tree plantations that have lower the environmental value<sup>10,37,57</sup>, while the 2008 Amazonian beef moratorium successfully slowed primary forest loss in Brazil but at the expense of naturally regenerating forests, which suffered a 280% increase in clearance rates<sup>58</sup>. Community participation and engagement in reforestation projects, as well as benefits sharing, is also critical in achieving socio-ecological benefits of restoration and long-term persistence59,60. Officially recognised indigenous territories have lower deforestation rates throughout the tropics<sup>61</sup>, while in the Brazilian Amazon collective property rights leads to larger, older

regenerating forest extents<sup>38</sup>. Engaging communities in restoration efforts and improving the security of land-tenure could therefore increase the long-term success of naturally regenerating forests in tropical landscapes.

Addressing the underlying societal causes of deforestation in regenerating forests would ensure greater longevity and provision of ecosystem services where natural regeneration occurs. Where smallholder cultivation is leading to losses of regenerating forest, programmes to improve yields<sup>62</sup> and diversify income streams for local communities (for example, through provision of timber or non-timber forest products or carbon-funded ANR)<sup>63</sup> could reduce deforestation pressure. At the global scale, closing yield gaps and implementing economically efficient land-use decisions could increase agricultural production by 79-148%<sup>64</sup>, while shifts away from land-intensive diets relying heavily on ruminant meats could reduce land requirements by up to 70%, opening up some of the ~3.7 billion hectares used for livestock production<sup>65</sup> to long-term natural regeneration. Expansion of naturally regenerating forests into abandoned crop and pastureland through natural colonisation and ANR<sup>49</sup> could then be strategically targeted to maximise the environmental and social values of natural regeneration<sup>2,7,16,57</sup>.

Overall, this study highlights that despite the existence of pockets of long-term natural regeneration in the tropics, the potential restoration value of naturally regenerating tropical forests is currently not being realised. Alongside the protection of remaining old-growth forests, preventing the continued cyclical deforestation of naturally regenerating forests through economic investment, community engagement, effective regulation and societal shifts in diet will be crucial in realising the restoration potential of naturally regenerating forests. Only through these pathways will regenerating forests transition from ephemeral ecosystems<sup>20,21</sup> into a key pillar of global restoration efforts.

#### Methods

#### Mapping regenerating TMF age

To create our map of tropical regenerating forest age, we used the TMF annual change dataset<sup>22</sup>. The TMF annual change product is a forest change dataset from the European Commision's Joint Research Centre that uses Landsat to track land cover change dynamics in humid TMFs (>90% canopy cover) at a resolution of 30 m between the years 1990 and 2023<sup>22</sup>. To create the map of regenerating TMF age in 2023, we first used the transition map and retained only pixels identified as 'regrowth forest', that is, pixels demonstrating naturally regrowing vegetation for 3 years or more after a deforestation event that resulted in a period of more than 2.5 years without detectable tree cover. After masking to retain only regenerating TMF pixels in the year 2023, we then stacked each year of the annual change data (1990-2023) together. Following a similar method to Heinrich et al.<sup>66</sup> and Silva et al.<sup>18</sup>, and starting from 1990, for every consecutive year where regenerating forest was detected in a pixel, the forest age value of that pixel increased by 1. If a different land-use classification was detected on a pixel (for example, owing to a deforestation event), the regenerating forest age for that pixel was reset to 0. The age of the pixel remained 0 until the next year that regenerating forest was detected in the pixel, at which point the age counter would start again from one. This process was repeated cumulatively for the years 1990 through to 2023, with the final layer representing the age of regenerating TMF (that is, how many cumulative years that regenerating forest had been detected in that pixel) in the year 2023. The requirement for 3 years of forest cover before designation as a 'forest regrowth' pixel and the earliest year of the annual change dataset (1990) thus give our regenerating forest age values a possible range of 3-34 years old.

To remove any potential oil palm plantations classified as regenerating forest in the dataset, we masked out all pixels mapped as oil palm plantations in the map produced by Descals et al.<sup>41</sup>. We also tested our map of regenerating forest age against the map of regenerating vegetation age for Brazil from the MapBiomas project<sup>46,47</sup> (Supplementary Note 2 and Supplementary Fig. 5) and the global map of forest management by Lesiv et al.<sup>42</sup> (2022) (Supplementary Note 1 and Supplementary Table 2).

The forest aging process was carried out in Google Earth Engine<sup>67</sup>, with all subsequent spatial analysis conducted in RStudio version 4.4.4<sup>68</sup>. We used the 'terra'<sup>69</sup> package to aggregate age maps for visualisation, identify pan-tropical hotspots of older regenerating TMF (defined as  $\geq$ 20 years old) and analyse regenerating forest extent and age profiles for each country on the basis of national boundaries from GADM<sup>70</sup>. Maps were plotted on the basis of national boundaries from 'rnaturalearth'<sup>71</sup>. To estimate the carbon accumulation potential of protecting all young ( $\leq$ 5 years old) naturally regenerating TMF for the next 25 years, we extracted carbon accumulation values from spatially explicit estimates of annual aboveground carbon accumulation in naturally regenerating forests<sup>7</sup> and multiplied these by 25.

#### Identifying the best predictors of regenerating forest age

To identify the most important variables in predicting the age of regenerating TMF pixels, we used random forest regression models to predict regenerating forest age on the basis of a number of input variables. Random forest models employ an ensemble of decision trees for predictions, and were used since they are generally robust to interactions between predictors and complex nonlinear and non-monotonic relationships<sup>72</sup>. To ensure that the models were computationally tractable across a sensible timeframe, we took a random-stratified sample of points across forest ages. For each possible value of forest age between 3 and 34 years we took a random sample of 10,000 points. For each point we then identified values for a suite of 32 spatially explicit predictors that could influence regenerating forest age and deforestation pressure, covering locational factors (see Supplementary Fig. 6 for a model without spatial coordinates), landscape level forest characteristics, environmental conditions and human-related deforestation pressures (see the Supplementary Information for more information). We removed any variables that were highly correlated (R > 0.9). This process was repeated for each tropical region (Americas, Africa and Asia-Pacific), and we fit a separate random forest model for each region.

Modelling was performed using the 'ranger' implementation of random forests<sup>73</sup> and trained using the 'caret' package<sup>74</sup>. We used 75% of the data for model training, and held 25% back for model testing. To ensure that we fit the most accurate model possible, we used 'caret' to tune important model hyperparameters to identify the most effective model. We tested a range of values for both 'ntree' (100-1500) and 'mtry' (3-29) under sevenfold cross validation, and selected the combination of 'ntree' and 'mtry' that delivered the highest level of accuracy in an acceptable timeframe for computation. The final model therefore used 500 trees and 18 'mtry' values. We calculated  $R^2$  scores on the basis of the predictive accuracy of the model on the held-back test data. To estimate the importance of each variable in predicting regenerating forest age, we used the corrected impurity measure which estimates mean decrease in accuracy and is unbiased in terms of the number of categories and category frequencies75,76, and created partial dependence plots for each variable using the *Dalex* package<sup>77</sup>.

#### **Reporting summary**

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

#### Data availability

The underlying TMF dataset used in this study is freely accessible and available via EU Science Hub at https://forobs.jrc.ec.europa.eu/TMF/ data. Additional data sources used for random forest modelling are also freely accessible and available for download in the links provided

in the Supplementary Information. The final map of regenerating tropical moist forest age for the year 2023 is available via Zenodo at https://doi.org/10.5281/zenodo.15120870 (ref. 78).

#### **Code availability**

The code used to generate the results are available via Zenodo at https://doi.org/10.5281/zenodo.15120870 (ref. 78).

#### References

- 1. Mo, L. et al. Integrated global assessment of the natural forest carbon potential. *Nature* **624**, 92–101 (2023).
- 2. Strassburg, B. B. N. et al. Global priority areas for ecosystem restoration. *Nature* **586**, 724–729 (2020).
- Lewis, S. L., Wheeler, C. E., Mitchard, E. T. A. & Koch, A. Restoring natural forests is the best way to remove atmospheric carbon. *Nature* 568, 25–28 (2019).
- 4. Press Release: New UN Decade on Ecosystem Restoration Offers Unparalleled Opportunity for Job Creation, Food Security and Addressing Climate Change (United Nations Environment Program, 2019).
- 5. First Draft of the Post-2020 Global Biodiversity Framework (CBD, 2021).
- 6. The Bonn Challenge (IUCN, 2020).
- 7. Cook-Patton, S. C. et al. Mapping carbon accumulation potential from global natural forest regrowth. *Nature* **585**, 545–550 (2020).
- Busch, J. et al. Cost-effectiveness of natural forest regeneration and plantations for climate mitigation. *Nat. Clim. Chang.* 14, 996–1002 (2024).
- Fagan, M. E., Reid, J. L., Holland, M. B., Drew, J. G. & Zahawi, R. A. How feasible are global forest restoration commitments? Conserv. Lett. 13, e12700 (2020).
- Hua, F. et al. The biodiversity and ecosystem service contributions and trade-offs of forest restoration approaches. Science 376, 839–844 (2022).
- 11. Poorter, L. et al. Biomass resilience of Neotropical secondary forests. *Nature* **530**, 211–214 (2016).
- 12. Rozendaal, D. M. A. et al. Biodiversity recovery of Neotropical secondary forests. *Sci. Adv.* **5**, eaau3114 (2019).
- Martin, P. A., Newton, A. C. & Bullock, J. M. Carbon pools recover more quickly than plant biodiversity in tropical secondary forests. *Proc. R. Soc. B* 280, 20132236 (2013).
- Edwards, D. P., Massam, M. R., Haugaasen, T. & Gilroy, J. J. Tropical secondary forest regeneration conserves high levels of avian phylogenetic diversity. *Biol. Conserv.* 209, 432–439 (2017).
- Matos, F. A. R. et al. Secondary forest fragments offer important carbon and biodiversity cobenefits. *Glob. Change Biol.* 26, 509–522 (2020).
- Williams, B. A. et al. Global potential for natural regeneration in deforested tropical regions. *Nature* https://doi.org/10.1038/ s41586-024-08106-4 (2024).
- Smith, C. C. et al. Secondary forests offset less than 10% of deforestation-mediated carbon emissions in the Brazilian Amazon. *Glob. Change Biol.* 26, 7006–7020 (2020).
- Silva Junior, C. H. L. et al. Benchmark maps of 33 years of secondary forest age for Brazil. Sci. Data 7, 269 (2020).
- Schwartz, N. B., Uriarte, M., DeFries, R., Gutierrez-Velez, V. H. & Pinedo-Vasquez, M. A. Land-use dynamics influence estimates of carbon sequestration potential in tropical second-growth forest. *Environ. Res. Lett.* 12, 074023 (2017).
- Reid, J. L., Fagan, M. E., Lucas, J., Slaughter, J. & Zahawi, R. A. The ephemerality of secondary forests in southern Costa Rica. *Conserv. Lett.* 12, e12607 (2019).
- 21. Piffer, P. R. et al. Ephemeral forest regeneration limits carbon sequestration potential in the Brazilian Atlantic Forest. *Glob. Change Biol.* **28**, 630–643 (2022).

- 22. Vancutsem, C. et al. Long-term (1990–2019) monitoring of forest cover changes in the humid tropics. *Sci. Adv.* **7**, eabe1603 (2021).
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., Da Fonseca, G. A. B. & Kent, J. Biodiversity hotspots for conservation priorities. *Nature* 403, 853–858 (2000).
- 24. Brancalion, P. H. S. et al. Global restoration opportunities in tropical rainforest landscapes. *Sci. Adv.* **5**, eaav3223 (2019).
- 25. Roy, E. D. et al. The phosphorus cost of agricultural intensification in the tropics. *Nat. Plants* **2**, 16043 (2016).
- 26. Montgomery, M. R. Theurban transformation of the developing world. Science **319**, 761–764 (2008).
- 27. Aide, T. M. et al. Woody vegetation dynamics in the tropical and subtropical Andes from 2001 to 2014: satellite image interpretation and expert validation. *Glob. Change Biol.* **25**, 2112–2126 (2019).
- 28. Estoque, R. C. et al. Spatiotemporal pattern of global forest change over the past 60 years and the forest transition theory. *Environ. Res. Lett.* **17**, 084022 (2022).
- Fagan, M. E. et al. Land cover dynamics following a deforestation ban in northern Costa Rica. *Environ. Res. Lett.* 8, 034017 (2013).
- Curtis, P. G., Slay, C. M., Harris, N. L., Tyukavina, A. & Hansen, M. C. Classifying drivers of global forest loss. *Science* **361**, 1108–1111 (2018).
- 31. Heinimann, A. et al. A global view of shifting cultivation: recent, current, and future extent. *PLoS One* **12**, e0184479 (2017).
- Borah, J. R., Gilroy, J. J., Evans, K. L. & Edwards, D. P. The value of shifting cultivation for biodiversity in Northeast India. *Divers. Distrib.* 28, 1979–1992 (2022).
- 33. Karp, D. S. et al. Intensive agriculture erodes  $\beta$ -diversity at large scales. *Ecol. Lett.* **15**, 963–970 (2012).
- Silva, J. M. N., Carreiras, J. M. B., Rosa, I. & Pereira, J. M. C. Greenhouse gas emissions from shifting cultivation in the tropics, including uncertainty and sensitivity analysis. *J. Geophys. Res.* 116, D20304 (2011).
- 35. Edwards, D. P. et al. Conservation of tropical forests in the anthropocene. *Curr. Biol.* **29**, R1008–R1020 (2019).
- 36. Grantham, H. S. et al. Anthropogenic modification of forests means only 40% of remaining forests have high ecosystem integrity. *Nat. Commun.* **11**, 5978 (2020).
- 37. Crouzeilles, R. et al. Ecological restoration success is higher for natural regeneration than for active restoration in tropical forests. *Sci. Adv.* **3**, e1701345 (2017).
- Baragwanath, K., Bayi, E. & Shinde, N. Collective property rights lead to secondary forest growth in the Brazilian Amazon. Proc. Natl Acad. Sci. USA 120, e2221346120 (2023).
- Chazdon, R. L. et al. The intervention continuum in restoration ecology: rethinking the active–passive dichotomy. *Restor. Ecol.* 32, e13535 (2024).
- 40. Fagan, M. E. et al. The expansion of tree plantations across tropical biomes. *Nat. Sustain.* **5**, 681–688 (2022).
- Descals, A. et al. High-resolution global map of smallholder and industrial closed-canopy oil palm plantations. *Earth Syst. Sci. Data* 13, 1211–1231 (2021).
- 42. Lesiv, M. et al. Global forest management data for 2015 at a 100 m resolution. *Sci. Data* **9**, 199 (2022).
- 43. Cook, R. L., Binkley, D. & Stape, J. L. Eucalyptus plantation effects on soil carbon after 20 years and three rotations in Brazil. *For. Ecol. Manag.* **359**, 92–98 (2016).
- 44. Nambiar, E. K. S. & Harwood, C. E. Productivity of acacia and eucalypt plantations in Southeast Asia. 1. Bio-physical determinants of production: opportunities and challenges. *Int. For. Rev.* **16**, 225–248 (2014).

- 45. Brown, H. C. A., Berninger, F. A., Larjavaara, M. & Appiah, M. Above-ground carbon stocks and timber value of old timber plantations, secondary and primary forests in southern Ghana. *For. Ecol. Manag.* **472**, 118236 (2020).
- 46. Souza, C. M. et al. Reconstructing three decades of land use and land cover changes in Brazilian biomes with Landsat Archive and Earth Engine. *Remote Sens.* **12**, 2735 (2020).
- Collection 9 of the Annual Land Use Land Cover Maps of Brazil. (MapBiomas Project, accessed 28 February, 2025); https://doi.org/ 10.58053/MapBiomas/XXUKA8
- Nunes, S., Oliveira, L., Siqueira, J., Morton, D. C. & Souza, C. M. Unmasking secondary vegetation dynamics in the Brazilian Amazon. *Environ. Res. Lett.* 15, 034057 (2020).
- Chazdon, R. L. & Guariguata, M. R. Natural regeneration as a tool for large-scale forest restoration in the tropics: prospects and challenges. *Biotropica* 48, 716–730 (2016).
- Morton, O., Borah, J. R. & Edwards, D. P. Economically viable forest restoration in shifting cultivation landscapes. *Environ. Res. Lett.* 15, 064017 (2020).
- Evans, M. C. et al. Carbon farming via assisted natural regeneration as a cost-effective mechanism for restoring biodiversity in agricultural landscapes. *Environ. Sci. Policy* 50, 114–129 (2015).
- 52. Crouzeilles, R. et al. Achieving cost-effective landscape-scale forest restoration through targeted natural regeneration. *Conserv. Lett.* **13**, e12709 (2020).
- 53. Brancalion, P. H. S. et al. A call to develop carbon credits for second-growth forests. *Nat. Ecol. Evol.* **8**, 179–180 (2024).
- 54. Rosa, M. R. et al. Hidden destruction of older forests threatens Brazil's Atlantic Forest and challenges restoration programs. *Sci. Adv.* **7**, eabc4547 (2021).
- Nanni, A. S. et al. The neotropical reforestation hotspots: a biophysical and socioeconomic typology of contemporary forest expansion. *Glob. Environ. Change* 54, 148–159 (2019).
- Vieira, I., Gardner, T., Ferreira, J., Lees, A. & Barlow, J. Challenges of governing second-growth forests: a case study from the Brazilian Amazonian state of Pará. *Forests* 5, 1737–1752 (2014).
- 57. Chazdon, R. L. et al. Fostering natural forest regeneration on former agricultural land through economic and policy interventions. *Environ. Res. Lett.* **15**, 043002 (2020).
- Wang, Y. et al. Upturn in secondary forest clearing buffers primary forest loss in the Brazilian Amazon. *Nat. Sustain.* 3, 290–295 (2020).
- 59. Di Sacco, A. et al. Ten golden rules for reforestation to optimize carbon sequestration, biodiversity recovery and livelihood benefits. *Glob. Change Biol.* **27**, 1328–1348 (2021).
- 60. Erbaugh, J. T. et al. Global forest restoration and the importance of prioritizing local communities. *Nat. Ecol. Evol.* **4**, 1472–1476 (2020).
- Sze, J. S., Carrasco, L. R., Childs, D. & Edwards, D. P. Reduced deforestation and degradation in indigenous lands pan-tropically. *Nat. Sustain.* 5, 123–130 (2021).
- Pelletier, J., Ngoma, H., Mason, N. M. & Barrett, C. B. Does smallholder maize intensification reduce deforestation? Evidence from Zambia. *Glob. Environ. Change* 63, 102127 (2020).
- Forero-Montaña, J., Marcano-Vega, H., Zimmerman, J. K. & Brandeis, T. J. Potential of second-growth Neotropical forests for forestry: the example of Puerto Rico. *For. Trees Livelihoods* 28, 126–141 (2019).
- 64. Mauser, W. et al. Global biomass production potentials exceed expected future demand without the need for cropland expansion. *Nat. Commun.* **6**, 8946 (2015).

- Foley, J. A. et al. Solutions for a cultivated planet. *Nature* **478**, 337–342 (2011).
- Heinrich, V. H. A. et al. The carbon sink of secondary and degraded humid tropical forests. *Nature* 615, 436–442 (2023).
- 67. Gorelick, N. et al. Google Earth Engine: planetary-scale geospatial analysis for everyone. *Remote Sens. Environ.* **202**, 18–27 (2017).
- R Core Team. R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing, 2013); http://www.R-project.org/
- 69. Hijmans, R. J. terra: spatial data analysis. R version 1.8–42 https://doi.org/10.32614/CRAN.package.terra (2020).
- 70. GADM. GADM database of global administrative areas. version 4.1. https://gadm.org/data.html (2022).
- 71. Massicotte, P. & South, A. rnaturalearth: world map data from Natural Earth. R version 1.0.1 https://doi.org/10.32614/CRAN. package.rnaturalearth (2017).
- 72. Breiman, L. Random forests. Mach. Learn. 45, 5–32 (2001).
- 73. Wright, M. N. & Ziegler, A. ranger: a fast implementation of random forests for high dimensional data in C++ and R. J. Stat. Soft. https://doi.org/10.18637/jss.v077.i01 (2017).
- 74. Kuhn, M. Building predictive models in R using the caret package. J. Stat. Soft. https://doi.org/10.18637/jss.v028.i05 (2008).
- 75. Greenwell, B. M. & Boehmke, B. C. Variable importance plots—an introduction to the vip package. *R J.* **12**, 343–366 (2020).
- 76. Nembrini, S., König, I. R. & Wright, M. N. The revival of the Gini importance? *Bioinformatics* **34**, 3711–3718 (2018).
- 77. Biecek, P. DALEX: explainers for complex predictive models in R. J. Mach. Learn. Res. **19**, 1–5 (2018).
- Bousfield, C. Data and code: the pan-tropical age distribution of regenerating tropical moist forest. *Zenodo* https://doi.org/ 10.5281/ZENODO.15120870 (2025).

#### **Author contributions**

C.G.B. and D.P.E. conceived the study. C.G.B. collected data, performed the analysis and drafted the initial paper. C.G.B. and D.P.E. edited the paper.

#### **Competing interests**

The authors declare no competing interests.

#### **Additional information**

**Supplementary information** The online version contains supplementary material available at https://doi.org/10.1038/s41559-025-02721-8.

**Correspondence and requests for materials** should be addressed to Christopher G. Bousfield.

**Peer review information** *Nature Ecology & Evolution* thanks Robin Chazdon and Edward Mitchard for their contribution to the peer review of this work.

**Reprints and permissions information** is available at www.nature.com/reprints.

**Publisher's note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate

#### Article

if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

© The Author(s) 2025

## nature portfolio

Corresponding author(s): Christopher Bousfield

Last updated by author(s): Apr 4, 2025

## **Reporting Summary**

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our <u>Editorial Policies</u> and the <u>Editorial Policy Checklist</u>.

#### **Statistics**

For	all st	atistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.
n/a	Cor	firmed
$\boxtimes$		The exact sample size $(n)$ for each experimental group/condition, given as a discrete number and unit of measurement
	$\square$	A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
	$\boxtimes$	The statistical test(s) used AND whether they are one- or two-sided Only common tests should be described solely by name; describe more complex techniques in the Methods section.
	$\boxtimes$	A description of all covariates tested
$\boxtimes$		A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
		A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
$\boxtimes$		For null hypothesis testing, the test statistic (e.g. F, t, r) with confidence intervals, effect sizes, degrees of freedom and P value noted Give P values as exact values whenever suitable.
$\boxtimes$		For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
$\boxtimes$		For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
$\boxtimes$		Estimates of effect sizes (e.g. Cohen's d, Pearson's r), indicating how they were calculated
		Our web collection on <u>statistics for biologists</u> contains articles on many of the points above.

#### Software and code

Policy information about availability of computer code						
Data collection	The code used to generate the results are available at https://doi.org/10.5281/zenodo.1512087078					
	The analysis was done using a combination of Google Earth Engine and RStudio (4.4)					
Data analysis	The code used to generate the results are available at https://doi.org/10.5281/zenodo.1512087078					
	The analysis was done using a combination of Google Earth Engine and RStudio (4.4)					

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio guidelines for submitting code & software for further information.

#### Data

#### Policy information about availability of data

All manuscripts must include a <u>data availability statement</u>. This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our policy

The underlying TMF dataset used in this study is freely accessible and available for download at https://forobs.jrc.ec.europa.eu/TMF/data. Additional data sources used for random forest modelling are also freely accessible and available for download in the links provided in the supplementary material. The final map of regenerating tropical moist forest age for the year 2023 is available for download at https://doi.org/10.5281/zenodo.1512087078

#### Research involving human participants, their data, or biological material

Policy information about studies with human participants or human data. See also policy information about sex, gender (identity/presentation), and sexual orientation and race, ethnicity and racism.

Reporting on sex and gender	NA
Reporting on race, ethnicity, or other socially relevant groupings	NA
Population characteristics	NA
Recruitment	NA
Ethics oversight	NA

Note that full information on the approval of the study protocol must also be provided in the manuscript.

### Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

Life sciences

Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see <u>nature.com/documents/nr-reporting-summary-flat.pdf</u>

No No

Yes

## Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	Open source remotely-sensed forest cover datasets were combined to produce a pan-tropical map of regenerating tropical forest age. Random forest models were also used to assess the importance of 32 predictors in predicting regenerating forest age.
Research sample	Overall dataset is based on all 2023 detections of 'regrowth forest' by the JRC TMF transition map. Sample for the random forest modeling is a randomly selected sample of 10,000 points for each potential age (3-34 years), which is aimed to represent the entire regenerating forest extent.
Sampling strategy	Random sample of 10,000 taken for each age group (3-34 years)
Data collection	NA
Timing and spatial scale	JRC TMF Annual change dataset covers the years 1990-2023, and covers the entire extent of tropical moist forest
Data exclusions	No data were excluded from the analysis
Reproducibility	Random forest models were rerun to check consistent fit
Randomization	Separate models were fit for each key tropical region (Americas, Africa, Asia-Pacific). Samples for models were taken randomly.
Blinding	Blinding was not relevant in this study.

Did the study involve field work?

## Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems	Methods	
n/a Involved in the study	n/a Involved in the study	
Antibodies	ChIP-seq	
Eukaryotic cell lines	Flow cytometry	
Palaeontology and archaeology	MRI-based neuroimaging	
Animals and other organisms	·	
🔀 🔲 Clinical data		
Dual use research of concern		
Plants		

#### Plants

Seed stocks	NA
Novel plant genotypes	NA
Noter plane generypes	
Authentication	NA