



Contents lists available at [ScienceDirect](#)

## Arthroplasty Today

journal homepage: <http://www.arthroplastytoday.org/>



### Corrigendum

## Corrigendum to “Revision Rates for Aseptic Loosening in the Obese Patient: A Comparison Between Stemmed, Uncemented, and Unstemmed Tibial Total Knee Arthroplasty Components” [Arthroplasty Today 32 (2025) 101,621]

Bernard H. van Duren, Amy M. Firth\*, Reshid Berber, Hosam E. Matar, Peter J. James, Benjamin V. Bloch

*Nottingham Elective Orthopaedic Services, Nottingham University Hospitals NHS Trust, Nottingham, United Kingdom*

The authors regret errors in the published article that should be corrected as follows.

In the “Methods” section of the article Abstract, the third sentence should read:

“Those who underwent cemented TKA using a tibial stem extension (Group 1) (n = 150) and those where cementless implants were used (Group 2) (n = 101) were compared to a control group (n = 1261) using a standard cemented implant.”

In the “Revisions” section within Results, the second sentence should read:

“Of the 1512 knees performed in patients with BMI in excess of 30, there were 37 all-cause revisions, of which 7 were for aseptic loosening (2 tibial, 1 femoral, and 4 involving both components).”

Finally, the figures in [Table 1](#) were not correctly adjusted. The full and correct [Table 1](#) is given here.

The authors would like to apologise for any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.artd.2025.101621>.

\* Corresponding author. Nottingham Elective Orthopaedics, Nottingham City Hospital, Hucknall Road, Nottingham NG5 1PB, United Kingdom. Tel.: +44 115 969 1169.

E-mail address: [a.firth1@nhs.net](mailto:a.firth1@nhs.net)

**Table 1**  
Overview of demographics preoperative characteristics and implant characteristics.

	Control (cemented primary Attune)		Cementless (Attune primary cementless)		Stemmed (Attune revision tray +50 × 14mm stem)	
Total No.	2443		221		162	
BMI > 30	1261	52%	101	46%	150	93%
BMI > 35	542	22%	37	17%	129	80%
BMI > 40	178	7%	8	4%	79	49%
Included in analysis: BMI > 30						
Total Prosthesis Years	8575		540		765	
Mean Follow Up (SD) (years)	6.8	5.3		5.1		
Range (years)	0-12.7		2.6-7.9		0-7.2	
Median (IQR) (years)	6.9 (5.1-8.6)		5.5 (4.2-6.6)		5.4 (4.6-6.2)	
Female	806	64%	41	41%	127	85%
Age (years)						
<65	437	35%	61	60%	82	55%
65-74	507	40%	33	33%	46	31%
≥75	317	25%	7	7%	22	15%
ASA						
I	55	4%	8	8%	2	1%
II	874	69%	83	82%	67	45%
III	325	26%	10	10%	80	53%
IV	7	1%	0	0%	1	1%
V	0	0%	0	0%	0	0%
Indication						
OA	1231	98%	99	98%	145	97%
Inflammatory Arthropathy	18	1%	1	1%	3	2%
AVN	2	0%	0	0%	0	0%
Trauma	8	1%	0	0%	1	1%
Other	2	0%	1	1%	1	1%
Cement						
Palacos R + G	378	30%	n/a		4	3%
Smartset GHV	883	70%	n/a		146	97%
Meniscal Constraint						
CR Mobile	755	60%	82	81%	10	7%
CR Fixed	387	31%	0	0%	70	47%
PS mobile	94	7%	19	19%	5	3%
PS Fixed	25	2%	0	0%	65	43%

ASA, American Society of Anesthesiologists; BMI, body mass index; CR, cruciate retaining; IQR, interquartile range; n/a, not applicable; PS, posterior stabilized; SD, standard deviation; AVN, avascular necrosis.