



This is a repository copy of *Search for long-lived charged particles using large specific ionisation loss and time of flight in 140 fb⁻¹ of pp collisions at = 13 TeV with the ATLAS detector*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/229454/>

Version: Published Version

Article:

Aad, G. orcid.org/0000-0002-6665-4934, Aakvaag, E. orcid.org/0000-0001-7616-1554, Abbott, B. orcid.org/0000-0002-5888-2734 et al. (2912 more authors) (2025) Search for long-lived charged particles using large specific ionisation loss and time of flight in 140 fb⁻¹ of pp collisions at = 13 TeV with the ATLAS detector. *Journal of High Energy Physics*, 2025 (7). 140. ISSN 1126-6708

[https://doi.org/10.1007/jhep07\(2025\)140](https://doi.org/10.1007/jhep07(2025)140)

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Search for long-lived charged particles using large specific ionisation loss and time of flight in 140 fb^{-1} of pp collisions at $\sqrt{s} = 13\text{ TeV}$ with the ATLAS detector



The ATLAS collaboration

E-mail: atlas.publications@cern.ch

ABSTRACT: This paper presents a search for massive, charged, long-lived particles with the ATLAS detector at the Large Hadron Collider using an integrated luminosity of 140 fb^{-1} of proton-proton collisions at $\sqrt{s} = 13\text{ TeV}$. These particles are expected to move significantly slower than the speed of light. In this paper, two signal regions provide complementary sensitivity. In one region, events are selected with at least one charged-particle track with high transverse momentum, large specific ionisation measured in the pixel detector, and time of flight to the hadronic calorimeter inconsistent with the speed of light. In the other region, events are selected with at least two tracks of opposite charge which both have a high transverse momentum and an anomalously large specific ionisation. The search is sensitive to particles with lifetimes greater than about 3 ns with masses ranging from 200 GeV to 3 TeV . The results are interpreted to set constraints on the supersymmetric pair production of long-lived R-hadrons, charginos and staus, with mass limits extending beyond those from previous searches in broad ranges of lifetime.

KEYWORDS: Beyond Standard Model, Hadron-Hadron Scattering, Supersymmetry

ARXIV EPRINT: [2502.06694](https://arxiv.org/abs/2502.06694)

Contents

1	Introduction	1
2	ATLAS detector	3
3	Data and Monte Carlo samples	5
4	Calibration	7
4.1	$\beta\gamma$ from ionisation energy loss	7
4.2	β from time of flight	8
5	Analysis	10
5.1	Overview	10
5.2	Event selection	11
5.3	Background estimation	14
5.4	Mass window definition	19
5.5	Uncertainties	22
6	Results	25
6.1	β -search results	25
6.2	Di-track search results	26
6.3	Lifetime-dependent mass limits	28
7	Conclusion	31
	The ATLAS collaboration	39

1 Introduction

A wide range of physics models that extend the Standard Model (SM) predict the existence of new, massive, charged long-lived particles (LLPs). These particles appear in certain dark matter models [1, 2] and in proposed solutions to the gauge hierarchy problem, including supersymmetric (SUSY) models [3–8] that either violate or conserve R -parity.¹ Particles can acquire macroscopic lifetimes in models of new physics via the same mechanisms that generate long-lived SM particles, with lifetimes depending on the mass hierarchies between new particles and/or the size of a new coupling.

A search is presented for particles that are massive, long-lived, and charged using 140 fb^{-1} of proton-proton collision data from the ATLAS experiment at the Large Hadron Collider (LHC) [9]. This analysis looks for a direct interaction of the LLPs with the ATLAS detector using the measurement of the ionisation energy loss (dE/dx) in the pixel detector and the

¹ R -parity is a quantum number defined as $(-1)^{3(B-L)+2S}$ where S is the particle spin and L and B are, respectively, its lepton and baryon number.




Figure 1. Representative production diagrams for (a) pair-produced gluinos which form R -hadrons decaying into neutralinos, (b) pair-produced charginos decaying into neutralinos, and (c) pair-produced staus decaying into gravitinos. The anti-particle labels are suppressed for simplicity.

time of flight (ToF) measured by the hadronic calorimeter. This search is, to first order, independent of the LLP decay mode and is therefore sensitive to many different models of new physics. The analysis is optimised for and interpreted in the context of several different long-lived, pair-produced, supersymmetric particles. Long-lived gluinos appear in models including mini-split SUSY [10, 11]; once produced, the gluinos hadronise with Standard Model quarks to produce R -hadrons [12]. Long-lived charginos are motivated by anomaly mediated supersymmetry-breaking (AMSB) models [13, 14]. Long-lived staus emerge in both co-annihilation dark matter models [15–17] and in gauge-mediated supersymmetry-breaking (GMSB) models [18–20]. Representative production and decay diagrams of the processes targeted by this search are shown in figure 1.

This search is designed to extend the reach of a previous paper [21], in which the primary signal selection requirement was an isolated, high-momentum track with large dE/dx . Two complementary search strategies extend the sensitivity with respect to the previous result for different signal topologies.

One analysis region (β -search) extends the sensitivity to higher masses for models with one or more charged, heavy LLPs by requiring that the tracks that pass signal candidate selections have both a large dE/dx and a ToF measurement consistent with a slow-moving particle. The additional ToF requirement reduces the contribution of background processes and therefore improves the sensitivity to models in which a large dE/dx in the pixel detector arises from a heavy, slow-moving particle, with a lifetime greater than about 10 ns. This analysis specifically targets heavy LLPs, including long-lived charginos and R -hadrons. By explicitly requiring $\beta \lesssim 0.8$, this channel is insensitive to models that predict large dE/dx from relativistic LLPs with an electric charge greater than one [22]. ATLAS reported on a search for long-lived multi-charged particles in ref. [23].

The second analysis region (di-track search) requires two signal tracks which both have significant dE/dx . As the requirement of a second signal track significantly reduces backgrounds, other selection requirements can be relaxed to enhance the signal significance, in particular for relatively light LLPs, which have a more modest ionisation signature. As this region is only sensitive to signatures with two charged LLPs, it does not add sensitivity for R -hadrons (which hadronise into a mix of charged and neutral states), nor for charginos produced in association with a neutralino. However, the large reduction in background and

enhanced acceptance for low-mass LLPs yields significant sensitivity gains for pair-produced sleptons with lifetimes greater than about 3 ns.

For both signal regions, the main observable used is the mass of the particle associated with the selected track(s). The candidate LLP mass is calculated directly via the relation $m \equiv p/\beta\gamma$ with two independent determinations of $\beta\gamma$, and p measured using the track curvature in the central magnetic field of ATLAS. In both regions, $\beta\gamma_{dE/dx}$ is extracted from a parameterisation of the Bethe-Bloch relationship between $\beta\gamma$ and dE/dx [24]. The di-track search obtains two independent $\beta\gamma_{dE/dx}$ measurements per event, one per track, while the β -search obtains both $\beta\gamma_{dE/dx}$ and $\beta\gamma_{\text{ToF}}$ for the same track. The $\beta\gamma_{\text{ToF}}$ is calculated from the ToF measured by a cluster of cells in the ATLAS calorimeter crossed by the candidate track and their distance from the proton-proton collision. Compatibility between the two mass measurements is finally required to maximise sensitivity.

This analysis uses the full Run 2 data sample and is an update of several previous searches performed by the ATLAS experiment in both Run 1 and Run 2 [21, 24–27]. The CMS experiment has also used a combination of dE/dx and ToF in previous searches [28–31]. ATLAS observed a 3.3 (3.6) global (local) Z significance excess at 1.4 TeV using only the pixel dE/dx , in 140 fb^{-1} of Run 2 collisions [21]. As reported in ref. [21], preliminary, uncalibrated ToF measurements of the calorimeter and muon systems for the tracks in the excess were not compatible with the hypothesis of slow massive particles. The search presented in this paper follows up on this excess by calibrating the ToF measurement in the calorimeter and by designing a signal region with enhanced sensitivity to heavy, charged, and slow LLPs with unit charge, while additionally extending the sensitivity to sleptons with moderate lifetimes with a new di-track region.

2 ATLAS detector

The ATLAS detector [32] is a general-purpose detector with a forward-backward-symmetric cylindrical layout² covering nearly 4π in solid angle. It consists of an inner detector (ID) tracking system which measures the trajectories of charged particles, surrounded by a 2 T solenoid, followed by calorimeters which measure the energy of particles that interact electromagnetically or hadronically, and a muon spectrometer (MS) inside toroidal magnets which provide additional tracking for muons. The detector is hermetic within its η acceptance and can therefore measure the missing transverse momentum ($\vec{p}_T^{\text{miss}}$, with magnitude E_T^{miss}) associated with each event. A two-level trigger system is used to select events [33]. The first-level trigger is hardware-based and uses a subset of detector information to accept events, produced by LHC at 40 MHz bunch crossing, at a rate below 100 kHz, which is the maximum detector readout rate. This is followed by a software-based high-level trigger, which runs calibration and prompt reconstruction algorithms, reducing the event recording rate to about 1 kHz. The events are eventually processed offline and reconstructed by making

²ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point in the centre of the detector and the z -axis coinciding with the axis of the beam pipe. The x -axis points from the interaction point to the centre of the LHC ring, and the y -axis points upward. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z -axis. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$.

use of a software suite [34], which also provides tools for data simulation, analysis, detector operations, trigger and data acquisition.

Two detectors, the pixel and calorimeter subsystems, are used to measure the $\beta\gamma$ of charged particles and are therefore described in more detail. The pixel detector [35–37] covers the innermost region of the ID and provides, on average, four precision measurements for each track in the region $|\eta| < 2.5$ at radial distances of 3.4 cm to 13 cm from the LHC beam line. These measurements determine both the track parameters and the charge released in each pixel. The charge is measured by digitising the time interval with the signal above a preset threshold (time-over-threshold or ToT), which is approximately proportional to the ionisation charge [38]. Compared with the other layers that can measure a charge corresponding to 10 minimum ionising particles (MIPs), the innermost pixel layer, called Insertable B-Layer (IBL) [36, 37] provides charge measurements with lower resolution and dynamic range. If the charge released in a pixel exceeds the IBL dynamic range (which is set at approximately two MIPs) an overflow bit is set. The charge released by a track crossing a layer of the pixel detector is rarely contained within just one pixel; neighbouring pixels registering hits are joined together using a connected component analysis [39] to form clusters. The charge of a cluster is calculated by summing the charges of all pixels belonging to the cluster. The dE/dx measurement assigned to each track is then calculated by averaging the ionisation measurements (charge collected in the cluster per unit track length in the sensor) of its individual clusters. To reduce the effect of the tails of the individual ionisation measurements on this distribution, a truncated average ($\langle dE/dx \rangle_{\text{trunc}}$) is evaluated after removing the highest dE/dx cluster, or the two highest dE/dx clusters in the rare case of more than four pixel clusters on a track. Clusters including pixels at the sensor edges and clusters with overflow ionisation in the IBL are excluded from the $\langle dE/dx \rangle_{\text{trunc}}$ calculation, as full charge detection is not guaranteed for these clusters. A track is considered for this analysis if the $\langle dE/dx \rangle_{\text{trunc}}$ is calculated using at least two clusters after removal of those meeting the criteria defined above. The average number of clusters used for the $\langle dE/dx \rangle_{\text{trunc}}$ calculation is approximately 2.7 per track. The $\langle dE/dx \rangle_{\text{trunc}}$ is then corrected for variations of the pixel detector conditions during the data-taking period (e.g. charge losses due to radiation damage) and for the residual η -dependence, as described in section 5.1. The output is the variable used in the signal selection for the search, and is further referred to simply as dE/dx . Like the *restricted energy loss* [40], this variable rejects high ionisation deposits.

Neither of these variables show a logarithmic rise at high values of $\beta\gamma$ nor sensitivity to radiative effects, which is expected based on the performance of the restricted energy loss in thin silicon sensors [40] and confirmed for the specific calculation of dE/dx in the ATLAS pixel detector with dedicated samples of electrons and muons from $Z \rightarrow ee$ and $Z \rightarrow \mu\mu$ events selected in data. The $\beta\gamma$ of a particle can be calculated from the dE/dx of its track using the Bethe-Bloch formula. A meaningful $\beta\gamma$ value can only be estimated in the range of $0.3 \lesssim \beta\gamma \lesssim 0.9$ using the pixel detector. The lower limit is a consequence of the ToT dynamic range, while the upper limit is due to the proximity of the MIP regime which begins at $\beta\gamma \approx 3$ and where dE/dx becomes quasi-independent of $\beta\gamma$.

A silicon microstrip track detector (SCT) [41] surrounds the pixel detector and contributes to the definition of an accepted track, which must reach a 45 cm radial distance from the colliding beams.




Figure 2. Schematic showing the TileCal cell layout and $|\eta|$ acceptance. The red dashed lines indicate where tracks from the origin with a given pseudorapidity will cross the calorimeter, the calorimeter cells A, (B, BC) and D belong to layers at increasing radius and all contribute to the ToF measurement. The special E-cells are not used in the analysis because their time resolution is poor. The calorimeter response is worse in the region $0.8 < |\eta| < 1.0$ (transition region between the barrel and the extended barrel).

The ATLAS calorimeter system is composed of two parts optimised to measure the energy of the particles interacting electromagnetically or hadronically. As the target LLPs are not expected to shower in the electromagnetic calorimeter and the timing resolution of the electromagnetic calorimeter decreases for small energy deposits, the ToF measurement is done with the central hadronic calorimeter (TileCal) [42] and uses, as a reference, the beam crossing time signal provided by the LHC. The TileCal is a barrel-shaped sampling device (made of steel plates acting as absorber and scintillator tiles as active medium) extending from a radius of 228 cm to 386.5 cm and covering the range $|\eta| < 1.6$ as shown in figure 2.

Wavelength-shifting fibres collect the light from scintillators and carry it to the photomultiplier tubes (PMTs). The analogue signals from the PMTs are amplified, shaped and digitised by sampling the signal every 25 ns and stored on detector until a trigger decision is received. The front-end electronics read out the signals produced by approximately 5000 cells organised into three radial layers. The tile calorimeter cells have a good signal time resolution (better than 1.0 ns (3.7 ns) when more than 5 GeV (500 MeV) are released in one cell) and can therefore determine β_{ToF} through a ToF measurement. Each calorimeter cell along the particle track contributes to the β_{ToF} measurement through a weighted average. This takes into account both the cell distance from the interaction point and the energy released in each cell, as the cell time resolution depends on the deposited energy [43].

3 Data and Monte Carlo samples

The analysis is conducted with 140 fb^{-1} of pp collision data that satisfy the ATLAS data quality requirements [44]. The data sample was taken during Run 2 of the LHC from 2015 to 2018, at a centre-of-mass energy of 13 TeV. The average number of collisions per bunch-crossing (pile-up, $\langle \mu \rangle$) is approximately 34. A dedicated sample of 21 pb^{-1} of low-pile-up data with $\langle \mu \rangle \sim 0.4$ taken in 2017 is used for the dE/dx -to- $\beta\gamma$ calibration. In this data sample,

tracks are reconstructed if they have a transverse momentum $p_T > 100$ MeV (while the minimum p_T requirement in the standard data sample is 500 MeV) allowing the measurement of low- $\beta\gamma$ pions, kaons and protons. Since LLPs are expected to behave similarly to muons in the calorimeter, data and high-statistic Monte Carlo (MC) samples of $Z \rightarrow \mu\mu$ are used to calibrate the ToF response of the calorimeter.

To optimise the analysis selection, MC samples were produced to simulate events containing long-lived gluinos, charginos, and staus, with lifetimes (τ) from 3 ns to stable, corresponding to the production diagrams shown in figure 1. These three signal models are complementary in this study. The gluino samples have large production cross-sections and are suited to probing the high-mass frontier beyond 2 TeV. The stau samples have production cross-sections that are several orders of magnitude smaller than for gluinos of the same mass, and are suited to probing the few-hundred-GeV mass range. The chargino sample cross-sections have intermediate values and are useful in probing the mass range around 1 TeV.

All signal samples were generated using MADGRAPH5_AMC@NLO 2.6.2 [45] with up to two additional partons at leading order, and interfaced to PYTHIA 8.240 [46] using the A14 set of tuned parameters (‘tune’) [47]. The R -hadron samples were generated with the NNPDF2.3LO [48] parton distribution function (PDF) set for parton showering and hadronisation, while for the chargino samples the CTEQ6.6 [49] and MSTW2008NLO90CL [50] PDF sets were used, with decays of bottom and charm hadrons performed by EVTGEN 1.6.0 [51]. The CKKW-L merging scheme [52, 53] was applied to combine the matrix element with the parton shower.

Gluino pair production was simulated for gluino masses ranging from 1.4 TeV to 2.4 TeV within a simplified model inspired by a split-SUSY scenario [10, 11]. The long-lived gluino, which carries colour charge, hadronises to form a colourless composite particle called an R -hadron. The details of the R -hadron simulation are given in ref. [54]. Each gluino decays into a stable neutralino and two quarks via a virtual squark at a very high mass scale in an R -parity conserving decay. The gluino acquires a long lifetime as the only decay channel available is via a massive virtual squark. To probe decays with different kinematics, two sets of samples were produced: one with a fixed neutralino mass of $m(\tilde{\chi}_1^0) = 100$ GeV, and the other one with a fixed mass splitting of $\Delta m(\tilde{g}, \tilde{\chi}_1^0) = 30$ GeV. These two series of mass parameters are complementary and illustrate that the search is open to various models as it does not require explicit decay properties of the charged LLP, like the visible mass of the decay. The nominal cross-section values were calculated at next-to-leading-order (NLO) with resummation of next-to-leading logarithms (NLL). Their uncertainties were taken from an envelope of predictions using different PDF sets and factorisation and renormalisation scales [55].

Samples with a combination of chargino-neutralino ($\tilde{\chi}_1^\pm \tilde{\chi}_1^0$) and chargino-chargino ($\tilde{\chi}_1^+ \tilde{\chi}_1^-$) events were generated with nearly degenerate chargino and neutralino masses, motivated by the ‘pure wino’ AMSB scenario [13, 14]. Each long-lived chargino decays into a stable neutralino and a pion, where the mass-splitting between the chargino and neutralino is set to 160 MeV. Although the AMSB model has a specific preference for the chargino’s lifetime ($\mathcal{O}(0.2)$ ns) and mass relation via the loop dynamics [56], this theoretical constraint was artificially loosened for experimental benchmarking, and charginos with higher lifetimes and masses ranging from 0.7 TeV to 1.4 TeV were examined. A 100% branching ratio for

$\tilde{\chi}_1^\pm \rightarrow \pi^\pm \tilde{\chi}_1^0$ is assumed. The production cross-sections are computed at NLO plus NLL precision in the limit of mass-degenerate $\tilde{\chi}_1^+$, $\tilde{\chi}_1^-$, and $\tilde{\chi}_1^0$, and with all the other sparticles assumed to be heavy and decoupled [57, 58].

Events with pair-produced staus, each of which decays into a τ -lepton and a stable gravitino, were produced in a simplified model motivated by the GMSB scenario [59–61]. The stau masses range from 200 GeV to 1 TeV and the small mass of the gravitino is neglected; the long stau lifetime is due to the small coupling to the gravitino. Signal cross-sections were calculated assuming direct $\tilde{\tau}$ production at NLO in α_s , with soft-gluon emission effects added at NLL accuracy, assuming mass-degenerate left- and right-handed staus ($\tilde{\tau}_{L,R}$) with no mixing [57, 58, 62–64].

Inelastic pp interactions were generated using PYTHIA 8.186 [65] and EVTGEN 1.6.0 with the NNPDF2.3LO PDF set and the A3 tune [66]. The inelastic collisions were overlaid onto the hard-scattering process to simulate the effect of multiple pp interactions. MC samples were reweighed to match the distribution of the mean number of interactions per bunch crossing observed in data.

The MC events were passed through a full detector simulation [67] based on GEANT4 [68]. The propagation and decays of charginos and staus were simulated within GEANT4, taking into account ionisation loss and interactions with the detector. The propagation of R -hadrons and their interactions were handled by GEANT4 until their decay, at which point the decay chains and subsequent hadronisation were simulated by PYTHIA 8; the information about the outgoing particles was then transferred back to GEANT4.

4 Calibration

This section describes the calibration of the $\beta\gamma$ measurements provided by the pixel detector and the tile calorimeter.

4.1 $\beta\gamma$ from ionisation energy loss

The most probable value (MPV) of the track $\langle dE/dx \rangle_{\text{trunc}}$ measured by the pixel detector varies as a function of the delivered luminosity and detector region. The radiation dose received, and the consequent charge trapping varies the track $\langle dE/dx \rangle_{\text{trunc}}$ by up to 40% in the data sample. These effects in combination with changing detector operating conditions require a data-derived set of run-by-run and $|\eta|$ -dependent corrections such as to equalise the most-probable value of $\langle dE/dx \rangle_{\text{trunc}}$ as a function of time and η and finally provide the dE/dx value. These corrections are the same as those used in ref. [21] where their detailed description is given. After these corrections are applied, samples of electrons and muons from selected $Z \rightarrow ee$ and $Z \rightarrow \mu\mu$ events in data show that the corrected dE/dx distribution has negligible dependence on the number of concurrent proton-proton collisions in the event.

The method used to associate a $\beta\gamma$ to a dE/dx value is based on the measurement of low-momentum SM particles. The correlation between $\beta\gamma$ and dE/dx is extracted from data by fitting a parameterisation of the Bethe-Bloch relation, with the assumptions that $\beta\gamma < 1$. Reconstructing tracks with momenta ranging from 100 MeV to a few GeV allows to resolve and identify electrons, pions, kaons, protons, and deuterons. The dE/dx spectrum is a superposition of Landau distributions of those particles. Fitting this spectrum extracts

the MPV of each particle for each momentum slice. The mapping of $(\beta\gamma, \text{MPV}_{dE/dx})$ is redundantly obtained for pion, kaon, proton (and deuteron when statistics are sufficient). While the proton data sample tends to cover the lower $\beta\gamma$ range down to $\beta\gamma \gtrsim 0.35$, the pion data sample can cover up to a MIP ($\beta\gamma \approx 3$). The kaon data sample overlaps between the two data samples. The mapping used is the one from ref. [21], where more details of the $\beta\gamma$ calibration process can be found.

The simulation of the dE/dx response of the pixel detector is based on a realistic charge-deposition model [69], but due to the sensitivity of the dE/dx measurement to detector conditions, including radiation damage, the simulated track dE/dx and especially the probability that a track has a hit in the IBL overflow do not describe the data accurately enough for this analysis. Hence, the dE/dx response for simulated events was modelled by replacing the simulated value with values from a data-driven template [21], derived from low-momentum events as a function of $\beta\gamma$.

4.2 β from time of flight

Each TileCal cell provides an independent measurement of β . The β_i measurement of the i -th calorimeter cell is obtained from the time measurement in the cell t_i (such that a particle travelling from the interaction point with the speed of light produces a signal at time $t = 0$, in each calorimeter cell), the distance of the cell's centre from the interaction point l_i and the speed of light c : $\beta_i = 1/(1 + \frac{ct_i}{l_i})$. Only cells with an energy deposition above 500 MeV are considered in order to minimise the effect of the noise contribution.

The final β_{ToF} exploits the average over $1/\beta_i$ values whose uncertainties are similar to a Gaussian distribution. Thus, $1/\beta_{\text{ToF}}$ is obtained as the average over $1/\beta_i$ weighted by $1/\sigma_i^2$, where σ_i is the time resolution in the i -th cell.

The β_{ToF} calibration is obtained in consecutive steps: the time offset correction in each cell, the correction based on the track pseudorapidity with respect to the cell's centre, and the determination of the cell time resolution σ_i . Isolated muons from $Z \rightarrow \mu\mu$ decays are used for this purpose.

The cell time calibration is performed separately for each data-taking year. The core of the time spectrum in each TileCal cell is approximately Gaussian. The mean value of the Gaussian fit in the $\pm 2\sigma$ region around the peak is taken as the calibration constant to be subtracted from the time measurement. A run-by-run correction did not show any effect on the measured β_{ToF} , hence it is not applied. No effect of the energy deposited in a cell on the mean value of the reconstructed time was visible. The reconstructed time in the cells shows a dependence on the distance $\Delta\eta$ of the track's impact point in the cell η_{track} to the cell centre η_{cell} ($\Delta\eta \equiv \eta_{\text{track}} - \eta_{\text{cell}}$). This effect is greater in cells spanning a larger pseudorapidity region. A correction is provided by a linear fit to the mean reconstructed time as a function of $\Delta\eta$ for each cell type.³

The cell time resolution σ_i improves with increasing deposited energy E as

$$\sigma_i = \sqrt{p_0^2 + \frac{p_1^2}{E} + \left(\frac{p_2}{E}\right)^2} \quad (4.1)$$

³The cell type is defined by the calorimeter layer and pseudorapidity (A1, BC2, etc), see figure 2.




Figure 3. (a) Distribution of β_{ToF} obtained with isolated muons from $Z \rightarrow \mu\mu$ decays (2018 data) with the default calibration (Default) and with the calibration illustrated in section 4 (Calibrated). (b) Dependence of the resolution of β_{ToF} on the pseudorapidity, using isolated muons from $Z \rightarrow \mu\mu$ decays (2018 data) with all the calibrations applied.

The parameters p_0, p_1 and p_2 are determined separately in each radial layer (cells A, (B,BC) and D in figure 2) by a fit of eq. (4.1) to the muons from $Z \rightarrow \mu\mu$ decays collected in the Run 2 dataset. For each radial layer, the energy spectrum is divided in 17 slices, and in each slice the time spectrum is fitted with a Gaussian distribution in the $\pm 2\sigma$ region around the peak. The standard deviation of the fit is taken as the time resolution σ_i for the slice with deposited energy E . Good agreement between data and the fitting curve is obtained for cells in the two outermost radial layers (BC and D), while worse agreement is obtained for cells in the innermost layer (A cells) due to shorter particle path length and lower energy deposits. For the same reason A cells also show worse time resolution.

The impact of the calibration described above on the β_{ToF} performance is checked with isolated muons originating from $Z \rightarrow \mu\mu$ decays. The performance before and after calibration are compared in figure 3(a) showing a 7% improvement on $\sigma(\beta_{\text{ToF}})$. The $|\eta|$ -dependence of the β_{ToF} resolution is shown in figure 3(b). The β_{ToF} resolution improves at larger $|\eta|$ because of the longer track path. This trend is counterbalanced in the barrel-endcap transition region and in the very high $|\eta|$ region as fewer calorimeter cells contribute to the ToF measurement.

To account for the difference between the reconstructed time between data and MC, a smearing of the cell time distribution is applied to Monte Carlo samples in such a way as to get the best matching with the $Z \rightarrow \mu\mu$ data taken in 2018. Figure 4 shows the agreement in the calorimeter β spectra between $Z \rightarrow \mu\mu$ data and Monte Carlo after the time smearing is applied. The residual difference between data and Monte Carlo is below 5% for $\beta_{\text{ToF}} < 1$.




Figure 4. Calorimeter β_{ToF} distribution for $Z \rightarrow \mu\mu$ events in 2018 data and in Monte Carlo after the time smearing procedure. The ratio of the MC to data is shown in the bottom of the plot where a solid black line is drawn at unity for reference.

5 Analysis

5.1 Overview

This analysis searches for heavy ($m > 200$ GeV) charged particles with a proper lifetime $\tau > 3$ ns.

Events are selected online using the lowest-threshold un-prescaled calorimetric $E_{\text{T}}^{\text{miss}}$ trigger, which is based on the vectorial energy sum measured in the calorimeters [70]. Further selections are applied to triggered events and candidate tracks as detailed in section 5.2. For the β -search, a sample of high-momentum isolated tracks with large dE/dx is identified and used to perform two independent measurements, $m_{dE/dx}$ and m_{ToF} . For the di-track search, events with two high-momentum, isolated tracks with large dE/dx are selected. For both regions, the search then consists in comparing the data and predicted background yields in trapezoidal mass windows in the $[m_{dE/dx}, m_{\text{ToF}}]$ or in the $[m_{dE/dx,1}, m_{dE/dx,2}]$ planes. This enforces that the two measurements in the event — either two measurements of the same track or of two tracks — are consistent with the same mass hypothesis. A trapezoidal shape is chosen to take into account the degradation of the mass resolution with increasing mass as well as decreased backgrounds at higher mass. As described in section 5.4, the trapezoidal mass windows are optimised separately for the β -search and the di-track search; the windows are common to LLPs of the same target mass.

Backgrounds can arise from instrumental effects and tails in the measurements of SM processes, which in the case of the dE/dx include the unavoidable Landau tails of the deposited ionisation energy. The background yield and its distribution in the reconstructed mass spectrum is estimated in a fully data-driven approach, as described in section 5.3. Data control samples are used to parameterise the momentum, dE/dx and, when necessary, β_{ToF} distributions and their interdependence, and then to generate pseudo-data that predict the background distribution in the $[m_{dE/dx,1}, m_{dE/dx,2}]$ and the $[m_{dE/dx}, m_{\text{ToF}}]$ planes. Potential

signal contamination is minimised in these background samples by inverting some of the selection criteria, and the background estimate method is validated in separate data samples called validation regions.

5.2 Event selection

Events are selected with a trigger based on E_T^{miss} , which is calculated online using energy measurements in the calorimeter with corrections for multiple pp interactions in each event [70]. The high-level E_T^{miss} trigger threshold varies from 70 GeV to 120 GeV during the data-taking period depending on the pile-up conditions. The efficiency of signal events to satisfy the trigger ranges from 20% to over 95% and depends on the lifetime, mass, and decay mode of the target LLP.

The E_T^{miss} computation is refined in the offline reconstruction after events are required to satisfy basic data quality selection to ensure all parts of the detector are working correctly [44]. The offline E_T^{miss} is built from calibrated muons [71, 72] and electrons [73, 74] that satisfy baseline selections, from calibrated jets [75] reconstructed at the electromagnetic scale with the anti- k_t jet clustering algorithm [76, 77] with radius parameter $R = 0.4$, and from a term that includes selected soft tracks not associated with any other objects in the event [78] but consistent with the primary vertex (PV). To remove beam-induced backgrounds and spurious calorimeter signals that could spoil the calculation of E_T^{miss} , events are rejected if they contain at least one jet tagged as *bad* [79] as determined from shower shape information.

In events where the signal LLPs are detector stable (i.e. they decay outside the ATLAS detector), the LLPs leave only modest energy depositions in the calorimeters, even in the R -hadron case [80], and only a fraction of them are reconstructed as a muon owing to their late arrival time in the muon spectrometer. Therefore, most of the momentum of each LLP is not accounted for in the calorimeter or muon system. Any reconstructed jets from initial-state radiation (ISR) or additional partons in the hard scatter provide a visible contribution that results in a measured imbalance of transverse momentum. In events with metastable LLPs (i.e. LLPs that decay inside the ATLAS detector), stable neutralinos or gravitinos will carry away unmeasured momentum that contributes to the measured E_T^{miss} and increases the trigger efficiency in the assumption that R -parity is conserved and the lightest stable sparticle is electrically neutral, as is in the models under consideration, if there is also significant energy deposited in the detector from SM decay products. The E_T^{miss} trigger and offline selection efficiency increases for metastable LLPs relative to detector-stable LLPs for signals with decay products with significant energy deposition. This includes R -hadrons with a light neutralino, but not R -hadrons with the fixed and small mass-splitting of $\Delta m(\tilde{g}, \tilde{\chi}_1^0) = 30$ GeV, nor the charginos considered here.

For the β -search, events are required to have $E_T^{\text{miss}} > 170$ GeV to enhance the signal sensitivity by removing background events from many SM processes. As the di-track search suppresses background with the requirement of two candidate tracks, the offline selection on E_T^{miss} is relaxed to $E_T^{\text{miss}} > 20$ GeV, increasing the acceptance for low-mass LLPs that may produce online E_T^{miss} due to the presence of an ISR jet in the event but can have low reconstructed offline E_T^{miss} if both LLP tracks are included in the soft-track term or muon term. The background estimate is fully data-driven and insensitive to the difference between

the online and offline E_T^{miss} thresholds, while the effect on signal efficiency is covered by a dedicated systematic uncertainty, as described in section 5.5.

After satisfying the trigger and the offline E_T^{miss} selections, events are required to have a hard-scatter PV with at least two associated reconstructed tracks and to contain at least one (β -search) or two (di-track search) candidate tracks that satisfy the track-level selections detailed below. In the unlikely case that there are more candidate tracks than required in an event after all selections, the candidate(s) with the highest track p_T is (are) selected. The β -search and the di-track search are independent and their results are not combined; and an event can then, in principle, be selected by both analyses.

To enrich the selected sample in potential signal events and to minimise background, candidate tracks are required to have $p_T > 120$ GeV. For masses above 200 GeV, the LLP production is central and therefore accepting only central- η tracks is implemented to maximise significance. For the β -search, candidate tracks must have $|\eta| < 1.6$ to match the η -acceptance of the central calorimeter, and for the di-track search, candidate tracks must have $|\eta| < 1.8$.

Reconstructed tracks must have at least eight clusters across the pixel and SCT detectors. To be considered a candidate, the track must be associated within tolerances with the primary vertex and have an associated cluster in the innermost pixel layer if it passes through an active detector module. Following the optimisation done in ref. [21] a momentum uncertainty requirement, linearly dependent on p_T and with an upper limit at 200%, is applied to ensure good mass resolution at low masses and to increase the signal acceptance at high masses, while discarding poorly reconstructed tracks. Similarly, an explicit requirement of at least five associated clusters in the SCT detector is imposed to reject further poorly reconstructed tracks. Additional requirements are applied to candidate tracks to ensure the robustness of the track ionisation computation: no clusters on the track should be consistent with any other track [39, 81] and at least two pixel clusters, after discarding the cluster with the highest ionisation, must be included in the dE/dx calculation. Moreover, as the signal is expected to generate isolated tracks, and background processes could acquire significant dE/dx from energy deposits from particles with overlapping trajectories, a track-based isolation requirement is applied. The scalar sum of the p_T of additional primary tracks, in a cone of size $\Delta R=0.3$ around the candidate track, must be less than 5 GeV. Although R -hadrons are hadrons, the very massive gluino parton produces an expected fragmentation that is very hard [82], and the isolation requirement is above 80% efficient for the signal models considered.

Additional criteria are applied to reject SM backgrounds from specific processes. To veto tracks from leptonic W decays, the transverse mass of the candidate track must be greater than 130 GeV. The transverse mass between the track momentum $\vec{p}_{\text{trk}}$ and $\vec{p}_T^{\text{miss}}$ is defined as

$$m_T(\vec{p}_{\text{trk}}, \vec{p}_T^{\text{miss}}) \equiv \sqrt{2p_T^{\text{trk}} E_T^{\text{miss}} \left(1 - \cos \Delta\phi(\vec{p}_T^{\text{miss}}, \vec{p}_{\text{trk}})\right)}.$$

For the di-track search, only one of the candidate tracks must satisfy the transverse mass selection.

Tracks from electrons are removed as in ref. [21] by rejecting a track if any jet with $p_T > 20$ GeV is found within a cone of $\Delta R = 0.05$, and has at least 95% of its energy deposited in the electromagnetic calorimeter. Similarly, SM hadrons are removed by excluding tracks

for which any associated jet within a cone of $\Delta R = 0.05$ with $p_T > 20$ GeV has a calibrated energy larger than the track momentum.

Additional selections specific to the β - and di-track searches are detailed below.

5.2.1 β -search selections

The β -search requires at least one candidate track that satisfies the selections described in section 5.2. The specific ionisation of the candidate track measured by the pixel detector must be larger than $1.8 \text{ MeV g}^{-1}\text{cm}^2$, while the most probable value for a MIP is $1.0 \text{ MeV g}^{-1}\text{cm}^2$ with a resolution of $0.13 \text{ MeV g}^{-1}\text{cm}^2$. The surviving background, in particular that arising from the fluctuations in the ionisation tails, is further suppressed by requiring that β_{ToF} is not compatible with one. This is implemented as $\beta_{\text{ToF}} < \beta_{\text{cut}}$, where $\beta_{\text{cut}} = 1 - 2\sigma_{\beta_{\text{ToF}}}$ varies per event and $\sigma_{\beta_{\text{ToF}}}$ is defined in 0.1-wide slices of $|\eta|$, as shown in figure 3(b). The selection has an efficiency for signal events that satisfy the dE/dx requirement ranging from about 80% at low masses to 95% at high masses. The efficiency is high because the dE/dx requirement selects signal particles with low β that are then likely to satisfy the β_{cut} . The analysis sensitivity is limited to LLPs with lifetimes $\tau \gtrsim 3\text{ ns}$ by the requirement that the track reaches the calorimeter. Conversely, the additional discrimination provided by the calorimeter reduces the background by a factor of about 20 relative to ref. [21], which results in an improvement of sensitivity for LLPs with $\tau \gtrsim 10\text{ ns}$.

The efficiency for signal events to satisfy all selections, including the trigger, ranges from 1.0% to 7.5% for the simulated LLP events described in section 3 and with lifetime exceeding 10 ns, and increases with the lifetime of the LLP and its mass. The signal region for the β -search (β -SR) is defined in table 1.

None of the seven tracks associated with the excess reported in ref. [21] satisfy the β -search selections. In particular, none of them have $\beta_{\text{ToF}} < \beta_{\text{cut}}$. This indicates that the excess identified in ref. [21] is not due to heavy, highly-ionising and slow particles reaching the hadronic calorimeter.

5.2.2 Di-track search selections

The di-track search requires at least two candidate tracks that satisfy the selections described in section 5.2. Additional selections are imposed on properties of the di-track system: the two tracks must have opposite electric charge, and their invariant mass, m_{inv} , calculated with the assumption that the track is a pion, must be larger than 200 GeV. The latter selection significantly reduces the contribution from Z boson decays, along with other SM sources.

Two signal regions in the di-track search are then defined, differing in the ionisation selections applied to both candidate tracks. In the Discovery Region (Discovery-SR), the potential signal significance is maximised by requiring both tracks to have $dE/dx > 1.7 \text{ MeV g}^{-1}\text{cm}^2$, which strongly rejects background. In the Exclusion Region (Exclusion-SR), the exclusion sensitivity is maximised by loosening the ionisation selections to increase the signal acceptance. In this region, one track is required to have $dE/dx > 1.6 \text{ MeV g}^{-1}\text{cm}^2$, while the ionisation requirement on the other track is relaxed to $dE/dx > 1.3 \text{ MeV g}^{-1}\text{cm}^2$. The efficiency for a signal model with 400 GeV staus with a lifetime of 10 ns to satisfy

all selections except the ionisation requirements is about 20%; in the Discovery-SR and Exclusion-SR, the total efficiency is about 2% and 5%, respectively.

5.3 Background estimation

The mass distribution of tracks from background processes in the signal regions is estimated by using a data-driven technique in which tracks are sampled from several control regions (CR). The mass of a track is defined by its momentum and $\beta\gamma$; therefore, the background mass distributions, $m_{dE/dx}$ and m_{ToF} , are constructed by sampling templates of these variables extracted from control regions. Validation regions (VR) are defined to verify the closure of the analysis method.

For a narrow $\Delta\eta$ slice, dE/dx and p_T , as well as β_{ToF} and p_T , are assumed to be uncorrelated for background tracks. In addition to the VRs, cross-checks on both assumptions are performed and described in section 5.5. The independence of dE/dx and p_T for minimum ionising particles is explicitly confirmed using muon tracks from an independent sample of selected $Z \rightarrow \mu\mu$ events.

The p_T distribution of background tracks, for each $|\eta|$ slice, is sampled from kinematic CRs and used as a template for the background track momentum in the signal and validation regions. The kinematic CRs are defined by inverting the dE/dx selection for candidate signal tracks. The dE/dx and the β_{ToF} distributions are sampled from additional CRs, as described in the sections below.

To generate a ‘pseudo-data’ background track, a pair of p_T and $|\eta|$ values is sampled from the kinematic CR template. A dE/dx (or a β_{ToF}) value is sampled from the corresponding $|\eta|$ bin of the dE/dx (or β_{ToF}) template. From these sampled values, the track mass, $m_{dE/dx}$ or m_{ToF} , is calculated using the dE/dx - $\beta\gamma$ calibration (or the β_{ToF}). Enough tracks are generated that the number of pseudo-data samples does not limit the accuracy of the predictions. The normalisation and validation of each background estimate is described in the following sections.

5.3.1 β -search background estimation and validation

Two control regions, kin-CR and $\beta\gamma$ -CR, are defined adjacent in phase space to the signal region (SR) (see table 1). The kin-CR is defined by inverting the dE/dx requirement and relaxing the β_{ToF} requirement used in the SR, and the $\beta\gamma$ -CR is defined by inverting the E_T^{miss} requirement used in the SR and removing the dE/dx and relaxing the β_{ToF} requirements. Data events in the $\beta\gamma$ -CR are reweighed with an E_T^{miss} trigger threshold weight to prevent the E_T^{miss} trigger differences causing a bias on the background estimate. The dE/dx and the β_{ToF} distributions in the $\beta\gamma$ -CR serve as the template distributions for the background mass prediction of $m_{dE/dx}$ and m_{ToF} , respectively. Only tracks with $\beta_{ToF} < 1$ are considered as this is the condition to calculate $\beta\gamma_{ToF}$ and then m_{ToF} .

Pseudo-data background tracks are simulated using a pair of $1/p_T$ and $|\eta|$ values sampled from the kinematic control region template and a dE/dx (or a β_{ToF}) value sampled from the corresponding $|\eta|$ bin of the dE/dx (or β_{ToF}) template. From these sampled values, the track mass $m_{dE/dx}$ and m_{ToF} are calculated. Finally, the pseudo-data samples are normalised to data in a sub-region of kin-CR that is expected to be fully depleted in signal, with $\beta_{ToF} > \beta_{cut}$ and both $m_{dE/dx}$ and m_{ToF} lower than 160 GeV.

β -search region	E_T^{miss} [GeV]	dE/dx [MeV g ⁻¹ cm ²]	β_{ToF}
β -SR	> 170	> 1.8	$< \beta_{\text{cut}}$
kin-CR	> 170	< 1.6	< 1.0
$\beta\gamma$ -CR	< 150	—	< 1.0
High- β_{ToF} -VR	> 170	> 1.8	$[\beta_{\text{cut}}, 1.0]$
High- β_{ToF} -VR kin-CR	> 170	< 1.6	$[\beta_{\text{cut}}, 1.0]$
High- β_{ToF} -VR $\beta\gamma$ -CR	< 150	—	$[\beta_{\text{cut}}, 1.0]$
Low- dE/dx -VR	> 170	$[1.05, 1.6]$	$< \beta_{\text{cut}}$
Low- dE/dx -VR kin-CR	> 170	< 1.05	< 1.0
Low- dE/dx -VR $\beta\gamma$ -CR	< 150	< 1.6	< 1.0

Table 1. Definitions of the signal, control and validation regions for the β -search. The β_{cut} varies per event and is defined as $\beta_{\text{cut}} = 1 - 2\sigma_{\beta_{\text{ToF}}}$. The estimation of the background for each signal and validation region requires two dedicated control regions.

The procedure for estimating both the normalisation and shape of the expected background is validated in two regions. One validation region is characterised by high- β_{ToF} (High- β_{ToF} -VR), i.e. it contains tracks that do not satisfy the β_{ToF} cut used for the signal selection (i.e. $\beta_{\text{ToF}} > \beta_{\text{cut}}$) but extends to high dE/dx . The other validation region is characterised by low- dE/dx (Low- dE/dx -VR), i.e. it contains tracks that do not satisfy the dE/dx requirement used for the signal selection (track ionisation in the range of $[1.05, 1.6]$ MeV g⁻¹cm²) but tests low values of β_{ToF} . The definition of these regions and the corresponding control regions used for the background estimation are shown in table 1. The validation regions are mutually exclusive and exclusive with the signal region by construction. The availability of the β_{ToF} measurement allows to define validation regions without a p_T upper limit and therefore allows the background estimate to be validated at high masses, which is an improvement with respect to the validation strategy used in the previous dE/dx analysis [21].

The contribution of possible signal contamination in each validation region was studied by comparing the number of signal events from various signal samples to the number of background tracks predicted by the background estimation procedure. It was found that the possible signal contamination was smaller than about 5% ($N_{\text{signal}}/\sqrt{N_{\text{background}}} < 0.9$) in the Low- dE/dx -VR and smaller than about 12% ($N_{\text{signal}}/\sqrt{N_{\text{background}}} < 1.2$) in the High- β_{ToF} -VR for all samples and masses not excluded by previous searches. The signal contamination is found to be negligible within all control regions and regions used for normalisation.

The predicted m_{ToF} ($m_{dE/dx}$) background distribution compared with the m_{ToF} ($m_{dE/dx}$) distribution in the data are shown in figure 5 and 6. There is good agreement between prediction and data both in the shape of the distributions and in the total yield. The predicted yield, including statistical and systematic uncertainties, is 316 ± 27 (91 ± 6) while the observed yield is 290 (93) for the Low- dE/dx -VR (High- β_{ToF} -VR).




Figure 5. Comparison of predicted (a) m_{ToF} and (b) $m_{\text{dE/dx}}$ background to data in the low- dE/dx validation region. The statistical and systematic uncertainty in the predicted background is calculated as indicated in section 5.5 and shown as a coloured band. The histogram overflow is added into the rightmost bin.




Figure 6. Comparison of predicted (a) m_{ToF} and (b) $m_{\text{dE/dx}}$ background to data in the high- β_{ToF} validation region. The statistical and systematic uncertainty in the predicted background is calculated as indicated in section 5.5 and shown as a coloured band. The histogram overflow is added into the rightmost bin.

5.3.2 Di-track background estimation and validation

The number of events from background processes with two tracks that satisfy all selections and the $m_{\text{dE/dx}}$ distribution of each track is predicted from pseudo-data events sampled from templates constructed from control regions. The kinematics of pairs of background tracks are extracted from a sample of events (kin-CR) that have two tracks that satisfy an inverted dE/dx selection. Sampling two tracks from the same event ensures that all kinematic correlations are retained. Inverting the dE/dx selection on both tracks ensures that there is no significant signal presence in the control region.

The dE/dx template is formed from events that have two tracks, at least one of which has a low p_{T} value to exclude possible signal (dE/dx -CR). As there is a small residual correlation

Di-track region	m_{inv} [GeV]	m_T [GeV]	Track 1 p_T [GeV]	Track 2 p_T [GeV]	Track 1 dE/dx [MeV g ⁻¹ cm ²]	Track 2 dE/dx [MeV g ⁻¹ cm ²]
Exclusion-SR	> 200	> 130	> 120	> 120	> 1.6	> 1.3
Discovery-SR			> 120	> 120	> 1.7	> 1.7
kin-CR			> 120	> 120	< 1.3	< 1.3
dE/dx -CR			> 50	[50, 120]	—	—
LowpT-VR	> 200	> 130	[70, 100]	[70, 100]	> 1.6	> 1.3
LowpT-VR kin-CR			[70, 100]	[70, 100]	< 1.3	< 1.3
LowpT-VR dE/dx -CR			> 10	[10, 70]	—	—
W-VR	< 200	—	[70, 100]	[70, 100]	> 1.6	> 1.3
W-VR kin-CR			[70, 100]	[70, 100]	< 1.3	< 1.3
W-VR dE/dx -CR			> 10	[10, 70]	—	—
InvMass-VR	< 200	> 130	> 120	> 120	> 1.6	> 1.3
InvMass-VR kin-CR			> 120	> 120	< 1.3	< 1.3
InvMass-VR dE/dx -CR			> 50	[50, 120]	—	—
Z-VR	[80, 100]	—	> 120	> 120	> 1.6	> 1.3
Z-VR kin-CR			> 120	> 120	< 1.3	< 1.3
Z-VR dE/dx -CR			> 50	[50, 120]	—	—

Table 2. Definitions of the signal, control and validation regions for the di-track search. Events must satisfy all event-level requirements and have at least two candidate tracks that satisfy all other selection requirements, as defined in the text. Tracks are ranked first by p_T and then by dE/dx ; Track 1 is the candidate track with the larger p_T or dE/dx , depending on the stage of the selection. The estimation of the background for each signal and validation region requires two dedicated control regions.

between dE/dx and $|\eta|$, the dE/dx template is binned in $|\eta|$. The definitions of the signal and control regions, as well as the validation regions defined below, are shown in table 2.

To form a pseudo-data event, an event from the kin-CR is assigned two independent dE/dx values sampled from the corresponding $|\eta|$ bins of the dE/dx template, and the track masses $m_{dE/dx,1}$ and $m_{dE/dx,2}$ are calculated. Events from the kin-CR can be reused with different sampled dE/dx values.

Enough pseudo-data events are generated so that the statistical uncertainty due to sampling is negligible. Treatment of the statistical correlation due to the kin-CR event reuse is discussed in section 5.5. Each pseudo-data event is normalised to data by the relation: $\frac{N_{\text{kin}}}{N_{\text{pseudo}}} \times \frac{1}{(1-f_1)(1-f_2)}$, where N_{kin} is the number of events in the kinematic template, N_{pseudo} is the number of pseudo-data events, and f_1 and f_2 are the fraction of tracks in the dE/dx template with high dE/dx in the η -bin corresponding to the first and second track of the pseudo-data event, respectively. The $(1-f_1)$ and $(1-f_2)$ factors correct for the exclusion of high dE/dx events from the kin-CR. Applying the signal dE/dx selection on the pseudo-data events produces a normalised distribution of $m_{dE/dx}$ for two-track events expected in the signal region.

The background estimation procedure is validated in several regions, in which the full pseudo-data method is repeated using dedicated control regions for each validation region, as defined in table 2. The inclusive LowpT-VR keeps all selections the same as the signal region,

Region	Predicted yield	Observed yield
LowpT-VR	13.0 ± 2.6	14
InvMass-VR	36 ± 1.9	35
W-VR	126 ± 4.0	138
Z-VR	71 ± 4.7	66

Table 3. The expected and observed yields in the di-track validation regions. Only the statistical uncertainty is included in the prediction. Not all regions are orthogonal.

except for the track p_T requirements, which are loosened so that each signal track is required to have p_T between 70 and 100 GeV. The dE/dx control region for LowpT-VR similarly loosens the lower track p_T to 10 GeV, in order to ensure sufficient template statistics. To ensure that the method closes at nominal values of track p_T , an independent InvMass-VR is constructed, in which the track p_T requirements are the same as the signal region, but the invariant mass requirement of the two tracks is inverted. Additionally, two higher statistics validation regions are formed that target background processes with W and Z bosons; both regions omit the one-track transverse mass selection, which dramatically enhances the number of W bosons relative to the InvMass-VR for the W-VR, and the Z-VR additionally omits the offline E_T^{miss} selection and requires that the two tracks each be matched to a reconstructed muon and have an invariant mass consistent with the Z boson. Good agreement between the pseudo-data prediction and the observed data is seen in all validation regions, as shown in table 3 and figure 7.

An additional validation of the expected yield in the signal and validation regions was performed using an $ABCD$ method [83] that requires the leading track to satisfy all signal selections and uses the p_T and dE/dx of the second track as the independent variables. Yields were found to be in agreement with both the pseudo-data model prediction and the observed data in the validation regions. To validate the background behaviour at higher dE/dx values needed for the Discovery-SR, the $ABCD$ estimate was tested in a modified InvMass-VR and Z-VR with symmetric dE/dx selections of 1.3, 1.4, 1.5, and 1.6 $\text{MeV g}^{-1}\text{cm}^2$, and agreement was found within statistical uncertainty in every test. To obtain enough statistics to test the behaviour at $dE/dx > 1.7 \text{ MeV g}^{-1}\text{cm}^2$, a low- p_T single-track validation region was developed, with a prediction of 1204 ± 38 events with 1246 observed in data.




Figure 7. Comparison of the predicted $m_{dE/dx}$ background to data in the di-track validation regions. The statistical and systematic uncertainty in the predicted background is calculated as indicated in section 5.5.

5.4 Mass window definition

The sensitivity to signal LLPs is measured comparing data, expected background, and potential signal in two-dimensional mass windows.

Two masses, $m_{dE/dx}$ and m_{ToF} , can be calculated for each track surviving the β -driven signal selection using $\beta\gamma_{dE/dx}$, $\beta\gamma_{\text{ToF}}$ and the momentum, and similarly two $m_{dE/dx}$ are calculated in each signal di-track event, one per track. The measurement of two masses allows to define surfaces in the $[m_{dE/dx}, m_{\text{ToF}}]$ or $[m_{dE/dx,1}, m_{dE/dx,2}]$ plane that are optimised for each signal mass. A trapezoidal shaped window is found to maximise sensitivity, reflecting the worsening mass resolution at higher mass, caused primarily by the worsening momentum resolution at higher momenta. The opening angle limiting the trapezoids is assumed to be symmetric around the line $m_{dE/dx} = m_{\text{ToF}}$ or $m_{dE/dx,1} = m_{dE/dx,2}$ and is sketched in figure 8.

5.4.1 β -search mass windows

The mass windows are constructed to contain about 70% of the expected signal. The selected surfaces are different for different signal masses. The trapezoidal mass windows are determined within an angle in the $[m_{dE/dx}, m_{\text{ToF}}]$ plane that is determined by the spread in




Figure 8. The trapezoidal mass windows are determined within an angle θ in the $[m_{\text{ToF}}, m_{\text{dE/dx}}]$ or $[m_{\text{dE/dx},1}, m_{\text{dE/dx},2}]$ plane. The mass points at the centre of the upper and lower edges of the trapezoids are shown as red dots and parameterise the mass window for each target mass. The target mass of the trapezoid is shown as a black dot.

$\beta\gamma_{\text{dE/dx}}$ relative to $\beta\gamma_{\text{ToF}}$. The spread in the difference between the two $\beta\gamma$ measurements translates into an expected spread in $m_{\text{dE/dx}}$ relative to m_{ToF} . As $m_{\text{dE/dx}}$ and m_{ToF} are both determined using the same momentum measurement, their difference for each track reflects only the difference in $\beta\gamma$. The trapezoid opening angle is set to 22 degrees around the bisector of the $[m_{\text{dE/dx}}, m_{\text{ToF}}]$ plane, approximately optimal for all mass points studied.

The trapezoids do not depend nor on the LLP lifetime nor on their identity, but only on their mass. The trapezoid lower and upper edges (see table 4) are defined through a procedure designed to optimise the signal sensitivity. First, the lower edge is scanned to find the optimal sensitivity with the upper edge fixed at 7 TeV. Then, the upper edge is defined by lowering it until the sensitivity is maximal while still keeping at least 70% of the signal events.

β -search	Trapezoid parameters, opening angle $\theta=22$ degrees	
Target signal mass [GeV]	Lower edge mass [GeV]	Upper edge mass [GeV]
150	120	210
200	160	290
250	210	380
300	250	490
350	270	640
400	320	680
450	370	700
500	400	810
550	470	930
600	480	1360
650	530	1360
700	530	1390
800	570	2380
900	620	5340
1000	720	6170
1100	780	6170
1200	860	6170
1300	860	6170
1400	950	7000
1600	950	7000
1800	950	7000
2000	1100	7000
2200	1200	7000
2400	1200	7000
2600	1660	7000

Table 4. The 2D mass window parameters for the β -driven signal region derived using the trapezoidal method, given an opening angle of 22 degrees. The lower edge mass identifies the mass point ($m_{dE/dx}$, m_{ToF}) at the centre of the trapezoidal lower edge. The upper edge mass locates the centre at point ($m_{dE/dx}$, m_{ToF}) of the trapezoid upper edge in the 2D mass plane.

5.4.2 Di-track search mass windows

Two masses, $m_{dE/dx,1}$ and $m_{dE/dx,2}$, can be calculated for the pair of tracks surviving the two-track signal selection using $\beta\gamma_{dE/dx}$ and the momentum of each track. Pair-produced sleptons should have the same mass, and are therefore more likely than the background to populate a trapezoid in the $[m_{dE/dx,1}, m_{dE/dx,2}]$ plane. The trapezoid window definition is optimised for the slepton mass region. Two cases are considered separately: the Discovery SR, with only two mass windows created to maximise the signal significance (see table 5) and the Exclusion SR, with mass windows optimised for the strongest signal exclusion (see table 6). There is no upper mass limit to the trapezoid in either the Discovery or Exclusion SR, as adding one was not found to improve sensitivity in the two-track analysis. The optimal angle of the trapezoid is found to increase slowly with target mass, as the momentum measurement

Di-track search Target signal mass [GeV]	Opening angle [Degrees]	Lower edge mass [GeV]
< 350	45	160
≥ 350	45	300

Table 5. The 2D mass window parameters for the two-track Discovery SR derived using the trapezoidal method. The angle identifies the region of compatibility between the mass measurements, and the lower edge mass identifies the mass point ($m_{dE/dx,1}$, $m_{dE/dx,2}$) at the centre of the trapezoidal lower edge.

Di-track search Target signal mass [GeV]	Opening angle [Degrees]	Lower edge mass [GeV]
200	20	160
300	20	260
400	26	340
500	28	430
600	28	520
700	28	600

Table 6. The 2D mass window parameters for the two-track Exclusion SR derived using the trapezoidal method. The angle identifies the region of compatibility between the mass measurements, and the lower edge mass identifies the mass point ($m_{dE/dx,1}$, $m_{dE/dx,2}$) at the centre of the trapezoidal lower edge.

dominates the reconstructed mass at moderate mass and above, and the reconstructed momentum of the tracks are independent (unlike in the β -SR.)

5.5 Uncertainties

Systematic uncertainties come from several sources: the data-driven background determination, corrections for the detector effects, and the experimental and theoretical signal modelling uncertainties.

The data-driven background estimate is based on a pseudo-data method, where the p_T , η , dE/dx and β_{ToF} variables are generated for each pseudo-data event. This generation method comes with the assumption that the dE/dx and p_T are uncorrelated while the η correlation is taken into account through the η -slicing of the samples. To evaluate the validity of this assumption, a closure test is implemented for the β -search where all kinematic, dE/dx , and β_{ToF} templates are extracted from the $\beta\gamma$ -CR of the SR to generate a background distribution and compared with data in a subset of this same region after applying the same dE/dx and β_{ToF} cuts used for the SR. Any observed non-closure might signify that there are correlations that are not taken into account, and a *template correlation uncertainty* must be assigned according to the size of this non-closure in each mass trapezoid. As an additional check the same closure test is repeated also in the $\beta\gamma$ -CR of each VR. This uncertainty ranges from 5% at 200 GeV to 23% at 700 GeV and is the largest single uncertainty in most mass windows.

For the di-track search, the dominant systematic uncertainty in the background estimate is also obtained from a closure test. The validation regions test the assumptions of the background estimation method, but the comparison between prediction and data is limited by the data statistics in the validation regions. The mass distributions from each of the validation regions is compared between the pseudo-data prediction and the observed data

and used to construct a ratio; a 0.68 confidence internal band is fit across all regions from the deviation from unity in this ratio. This uncertainty is calculated and applied per mass bin; it increases with the target mass from about 10% in the lowest mass windows to about 40% in the highest mass window.

The choice of the η binning has a direct effect on the background template shape, and a different η slicing choice could impact the final results. For this reason, an alternative η slicing⁴ is used to generate an alternative data-driven background and then calculate an *alternative η -slice uncertainty*, which is below 5% in all mass windows. Another concern with the dE/dx template is the lack of data at large dE/dx values, which can result in the large dE/dx values to be over or under represented in the data-driven background estimate. To circumvent this issue, alternative dE/dx templates are generated by fitting the original dE/dx templates with a Crystal Ball function [21] that models the dE/dx template with high accuracy for small dE/dx values and does not suffer from statistical uncertainty issues at large dE/dx values. The difference between these templates for large dE/dx values allows to calculate a *dE/dx tail uncertainty*, which is everywhere below 1%. For the di-track analysis, these two uncertainties are found to be negligible.

The data-driven background mass distributions in the signal regions are simulated using pseudo-data events extracted from dE/dx , β_{ToF} and kinematics templates. The statistical precision of the pseudo-data experiments is limited by the statistical uncertainty of the input templates. To compute a correct statistical uncertainty of the background templates, the statistical uncertainties of dE/dx , β_{ToF} and kinematic templates need to be propagated to the final background distributions that are generated by pseudo-data events. To calculate the statistical uncertainty due to the background templates for the β -search, the input kinematic templates are first smoothed to remove empty bins, then these templates are randomly fluctuated assuming Poisson distributions. The newly generated templates are then used to throw new pseudo-data generating alternative background mass distributions. The root mean square difference between the alternative mass distributions is then used as the *statistical uncertainty*, with a maximum value of 5%. For the β -search, an additional percent-level *normalisation uncertainty* is computed by propagating the statistical uncertainties of the generated background and data (which dominates this uncertainty) through the normalisation method and combining them quadratically. For the di-track search, the dE/dx template uncertainties are handled similarly, but the kinematic CR statistical uncertainty is different because the template consists of unbinned events. For the statistical uncertainty related to the finite number of events in the kinematic CR, each event is Poisson fluctuated from a nominal expectation value of one event in the sample; i.e. some events do not appear in a fluctuated sample while others appear multiple times. This is known as a Bootstrap procedure.

An E_T^{miss} *trigger uncertainty* in the background estimate is evaluated by re-generating the background without reweighing the events in the $\beta\gamma$ -CR and comparing with the same events reweighed with an E_T^{miss} trigger threshold weight. The E_T^{miss} *trigger uncertainty* is below 2%. There is no E_T^{miss} reweighing of the data for the di-track search, and as E_T^{miss} does not play a role in the di-track background estimate, there is no associated uncertainty.

⁴The edges of the baseline η -binning are [0, 0.1, 0.2, 0.3, ..., 1.3, 1.4, 1.5, 1.6] while the edges of the alternative η -binning are [0, 0.15, 0.25, 0.35, ..., 1.35, 1.45, 1.6].

For the β -search, two systematic uncertainties are considered that evaluate the effect on the background estimate of choices made in treating the β_{ToF} variable. The β_{ToF} η *slicing uncertainty* evaluates the effect of the choice of η -binning of the β_{ToF} templates, and is calculated similarly to the dE/dx η slicing uncertainty described above. The β_{ToF} η *slicing uncertainty* amounts to roughly 10% in all mass windows. The other β_{ToF} uncertainty is similar to the dE/dx tail uncertainty. Also in this case it is necessary to evaluate how well the background estimate models the low tail of the β_{ToF} distribution. This tail is not well-populated in the η -sliced β_{ToF} templates, and so not all η -dependent effects might be captured in the background estimate. To account for this, a new background estimate is created where β_{ToF} is drawn from a Crystal Ball function that is fitted to the data rather than from the low tails of the β_{ToF} templates. The difference between this new background estimate and the nominal background estimate defines the β_{ToF} *tail uncertainty*, which ranges from 3% to 10%.

The signal cross-section uncertainties are displayed in the final limit plots as theoretical uncertainties in the excluded cross-section. Signals with long lifetimes that are detector stable and signal models with a small mass splitting between the invisible decay product and the LLP parent mass mostly rely on the presence of an ISR jet to satisfy the online $E_{\text{T}}^{\text{miss}}$ trigger. For these signal models, the ISR modelling is expected to be the largest signal systematic uncertainty. To estimate this uncertainty, alternative generator-level signal MC samples were generated with factorisation, renormalisation, and merging scales varied from the nominal values separately by 0.5 and 2.0, as well as with five variations concerning parton shower tuning and radiation uncertainty [47]. The p_{T} of the sparticle and signal-mass dependent weights are then extracted from these samples to parameterise the differences between the samples with scale variations. The newly acquired weights are then applied to fully reconstructed signal MC samples. The differences between the reweighted MC samples and the nominal samples is used as a systematic uncertainty. The majority of these uncertainties are small; the leading ISR jet uncertainty is around 10%.

Additional uncertainties in the signal selection acceptance and efficiency associated with the simulation modelling related to the modelling of the pile-up distribution, the calculation of $E_{\text{T}}^{\text{miss}}$, and track-level quantities are assessed; no individual uncertainty due to the modelling of these quantities has an impact larger than a few percent on the signal yield. The modelling of the $E_{\text{T}}^{\text{miss}}$ trigger efficiency in simulation is validated and an uncertainty derived by comparing the trigger efficiency, as a function of $E_{\text{T}}^{\text{miss}}$, between data and simulation for a sample of $Z \rightarrow \mu\mu$ events with similar track-level requirements as the tracks in the search. The difference between the observed and simulated trigger efficiency measured in $Z \rightarrow \mu\mu$ events is used to correct the signal trigger efficiency based on the reconstructed offline $E_{\text{T}}^{\text{miss}}$, neglecting both muons and the soft track term. The difference between this shifted value and the nominal yield in simulation is used as an additional uncertainty in the signal yield, which is about 15% or lower for all stau signals.

The uncertainty in the combined 2015–2018 integrated luminosity is 0.83% [84], obtained using the LUCID-2 detector [85] for the primary luminosity measurements, complemented by measurements using the inner detector and calorimeters.




Figure 9. The distribution of data and predicted background in the β -search signal region. The observed data events are indicated as magenta circles if they are inside the mass-compatibility angle (shown as grey lines) and as orange diamonds if they are outside, while the blue area is the mass distribution of the background. The overflow is included in the $900 < m_{dE/dx} < 1000$ GeV and in the $900 < m_{ToF} < 1000$ GeV regions.

6 Results

Results are presented separately for the β -search and the two-track search. The statistical analysis and likelihood construction were implemented in the `pyhf` software framework [86]. For each trapezoidal mass window, the likelihood of the background-only hypothesis given the observed data was constructed from the background prediction and the associated systematic uncertainties. The effect of the systematic uncertainties is incorporated through nuisance parameters that are constrained to be Gaussian-distributed. Using a profile-likelihood-based test statistic [87], independent p_0 -values quantifying the level of agreement between the observed data and the background prediction were calculated for each of these windows.

6.1 β -search results

Without any mass-compatibility requirement, nine events are observed in the signal region while the background expectation is of 5.1 ± 0.5 events. The mass values of those events are shown as circles in the $[m_{dE/dx}, m_{ToF}]$ plane in figure 9.

Restricting the observation to be in the 22 degree mass-compatibility angle, the observed events are reduced to six and the background expectation to 3.7 ± 0.4 . The observed data and the expected background in each mass window are shown in table 7. The distribution of the average of $m_{dE/dx}$ and m_{ToF} is shown in figure 10 for the events inside the mass-compatibility angle.

The lowest p -value of 3.3×10^{-2} is measured in the 200 GeV mass window.

Target signal mass [GeV]	Expected background	Observed data
150	1.73 ± 0.17	2
200	1.89 ± 0.20	5
250	1.40 ± 0.17	4
300	1.24 ± 0.17	1
350	1.23 ± 0.18	0
400	0.88 ± 0.14	0
> 400	0.80 ± 0.12	0

Table 7. Data and background yields in the trapezoids defined for different masses in the β -driven analysis. The table extends up to 400 GeV covering the mass region where there are data entries and beyond, including an overflow bin. The regions are not orthogonal.




Figure 10. The distribution of the average of m_{ToF} and $m_{\text{dE/dx}}$ compared with the expected background in the β -search signal region. The systematic uncertainty in the predicted background is calculated as indicated in section 5.5. Only events inside the 22 degrees mass-compatibility angle are considered.

6.2 Di-track search results

There are 15 events observed in the inclusive Exclusion-SR region, with a background expectation of 20.7 ± 4.5 events. The distribution of $m_{\text{dE/dx},1}$ and $m_{\text{dE/dx},2}$ for each event in the inclusive Exclusion-SR is shown in figure 11. There are five observed events that fall into the union of all mass windows; the distribution of the average $m_{\text{dE/dx}}$ of both tracks in these 5 events is shown in figure 12. The observed data and expected background in each mass window in the Exclusion-SR are shown in table 8.

There are zero events observed in the Discovery-SR, with an inclusive background expectation of 0.79 ± 0.19 events. The expected background in both Discovery-SR mass windows is shown in table 9, along with observed and expected model-independent 95% confidence level (CL) upper limits on the number of signal events and the observed 95% CL upper limit on the visible cross-section. Also shown is the discovery p-value, which




Figure 11. The distribution of data and predicted background in the di-track Exclusion-SR. The observed data events are indicated as magenta circles if they are inside the mass-compatibility angle (shown as grey lines) and as orange diamonds if they are outside, while the blue area is the mass distribution of the expected background. The last bins include overflow events.




Figure 12. The distribution of the average $m_{dE/dx}$ of both tracks in each event in the union of all the mass windows in the di-track Exclusion-SR, compared with the expected background. The systematic uncertainty in the predicted background is calculated as indicated in section 5.5. The yield for a 500 GeV stau signal with a lifetime of 10 ns is also shown.

Target signal mass [GeV]	Expected background	Observed data
200	7.93 ± 1.56	4
300	3.49 ± 0.89	1
400	2.09 ± 0.74	1
500	1.07 ± 0.49	0
600	0.59 ± 0.32	0
700	0.35 ± 0.20	0

Table 8. The observed data and expected background in each mass window of the di-track Exclusion-SR. Both systematic and statistical uncertainties in the expected background are included. The regions are not orthogonal.

Lower mass edge [GeV]	Expected background	Observed data	S_{obs}^{95}	S_{exp}^{95}	σ_{vis} [fb]	$P(s = 0)$
160	0.67 ± 0.17	0	3.0	3.1	0.02	0.5
300	0.29 ± 0.10	0	3.0	3.0	0.02	0.5

Table 9. The observed data and expected background in both mass windows of the di-track Discovery-SR. Both systematic and statistical uncertainties in the expected background are included. The regions are not orthogonal. Also shown are model-independent observed and expected 95% CL upper limits on the number of signal events, S_{obs}^{95} and S_{exp}^{95} , the observed 95% CL upper limit on the visible cross-section, σ_{vis} , and the discovery p-value, $P(s = 0)$. The p-value is capped at 0.5.

measures the compatibility of the observed data with the background-only hypothesis relative to fluctuations of the background.

6.3 Lifetime-dependent mass limits

The results of this study are interpreted for the benchmark signal models considered, and the 95% CL upper limit on the cross-section is extracted using pseudo-experiments and the CL_s prescription [88] for each signal mass and lifetime hypothesis. Cross-section limits as a function of lifetime, for varying signal masses, and cross-sections limits as a function of signal mass, for varying lifetimes, are provided for all benchmark signal models in ref. [89].

To more accurately probe the sensitivity of the analysis to LLP lifetimes other than those used in the generation of the signal samples, the same samples are reinterpreted for intermediate lifetime values by reweighing the LLP particle decay spectra. Intermediate lifetimes are modelled by reweighing the closest longer-lifetime sample to shorter lifetimes, except for $\tau > 30$ ns. The choice of target lifetimes for $\tau > 30$ ns is limited by the reduced size of the reweighed sample.

The di-track search is optimised for the stau scenario. It has higher sensitivity than the β -search in this domain, but lower sensitivity for charginos and R -hadrons. The mass limits reported below are therefore obtained with the di-track analysis for staus and with the β -driven analysis for charginos and R -hadrons.

The limits in the stau scenario are shown in figure 13(a). While the interpretation is done only for staus, the analysis has similar sensitivity for selectrons and smuons. Differences between signal selection efficiency arise only from second-order effects on $E_{\text{T}}^{\text{miss}}$ and isolation criteria from the different interactions of the slepton decay products with the detector. The

sensitivity for staus peaks at around 30 ns for two reasons: at lower lifetimes, the LLPs do not travel far enough to be reconstructed as tracks, while at higher lifetimes, the E_T^{miss} trigger efficiency drops, as discussed in section 5.2.

The mass range 200–560 GeV is excluded for mass-degenerate $\tilde{\tau}_L$ and $\tilde{\tau}_R$ with lifetimes $\tau = 10$ ns, while the corresponding expected exclusion is 200–550 GeV. This search is not sensitive to masses below 200 GeV as lighter LLPs have lower p_T and lower dE/dx which do not allow for reasonable background discrimination. These results are the most sensitive to date for detector-unstable $\tilde{\tau}_{L,R}$ with lifetimes above 3 ns. At lower lifetimes, the ATLAS search for displaced leptons provides exclusions for $\tilde{\tau}_{L,R}$ with $\tau = 0.3$ ns up to 380 GeV [90]. Searches for detector-stable LLPs, exploiting the muon system as a trigger, have previously been performed, including an ATLAS search for detector-stable LLPs with dE/dx and ToF using 36 fb^{-1} of data that excluded stable nearly pure $\tilde{\tau}_R$ with masses up to 430 GeV [28], and a CMS result using 101 fb^{-1} of data that excluded detector-stable $\tilde{\tau}_R$ up to 520 GeV and $\tilde{\tau}_{L,R}$ with masses up to 730 GeV [31].

Figure 13(b) shows the mass limits for sum of the $\tilde{\chi}_1^\pm \tilde{\chi}_1^0$ and $\tilde{\chi}_1^+ \tilde{\chi}_1^-$ production. The chargino mass limit is ≈ 1.3 TeV for lifetimes $\tau > 100$ ns. These results provide the most stringent limits to date for detector-unstable charginos in the lifetime range above 10 ns. The previous ATLAS search that selected a single track with significant dE/dx had greater sensitivity for charginos with lifetimes from 3 ns to 10 ns [21], as the requirement for the LLPs to travel to the hadronic calorimeter is less efficient for signals with shorter lifetimes.

Figure 14 shows the mass limits for gluino R -hadron pair production for both the $m(\tilde{\chi}_1^0) = 100$ GeV and $\Delta m(\tilde{g}, \tilde{\chi}_1^0) = 30$ GeV cases. The sensitivity for R -hadrons with $m(\tilde{\chi}_1^0) = 100$ GeV falls for lifetimes above 30 ns due to a loss of efficiency for the E_T^{miss} trigger, as discussed in section 5.2. The charginos and R -hadrons with $\Delta m(\tilde{g}, \tilde{\chi}_1^0) = 30$ GeV do not have decay products that interact significantly with the detector, so their E_T^{miss} trigger efficiency is flat as a function of lifetime. The highest observed lower limit on the mass is 2.27 TeV (2.20 TeV) and is obtained at $\tau = 30$ ns ($\tau > 200$ ns) for $m(\tilde{\chi}_1^0) = 100$ GeV ($\Delta m(\tilde{g}, \tilde{\chi}_1^0) = 30$ GeV), while the corresponding expected limit matches the observed limit. These results provide the most stringent limits to date for detector-unstable LLPs in the lifetime range above 10 ns. At lifetimes lower than 10 ns, other strategies including searches for disappearing tracks [91, 92] and searches for displaced vertices [93, 94] provide better sensitivity.




Figure 13. (a) The contour for the excluded mas5-2ifetime region for stau pair production obtained with the di-track search. All masses and lifetimes shown that are below the curve and above 200 GeV are excluded by the observed data (while the expected exclusion is between the upper curve down to 210 GeV for lifetimes above 3000 ns). The sensitivity extends indefinitely to longer lifetimes. (b) Upper limits on the chargino mass, assuming the $\tilde{\chi}_1^\pm\tilde{\chi}_1^0$ and $\tilde{\chi}_1^\pm\tilde{\chi}_1^\mp$ contributions, versus lifetime obtained with the β -search. Lifetimes below 3 ns and masses below 200 GeV are not probed by these analyses. Observed limits are indicated as solid blue lines and expected limits are indicated by dotted black lines. The shaded band around the expected limit indicate the $1\sigma_{\text{exp}}$ (and $2\sigma_{\text{exp}}$) uncertainty range, derived as explained in section 5.5.




Figure 14. Upper limits on the gluino mass, from gluino R -hadron pair production, as a function of the gluino lifetime for two neutralino mass assumptions of (a) $m(\tilde{\chi}_1^0) = 100$ GeV and (b) $\Delta m(\tilde{g}, \tilde{\chi}_1^0) = 30$ GeV obtained with the β -search. Observed limits are indicated as solid blue lines and expected limits are indicated by dotted black lines. The shaded band around the expected limit indicate the $1\sigma_{\text{exp}}$ (and $2\sigma_{\text{exp}}$) uncertainty range, derived as explained in section 5.5. The upper $1\sigma_{\text{exp}}$ expected bound is very close to the expected limit for some lifetime values due to the expected background approaching zero events. For a given lifetime, the mass values below the curve are excluded.

7 Conclusion

A search is performed for heavy charged LLPs of lifetime exceeding 3 ns produced at the LHC in 140 fb^{-1} of pp collisions at $\sqrt{s} = 13 \text{ TeV}$. This search is based on two independent and compatible measurements of the LLP mass compared to a data-driven background estimate. The two mass measurements are obtained either for two heavily-ionising opposite-sign particles or for one heavily ionising particle that is also measured to be slow-moving. The mass values are determined by the $\beta\gamma$ measurements obtained either through the specific ionisation in the pixel detector or through the ToF measured by the hadronic calorimeter. Two independent determinations of $\beta\gamma$ minimise the effect of the fluctuations that happen in the far tails of those distributions.

Observed yields and distributions agree with the SM background expectations and limits are placed on several simplified SUSY models. The highest sensitivity is reached for LLPs with lifetimes exceeding 10 ns. Masses smaller than 2.27 TeV are excluded at the 95% confidence level for gluino R -hadrons with a lifetime of 30 ns and $m(\tilde{\chi}_1^0) = 100 \text{ GeV}$. The mass limit for compressed-scenario R -hadrons, with $\Delta m(\tilde{g}, \tilde{\chi}_1^0) = 30 \text{ GeV}$ and a lifetime $> 200 \text{ ns}$, is 2.20 TeV. Charginos with masses smaller than 1.3 TeV and lifetime $> 100 \text{ ns}$ are excluded. Masses in the range of 200–560 GeV for staus are excluded for lifetimes of 10 ns.

The limits for detector unstable LLPs in the mas5-2ifetime plane are the most stringent to date in the lifetime domain exceeding 10 ns and provide further constraints on the R -hadron, chargino and stau production models considered.

The seven events in the 3.3 Z significance excess observed in the signal region defined by ref. [21] are excluded by the β_{ToF} selection. This indicates that this excess is not due to heavy, highly-ionising and slow particles reaching the hadronic calorimeter.

Acknowledgments

We thank CERN for the very successful operation of the LHC and its injectors, as well as the support staff at CERN and at our institutions worldwide without whom ATLAS could not be operated efficiently.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF/SFU (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), RAL (U.K.) and BNL (U.S.A.), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in ref. [95].

We gratefully acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; Minciencias, Colombia; MEYS CR, Czech Republic; DNRF and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSRI, Greece; RGC and Hong Kong SAR, China; ICHEP and Academy of Sciences and Humanities, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MNiSW, Poland; FCT, Portugal; MNE/IFA, Romania; MSTDI,

Serbia; MSSR, Slovakia; ARIS and MVZI, Slovenia; DSI/NRF, South Africa; MICIU/AEI, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; NSTC, Taipei; TENMAK, Türkiye; STFC/UKRI, United Kingdom; DOE and NSF, United States of America.

Individual groups and members have received support from BCKDF, CANARIE, CRC and DRAC, Canada; CERN-CZ, FORTE and PRIMUS, Czech Republic; COST, ERC, ERDF, Horizon 2020, ICSC-NextGenerationEU and Marie Skłodowska-Curie Actions, European Union; Investissements d’Avenir Labex, Investissements d’Avenir Idex and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and MINERVA, Israel; NCN and NAWA, Poland; La Caixa Banking Foundation, CERCA Programme Generalitat de Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; The Royal Society and Leverhulme Trust, United Kingdom.

In addition, individual members wish to acknowledge support from Armenia: Yerevan Physics Institute (FAPERJ); CERN: European Organization for Nuclear Research (CERN DOCT); Chile: Agencia Nacional de Investigación y Desarrollo (FONDECYT 1230812, FONDECYT 1230987, FONDECYT 1240864); China: Chinese Ministry of Science and Technology (MOST-2023YFA1605700, MOST-2023YFA1609300), National Natural Science Foundation of China (NSFC — 12175119, NSFC 12275265, NSFC-12075060); Czech Republic: Czech Science Foundation (GACR — 24-11373S), Ministry of Education Youth and Sports (ERC-CZ-LL2327, FORTE CZ.02.01.01/00/22_008/0004632), PRIMUS Research Programme (PRIMUS/21/SCI/017); EU: H2020 European Research Council (ERC — 101002463); European Union: European Research Council (ERC — 948254, ERC 101089007, ERC, BARD, 101116429), Horizon 2020 Framework Programme (MUCCA — CHIST-ERA-19-XAI-00), European Union, Future Artificial Intelligence Research (FAIR-NextGenerationEU PE00000013), Italian Center for High Performance Computing, Big Data and Quantum Computing (ICSC, NextGenerationEU); France: Agence Nationale de la Recherche (ANR-20-CE31-0013, ANR-21-CE31-0013, ANR-21-CE31-0022, ANR-22-EDIR-0002); Germany: Baden-Württemberg Stiftung (BW Stiftung-Postdoc Eliteprogramme), Deutsche Forschungsgemeinschaft (DFG — 469666862, DFG — CR 312/5-2); Italy: Istituto Nazionale di Fisica Nucleare (ICSC, NextGenerationEU), Ministero dell’Università e della Ricerca (NextGenEU I53D23000820006 M4C2.1.1); Japan: Japan Society for the Promotion of Science (JSPS KAKENHI JP22H01227, JSPS KAKENHI JP22H04944, JSPS KAKENHI JP22KK0227, JSPS KAKENHI JP23KK0245); Norway: Research Council of Norway (RCN-314472); Poland: Ministry of Science and Higher Education (IDUB AGH, POB8, D4 no 9722), Polish National Agency for Academic Exchange (PPN/PPO/2020/1/00002/U/00001), Polish National Science Centre (NCN 2021/42/E/ST2/00350, NCN OPUS 2023/51/B/ST2/02507, NCN OPUS nr 2022/47/B/ST2/03059, NCN UMO-2019/34/E/ST2/00393, UMO-2020/37/B/ST2/01043, UMO-2021/40/C/ST2/00187, UMO-2022/47/O/ST2/00148, UMO-2023/49/B/ST2/04085, UMO-2023/51/B/ST2/00920); Portugal: Foundation for Science and Technology (FCT); Spain: Generalitat Valenciana (Artemisa, FEDER, IDIFEDER/2018/048), Ministry of Science and Innovation (MCIN & NextGenEU PCI2022-135018-2, MICIN & FEDER PID2021-

125273NB, RYC2019-028510-I, RYC2020-030254-I, RYC2021-031273-I, RYC2022-038164-I); Sweden: Carl Trygger Foundation (Carl Trygger Foundation CTS 22:2312), Swedish Research Council (Swedish Research Council 2023-04654, VR 2018-00482, VR 2021-03651, VR 2022-03845, VR 2022-04683, VR 2023-03403), Knut and Alice Wallenberg Foundation (KAW 2018.0458, KAW 2019.0447, KAW 2022.0358); Switzerland: Swiss National Science Foundation (SNSF — PCEFP2_194658); United Kingdom: Leverhulme Trust (Leverhulme Trust RPG-2020-004), Royal Society (NIF-R1-231091); United States of America: U.S. Department of Energy (ECA DE-AC02-76SF00515), Neubauer Family Foundation.

Data Availability Statement. The public release of data supporting the findings of this article will follow the CERN Open Data Policy [96]. Inquiries about plots and tables associated with this article can be addressed to atlas.publications@cern.ch.

Code Availability Statement. The ATLAS collaboration’s Athena software, including the configuration of the event generators, is open source (<http://gitlab.cern.ch/atlas/athena>).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] H. Acaroglu, P. Agrawal and M. Blanke, *Lepton-flavoured scalar dark matter in dark minimal flavour violation*, *JHEP* **05** (2023) 106 [[arXiv:2211.03809](https://arxiv.org/abs/2211.03809)] [[INSPIRE](#)].
- [2] P. Asadi, A. Radick and T.-T. Yu, *Interplay of freeze-in and freeze-out: lepton-flavored dark matter and muon colliders*, *Phys. Rev. D* **110** (2024) 035022 [[arXiv:2312.03826](https://arxiv.org/abs/2312.03826)] [[INSPIRE](#)].
- [3] Y.A. Golfand and E.P. Likhtman, *Extension of the algebra of Poincaré group generators and violation of p invariance*, *JETP Lett.* **13** (1971) 323 [[INSPIRE](#)].
- [4] D.V. Volkov and V.P. Akulov, *Is the neutrino a Goldstone particle?*, *Phys. Lett. B* **46** (1973) 109 [[INSPIRE](#)].
- [5] J. Wess and B. Zumino, *Supergauge transformations in four-dimensions*, *Nucl. Phys. B* **70** (1974) 39 [[INSPIRE](#)].
- [6] J. Wess and B. Zumino, *Supergauge invariant extension of quantum electrodynamics*, *Nucl. Phys. B* **78** (1974) 1 [[INSPIRE](#)].
- [7] S. Ferrara and B. Zumino, *Supergauge invariant Yang-Mills theories*, *Nucl. Phys. B* **79** (1974) 413 [[INSPIRE](#)].
- [8] A. Salam and J.A. Strathdee, *Supersymmetry and non-Abelian gauges*, *Phys. Lett. B* **51** (1974) 353 [[INSPIRE](#)].
- [9] L. Evans and P. Bryant, *LHC machine*, 2008 *JINST* **3** S08001 [[INSPIRE](#)].
- [10] G.F. Giudice and A. Romanino, *Split supersymmetry*, *Nucl. Phys. B* **699** (2004) 65 [[hep-ph/0406088](https://arxiv.org/abs/hep-ph/0406088)] [[INSPIRE](#)].
- [11] N. Arkani-Hamed and S. Dimopoulos, *Supersymmetric unification without low energy supersymmetry and signatures for fine-tuning at the LHC*, *JHEP* **06** (2005) 073 [[hep-th/0405159](https://arxiv.org/abs/hep-th/0405159)] [[INSPIRE](#)].

- [12] M. Fairbairn et al., *Stable massive particles at colliders*, *Phys. Rept.* **438** (2007) 1 [[hep-ph/0611040](#)] [[INSPIRE](#)].
- [13] G.F. Giudice, M.A. Luty, H. Murayama and R. Rattazzi, *Gaugino mass without singlets*, *JHEP* **12** (1998) 027 [[hep-ph/9810442](#)] [[INSPIRE](#)].
- [14] L. Randall and R. Sundrum, *Out of this world supersymmetry breaking*, *Nucl. Phys. B* **557** (1999) 79 [[hep-th/9810155](#)] [[INSPIRE](#)].
- [15] J.R. Ellis, T. Falk and K.A. Olive, *Neutralino-stau coannihilation and the cosmological upper limit on the mass of the lightest supersymmetric particle*, *Phys. Lett. B* **444** (1998) 367 [[hep-ph/9810360](#)] [[INSPIRE](#)].
- [16] T. Jittoh, J. Sato, T. Shimomura and M. Yamanaka, *Long life stau in the minimal supersymmetric standard model*, *Phys. Rev. D* **73** (2006) 055009 [Erratum *ibid.* **87** (2013) 019901] [[hep-ph/0512197](#)] [[INSPIRE](#)].
- [17] S. Kaneko et al., *Measuring lepton flavor violation at LHC with a long-lived slepton in the coannihilation region*, *Phys. Rev. D* **78** (2008) 116013 [Erratum *ibid.* **87** (2013) 039904] [[arXiv:0811.0703](#)] [[INSPIRE](#)].
- [18] M. Dine and W. Fischler, *A phenomenological model of particle physics based on supersymmetry*, *Phys. Lett. B* **110** (1982) 227 [[INSPIRE](#)].
- [19] L. Alvarez-Gaume, M. Claudson and M.B. Wise, *Low-energy supersymmetry*, *Nucl. Phys. B* **207** (1982) 96 [[INSPIRE](#)].
- [20] C.R. Nappi and B.A. Ovrut, *Supersymmetric extension of the $SU(3) \times SU(2) \times U(1)$ model*, *Phys. Lett. B* **113** (1982) 175 [[INSPIRE](#)].
- [21] ATLAS collaboration, *Search for heavy, long-lived, charged particles with large ionisation energy loss in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS experiment and the full Run 2 dataset*, *JHEP* **06** (2023) 158 [[arXiv:2205.06013](#)] [[INSPIRE](#)].
- [22] G.F. Giudice, M. McCullough and D. Teresi, *dE/dx from boosted long-lived particles*, *JHEP* **08** (2022) 012 [[arXiv:2205.04473](#)] [[INSPIRE](#)].
- [23] ATLAS collaboration, *Search for heavy long-lived multi-charged particles in the full LHC run 2 pp collision data at $\sqrt{s} = 13$ TeV using the ATLAS detector*, *Phys. Lett. B* **847** (2023) 138316 [[arXiv:2303.13613](#)] [[INSPIRE](#)].
- [24] ATLAS collaboration, *Search for heavy charged long-lived particles in proton-proton collisions at $\sqrt{s} = 13$ TeV using an ionisation measurement with the ATLAS detector*, *Phys. Lett. B* **788** (2019) 96 [[arXiv:1808.04095](#)] [[INSPIRE](#)].
- [25] ATLAS collaboration, *Search for heavy long-lived charged particles with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV*, *Phys. Lett. B* **703** (2011) 428 [[arXiv:1106.4495](#)] [[INSPIRE](#)].
- [26] ATLAS collaboration, *Search for metastable heavy charged particles with large ionisation energy loss in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS experiment*, *Eur. Phys. J. C* **75** (2015) 407 [[arXiv:1506.05332](#)] [[INSPIRE](#)].
- [27] ATLAS collaboration, *Search for metastable heavy charged particles with large ionization energy loss in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS experiment*, *Phys. Rev. D* **93** (2016) 112015 [[arXiv:1604.04520](#)] [[INSPIRE](#)].
- [28] ATLAS collaboration, *Search for heavy charged long-lived particles in the ATLAS detector in 36.1 fb^{-1} of proton-proton collision data at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **99** (2019) 092007 [[arXiv:1902.01636](#)] [[INSPIRE](#)].

- [29] ATLAS collaboration, *Search for heavy long-lived charged R-hadrons with the ATLAS detector in 3.2fb^{-1} of proton-proton collision data at $\sqrt{s} = 13\text{ TeV}$* , *Phys. Lett. B* **760** (2016) 647 [[arXiv:1606.05129](#)] [[INSPIRE](#)].
- [30] CMS collaboration, *Search for long-lived charged particles in proton-proton collisions at $\sqrt{s} = 13\text{ TeV}$* , *Phys. Rev. D* **94** (2016) 112004 [[arXiv:1609.08382](#)] [[INSPIRE](#)].
- [31] CMS collaboration, *Search for heavy long-lived charged particles with large ionization energy loss in proton-proton collisions at $\sqrt{s} = 13\text{ TeV}$* , *JHEP* **04** (2025) 109 [[arXiv:2410.09164](#)] [[INSPIRE](#)].
- [32] ATLAS collaboration, *The ATLAS experiment at the CERN Large Hadron Collider*, 2008 *JINST* **3** S08003 [[INSPIRE](#)].
- [33] ATLAS collaboration, *Performance of the ATLAS trigger system in 2015*, *Eur. Phys. J. C* **77** (2017) 317 [[arXiv:1611.09661](#)] [[INSPIRE](#)].
- [34] ATLAS collaboration, *Software and computing for run 3 of the ATLAS experiment at the LHC*, *Eur. Phys. J. C* **85** (2025) 234 [[arXiv:2404.06335](#)] [[INSPIRE](#)].
- [35] ATLAS collaboration, *ATLAS pixel detector*, *ATLAS-TDR-11*, CERN, Geneva, Switzerland (1998).
- [36] ATLAS collaboration, *ATLAS Insertable B-Layer technical design report*, *CERN-LHCC-2010-013*, CERN, Geneva, Switzerland (2010).
- [37] ATLAS IBL collaboration, *Production and integration of the ATLAS Insertable B-Layer*, 2018 *JINST* **13** T05008 [[arXiv:1803.00844](#)] [[INSPIRE](#)].
- [38] BABAR collaboration, *The BaBar detector*, *Nucl. Instrum. Meth. A* **479** (2002) 1 [[hep-ex/0105044](#)] [[INSPIRE](#)].
- [39] ATLAS collaboration, *A neural network clustering algorithm for the ATLAS silicon pixel detector*, 2014 *JINST* **9** P09009 [[arXiv:1406.7690](#)] [[INSPIRE](#)].
- [40] PARTICLE DATA GROUP collaboration, *Review of particle physics*, *PTEP* **2020** (2020) 083C01 [[INSPIRE](#)].
- [41] ATLAS collaboration, *Operation and performance of the ATLAS semiconductor tracker in LHC run 2*, 2022 *JINST* **17** P01013 [[arXiv:2109.02591](#)] [[INSPIRE](#)].
- [42] ATLAS collaboration, *Operation and performance of the ATLAS tile calorimeter in run 1*, *Eur. Phys. J. C* **78** (2018) 987 [[arXiv:1806.02129](#)] [[INSPIRE](#)].
- [43] ATLAS collaboration, *Performance and calibration of the ATLAS tile calorimeter*, *Instruments* **6** (2022) 25 [[INSPIRE](#)].
- [44] ATLAS collaboration, *ATLAS data quality operations and performance for 2015–2018 data-taking*, 2020 *JINST* **15** P04003 [[arXiv:1911.04632](#)] [[INSPIRE](#)].
- [45] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].
- [46] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[arXiv:1410.3012](#)] [[INSPIRE](#)].
- [47] ATLAS collaboration, *ATLAS Pythia 8 tunes to 7 TeV data*, *ATL-PHYS-PUB-2014-021*, CERN, Geneva, Switzerland (2014).
- [48] R.D. Ball et al., *Parton distributions with LHC data*, *Nucl. Phys. B* **867** (2013) 244 [[arXiv:1207.1303](#)] [[INSPIRE](#)].

- [49] J. Pumplin et al., *New generation of parton distributions with uncertainties from global QCD analysis*, *JHEP* **07** (2002) 012 [[hep-ph/0201195](#)] [[INSPIRE](#)].
- [50] A.D. Martin, W.J. Stirling, R.S. Thorne and G. Watt, *Parton distributions for the LHC*, *Eur. Phys. J. C* **63** (2009) 189 [[arXiv:0901.0002](#)] [[INSPIRE](#)].
- [51] D.J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152 [[INSPIRE](#)].
- [52] S. Catani, F. Krauss, R. Kuhn and B.R. Webber, *QCD matrix elements + parton showers*, *JHEP* **11** (2001) 063 [[hep-ph/0109231](#)] [[INSPIRE](#)].
- [53] S. Höche, F. Krauss, S. Schumann and F. Siegert, *QCD matrix elements and truncated showers*, *JHEP* **05** (2009) 053 [[arXiv:0903.1219](#)] [[INSPIRE](#)].
- [54] ATLAS collaboration, *Generation and simulation of R-hadrons in the ATLAS experiment*, [ATL-PHYS-PUB-2019-019](#), CERN, Geneva, Switzerland (2019).
- [55] C. Borschensky et al., *Squark and gluino production cross sections in pp collisions at $\sqrt{s} = 13, 14, 33$ and 100 TeV*, *Eur. Phys. J. C* **74** (2014) 3174 [[arXiv:1407.5066](#)] [[INSPIRE](#)].
- [56] M. Ibe, S. Matsumoto and R. Sato, *Mass splitting between charged and neutral winos at two-loop level*, *Phys. Lett. B* **721** (2013) 252 [[arXiv:1212.5989](#)] [[INSPIRE](#)].
- [57] B. Fuks, M. Klasen, D.R. Lamprea and M. Rothering, *Gaugino production in proton-proton collisions at a center-of-mass energy of 8 TeV*, *JHEP* **10** (2012) 081 [[arXiv:1207.2159](#)] [[INSPIRE](#)].
- [58] B. Fuks, M. Klasen, D.R. Lamprea and M. Rothering, *Precision predictions for electroweak superpartner production at hadron colliders with Resummino*, *Eur. Phys. J. C* **73** (2013) 2480 [[arXiv:1304.0790](#)] [[INSPIRE](#)].
- [59] J. Alwall, M.-P. Le, M. Lisanti and J.G. Wacker, *Searching for directly decaying gluinos at the Tevatron*, *Phys. Lett. B* **666** (2008) 34 [[arXiv:0803.0019](#)] [[INSPIRE](#)].
- [60] J. Alwall, P. Schuster and N. Toro, *Simplified models for a first characterization of new physics at the LHC*, *Phys. Rev. D* **79** (2009) 075020 [[arXiv:0810.3921](#)] [[INSPIRE](#)].
- [61] LHC NEW PHYSICS WORKING GROUP collaboration, *Simplified models for LHC new physics searches*, *J. Phys. G* **39** (2012) 105005 [[arXiv:1105.2838](#)] [[INSPIRE](#)].
- [62] W. Beenakker et al., *The production of charginos/neutralinos and sleptons at hadron colliders*, *Phys. Rev. Lett.* **83** (1999) 3780 [Erratum *ibid.* **100** (2008) 029901] [[hep-ph/9906298](#)] [[INSPIRE](#)].
- [63] J. Debove, B. Fuks and M. Klasen, *Threshold resummation for gaugino pair production at hadron colliders*, *Nucl. Phys. B* **842** (2011) 51 [[arXiv:1005.2909](#)] [[INSPIRE](#)].
- [64] J. Fiaschi and M. Klasen, *Neutralino-chargino pair production at NLO+NLL with resummation-improved parton density functions for LHC run II*, *Phys. Rev. D* **98** (2018) 055014 [[arXiv:1805.11322](#)] [[INSPIRE](#)].
- [65] T. Sjöstrand, S. Mrenna and P.Z. Skands, *A brief introduction to PYTHIA 8.1*, *Comput. Phys. Commun.* **178** (2008) 852 [[arXiv:0710.3820](#)] [[INSPIRE](#)].
- [66] ATLAS collaboration, *The Pythia 8 A3 tune description of ATLAS minimum bias and inelastic measurements incorporating the Donnachie-Landshoff diffractive model*, [ATL-PHYS-PUB-2016-017](#), CERN, Geneva, Switzerland (2016).
- [67] ATLAS collaboration, *The ATLAS simulation infrastructure*, *Eur. Phys. J. C* **70** (2010) 823 [[arXiv:1005.4568](#)] [[INSPIRE](#)].

- [68] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [INSPIRE].
- [69] F. Wang et al., *The impact of incorporating shell-corrections to energy loss in silicon*, *Nucl. Instrum. Meth. A* **899** (2018) 1 [arXiv:1711.05465] [INSPIRE].
- [70] ATLAS collaboration, *Performance of the missing transverse momentum triggers for the ATLAS detector during run-2 data taking*, *JHEP* **08** (2020) 080 [arXiv:2005.09554] [INSPIRE].
- [71] ATLAS collaboration, *Muon reconstruction and identification efficiency in ATLAS using the full run 2 pp collision data set at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **81** (2021) 578 [arXiv:2012.00578] [INSPIRE].
- [72] ATLAS collaboration, *Studies of the muon momentum calibration and performance of the ATLAS detector with pp collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **83** (2023) 686 [arXiv:2212.07338] [INSPIRE].
- [73] ATLAS collaboration, *Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton-proton collision data*, *2019 JINST* **14** P12006 [arXiv:1908.00005] [INSPIRE].
- [74] ATLAS collaboration, *Electron and photon energy calibration with the ATLAS detector using LHC run 2 data*, *2024 JINST* **19** P02009 [arXiv:2309.05471] [INSPIRE].
- [75] ATLAS collaboration, *Jet energy scale and resolution measured in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **81** (2021) 689 [arXiv:2007.02645] [INSPIRE].
- [76] M. Cacciari, G.P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063 [arXiv:0802.1189] [INSPIRE].
- [77] M. Cacciari, G.P. Salam and G. Soyez, *FastJet user manual*, *Eur. Phys. J. C* **72** (2012) 1896 [arXiv:1111.6097] [INSPIRE].
- [78] ATLAS collaboration, *The performance of missing transverse momentum reconstruction and its significance with the ATLAS detector using 140fb^{-1} of $\sqrt{s} = 13$ TeV pp collisions*, *Eur. Phys. J. C* **85** (2025) 606 [arXiv:2402.05858] [INSPIRE].
- [79] ATLAS collaboration, *Selection of jets produced in 13 TeV proton-proton collisions with the ATLAS detector*, *ATLAS-CONF-2015-029*, CERN, Geneva, Switzerland (2015).
- [80] R. Mackeprang and D. Milstead, *An updated description of heavy-hadron interactions in GEANT4*, *Eur. Phys. J. C* **66** (2010) 493 [arXiv:0908.1868] [INSPIRE].
- [81] ATLAS collaboration, *Performance of the ATLAS track reconstruction algorithms in dense environments in LHC run 2*, *Eur. Phys. J. C* **77** (2017) 673 [arXiv:1704.07983] [INSPIRE].
- [82] M. Fairbairn et al., *Stable massive particles at colliders*, *Phys. Rept.* **438** (2007) 1 [hep-ph/0611040] [INSPIRE].
- [83] ATLAS collaboration, *Search for long-lived, multi-charged particles in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector*, *Phys. Lett. B* **722** (2013) 305 [arXiv:1301.5272] [INSPIRE].
- [84] ATLAS collaboration, *Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC*, *Eur. Phys. J. C* **83** (2023) 982 [arXiv:2212.09379] [INSPIRE].
- [85] G. Avoni et al., *The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS*, *2018 JINST* **13** P07017 [INSPIRE].

- [86] M. Feickert, L. Heinrich and G. Stark, *pyhf: a pure-python statistical fitting library with tensors and automatic differentiation*, *PoS ICHEP2022* (2022) 245 [[arXiv:2211.15838](#)] [[INSPIRE](#)].
- [87] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [Erratum *ibid.* **73** (2013) 2501] [[arXiv:1007.1727](#)] [[INSPIRE](#)].
- [88] A.L. Read, *Presentation of search results: the CL_s technique*, *J. Phys. G* **28** (2002) 2693 [[INSPIRE](#)].
- [89] ATLAS collaboration, *HepData record for this publication*, <https://www.hepdata.net/record/ins2878503>.
- [90] ATLAS collaboration, *Search for displaced leptons in $\sqrt{s} = 13$ TeV and 13.6 TeV pp collisions with the ATLAS detector*, [arXiv:2410.16835](#) [[INSPIRE](#)].
- [91] CMS collaboration, *Search for supersymmetry in final states with disappearing tracks in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **109** (2024) 072007 [[arXiv:2309.16823](#)] [[INSPIRE](#)].
- [92] ATLAS collaboration, *Search for long-lived charginos based on a disappearing-track signature using 136 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **82** (2022) 606 [[arXiv:2201.02472](#)] [[INSPIRE](#)].
- [93] CMS collaboration, *Search for long-lived particles using displaced vertices and missing transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Phys. Rev. D* **109** (2024) 112005 [[arXiv:2402.15804](#)] [[INSPIRE](#)].
- [94] ATLAS collaboration, *Search for long-lived, massive particles in events with displaced vertices and multiple jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **06** (2023) 200 [[arXiv:2301.13866](#)] [[INSPIRE](#)].
- [95] ATLAS collaboration, *ATLAS computing acknowledgements*, [ATL-SOFT-PUB-2025-001](#), CERN, Geneva, Switzerland (2025).
- [96] CERN, *CERN open data policy for the LHC experiments*, [CERN-OPEN-2020-013](#), CERN, Geneva, Switzerland (2020) [[DOI:10.17181/CERN.QXNK.8L2G](#)].

The ATLAS collaboration

G. Aad ^{[105](#)}, E. Aakvaag ^{[17](#)}, B. Abbott ^{[124](#)}, S. Abdelhameed ^{[120a](#)}, K. Abeling ^{[56](#)},
N.J. Abicht ^{[50](#)}, S.H. Abidi ^{[30](#)}, M. Aboelela ^{[46](#)}, A. Aboulhorma ^{[36e](#)}, H. Abramowicz ^{[158](#)},
Y. Abulaiti ^{[121](#)}, B.S. Acharya ^{[70a,70b,o](#)}, A. Ackermann ^{[64a](#)}, C. Adam Bourdarios ^{[4](#)},
L. Adamczyk ^{[87a](#)}, S.V. Addepalli ^{[150](#)}, M.J. Addison ^{[104](#)}, J. Adelman ^{[119](#)}, A. Adiguzel ^{[22c](#)},
T. Adye ^{[138](#)}, A.A. Affolder ^{[140](#)}, Y. Afik ^{[41](#)}, M.N. Agaras ^{[13](#)}, A. Aggarwal ^{[103](#)},
C. Agheorghiesei ^{[28c](#)}, F. Ahmadov ^{[40,ae](#)}, S. Ahuja ^{[98](#)}, X. Ai ^{[144b](#)}, G. Aielli ^{[77a,77b](#)},
A. Aikot ^{[170](#)}, M. Ait Tamlihat ^{[36e](#)}, B. Aitbenkikh ^{[36a](#)}, M. Akbiyik ^{[103](#)}, T.P.A. Åkesson ^{[101](#)},
A.V. Akimov ^{[152](#)}, D. Akiyama ^{[175](#)}, N.N. Akolkar ^{[25](#)}, S. Aktas ^{[22a](#)}, G.L. Alberghi ^{[24b](#)},
J. Albert ^{[172](#)}, P. Albicocco ^{[54](#)}, G.L. Albouy ^{[61](#)}, S. Alderweireldt ^{[53](#)}, Z.L. Alegria ^{[125](#)},
M. Aleksa ^{[37](#)}, I.N. Aleksandrov ^{[40](#)}, C. Alexa ^{[28b](#)}, T. Alexopoulos ^{[10](#)}, F. Alfonsi ^{[24b](#)},
M. Algren ^{[57](#)}, M. Alhroob ^{[174](#)}, B. Ali ^{[136](#)}, H.M.J. Ali ^{[94,x](#)}, S. Ali ^{[32](#)}, S.W. Alibocus ^{[95](#)},
M. Aliev ^{[34c](#)}, G. Alimonti ^{[72a](#)}, W. Alkakh ^{[56](#)}, C. Allaire ^{[67](#)}, B.M.M. Allbrooke ^{[153](#)},
J.S. Allen ^{[104](#)}, J.F. Allen ^{[53](#)}, P.P. Allport ^{[21](#)}, A. Aloisio ^{[73a,73b](#)}, F. Alonso ^{[93](#)}, C. Alpigiani ^{[143](#)},
Z.M.K. Alsolami ^{[94](#)}, A. Alvarez Fernandez ^{[103](#)}, M. Alves Cardoso ^{[57](#)}, M.G. Alviggi ^{[73a,73b](#)},
M. Aly ^{[104](#)}, Y. Amaral Coutinho ^{[84b](#)}, A. Ambler ^{[107](#)}, C. Amelung ^{[37](#)}, M. Amerl ^{[104](#)},
C.G. Ames ^{[112](#)}, D. Amidei ^{[109](#)}, B. Amini ^{[55](#)}, K. Amirie ^{[162](#)}, A. Amirkhanov ^{[40](#)},
S.P. Amor Dos Santos ^{[134a](#)}, K.R. Amos ^{[170](#)}, D. Amperiadou ^{[159](#)}, S. An ^{[85](#)}, V. Ananiev ^{[129](#)},
C. Anastopoulos ^{[146](#)}, T. Andeen ^{[11](#)}, J.K. Anders ^{[95](#)}, A.C. Anderson ^{[60](#)}, A. Andreazza ^{[72a,72b](#)},
S. Angelidakis ^{[9](#)}, A. Angerami ^{[43](#)}, A.V. Anisenkov ^{[40](#)}, A. Annovi ^{[75a](#)}, C. Antel ^{[57](#)},
E. Antipov ^{[152](#)}, M. Antonelli ^{[54](#)}, F. Anulli ^{[76a](#)}, M. Aoki ^{[85](#)}, T. Aoki ^{[160](#)}, M.A. Aparo ^{[153](#)},
L. Aperio Bella ^{[49](#)}, C. Appelt ^{[158](#)}, A. Apyan ^{[27](#)}, S.J. Arbiol Val ^{[88](#)}, C. Arcangeletti ^{[54](#)},
A.T.H. Arce ^{[52](#)}, J-F. Arguin ^{[111](#)}, S. Argyropoulos ^{[159](#)}, J.-H. Arling ^{[49](#)}, O. Arnaez ^{[4](#)},
H. Arnold ^{[152](#)}, G. Artoni ^{[76a,76b](#)}, H. Asada ^{[114](#)}, K. Asai ^{[122](#)}, S. Asai ^{[160](#)}, N.A. Asbah ^{[37](#)},
R.A. Ashby Pickering ^{[174](#)}, A.M. Aslam ^{[98](#)}, K. Assamagan ^{[30](#)}, R. Astalos ^{[29a](#)},
K.S.V. Astrand ^{[101](#)}, S. Atashi ^{[166](#)}, R.J. Atkin ^{[34a](#)}, H. Atmani ^{[36f](#)}, P.A. Atmasiddha ^{[132](#)},
K. Augsten ^{[136](#)}, A.D. Aurio ^{[42](#)}, V.A. Austrup ^{[104](#)}, G. Avolio ^{[37](#)}, K. Axiotis ^{[57](#)},
G. Azuelos ^{[111,aj](#)}, D. Babal ^{[29b](#)}, H. Bachacou ^{[139](#)}, K. Bachas ^{[159,s](#)}, A. Bachi ^{[35](#)},
E. Bachmann ^{[51](#)}, M.J. Backes ^{[64a](#)}, A. Badea ^{[41](#)}, T.M. Baer ^{[109](#)}, P. Bagnaia ^{[76a,76b](#)},
M. Bahmani ^{[19](#)}, D. Bahner ^{[55](#)}, K. Bai ^{[127](#)}, J.T. Baines ^{[138](#)}, L. Baines ^{[97](#)}, O.K. Baker ^{[179](#)},
E. Bakos ^{[16](#)}, D. Bakshi Gupta ^{[8](#)}, L.E. Balabram Filho ^{[84b](#)}, V. Balakrishnan ^{[124](#)},
R. Balasubramanian ^{[4](#)}, E.M. Baldin ^{[39](#)}, P. Balek ^{[87a](#)}, E. Ballabene ^{[24b,24a](#)}, F. Balli ^{[139](#)},
L.M. Baltes ^{[64a](#)}, W.K. Balunas ^{[33](#)}, J. Balz ^{[103](#)}, I. Bamwidhi ^{[120b](#)}, E. Banas ^{[88](#)},
M. Bandieramonte ^{[133](#)}, A. Bandyopadhyay ^{[25](#)}, S. Bansal ^{[25](#)}, L. Barak ^{[158](#)}, M. Barakat ^{[49](#)},
E.L. Barberio ^{[108](#)}, D. Barberis ^{[18b](#)}, M. Barbero ^{[105](#)}, M.Z. Barel ^{[118](#)}, T. Barillari ^{[113](#)},
M.-S. Barisits ^{[37](#)}, T. Barklow ^{[150](#)}, P. Baron ^{[126](#)}, D.A. Baron Moreno ^{[104](#)}, A. Baroncelli ^{[63](#)},
A.J. Barr ^{[130](#)}, J.D. Barr ^{[99](#)}, F. Barreiro ^{[102](#)}, J. Barreiro Guimarães da Costa ^{[14](#)},
M.G. Barros Teixeira ^{[134a](#)}, S. Barsov ^{[39](#)}, F. Bartels ^{[64a](#)}, R. Bartoldus ^{[150](#)}, A.E. Barton ^{[94](#)},
P. Bartos ^{[29a](#)}, A. Basan ^{[103](#)}, M. Baselga ^{[50](#)}, S. Bashiri ^{[88](#)}, A. Bassalat ^{[67,b](#)}, M.J. Basso ^{[163a](#)},
S. Bataju ^{[46](#)}, R. Bate ^{[171](#)}, R.L. Bates ^{[60](#)}, S. Batlamous ^{[102](#)}, M. Battaglia ^{[140](#)}, D. Battulga ^{[19](#)},
M. Baue ^{[76a,76b](#)}, M. Bauer ^{[80](#)}, P. Bauer ^{[25](#)}, L.T. Bayer ^{[49](#)}, L.T. Bazzano Hurrell ^{[31](#)},
J.B. Beacham ^{[113](#)}, T. Beau ^{[131](#)}, J.Y. Beaucamp ^{[93](#)}, P.H. Beauchemin ^{[165](#)}, P. Bechtle ^{[25](#)},
H.P. Beck ^{[20,r](#)}, K. Becker ^{[174](#)}, A.J. Beddall ^{[83](#)}, V.A. Bednyakov ^{[40](#)}, C.P. Bee ^{[152](#)},

L.J. Beemster¹⁶, M. Begalli^{84d}, M. Begel³⁰, J.K. Behr⁴⁹, J.F. Beirer³⁷, F. Beisiegel²⁵,
 M. Belfkir^{120b}, G. Bella¹⁵⁸, L. Bellagamba^{24b}, A. Bellerive³⁵, P. Bellos²¹,
 K. Beloborodov³⁹, D. Benckekroun^{36a}, F. Bendebba^{36a}, Y. Benhammou¹⁵⁸,
 K.C. Benkendorfer⁶², L. Beresford⁴⁹, M. Beretta⁵⁴, E. Bergeaas Kuutmann¹⁶⁸, N. Berger⁴,
 B. Bergmann¹³⁶, J. Beringer^{18a}, G. Bernardi⁵, C. Bernius¹⁵⁰, F.U. Bernlochner²⁵,
 F. Bernon³⁷, A. Berrocal Guardia¹³, T. Berry⁹⁸, P. Berta¹³⁷, A. Berthold⁵¹,
 R. Bertrand¹⁰⁵, S. Bethke¹¹³, A. Betti^{76a,76b}, A.J. Bevan⁹⁷, L. Bezio⁵⁷, N.K. Bhalla⁵⁵,
 S. Bharghava¹¹³, S. Bhatta¹⁵², D.S. Bhattacharya¹⁷³, P. Bhattarai¹⁵⁰, Z.M. Bhatti¹²¹,
 K.D. Bhide⁵⁵, V.S. Bhopatkar¹²⁵, R.M. Bianchi¹³³, G. Bianco^{24b,24a}, O. Biebel¹¹²,
 M. Biglietti^{78a}, C.S. Billingsley⁴⁶, Y. Bimgdi^{36f}, M. Bindi⁵⁶, A. Bingham¹⁷⁸, A. Bingul^{22b},
 C. Bini^{76a,76b}, G.A. Bird³³, M. Birman¹⁷⁶, M. Biros¹³⁷, S. Biryukov¹⁵³, T. Bisanz⁵⁰,
 E. Bisceglie^{24b,24a}, J.P. Biswal¹³⁸, D. Biswas¹⁴⁸, I. Bloch⁴⁹, A. Blue⁶⁰,
 U. Blumenschein⁹⁷, J. Blumenthal¹⁰³, V.S. Bobrovnikov⁴⁰, M. Boehler⁵⁵, B. Boehm¹⁷³,
 D. Bogavac³⁷, A.G. Bogdanchikov³⁹, L.S. Boggia¹³¹, V. Boisvert⁹⁸, P. Bokan³⁷,
 T. Bold^{87a}, M. Bomben⁵, M. Bona⁹⁷, M. Boonekamp¹³⁹, A.G. Borbély⁶⁰,
 I.S. Bordulev³⁹, G. Borissov⁹⁴, D. Bortoletto¹³⁰, D. Boscherini^{24b}, M. Bosman¹³,
 K. Bouaouda^{36a}, N. Bouchhar¹⁷⁰, L. Boudet⁴, J. Boudreau¹³³, E.V. Bouhova-Thacker⁹⁴,
 D. Boumediene⁴², R. Bouquet^{58b,58a}, A. Boveia¹²³, J. Boyd³⁷, D. Boye³⁰, I.R. Boyko⁴⁰,
 L. Bozianu⁵⁷, J. Bracinik²¹, N. Brahimi⁴, G. Brandt¹⁷⁸, O. Brandt³³, B. Brau¹⁰⁶,
 J.E. Brau¹²⁷, R. Brenner¹⁷⁶, L. Brenner¹¹⁸, R. Brenner¹⁶⁸, S. Bressler¹⁷⁶,
 G. Brianti^{79a,79b}, D. Britton⁶⁰, D. Britzger¹¹³, I. Brock²⁵, R. Brock¹¹⁰, G. Brooijmans⁴³,
 A.J. Brooks⁶⁹, E.M. Brooks^{163b}, E. Brost³⁰, L.M. Brown^{172,163a}, L.E. Bruce⁶²,
 T.L. Bruckler¹³⁰, P.A. Bruckman de Renstrom⁸⁸, B. Brüers⁴⁹, A. Bruni^{24b}, G. Bruni^{24b},
 D. Brunner^{48a,48b}, M. Bruschi^{24b}, N. Bruscino^{76a,76b}, T. Buanes¹⁷, Q. Buat¹⁴³,
 D. Buchin¹¹³, A.G. Buckley⁶⁰, O. Bulekov³⁹, B.A. Bullard¹⁵⁰, S. Burdin⁹⁵,
 C.D. Burgard⁵⁰, A.M. Burger³⁷, B. Burghgrave⁸, O. Burlayenko⁵⁵, J. Burleson¹⁶⁹,
 J.T.P. Burr³³, J.C. Burzynski¹⁴⁹, E.L. Busch⁴³, V. Büscher¹⁰³, P.J. Bussey⁶⁰,
 J.M. Butler²⁶, C.M. Buttar⁶⁰, J.M. Butterworth⁹⁹, W. Buttinger¹³⁸,
 C.J. Buxo Vazquez¹¹⁰, A.R. Buzykaev⁴⁰, S. Cabrera Urbán¹⁷⁰, L. Cadamuro⁶⁷,
 D. Caforio⁵⁹, H. Cai¹³³, Y. Cai^{24b,115c,24a}, Y. Cai^{115a}, V.M.M. Cairo³⁷, O. Cakir^{3a},
 N. Calace³⁷, P. Calafiura^{18a}, G. Calderini¹³¹, P. Calfayan³⁵, G. Callea⁶⁰, L.P. Caloba^{84b},
 D. Calvet⁴², S. Calvet⁴², R. Camacho Toro¹³¹, S. Camarda³⁷, D. Camarero Munoz²⁷,
 P. Camarri^{77a,77b}, M.T. Camerlingo^{73a,73b}, C. Camincher¹⁷², M. Campanelli⁹⁹,
 A. Camplani⁴⁴, V. Canale^{73a,73b}, A.C. Canbay^{3a}, E. Canonero⁹⁸, J. Cantero¹⁷⁰,
 Y. Cao¹⁶⁹, F. Capocasa²⁷, M. Capua^{45b,45a}, A. Carbone^{72a,72b}, R. Cardarelli^{77a},
 J.C.J. Cardenas⁸, M.P. Cardiff²⁷, G. Carducci^{45b,45a}, T. Carli³⁷, G. Carlino^{73a},
 J.I. Carlotto¹³, B.T. Carlson^{133,t}, E.M. Carlson¹⁷², J. Carmignani⁹⁵, L. Carminati^{72a,72b},
 A. Carnelli⁴, M. Carnesale³⁷, S. Caron¹¹⁷, E. Carquin^{141f}, I.B. Carr¹⁰⁸, S. Carrá^{72a},
 G. Carratta^{24b,24a}, A.M. Carroll¹²⁷, M.P. Casado^{13,i}, M. Caspar⁴⁹, F.L. Castillo⁴,
 L. Castillo Garcia¹³, V. Castillo Gimenez¹⁷⁰, N.F. Castro^{134a,134e}, A. Catinaccio³⁷,
 J.R. Catmore¹²⁹, T. Cavaliere⁴, V. Cavaliere³⁰, L.J. Caviedes Betancourt^{23b},
 Y.C. Cekmecelioglu⁴⁹, E. Celebi⁸³, S. Cella³⁷, V. Cepaitis⁵⁷, K. Cerny¹²⁶,
 A.S. Cerqueira^{84a}, A. Cerri^{75a,75b}, L. Cerrito^{77a,77b}, F. Cerutti^{18a}, B. Cervato^{72a,72b},

A. Cervelli [ID](#)^{24b}, G. Cesarini [ID](#)⁵⁴, S.A. Cetin [ID](#)⁸³, P.M. Chabrilat [ID](#)¹³¹, J. Chan [ID](#)^{18a},
W.Y. Chan [ID](#)¹⁶⁰, J.D. Chapman [ID](#)³³, E. Chapon [ID](#)¹³⁹, B. Chargeishvili [ID](#)^{156b}, D.G. Charlton [ID](#)²¹,
C. Chauhan [ID](#)¹³⁷, Y. Che [ID](#)^{115a}, S. Chekanov [ID](#)⁶, S.V. Chekulaev [ID](#)^{163a}, G.A. Chelkov [ID](#)^{40,a},
B. Chen [ID](#)¹⁵⁸, B. Chen [ID](#)¹⁷², H. Chen [ID](#)^{115a}, H. Chen [ID](#)³⁰, J. Chen [ID](#)^{145a}, J. Chen [ID](#)¹⁴⁹,
M. Chen [ID](#)¹³⁰, S. Chen [ID](#)⁹⁰, S.J. Chen [ID](#)^{115a}, X. Chen [ID](#)^{145a}, X. Chen [ID](#)^{15,ai}, Z. Chen [ID](#)⁶³,
C.L. Cheng [ID](#)¹⁷⁷, H.C. Cheng [ID](#)^{65a}, S. Cheong [ID](#)¹⁵⁰, A. Cheplakov [ID](#)⁴⁰, E. Cheremushkina [ID](#)⁴⁹,
E. Cherepanova [ID](#)¹¹⁸, R. Cherkaoui El Moursli [ID](#)^{36e}, E. Cheu [ID](#)⁷, K. Cheung [ID](#)⁶⁶, L. Chevalier [ID](#)¹³⁹,
V. Chiarella [ID](#)⁵⁴, G. Chiarelli [ID](#)^{75a}, N. Chiedde [ID](#)¹⁰⁵, G. Chiodini [ID](#)^{71a}, A.S. Chisholm [ID](#)²¹,
A. Chitan [ID](#)^{28b}, M. Chitishvili [ID](#)¹⁷⁰, M.V. Chizhov [ID](#)^{40,u}, K. Choi [ID](#)¹¹, Y. Chou [ID](#)¹⁴³,
E.Y.S. Chow [ID](#)¹¹⁷, K.L. Chu [ID](#)¹⁷⁶, M.C. Chu [ID](#)^{65a}, X. Chu [ID](#)^{14,115c}, Z. Chubinidze [ID](#)⁵⁴,
J. Chudoba [ID](#)¹³⁵, J.J. Chwastowski [ID](#)³⁸, D. Cieri [ID](#)¹¹³, K.M. Ciesla [ID](#)^{87a}, V. Cindro [ID](#)⁹⁶,
A. Ciocio [ID](#)^{18a}, F. Ciotto [ID](#)^{73a,73b}, Z.H. Citron [ID](#)¹⁷⁶, M. Citterio [ID](#)^{72a}, D.A. Ciubotaru [ID](#)^{28b},
A. Clark [ID](#)⁵⁷, P.J. Clark [ID](#)⁵³, N. Clarke Hall [ID](#)⁹⁹, C. Clarry [ID](#)¹⁶², S.E. Clawson [ID](#)⁴⁹,
C. Clement [ID](#)^{48a,48b}, Y. Coadou [ID](#)¹⁰⁵, M. Cobal [ID](#)^{70a,70c}, A. Coccaro [ID](#)^{58b}, R.F. Coelho Barrue [ID](#)^{134a},
R. Coelho Lopes De Sa [ID](#)¹⁰⁶, S. Coelli [ID](#)^{72a}, L.S. Colangeli [ID](#)¹⁶², B. Cole [ID](#)⁴³, P. Collado Soto [ID](#)¹⁰²,
J. Collot [ID](#)⁶¹, P. Conde Muiño [ID](#)^{134a,134g}, M.P. Connell [ID](#)^{34c}, S.H. Connell [ID](#)^{34c}, E.I. Conroy [ID](#)¹³⁰,
F. Conventi [ID](#)^{73a,ak}, H.G. Cooke [ID](#)²¹, A.M. Cooper-Sarkar [ID](#)¹³⁰, F.A. Corchia [ID](#)^{24b,24a},
A. Cordeiro Oudot Choi [ID](#)¹³¹, L.D. Corpe [ID](#)⁴², M. Corradi [ID](#)^{76a,76b}, F. Corriveau [ID](#)^{107,ac},
A. Cortes-Gonzalez [ID](#)¹⁹, M.J. Costa [ID](#)¹⁷⁰, F. Costanza [ID](#)⁴, D. Costanzo [ID](#)¹⁴⁶, B.M. Cote [ID](#)¹²³,
J. Couthures [ID](#)⁴, G. Cowan [ID](#)⁹⁸, K. Cranmer [ID](#)¹⁷⁷, L. Cremer [ID](#)⁵⁰, D. Cremonini [ID](#)^{24b,24a},
S. Crépé-Renaudin [ID](#)⁶¹, F. Crescioli [ID](#)¹³¹, T. Cresta [ID](#)^{74a,74b}, M. Cristinziani [ID](#)¹⁴⁸,
M. Cristoforetti [ID](#)^{79a,79b}, V. Croft [ID](#)¹¹⁸, J.E. Crosby [ID](#)¹²⁵, G. Crosetti [ID](#)^{45b,45a}, A. Cueto [ID](#)¹⁰²,
H. Cui [ID](#)⁹⁹, Z. Cui [ID](#)⁷, W.R. Cunningham [ID](#)⁶⁰, F. Curcio [ID](#)¹⁷⁰, J.R. Curran [ID](#)⁵³, P. Czodrowski [ID](#)³⁷,
M.J. Da Cunha Sargedas De Sousa [ID](#)^{58b,58a}, J.V. Da Fonseca Pinto [ID](#)^{84b}, C. Da Via [ID](#)¹⁰⁴,
W. Dabrowski [ID](#)^{87a}, T. Dado [ID](#)³⁷, S. Dahbi [ID](#)¹⁵⁵, T. Dai [ID](#)¹⁰⁹, D. Dal Santo [ID](#)²⁰, C. Dallapiccola [ID](#)¹⁰⁶,
M. Dam [ID](#)⁴⁴, G. D’amen [ID](#)³⁰, V. D’Amico [ID](#)¹¹², J. Damp [ID](#)¹⁰³, J.R. Dandoy [ID](#)³⁵, D. Dannheim [ID](#)³⁷,
M. Danninger [ID](#)¹⁴⁹, V. Dao [ID](#)¹⁵², G. Darbo [ID](#)^{58b}, S.J. Das [ID](#)³⁰, F. Dattola [ID](#)⁴⁹, S. D’Auria [ID](#)^{72a,72b},
A. D’Avanzo [ID](#)^{73a,73b}, T. Davidek [ID](#)¹³⁷, I. Dawson [ID](#)⁹⁷, H.A. Day-hall [ID](#)¹³⁶, K. De [ID](#)⁸,
C. De Almeida Rossi [ID](#)¹⁶², R. De Asmundis [ID](#)^{73a}, N. De Biase [ID](#)⁴⁹, S. De Castro [ID](#)^{24b,24a},
N. De Groot [ID](#)¹¹⁷, P. de Jong [ID](#)¹¹⁸, H. De la Torre [ID](#)¹¹⁹, A. De Maria [ID](#)^{115a}, A. De Salvo [ID](#)^{76a},
U. De Sanctis [ID](#)^{77a,77b}, F. De Santis [ID](#)^{71a,71b}, A. De Santo [ID](#)¹⁵³, J.B. De Vivie De Regie [ID](#)⁶¹,
J. Debevc [ID](#)⁹⁶, D.V. Dedovich [ID](#)⁴⁰, J. Degens [ID](#)⁹⁵, A.M. Deiana [ID](#)⁴⁶, J. Del Peso [ID](#)¹⁰²,
L. Delagrangé [ID](#)¹³¹, F. Deliot [ID](#)¹³⁹, C.M. Delitzsch [ID](#)⁵⁰, M. Della Pietra [ID](#)^{73a,73b}, D. Della Volpe [ID](#)⁵⁷,
A. Dell’Acqua [ID](#)³⁷, L. Dell’Asta [ID](#)^{72a,72b}, M. Delmastro [ID](#)⁴, C.C. Delogu [ID](#)¹⁰³, P.A. Delsart [ID](#)⁶¹,
S. Demers [ID](#)¹⁷⁹, M. Demichev [ID](#)⁴⁰, S.P. Denisov [ID](#)³⁹, H. Denizli [ID](#)^{22a,m}, L. D’Eramo [ID](#)⁴²,
D. Derendarz [ID](#)⁸⁸, F. Derue [ID](#)¹³¹, P. Dervan [ID](#)⁹⁵, K. Desch [ID](#)²⁵, C. Deutsch [ID](#)²⁵,
F.A. Di Bello [ID](#)^{58b,58a}, A. Di Ciaccio [ID](#)^{77a,77b}, L. Di Ciaccio [ID](#)⁴, A. Di Domenico [ID](#)^{76a,76b},
C. Di Donato [ID](#)^{73a,73b}, A. Di Girolamo [ID](#)³⁷, G. Di Gregorio [ID](#)³⁷, A. Di Luca [ID](#)^{79a,79b},
B. Di Micco [ID](#)^{78a,78b}, R. Di Nardo [ID](#)^{78a,78b}, K.F. Di Petrillo [ID](#)⁴¹, M. Diamantopoulou [ID](#)³⁵,
F.A. Dias [ID](#)¹¹⁸, T. Dias Do Vale [ID](#)¹⁴⁹, M.A. Diaz [ID](#)^{141a,141b}, A.R. Didenko [ID](#)⁴⁰, M. Didenko [ID](#)¹⁷⁰,
E.B. Diehl [ID](#)¹⁰⁹, S. Díez Cornell [ID](#)⁴⁹, C. Díez Pardos [ID](#)¹⁴⁸, C. Dimitriadi [ID](#)¹⁵¹, A. Dimitrievska [ID](#)²¹,
A. Dimri [ID](#)¹⁵², J. Dingfelder [ID](#)²⁵, T. Dingley [ID](#)¹³⁰, I-M. Dinu [ID](#)^{28b}, S.J. Dittmeier [ID](#)^{64b}, F. Dittus [ID](#)³⁷,
M. Divisek [ID](#)¹³⁷, B. Dixit [ID](#)⁹⁵, F. Djama [ID](#)¹⁰⁵, T. Djobava [ID](#)^{156b}, C. Doglioni [ID](#)^{104,101},

A. Dohnalova ^{[129a](#)}, Z. Dolezal ^{[137](#)}, K. Domijan ^{[187a](#)}, K.M. Dona ^{[41](#)}, M. Donadelli ^{[184d](#)},
B. Dong ^{[110](#)}, J. Donini ^{[42](#)}, A. D’Onofrio ^{[73a,73b](#)}, M. D’Onofrio ^{[95](#)}, J. Dopke ^{[138](#)}, A. Doria ^{[73a](#)},
N. Dos Santos Fernandes ^{[134a](#)}, P. Dougan ^{[104](#)}, M.T. Dova ^{[93](#)}, A.T. Doyle ^{[60](#)}, M.A. Draguet ^{[130](#)},
M.P. Drescher ^{[56](#)}, E. Dreyer ^{[176](#)}, I. Drivas-koulouris ^{[10](#)}, M. Drnevich ^{[121](#)}, M. Drozdova ^{[57](#)},
D. Du ^{[63](#)}, T.A. du Pree ^{[118](#)}, F. Dubinin ^{[40](#)}, M. Dubovsky ^{[29a](#)}, E. Duchovni ^{[176](#)},
G. Duckeck ^{[112](#)}, P.K. Duckett ^{[99](#)}, O.A. Ducu ^{[28b](#)}, D. Duda ^{[53](#)}, A. Dudarev ^{[37](#)}, E.R. Duden ^{[27](#)},
M. D’uffizi ^{[104](#)}, L. Dufлот ^{[67](#)}, M. Dührssen ^{[37](#)}, I. Duminica ^{[28g](#)}, A.E. Dumitriu ^{[28b](#)},
M. Dunford ^{[64a](#)}, S. Dungs ^{[50](#)}, K. Dunne ^{[48a,48b](#)}, A. Duperrin ^{[105](#)}, H. Duran Yildiz ^{[3a](#)},
M. Düren ^{[59](#)}, A. Durglishvili ^{[156b](#)}, D. Duvnjak ^{[35](#)}, B.L. Dwyer ^{[119](#)}, G.I. Dyckes ^{[18a](#)},
M. Dyndal ^{[87a](#)}, B.S. Dziedzic ^{[37](#)}, Z.O. Earnshaw ^{[153](#)}, G.H. Eberwein ^{[130](#)}, B. Eckerova ^{[29a](#)},
S. Eggebrecht ^{[56](#)}, E. Egidio Purcino De Souza ^{[84e](#)}, G. Eigen ^{[17](#)}, K. Einsweiler ^{[18a](#)},
T. Ekelof ^{[168](#)}, P.A. Ekman ^{[101](#)}, S. El Farkh ^{[36b](#)}, Y. El Ghazali ^{[63](#)}, H. El Jarrari ^{[37](#)},
A. El Moussaouy ^{[36a](#)}, V. Ellajosyula ^{[168](#)}, M. Ellert ^{[168](#)}, F. Ellinghaus ^{[178](#)}, N. Ellis ^{[37](#)},
J. Elmsheuser ^{[30](#)}, M. Elsayy ^{[120a](#)}, M. Elsing ^{[37](#)}, D. Emelianov ^{[138](#)}, Y. Enari ^{[85](#)}, I. Ene ^{[18a](#)},
S. Epari ^{[111](#)}, D. Ernani Martins Neto ^{[88](#)}, M. Errenst ^{[178](#)}, M. Escalier ^{[67](#)}, C. Escobar ^{[170](#)},
E. Etzion ^{[158](#)}, G. Evans ^{[134a,134b](#)}, H. Evans ^{[69](#)}, L.S. Evans ^{[98](#)}, A. Ezhilov ^{[39](#)},
S. Ezzarqtouni ^{[36a](#)}, F. Fabbri ^{[24b,24a](#)}, L. Fabbri ^{[24b,24a](#)}, G. Facini ^{[99](#)}, V. Fadeyev ^{[140](#)},
R.M. Fakhrutdinov ^{[39](#)}, D. Fakoudis ^{[103](#)}, S. Falciano ^{[76a](#)}, L.F. Falda Ulhoa Coelho ^{[134a](#)},
F. Fallavollita ^{[113](#)}, G. Falsetti ^{[45b,45a](#)}, J. Faltova ^{[137](#)}, C. Fan ^{[169](#)}, K.Y. Fan ^{[65b](#)}, Y. Fan ^{[14](#)},
Y. Fang ^{[14,115c](#)}, M. Fanti ^{[72a,72b](#)}, M. Faraj ^{[70a,70b](#)}, Z. Farazpay ^{[100](#)}, A. Farbin ^{[8](#)}, A. Farilla ^{[78a](#)},
T. Farooque ^{[110](#)}, J.N. Farr ^{[179](#)}, S.M. Farrington ^{[138,53](#)}, F. Fassi ^{[36e](#)}, D. Fassouliotis ^{[9](#)},
L. Fayard ^{[67](#)}, P. Federic ^{[137](#)}, P. Federicova ^{[135](#)}, O.L. Fedin ^{[39,a](#)}, M. Feickert ^{[177](#)},
L. Feligioni ^{[105](#)}, D.E. Fellers ^{[18a](#)}, C. Feng ^{[144a](#)}, Z. Feng ^{[118](#)}, M.J. Fenton ^{[166](#)}, L. Ferencz ^{[49](#)},
B. Fernandez Barbadillo ^{[94](#)}, P. Fernandez Martinez ^{[68](#)}, M.J.V. Fernoux ^{[105](#)}, J. Ferrando ^{[94](#)},
A. Ferrari ^{[168](#)}, P. Ferrari ^{[118,117](#)}, R. Ferrari ^{[74a](#)}, D. Ferrere ^{[57](#)}, C. Ferretti ^{[109](#)}, M.P. Fewell ^{[1](#)},
D. Fiacco ^{[76a,76b](#)}, F. Fiedler ^{[103](#)}, P. Fiedler ^{[136](#)}, S. Filimonov ^{[40](#)}, A. Filipčić ^{[96](#)},
E.K. Filmer ^{[163a](#)}, F. Filthaut ^{[117](#)}, M.C.N. Fiolhais ^{[134a,134c,c](#)}, L. Fiorini ^{[170](#)}, W.C. Fisher ^{[110](#)},
T. Fitschen ^{[104](#)}, P.M. Fitzhugh ^{[139](#)}, I. Fleck ^{[148](#)}, P. Fleischmann ^{[109](#)}, T. Flick ^{[178](#)},
M. Flores ^{[34d,ag](#)}, L.R. Flores Castillo ^{[65a](#)}, L. Flores Sanz De Acedo ^{[37](#)}, F.M. Follega ^{[79a,79b](#)},
N. Fomin ^{[33](#)}, J.H. Foo ^{[162](#)}, A. Formica ^{[139](#)}, A.C. Forti ^{[104](#)}, E. Fortin ^{[37](#)}, A.W. Fortman ^{[18a](#)},
L. Fountas ^{[9,j](#)}, D. Fournier ^{[67](#)}, H. Fox ^{[94](#)}, P. Francavilla ^{[75a,75b](#)}, S. Francescato ^{[62](#)},
S. Franchellucci ^{[57](#)}, M. Franchini ^{[24b,24a](#)}, S. Franchino ^{[64a](#)}, D. Francis ^{[37](#)}, L. Franco ^{[117](#)},
V. Franco Lima ^{[37](#)}, L. Franconi ^{[49](#)}, M. Franklin ^{[62](#)}, G. Frattari ^{[27](#)}, Y.Y. Frid ^{[158](#)}, J. Friend ^{[60](#)},
N. Fritzsche ^{[37](#)}, A. Froch ^{[57](#)}, D. Froidevaux ^{[37](#)}, J.A. Frost ^{[130](#)}, Y. Fu ^{[110](#)},
S. Fuenzalida Garrido ^{[141f](#)}, M. Fujimoto ^{[105](#)}, K.Y. Fung ^{[65a](#)}, E. Furtado De Simas Filho ^{[84e](#)},
M. Furukawa ^{[160](#)}, J. Fuster ^{[170](#)}, A. Gaa ^{[56](#)}, A. Gabrielli ^{[24b,24a](#)}, A. Gabrielli ^{[162](#)}, P. Gadow ^{[37](#)},
G. Gagliardi ^{[58b,58a](#)}, L.G. Gagnon ^{[18a](#)}, S. Gaid ^{[89b](#)}, S. Galantzan ^{[158](#)}, J. Gallagher ^{[1](#)},
E.J. Gallas ^{[130](#)}, A.L. Gallen ^{[168](#)}, B.J. Gallop ^{[138](#)}, K.K. Gan ^{[123](#)}, S. Ganguly ^{[160](#)}, Y. Gao ^{[53](#)},
A. Garabaglu ^{[143](#)}, F.M. Garay Walls ^{[141a,141b](#)}, B. Garcia ^{[30](#)}, C. García ^{[170](#)}, A. Garcia Alonso ^{[118](#)},
A.G. Garcia Caffaro ^{[179](#)}, J.E. García Navarro ^{[170](#)}, M. Garcia-Sciveres ^{[18a](#)}, G.L. Gardner ^{[132](#)},
R.W. Gardner ^{[41](#)}, N. Garelli ^{[165](#)}, R.B. Garg ^{[150](#)}, J.M. Gargan ^{[53](#)}, C.A. Garner ^{[162](#)},
C.M. Garvey ^{[34a](#)}, V.K. Gassmann ^{[165](#)}, G. Gaudio ^{[74a](#)}, V. Gautam ^{[13](#)}, P. Gauzzi ^{[76a,76b](#)},
J. Gavranovic ^{[96](#)}, I.L. Gavrilenko ^{[134a](#)}, A. Gavriluk ^{[39](#)}, C. Gay ^{[171](#)}, G. Gaycken ^{[127](#)},

E.N. Gaziz ¹⁰, A. Gekow¹²³, C. Gemme ^{58b}, M.H. Genest ⁶¹, A.D. Gentry ¹¹⁶, S. George ⁹⁸, W.F. George ²¹, T. Geralis ⁴⁷, A.A. Gerwin ¹²⁴, P. Gessinger-Befurt ³⁷, M.E. Geyik ¹⁷⁸, M. Ghani ¹⁷⁴, K. Ghorbanian ⁹⁷, A. Ghosal ¹⁴⁸, A. Ghosh ¹⁶⁶, A. Ghosh ⁷, B. Giacobbe ^{24b}, S. Giagu ^{76a,76b}, T. Giani ¹¹⁸, A. Giannini ⁶³, S.M. Gibson ⁹⁸, M. Gignac ¹⁴⁰, D.T. Gil ^{87b}, A.K. Gilbert ^{87a}, B.J. Gilbert ⁴³, D. Gillberg ³⁵, G. Gilles ¹¹⁸, L. Ginabat ¹³¹, D.M. Gingrich ^{2,aj}, M.P. Giordani ^{70a,70c}, P.F. Giraud ¹³⁹, G. Giugliarelli ^{70a,70c}, D. Giugni ^{72a}, F. Giuli ^{77a,77b}, I. Gkialas ^{9,j}, L.K. Gladilin ³⁹, C. Glasman ¹⁰², G. Glemža ⁴⁹, M. Glisic ¹²⁷, I. Gnesi ^{45b}, Y. Go ³⁰, M. Goblirsch-Kolb ³⁷, B. Gocke ⁵⁰, D. Godin¹¹¹, B. Gokturk ^{22a}, S. Goldfarb ¹⁰⁸, T. Golling ⁵⁷, M.G.D. Gololo ^{34c}, D. Golubkov ³⁹, J.P. Gombas ¹¹⁰, A. Gomes ^{134a,134b}, G. Gomes Da Silva ¹⁴⁸, A.J. Gomez Delegido ¹⁷⁰, R. Gonçalo ^{134a}, L. Gonella ²¹, A. Gongadze ^{156c}, F. Gonnella ²¹, J.L. Gonski ¹⁵⁰, R.Y. González Andana ⁵³, S. González de la Hoz ¹⁷⁰, C. Gonzalez Renteria ^{18a}, M.V. Gonzalez Rodrigues ⁴⁹, R. Gonzalez Suarez ¹⁶⁸, S. Gonzalez-Sevilla ⁵⁷, L. Goossens ³⁷, B. Gorini ³⁷, E. Gorini ^{71a,71b}, A. Gorišek ⁹⁶, T.C. Gosart ¹³², A.T. Goshaw ⁵², M.I. Gostkin ⁴⁰, S. Goswami ¹²⁵, C.A. Gottardo ³⁷, S.A. Gotz ¹¹², M. Goughri ^{36b}, A.G. Goussiou ¹⁴³, N. Govender ^{34c}, R.P. Grabarczyk ¹³⁰, I. Grabowska-Bold ^{87a}, K. Graham ³⁵, E. Gramstad ¹²⁹, S. Grancagnolo ^{71a,71b}, C.M. Grant^{1,139}, P.M. Gravila ^{28f}, F.G. Gravili ^{71a,71b}, H.M. Gray ^{18a}, M. Greco ¹¹³, M.J. Green ¹, C. Grefe ²⁵, A.S. Grefsrud ¹⁷, I.M. Gregor ⁴⁹, K.T. Greif ¹⁶⁶, P. Grenier ¹⁵⁰, S.G. Grewe¹¹³, A.A. Grillo ¹⁴⁰, K. Grimm ³², S. Grinstein ^{13,y}, J.-F. Grivaz ⁶⁷, E. Gross ¹⁷⁶, J. Grosse-Knetter ⁵⁶, L. Guan ¹⁰⁹, G. Guerrieri ³⁷, R. Guevara ¹²⁹, R. Gugel ¹⁰³, J.A.M. Guhit ¹⁰⁹, A. Guida ¹⁹, E. Guilloton ¹⁷⁴, S. Guindon ³⁷, F. Guo ^{14,115c}, J. Guo ^{145a}, L. Guo ⁴⁹, L. Guo ^{115b,w}, Y. Guo ¹⁰⁹, A. Gupta ⁵⁰, R. Gupta ¹³³, S. Gurbuz ²⁵, S.S. Gurdasani ⁴⁹, G. Gustavino ^{76a,76b}, P. Gutierrez ¹²⁴, L.F. Gutierrez Zagazeta ¹³², M. Gutsche ⁵¹, C. Gutscheow ⁹⁹, C. Gwenlan ¹³⁰, C.B. Gwilliam ⁹⁵, E.S. Haaland ¹²⁹, A. Haas ¹²¹, M. Habedank ⁶⁰, C. Haber ^{18a}, H.K. Hadavand ⁸, A. Haddad ⁴², A. Hadeef ⁵¹, A.I. Hagan ⁹⁴, J.J. Hahn ¹⁴⁸, E.H. Haines ⁹⁹, M. Haleem ¹⁷³, J. Haley ¹²⁵, G.D. Hallewell ¹⁰⁵, L. Halser ²⁰, K. Hamano ¹⁷², M. Hamer ²⁵, S.E.D. Hammoud ⁶⁷, E.J. Hampshire ⁹⁸, J. Han ^{144a}, L. Han ^{115a}, L. Han ⁶³, S. Han ^{18a}, K. Hanagaki ⁸⁵, M. Hance ¹⁴⁰, D.A. Hangal ⁴³, H. Hanif ¹⁴⁹, M.D. Hank ¹³², J.B. Hansen ⁴⁴, P.H. Hansen ⁴⁴, D. Harada ⁵⁷, T. Harenberg ¹⁷⁸, S. Harkusha ¹⁸⁰, M.L. Harris ¹⁰⁶, Y.T. Harris ²⁵, J. Harrison ¹³, N.M. Harrison ¹²³, P.F. Harrison¹⁷⁴, N.M. Hartman ¹¹³, N.M. Hartmann ¹¹², R.Z. Hasan ^{98,138}, Y. Hasegawa ¹⁴⁷, F. Haslbeck ¹³⁰, S. Hassan ¹⁷, R. Hauser ¹¹⁰, M. Haviernik ¹³⁷, C.M. Hawkes ²¹, R.J. Hawkins ³⁷, Y. Hayashi ¹⁶⁰, D. Hayden ¹¹⁰, C. Hayes ¹⁰⁹, R.L. Hayes ¹¹⁸, C.P. Hays ¹³⁰, J.M. Hays ⁹⁷, H.S. Hayward ⁹⁵, F. He ⁶³, M. He ^{14,115c}, Y. He ⁴⁹, Y. He ⁹⁹, N.B. Heatley ⁹⁷, V. Hedberg ¹⁰¹, A.L. Heggelund ¹²⁹, C. Heidegger ⁵⁵, K.K. Heidegger ⁵⁵, J. Heilman ³⁵, S. Heim ⁴⁹, T. Heim ^{18a}, J.G. Heinlein ¹³², J.J. Heinrich ¹²⁷, L. Heinrich ^{113,ah}, J. Hejbal ¹³⁵, A. Held ¹⁷⁷, S. Hellesund ¹⁷, C.M. Helling ¹⁷¹, S. Hellman ^{48a,48b}, L. Henkelmann ³³, A.M. Henriques Correia³⁷, H. Herde ¹⁰¹, Y. Hernández Jiménez ¹⁵², L.M. Herrmann ²⁵, T. Herrmann ⁵¹, G. Herten ⁵⁵, R. Hertenberger ¹¹², L. Hervas ³⁷, M.E. Hespings ¹⁰³, N.P. Hessey ^{163a}, J. Hessler ¹¹³, M. Hidaoui ^{36b}, N. Hidic ¹³⁷, E. Hill ¹⁶², S.J. Hillier ²¹, J.R. Hinds ¹¹⁰, F. Hinterkeuser ²⁵, M. Hirose ¹²⁸, S. Hirose ¹⁶⁴, D. Hirschbuehl ¹⁷⁸,

T.G. Hitchings [104](#), B. Hiti [96](#), J. Hobbs [152](#), R. Hobincu [28e](#), N. Hod [176](#),
M.C. Hodgkinson [146](#), B.H. Hodgkinson [130](#), A. Hoecker [37](#), D.D. Hofer [109](#), J. Hofer [170](#),
M. Holzbock [37](#), L.B.A.H. Hommels [33](#), V. Homsak [130](#), B.P. Honan [104](#), J.J. Hong [69](#),
J. Hong [145a](#), T.M. Hong [133](#), B.H. Hooberman [169](#), W.H. Hopkins [6](#), M.C. Hoppesch [169](#),
Y. Horii [114](#), M.E. Horstmann [113](#), S. Hou [155](#), M.R. Housenga [169](#), A.S. Howard [96](#),
J. Howarth [60](#), J. Hoya [6](#), M. Hrabovsky [126](#), T. Hryn'ova [4](#), P.J. Hsu [66](#), S.-C. Hsu [143](#),
T. Hsu [67](#), M. Hu [18a](#), Q. Hu [63](#), S. Huang [33](#), X. Huang [14,115c](#), Y. Huang [137](#),
Y. Huang [115b](#), Y. Huang [103](#), Y. Huang [14](#), Z. Huang [104](#), Z. Hubacek [136](#), M. Huebner [25](#),
F. Huegging [25](#), T.B. Huffman [130](#), M. Hufnagel Maranha De Faria [84a](#), C.A. Hugli [49](#),
M. Huhtinen [37](#), S.K. Huiberts [17](#), R. Hulsken [107](#), C.E. Hultquist [18a](#), N. Huseynov [12,g](#),
J. Huston [110](#), J. Huth [62](#), R. Hyneman [7](#), G. Iacobucci [57](#), G. Iakovidis [30](#),
L. Iconomidou-Fayard [67](#), J.P. Iddon [37](#), P. Iengo [73a,73b](#), R. Iguchi [160](#), Y. Iiyama [160](#),
T. Iizawa [130](#), Y. Ikegami [85](#), D. Iliadis [159](#), N. Ilic [162](#), H. Imam [84c](#), G. Inacio Goncalves [84d](#),
S.A. Infante Cabanas [141c](#), T. Ingebretsen Carlson [48a,48b](#), J.M. Inglis [97](#), G. Introzzi [74a,74b](#),
M. Iodice [78a](#), V. Ippolito [76a,76b](#), R.K. Irwin [95](#), M. Ishino [160](#), W. Islam [177](#), C. Issever [19](#),
S. Istin [22a,ao](#), K. Itabashi [85](#), H. Ito [175](#), R. Iuppa [79a,79b](#), A. Ivina [176](#), V. Izzo [73a](#),
P. Jacka [135](#), P. Jackson [1](#), P. Jain [49](#), K. Jakobs [55](#), T. Jakoubek [176](#), J. Jamieson [60](#),
W. Jang [160](#), S. Jankovych [137](#), M. Javurkova [106](#), P. Jawahar [104](#), L. Jeanty [127](#),
J. Jejelava [156a,af](#), P. Jenni [55,f](#), C.E. Jessiman [35](#), C. Jia [144a](#), H. Jia [171](#), J. Jia [152](#),
X. Jia [14,115c](#), Z. Jia [115a](#), C. Jiang [53](#), Q. Jiang [65b](#), S. Jiggins [49](#), J. Jimenez Pena [13](#),
S. Jin [115a](#), A. Jinaru [28b](#), O. Jinnouchi [142](#), P. Johansson [146](#), K.A. Johns [7](#),
J.W. Johnson [140](#), F.A. Jolly [49](#), D.M. Jones [153](#), E. Jones [49](#), K.S. Jones [8](#), P. Jones [33](#),
R.W.L. Jones [94](#), T.J. Jones [95](#), H.L. Joos [56,37](#), R. Joshi [123](#), J. Jovicevic [16](#), X. Ju [18a](#),
J.J. Junggeburth [37](#), T. Junkermann [64a](#), A. Juste Rozas [13,y](#), M.K. Juzek [88](#), S. Kabana [141e](#),
A. Kaczmarska [88](#), M. Kado [113](#), H. Kagan [123](#), M. Kagan [150](#), A. Kahn [132](#), C. Kahra [103](#),
T. Kaji [160](#), E. Kajomovitz [157](#), N. Kakati [176](#), N. Kakoty [13](#), I. Kalaitzidou [55](#), S. Kandel [8](#),
N.J. Kang [140](#), D. Kar [34g](#), K. Karava [130](#), E. Karentzos [25](#), O. Karkout [118](#), S.N. Karpov [40](#),
Z.M. Karpova [40](#), V. Kartvelishvili [94](#), A.N. Karyukhin [39](#), E. Kasimi [159](#), J. Katzy [49](#),
S. Kaur [35](#), K. Kawade [147](#), M.P. Kawale [124](#), C. Kawamoto [90](#), T. Kawamoto [63](#), E.F. Kay [37](#),
F.I. Kaya [165](#), S. Kazakos [110](#), V.F. Kazanin [39](#), Y. Ke [152](#), J.M. Keaveney [34a](#), R. Keeler [172](#),
G.V. Kehris [62](#), J.S. Keller [35](#), J.J. Kempster [153](#), O. Kepka [135](#), J. Kerr [163b](#),
B.P. Kerridge [138](#), B.P. Kerševan [96](#), L. Keszeghova [29a](#), R.A. Khan [133](#), A. Khanov [125](#),
A.G. Kharlamov [39](#), T. Kharlamova [39](#), E.E. Khoda [143](#), M. Kholodenko [134a](#), T.J. Khoo [19](#),
G. Khorauli [173](#), J. Khubua [156b,*](#), Y.A.R. Khwaira [131](#), B. Kibirige [34g](#), D. Kim [6](#),
D.W. Kim [48a,48b](#), Y.K. Kim [41](#), N. Kimura [99](#), M.K. Kingston [56](#), A. Kirchhoff [56](#),
C. Kirfel [25](#), F. Kirfel [25](#), J. Kirk [138](#), A.E. Kiryunin [113](#), S. Kita [164](#), C. Kitsaki [10](#),
O. Kivernyk [25](#), M. Klassen [165](#), C. Klein [35](#), L. Klein [173](#), M.H. Klein [46](#), S.B. Klein [57](#),
U. Klein [95](#), A. Klimentov [30](#), T. Klioutchnikova [37](#), P. Kluit [118](#), S. Kluth [113](#),
E. Kneringer [80](#), T.M. Knight [162](#), A. Knue [50](#), M. Kobel [51](#), D. Kobylanskii [176](#),
S.F. Koch [130](#), M. Kocian [150](#), P. Kodyš [137](#), D.M. Koeck [127](#), P.T. Koenig [25](#), T. Koffas [35](#),
O. Kolay [51](#), I. Koletsou [4](#), T. Komarek [88](#), K. Köneke [56](#), A.X.Y. Kong [1](#), T. Kono [122](#),
N. Konstantinidis [99](#), P. Kontaxakis [57](#), B. Konya [101](#), R. Kopeliansky [43](#), S. Koperny [87a](#),
K. Korcyl [88](#), K. Kordas [159,e](#), A. Korn [99](#), S. Korn [56](#), I. Korolkov [13](#), N. Korotkova [39](#),

B. Kortman ^{[118](#)}, O. Kortner ^{[113](#)}, S. Kortner ^{[113](#)}, W.H. Kostecka ^{[119](#)}, M. Kostov ^{[29a](#)},
V.V. Kostyukhin ^{[148](#)}, A. Kotskechagia ^{[37](#)}, A. Kotwal ^{[52](#)}, A. Koulouris ^{[37](#)},
A. Kourkumeli-Charalampidi ^{[74a,74b](#)}, C. Kourkumelis ^{[9](#)}, E. Kourlitis ^{[113](#)}, O. Kovanda ^{[127](#)},
R. Kowalewski ^{[172](#)}, W. Kozanecki ^{[127](#)}, A.S. Kozhin ^{[39](#)}, V.A. Kramarenko ^{[39](#)}, G. Kramberger ^{[96](#)},
P. Kramer ^{[25](#)}, M.W. Krasny ^{[131](#)}, A. Krasznahorkay ^{[106](#)}, A.C. Kraus ^{[119](#)}, J.W. Kraus ^{[178](#)},
J.A. Kremer ^{[49](#)}, N.B. Krengel ^{[148](#)}, T. Kresse ^{[51](#)}, L. Kretschmann ^{[178](#)}, J. Kretzschmar ^{[95](#)},
K. Kreul ^{[19](#)}, P. Krieger ^{[162](#)}, K. Krizka ^{[21](#)}, K. Kroeninger ^{[50](#)}, H. Kroha ^{[113](#)}, J. Kroll ^{[135](#)},
J. Kroll ^{[132](#)}, K.S. Krowpman ^{[110](#)}, U. Kruchonak ^{[40](#)}, H. Krüger ^{[25](#)}, N. Krumnack ^{[82](#)},
M.C. Kruse ^{[52](#)}, O. Kuchinskaia ^{[40](#)}, S. Kuday ^{[3a](#)}, S. Kuehn ^{[37](#)}, R. Kuesters ^{[55](#)}, T. Kuhl ^{[49](#)},
V. Kukhtin ^{[40](#)}, Y. Kulchitsky ^{[40](#)}, S. Kuleshov ^{[141d,141b](#)}, M. Kumar ^{[34g](#)}, N. Kumari ^{[49](#)},
P. Kumari ^{[163b](#)}, A. Kupco ^{[135](#)}, T. Kupfer ^{[50](#)}, A. Kupich ^{[39](#)}, O. Kuprash ^{[55](#)}, H. Kurashige ^{[86](#)},
L.L. Kurchaninov ^{[163a](#)}, O. Kurdysh ^{[4](#)}, Y.A. Kurochkin ^{[38](#)}, A. Kurova ^{[39](#)}, M. Kuze ^{[142](#)},
A.K. Kvam ^{[106](#)}, J. Kvita ^{[126](#)}, N.G. Kyriacou ^{[109](#)}, L.A.O. Laatu ^{[105](#)}, C. Lacasta ^{[170](#)},
F. Lacava ^{[76a,76b](#)}, H. Lacker ^{[19](#)}, D. Lacour ^{[131](#)}, N.N. Lad ^{[99](#)}, E. Ladygin ^{[40](#)}, A. Lafarge ^{[42](#)},
B. Laforge ^{[131](#)}, T. Lagouri ^{[179](#)}, F.Z. Lahbabi ^{[36a](#)}, S. Lai ^{[56](#)}, J.E. Lambert ^{[172](#)}, S. Lammers ^{[69](#)},
W. Lampl ^{[7](#)}, C. Lampoudis ^{[159,e](#)}, G. Lamprinoudis ^{[103](#)}, A.N. Lancaster ^{[119](#)}, E. Lançon ^{[30](#)},
U. Landgraf ^{[55](#)}, M.P.J. Landon ^{[97](#)}, V.S. Lang ^{[55](#)}, O.K.B. Langrekken ^{[129](#)}, A.J. Lankford ^{[166](#)},
F. Lanni ^{[37](#)}, K. Lantzsch ^{[25](#)}, A. Lanza ^{[74a](#)}, M. Lanzac Berrocal ^{[170](#)}, J.F. Laporte ^{[139](#)},
T. Lari ^{[72a](#)}, D. Larsen ^{[17](#)}, F. Lasagni Manghi ^{[24b](#)}, M. Lassnig ^{[37](#)}, V. Latonova ^{[135](#)},
S.D. Lawlor ^{[146](#)}, Z. Lawrence ^{[104](#)}, R. Lazaridou ^{[174](#)}, M. Lazzaroni ^{[72a,72b](#)}, H.D.M. Le ^{[110](#)},
E.M. Le Boulicaut ^{[179](#)}, L.T. Le Pottier ^{[18a](#)}, B. Leban ^{[24b,24a](#)}, M. LeBlanc ^{[104](#)},
F. Ledroit-Guillon ^{[61](#)}, S.C. Lee ^{[155](#)}, T.F. Lee ^{[95](#)}, L.L. Leeuw ^{[34c,am](#)}, M. Lefebvre ^{[172](#)},
C. Leggett ^{[18a](#)}, G. Lehmann Miotto ^{[37](#)}, M. Leigh ^{[57](#)}, W.A. Leight ^{[106](#)}, W. Leinonen ^{[117](#)},
A. Leisos ^{[159,v](#)}, M.A.L. Leite ^{[84c](#)}, C.E. Leitgeb ^{[19](#)}, R. Leitner ^{[137](#)}, K.J.C. Leney ^{[46](#)}, T. Lenz ^{[25](#)},
S. Leone ^{[75a](#)}, C. Leonidopoulos ^{[53](#)}, A. Leopold ^{[151](#)}, J.H. Lepage Bourbonnais ^{[35](#)}, R. Les ^{[110](#)},
C.G. Lester ^{[33](#)}, M. Levchenko ^{[39](#)}, J. Levêque ^{[4](#)}, L.J. Levinson ^{[176](#)}, G. Levrini ^{[24b,24a](#)},
M.P. Lewicki ^{[88](#)}, C. Lewis ^{[143](#)}, D.J. Lewis ^{[4](#)}, L. Lewitt ^{[146](#)}, A. Li ^{[30](#)}, B. Li ^{[144a](#)}, C. Li ^{[109](#)},
C-Q. Li ^{[113](#)}, H. Li ^{[63](#)}, H. Li ^{[144a](#)}, H. Li ^{[104](#)}, H. Li ^{[15](#)}, H. Li ^{[63](#)}, H. Li ^{[144a](#)}, J. Li ^{[145a](#)}, K. Li ^{[14](#)},
L. Li ^{[145a](#)}, R. Li ^{[179](#)}, S. Li ^{[14,115c](#)}, S. Li ^{[145b,145a,d](#)}, T. Li ^{[5](#)}, X. Li ^{[107](#)}, Z. Li ^{[160](#)}, Z. Li ^{[14,115c](#)},
Z. Li ^{[63](#)}, S. Liang ^{[14,115c](#)}, Z. Liang ^{[14](#)}, M. Liberatore ^{[139](#)}, B. Liberti ^{[77a](#)}, K. Lie ^{[65c](#)},
J. Lieber Marin ^{[84e](#)}, H. Lien ^{[69](#)}, H. Lin ^{[109](#)}, L. Linden ^{[112](#)}, R.E. Lindley ^{[7](#)}, J.H. Lindon ^{[2](#)},
J. Ling ^{[62](#)}, E. Lipeles ^{[132](#)}, A. Lipniacka ^{[17](#)}, A. Lister ^{[171](#)}, J.D. Little ^{[69](#)}, B. Liu ^{[14](#)},
B.X. Liu ^{[115b](#)}, D. Liu ^{[145b,145a](#)}, E.H.L. Liu ^{[21](#)}, J.K.K. Liu ^{[33](#)}, K. Liu ^{[145b](#)}, K. Liu ^{[145b,145a](#)},
M. Liu ^{[63](#)}, M.Y. Liu ^{[63](#)}, P. Liu ^{[14](#)}, Q. Liu ^{[145b,143,145a](#)}, X. Liu ^{[63](#)}, X. Liu ^{[144a](#)},
Y. Liu ^{[115b,115c](#)}, Y.L. Liu ^{[144a](#)}, Y.W. Liu ^{[63](#)}, Z. Liu ^{[67,l](#)}, S.L. Lloyd ^{[97](#)}, E.M. Lobodzinska ^{[49](#)},
P. Loch ^{[7](#)}, E. Lodhi ^{[162](#)}, T. Lohse ^{[19](#)}, K. Lohwasser ^{[146](#)}, E. Loiacono ^{[49](#)}, J.D. Lomas ^{[21](#)},
J.D. Long ^{[43](#)}, I. Longarini ^{[166](#)}, R. Longo ^{[169](#)}, A. Lopez Solis ^{[49](#)}, N.A. Lopez-canelas ^{[7](#)},
N. Lorenzo Martinez ^{[4](#)}, A.M. Lory ^{[112](#)}, M. Losada ^{[120a](#)}, G. Löschcke Centeno ^{[153](#)}, O. Loseva ^{[39](#)},
X. Lou ^{[48a,48b](#)}, X. Lou ^{[14,115c](#)}, A. Lounis ^{[67](#)}, P.A. Love ^{[94](#)}, G. Lu ^{[14,115c](#)}, M. Lu ^{[67](#)}, S. Lu ^{[132](#)},
Y.J. Lu ^{[155](#)}, H.J. Lubatti ^{[143](#)}, C. Luci ^{[76a,76b](#)}, F.L. Lucio Alves ^{[115a](#)}, F. Luehring ^{[69](#)},
B.S. Lunday ^{[132](#)}, O. Lundberg ^{[151](#)}, B. Lund-Jensen ^{[151,*](#)}, N.A. Luongo ^{[6](#)}, M.S. Lutz ^{[37](#)},
A.B. Lux ^{[26](#)}, D. Lynn ^{[30](#)}, R. Lysak ^{[135](#)}, V. Lysenko ^{[136](#)}, E. Lytken ^{[101](#)}, V. Lyubushkin ^{[40](#)},
T. Lyubushkina ^{[40](#)}, M.M. Lyukova ^{[152](#)}, M.Firdaus M. Soberi ^{[53](#)}, H. Ma ^{[30](#)}, K. Ma ^{[63](#)},

L.L. Ma ^{[144a](#)}, W. Ma ^{[63](#)}, Y. Ma ^{[125](#)}, J.C. MacDonald ^{[103](#)}, P.C. Machado De Abreu Farias ^{[84e](#)}, R. Madar ^{[42](#)}, T. Madula ^{[99](#)}, J. Maeda ^{[86](#)}, T. Maeno ^{[30](#)}, P.T. Mafa ^{[34c,k](#)}, H. Maguire ^{[146](#)}, V. Maiboroda ^{[67](#)}, A. Maio ^{[134a,134b,134d](#)}, K. Maj ^{[87a](#)}, O. Majersky ^{[49](#)}, S. Majewski ^{[127](#)}, R. Makhmanazarov ^{[39](#)}, N. Makovec ^{[67](#)}, V. Maksimovic ^{[16](#)}, B. Malaescu ^{[131](#)}, J. Malamant ^{[129](#)}, Pa. Malecki ^{[88](#)}, V.P. Maleev ^{[39](#)}, F. Malek ^{[61,q](#)}, M. Mali ^{[96](#)}, D. Malito ^{[98](#)}, U. Mallik ^{[81,*](#)}, S. Maltezos ^{[10](#)}, A. Malvezzi Lopes ^{[84d](#)}, S. Malyukov ^{[40](#)}, J. Mamuzic ^{[13](#)}, G. Mancini ^{[54](#)}, M.N. Mancini ^{[27](#)}, G. Manco ^{[74a,74b](#)}, J.P. Mandalia ^{[97](#)}, S.S. Mandarry ^{[153](#)}, I. Mandić ^{[96](#)}, L. Manhaes de Andrade Filho ^{[84a](#)}, I.M. Maniatis ^{[176](#)}, J. Manjarres Ramos ^{[92](#)}, D.C. Mankad ^{[176](#)}, A. Mann ^{[112](#)}, T. Manoussos ^{[37](#)}, S. Manzoni ^{[37](#)}, L. Mao ^{[145a](#)}, X. Mapekula ^{[34c](#)}, A. Marantis ^{[159,v](#)}, R.R. Marcelo Gregorio ^{[97](#)}, G. Marchiori ^{[5](#)}, M. Marcisovsky ^{[135](#)}, C. Marcon ^{[72a](#)}, M. Marinescu ^{[21](#)}, S. Marium ^{[49](#)}, M. Marjanovic ^{[124](#)}, A. Markhoos ^{[55](#)}, M. Markovitch ^{[67](#)}, M.K. Maroun ^{[106](#)}, E.J. Marshall ^{[94](#)}, Z. Marshall ^{[18a](#)}, S. Marti-Garcia ^{[170](#)}, J. Martin ^{[99](#)}, T.A. Martin ^{[138](#)}, V.J. Martin ^{[53](#)}, B. Martin dit Latour ^{[17](#)}, L. Martinelli ^{[76a,76b](#)}, M. Martinez ^{[13,y](#)}, P. Martinez Agullo ^{[170](#)}, V.I. Martinez Outschoorn ^{[106](#)}, P. Martinez Suarez ^{[13](#)}, S. Martin-Haugh ^{[138](#)}, G. Martinovicova ^{[137](#)}, V.S. Martoiu ^{[28b](#)}, A.C. Martyniuk ^{[99](#)}, A. Marzin ^{[37](#)}, D. Mascione ^{[79a,79b](#)}, L. Masetti ^{[103](#)}, J. Masik ^{[104](#)}, A.L. Maslennikov ^{[40](#)}, S.L. Mason ^{[43](#)}, P. Massarotti ^{[73a,73b](#)}, P. Mastrandrea ^{[75a,75b](#)}, A. Mastroberardino ^{[45b,45a](#)}, T. Masubuchi ^{[128](#)}, T.T. Mathew ^{[127](#)}, J. Matousek ^{[137](#)}, D.M. Mattern ^{[50](#)}, J. Maurer ^{[28b](#)}, T. Maurin ^{[60](#)}, A.J. Maury ^{[67](#)}, B. Maček ^{[96](#)}, D.A. Maximov ^{[39](#)}, A.E. May ^{[104](#)}, E. Mayer ^{[42](#)}, R. Mazini ^{[34g](#)}, I. Maznas ^{[119](#)}, M. Mazza ^{[110](#)}, S.M. Mazza ^{[140](#)}, E. Mazzeo ^{[72a,72b](#)}, J.P. Mc Gowan ^{[172](#)}, S.P. Mc Kee ^{[109](#)}, C.A. Mc Lean ^{[6](#)}, C.C. McCracken ^{[171](#)}, E.F. McDonald ^{[108](#)}, A.E. McDougall ^{[118](#)}, L.F. Mcelhinney ^{[94](#)}, J.A. Mcfayden ^{[153](#)}, R.P. McGovern ^{[132](#)}, R.P. Mckenzie ^{[34g](#)}, T.C. McLachlan ^{[49](#)}, D.J. McLaughlin ^{[99](#)}, S.J. McMahon ^{[138](#)}, C.M. Mcpartland ^{[95](#)}, R.A. McPherson ^{[172,ac](#)}, S. Mehlhase ^{[112](#)}, A. Mehta ^{[95](#)}, D. Melini ^{[170](#)}, B.R. Mellado Garcia ^{[34g](#)}, A.H. Melo ^{[56](#)}, F. Meloni ^{[49](#)}, A.M. Mendes Jacques Da Costa ^{[104](#)}, H.Y. Meng ^{[162](#)}, L. Meng ^{[94](#)}, S. Menke ^{[113](#)}, M. Mentink ^{[37](#)}, E. Meoni ^{[45b,45a](#)}, G. Mercado ^{[119](#)}, S. Merianos ^{[159](#)}, C. Merlassino ^{[70a,70c](#)}, C. Meroni ^{[72a,72b](#)}, J. Metcalfe ^{[6](#)}, A.S. Mete ^{[6](#)}, E. Meuser ^{[103](#)}, C. Meyer ^{[69](#)}, J-P. Meyer ^{[139](#)}, R.P. Middleton ^{[138](#)}, M. Mihovilovic ^{[67](#)}, L. Mijović ^{[53](#)}, G. Mikenberg ^{[176](#)}, M. Mikestikova ^{[135](#)}, M. Mikuž ^{[96](#)}, H. Mildner ^{[103](#)}, A. Milic ^{[37](#)}, D.W. Miller ^{[41](#)}, E.H. Miller ^{[150](#)}, L.S. Miller ^{[35](#)}, A. Milov ^{[176](#)}, D.A. Milstead ^{[48a,48b](#)}, T. Min ^{[115a](#)}, A.A. Minaenko ^{[39](#)}, I.A. Minashvili ^{[156b](#)}, A.I. Mincer ^{[121](#)}, B. Mindur ^{[87a](#)}, M. Mineev ^{[40](#)}, Y. Mino ^{[90](#)}, L.M. Mir ^{[13](#)}, M. Miralles Lopez ^{[60](#)}, M. Mironova ^{[18a](#)}, M.C. Missio ^{[117](#)}, A. Mitra ^{[174](#)}, V.A. Mitsou ^{[170](#)}, Y. Mitsumori ^{[114](#)}, O. Miu ^{[162](#)}, P.S. Miyagawa ^{[97](#)}, T. Mkrtchyan ^{[64a](#)}, M. Mlinarevic ^{[99](#)}, T. Mlinarevic ^{[99](#)}, M. Mlynarikova ^{[37](#)}, S. Mobius ^{[20](#)}, P. Mogg ^{[112](#)}, M.H. Mohamed Farook ^{[116](#)}, A.F. Mohammed ^{[14,115c](#)}, S. Mohapatra ^{[43](#)}, S. Mohiuddin ^{[125](#)}, G. Mokgatitswane ^{[34g](#)}, L. Moleri ^{[176](#)}, B. Mondal ^{[148](#)}, S. Mondal ^{[136](#)}, K. Mönig ^{[49](#)}, E. Monnier ^{[105](#)}, L. Monsonis Romero ^{[170](#)}, J. Montejo Berlingen ^{[13](#)}, A. Montella ^{[48a,48b](#)}, M. Montella ^{[123](#)}, F. Montekali ^{[78a,78b](#)}, F. Monticelli ^{[93](#)}, S. Monzani ^{[70a,70c](#)}, A. Morancho Tarda ^{[44](#)}, N. Morange ^{[67](#)}, A.L. Moreira De Carvalho ^{[49](#)}, M. Moreno Llácer ^{[170](#)}, C. Moreno Martinez ^{[57](#)}, J.M. Moreno Perez ^{[23b](#)}, P. Morettini ^{[58b](#)}, S. Morgenstern ^{[37](#)}, M. Morii ^{[62](#)}, M. Morinaga ^{[160](#)}, M. Moritsu ^{[91](#)}, F. Morodei ^{[76a,76b](#)}, P. Moschovakos ^{[37](#)}, B. Moser ^{[130](#)}, M. Mosidze ^{[156b](#)}, T. Moskalets ^{[46](#)}, P. Moskvitina ^{[117](#)}, J. Moss ^{[32,n](#)}, P. Moszkowicz ^{[87a](#)}, A. Moussa ^{[36d](#)},

Y. Moyal ¹⁷⁶, E.J.W. Moyse ¹⁰⁶, O. Mtintsilana ^{34g}, S. Muanza ¹⁰⁵, J. Mueller ¹³³,
R. Müller ³⁷, G.A. Mullier ¹⁶⁸, A.J. Mullin ³³, J.J. Mullin ⁵², A.E. Mulski ⁶², D.P. Mungo ¹⁶²,
D. Munoz Perez ¹⁷⁰, F.J. Munoz Sanchez ¹⁰⁴, M. Murin ¹⁰⁴, W.J. Murray ^{174,138},
M. Muškinja ⁹⁶, C. Mwewa ³⁰, A.G. Myagkov ^{39,a}, A.J. Myers ⁸, G. Myers ¹⁰⁹, M. Myska ¹³⁶,
B.P. Nachman ^{18a}, K. Nagai ¹³⁰, K. Nagano ⁸⁵, R. Nagasaka ¹⁶⁰, J.L. Nagle ^{30,al}, E. Nagy ¹⁰⁵,
A.M. Nairz ³⁷, Y. Nakahama ⁸⁵, K. Nakamura ⁸⁵, K. Nakkalil ⁵, H. Nanjo ¹²⁸,
E.A. Narayanan ⁴⁶, Y. Narukawa ¹⁶⁰, I. Naryshkin ³⁹, L. Nasella ^{72a,72b}, S. Nasri ^{120b},
C. Nass ²⁵, G. Navarro ^{23a}, J. Navarro-Gonzalez ¹⁷⁰, A. Nayaz ¹⁹, P.Y. Nechaeva ³⁹,
S. Nechaeva ^{24b,24a}, F. Nechansky ¹³⁵, L. Nedic ¹³⁰, T.J. Neep ²¹, A. Negri ^{74a,74b},
M. Negrini ^{24b}, C. Nellist ¹¹⁸, C. Nelson ¹⁰⁷, K. Nelson ¹⁰⁹, S. Nemecek ¹³⁵, M. Nessi ^{37,h},
M.S. Neubauer ¹⁶⁹, J. Newell ⁹⁵, P.R. Newman ²¹, Y.W.Y. Ng ¹⁶⁹, B. Ngair ^{120a},
H.D.N. Nguyen ¹¹¹, R.B. Nickerson ¹³⁰, R. Nicolaidou ¹³⁹, J. Nielsen ¹⁴⁰, M. Niemeyer ⁵⁶,
J. Niermann ³⁷, N. Nikiforou ³⁷, V. Nikolaenko ^{39,a}, I. Nikolic-Audit ¹³¹, P. Nilsson ³⁰,
I. Ninca ⁴⁹, G. Ninio ¹⁵⁸, A. Nisati ^{76a}, N. Nishu ², R. Nisius ¹¹³, N. Nitika ^{70a,70c},
J-E. Nitschke ⁵¹, E.K. Nkadimeng ^{34g}, T. Nobe ¹⁶⁰, T. Nommensen ¹⁵⁴, M.B. Norfolk ¹⁴⁶,
B.J. Norman ³⁵, M. Noury ^{36a}, J. Novak ⁹⁶, T. Novak ⁹⁶, R. Novotny ¹¹⁶, L. Nozka ¹²⁶,
K. Ntekas ¹⁶⁶, N.M.J. Nunes De Moura Junior ^{84b}, J. Ocariz ¹³¹, A. Ochi ⁸⁶, I. Ochoa ^{134a},
S. Oerdek ^{49,z}, J.T. Offermann ⁴¹, A. Ogrodnik ¹³⁷, A. Oh ¹⁰⁴, C.C. Ohm ¹⁵¹, H. Oide ⁸⁵,
R. Oishi ¹⁶⁰, M.L. Ojeda ³⁷, Y. Okumura ¹⁶⁰, L.F. Oleiro Seabra ^{134a}, I. Oleksiyuk ⁵⁷,
S.A. Olivares Pino ^{141d}, G. Oliveira Correa ¹³, D. Oliveira Damazio ³⁰, J.L. Oliver ¹⁶⁶,
Ö.O. Öncel ⁵⁵, A.P. O'Neill ²⁰, A. Onofre ^{134a,134e}, P.U.E. Onyisi ¹¹, M.J. Oreglia ⁴¹,
D. Orestano ^{78a,78b}, R. Orlandini ^{78a,78b}, R.S. Orr ¹⁶², L.M. Osojnak ¹³², Y. Osumi ¹¹⁴,
G. Otero y Garzon ³¹, H. Otono ⁹¹, G.J. Ottino ^{18a}, M. Ouchrif ^{36d}, F. Ould-Saada ¹²⁹,
T. Ovsianikova ¹⁴³, M. Owen ⁶⁰, R.E. Owen ¹³⁸, V.E. Ozcan ^{22a}, F. Ozturk ⁸⁸, N. Ozturk ⁸,
S. Ozturk ⁸³, H.A. Pacey ¹³⁰, K. Pachal ^{163a}, A. Pacheco Pages ¹³, C. Padilla Aranda ¹³,
G. Padovano ^{76a,76b}, S. Pagan Griso ^{18a}, G. Palacino ⁶⁹, A. Palazzo ^{71a,71b}, J. Pampel ²⁵,
J. Pan ¹⁷⁹, T. Pan ^{65a}, D.K. Panchal ¹¹, C.E. Pandini ¹¹⁸, J.G. Panduro Vazquez ¹³⁸,
H.D. Pandya ¹, H. Pang ¹³⁹, P. Pani ⁴⁹, G. Panizzo ^{70a,70c}, L. Panwar ¹³¹, L. Paolozzi ⁵⁷,
S. Parajuli ¹⁶⁹, A. Paramonov ⁶, C. Paraskevopoulos ⁵⁴, D. Paredes Hernandez ^{65b},
A. Pareti ^{74a,74b}, K.R. Park ⁴³, T.H. Park ¹¹³, F. Parodi ^{58b,58a}, J.A. Parsons ⁴³,
U. Parzefall ⁵⁵, B. Pascual Dias ⁴², L. Pascual Dominguez ¹⁰², E. Pasqualucci ^{76a},
S. Passaggio ^{58b}, F. Pastore ⁹⁸, P. Patel ⁸⁸, U.M. Patel ⁵², J.R. Pater ¹⁰⁴, T. Pauly ³⁷,
F. Pauwels ¹³⁷, C.I. Pazos ¹⁶⁵, M. Pedersen ¹²⁹, R. Pedro ^{134a}, S.V. Peleganchuk ³⁹,
O. Penc ³⁷, E.A. Pender ⁵³, S. Peng ¹⁵, G.D. Penn ¹⁷⁹, K.E. Pensi ¹¹², M. Penzin ³⁹,
B.S. Peralva ^{84d}, A.P. Pereira Peixoto ¹⁴³, L. Pereira Sanchez ¹⁵⁰, D.V. Perepelitsa ^{30,al},
G. Perera ¹⁰⁶, E. Perez Codina ^{163a}, M. Perganti ¹⁰, H. Pernegger ³⁷, S. Perrella ^{76a,76b},
O. Perrin ⁴², K. Peters ⁴⁹, R.F.Y. Peters ¹⁰⁴, B.A. Petersen ³⁷, T.C. Petersen ⁴⁴, E. Petit ¹⁰⁵,
V. Petousis ¹³⁶, A.R. Petri ^{72a,72b}, C. Petridou ^{159,e}, T. Petru ¹³⁷, A. Petrukhin ¹⁴⁸,
M. Pettee ^{18a}, A. Petukhov ⁸³, K. Petukhova ³⁷, R. Pezoa ^{141f}, L. Pezzotti ^{24b,24a},
G. Pezzullo ¹⁷⁹, L. Pfaffenbichler ³⁷, A.J. Pflieger ³⁷, T.M. Pham ¹⁷⁷, T. Pham ¹⁰⁸,
P.W. Phillips ¹³⁸, G. Piacquadio ¹⁵², E. Pianori ^{18a}, F. Piazza ¹²⁷, R. Piegai ³¹,
D. Pietreanu ^{28b}, A.D. Pilkington ¹⁰⁴, M. Pinamonti ^{70a,70c}, J.L. Pinfold ²,
B.C. Pinheiro Pereira ^{134a}, J. Pinol Bel ¹³, A.E. Pinto Pinoargote ¹³¹, L. Pintucci ^{70a,70c},

K.M. Piper ^{[153](#)}, A. Pirttikoski ^{[57](#)}, D.A. Pizzi ^{[35](#)}, L. Pizzimento ^{[65b](#)}, A. Plebani ^{[33](#)},
M.-A. Pleier ^{[30](#)}, V. Pleskot ^{[137](#)}, E. Plotnikova ^{[40](#)}, G. Poddar ^{[97](#)}, R. Poettgen ^{[101](#)},
L. Poggioli ^{[131](#)}, S. Polacek ^{[137](#)}, G. Polesello ^{[74a](#)}, A. Poley ^{[149,163a](#)}, A. Polini ^{[24b](#)},
C.S. Pollard ^{[174](#)}, Z.B. Pollock ^{[123](#)}, E. Pompa Pacchi ^{[124](#)}, N.I. Pond ^{[99](#)}, D. Ponomarenko ^{[69](#)},
L. Pontecorvo ^{[37](#)}, S. Popa ^{[28a](#)}, G.A. Popeneciu ^{[28d](#)}, A. Poreba ^{[37](#)}, D.M. Portillo Quintero ^{[163a](#)},
S. Pospisil ^{[136](#)}, M.A. Postill ^{[146](#)}, P. Postolache ^{[28c](#)}, K. Potamianos ^{[174](#)}, P.A. Potepa ^{[87a](#)},
I.N. Potrap ^{[40](#)}, C.J. Potter ^{[33](#)}, H. Potti ^{[154](#)}, J. Poveda ^{[170](#)}, M.E. Pozo Astigarraga ^{[37](#)},
A. Prades Ibanez ^{[77a,77b](#)}, J. Pretel ^{[172](#)}, D. Price ^{[104](#)}, M. Primavera ^{[71a](#)}, L. Primomo ^{[70a,70c](#)},
M.A. Principe Martin ^{[102](#)}, R. Privara ^{[126](#)}, T. Procter ^{[60](#)}, M.L. Proffitt ^{[143](#)}, N. Proklova ^{[132](#)},
K. Prokofiev ^{[65c](#)}, G. Proto ^{[113](#)}, J. Proudfoot ^{[6](#)}, M. Przybycien ^{[87a](#)}, W.W. Przygoda ^{[87b](#)},
A. Psallidas ^{[47](#)}, J.E. Puddefoot ^{[146](#)}, D. Pudzha ^{[55](#)}, D. Pyatiizbyantseva ^{[117](#)}, J. Qian ^{[109](#)},
R. Qian ^{[110](#)}, D. Qichen ^{[104](#)}, Y. Qin ^{[13](#)}, T. Qiu ^{[53](#)}, A. Quadt ^{[56](#)}, M. Queitsch-Maitland ^{[104](#)},
G. Quetant ^{[57](#)}, R.P. Quinn ^{[171](#)}, G. Rabanal Bolanos ^{[62](#)}, D. Rafanoharana ^{[55](#)}, F. Raffaelli ^{[77a,77b](#)},
F. Ragusa ^{[72a,72b](#)}, J.L. Rainbolt ^{[41](#)}, J.A. Raine ^{[57](#)}, S. Rajagopalan ^{[30](#)}, E. Ramakoti ^{[39](#)},
L. Rambelli ^{[58b,58a](#)}, I.A. Ramirez-Berend ^{[35](#)}, K. Ran ^{[49,115c](#)}, D.S. Rankin ^{[132](#)},
N.P. Rapheeha ^{[34g](#)}, H. Rasheed ^{[28b](#)}, V. Raskina ^{[131](#)}, D.F. Rassloff ^{[64a](#)}, A. Rastogi ^{[18a](#)},
S. Rave ^{[103](#)}, S. Ravera ^{[58b,58a](#)}, B. Ravina ^{[37](#)}, I. Ravinovich ^{[176](#)}, M. Raymond ^{[37](#)},
A.L. Read ^{[129](#)}, N.P. Readioff ^{[146](#)}, D.M. Rebuzzi ^{[74a,74b](#)}, A.S. Reed ^{[113](#)}, K. Reeves ^{[27](#)},
J.A. Reidelsturz ^{[178](#)}, D. Reikher ^{[127](#)}, A. Rej ^{[50](#)}, C. Rembser ^{[37](#)}, H. Ren ^{[63](#)}, M. Renda ^{[28b](#)},
F. Renner ^{[49](#)}, A.G. Rennie ^{[166](#)}, A.L. Rescia ^{[49](#)}, S. Resconi ^{[72a](#)}, M. Ressegotti ^{[58b,58a](#)},
S. Rettie ^{[37](#)}, W.F. Rettie ^{[35](#)}, J.G. Reyes Rivera ^{[110](#)}, E. Reynolds ^{[18a](#)}, O.L. Rezanova ^{[40](#)},
P. Reznicek ^{[137](#)}, H. Riani ^{[36d](#)}, N. Ribaric ^{[52](#)}, E. Ricci ^{[79a,79b](#)}, R. Richter ^{[113](#)}, S. Richter ^{[48a,48b](#)},
E. Richter-Was ^{[87b](#)}, M. Ridel ^{[131](#)}, S. Ridouani ^{[36d](#)}, P. Rieck ^{[121](#)}, P. Riedler ^{[37](#)},
E.M. Riefel ^{[48a,48b](#)}, J.O. Rieger ^{[118](#)}, M. Rijssenbeek ^{[152](#)}, M. Rimoldi ^{[37](#)}, L. Rinaldi ^{[24b,24a](#)},
P. Rincke ^{[56](#)}, G. Ripellino ^{[168](#)}, I. Riu ^{[13](#)}, J.C. Rivera Vergara ^{[172](#)}, F. Rizatdinova ^{[125](#)},
E. Rizvi ^{[97](#)}, B.R. Roberts ^{[18a](#)}, S.S. Roberts ^{[140](#)}, D. Robinson ^{[33](#)}, M. Robles Manzano ^{[103](#)},
A. Robson ^{[60](#)}, A. Rocchi ^{[77a,77b](#)}, C. Roda ^{[75a,75b](#)}, S. Rodriguez Bosca ^{[37](#)},
Y. Rodriguez Garcia ^{[23a](#)}, A.M. Rodríguez Vera ^{[119](#)}, S. Roe ^{[37](#)}, J.T. Roemer ^{[37](#)}, O. Röhne ^{[129](#)},
R.A. Rojas ^{[37](#)}, C.P.A. Roland ^{[131](#)}, J. Roloff ^{[30](#)}, A. Romanouk ^{[80](#)}, E. Romano ^{[74a,74b](#)},
M. Romano ^{[24b](#)}, A.C. Romero Hernandez ^{[169](#)}, N. Rompotis ^{[95](#)}, L. Roos ^{[131](#)}, S. Rosati ^{[76a](#)},
B.J. Rosser ^{[41](#)}, E. Rossi ^{[130](#)}, E. Rossi ^{[73a,73b](#)}, L.P. Rossi ^{[62](#)}, L. Rossini ^{[55](#)}, R. Rosten ^{[123](#)},
M. Rotaru ^{[28b](#)}, B. Rottler ^{[55](#)}, D. Rousseau ^{[67](#)}, D. Rouso ^{[49](#)}, S. Roy-Garand ^{[162](#)},
A. Rozanov ^{[105](#)}, Z.M.A. Rozario ^{[60](#)}, Y. Rozen ^{[157](#)}, A. Rubio Jimenez ^{[170](#)},
V.H. Ruelas Rivera ^{[19](#)}, T.A. Ruggeri ^{[1](#)}, A. Ruggiero ^{[130](#)}, A. Ruiz-Martinez ^{[170](#)}, A. Rummler ^{[37](#)},
Z. Rurikova ^{[55](#)}, N.A. Rusakovich ^{[40](#)}, H.L. Russell ^{[172](#)}, G. Russo ^{[76a,76b](#)}, J.P. Rutherford ^{[7](#)},
S. Rutherford Colmenares ^{[33](#)}, M. Rybar ^{[137](#)}, P. Rybczynski ^{[87a](#)}, E.B. Rye ^{[129](#)}, A. Ryzhov ^{[46](#)},
J.A. Sabater Iglesias ^{[57](#)}, H.F-W. Sadrozinski ^{[140](#)}, F. Safai Tehrani ^{[76a](#)}, S. Saha ^{[1](#)},
M. Sahinsoy ^{[83](#)}, A. Saibel ^{[170](#)}, B.T. Saifuddin ^{[124](#)}, M. Saimpert ^{[139](#)}, M. Saito ^{[160](#)}, T. Saito ^{[160](#)},
A. Sala ^{[72a,72b](#)}, D. Salamani ^{[37](#)}, A. Salmikov ^{[150](#)}, J. Salt ^{[170](#)}, A. Salvador Salas ^{[158](#)},
D. Salvatore ^{[45b,45a](#)}, F. Salvatore ^{[153](#)}, A. Salzburger ^{[37](#)}, D. Sammel ^{[55](#)}, E. Sampson ^{[94](#)},
D. Sampsonidis ^{[159,e](#)}, D. Sampsonidou ^{[127](#)}, J. Sánchez ^{[170](#)}, V. Sanchez Sebastian ^{[170](#)},
H. Sandaker ^{[129](#)}, C.O. Sander ^{[49](#)}, J.A. Sandesara ^{[106](#)}, M. Sandhoff ^{[178](#)}, C. Sandoval ^{[23b](#)},
L. Sanfilippo ^{[64a](#)}, D.P.C. Sankey ^{[138](#)}, T. Sano ^{[90](#)}, A. Sansoni ^{[54](#)}, L. Santi ^{[37](#)}, C. Santoni ^{[42](#)},

H. Santos [ID](#)^{134a,134b}, A. Santra [ID](#)¹⁷⁶, E. Sanzani [ID](#)^{24b,24a}, K.A. Saoucha [ID](#)^{89b}, J.G. Saraiva [ID](#)^{134a,134d}, J. Sardain [ID](#)⁷, O. Sasaki [ID](#)⁸⁵, K. Sato [ID](#)¹⁶⁴, C. Sauer³⁷, E. Sauvan [ID](#)⁴, P. Savard [ID](#)^{162,aj}, R. Sawada [ID](#)¹⁶⁰, C. Sawyer [ID](#)¹³⁸, L. Sawyer [ID](#)¹⁰⁰, C. Sbarra [ID](#)^{24b}, A. Sbrizzi [ID](#)^{24b,24a}, T. Scanlon [ID](#)⁹⁹, J. Schaarschmidt [ID](#)¹⁴³, U. Schäfer [ID](#)¹⁰³, A.C. Schaffer [ID](#)^{67,46}, D. Schaile [ID](#)¹¹², R.D. Schamberger [ID](#)¹⁵², C. Scharf [ID](#)¹⁹, M.M. Schefer [ID](#)²⁰, V.A. Schegelsky [ID](#)³⁹, D. Scheirich [ID](#)¹³⁷, M. Schernau [ID](#)^{141e}, C. Scheulen [ID](#)⁵⁷, C. Schiavi [ID](#)^{58b,58a}, M. Schioppa [ID](#)^{45b,45a}, B. Schlag [ID](#)¹⁵⁰, S. Schlenker [ID](#)³⁷, J. Schmeing [ID](#)¹⁷⁸, M.A. Schmidt [ID](#)¹⁷⁸, K. Schmieden [ID](#)¹⁰³, C. Schmitt [ID](#)¹⁰³, N. Schmitt [ID](#)¹⁰³, S. Schmitt [ID](#)⁴⁹, L. Schoeffel [ID](#)¹³⁹, A. Schoening [ID](#)^{64b}, P.G. Scholer [ID](#)³⁵, E. Schopf [ID](#)¹⁴⁸, M. Schott [ID](#)²⁵, S. Schramm [ID](#)⁵⁷, T. Schroer [ID](#)⁵⁷, H-C. Schultz-Coulon [ID](#)^{64a}, M. Schumacher [ID](#)⁵⁵, B.A. Schumm [ID](#)¹⁴⁰, Ph. Schune [ID](#)¹³⁹, H.R. Schwartz [ID](#)¹⁴⁰, A. Schwartzman [ID](#)¹⁵⁰, T.A. Schwarz [ID](#)¹⁰⁹, Ph. Schwemling [ID](#)¹³⁹, R. Schwienhorst [ID](#)¹¹⁰, F.G. Sgacca [ID](#)²⁰, A. Sciandra [ID](#)³⁰, G. Sciolla [ID](#)²⁷, F. Scuri [ID](#)^{75a}, C.D. Sebastiani [ID](#)³⁷, K. Sedlaczek [ID](#)¹¹⁹, S.C. Seidel [ID](#)¹¹⁶, A. Seiden [ID](#)¹⁴⁰, B.D. Seidlitz [ID](#)⁴³, C. Seitz [ID](#)⁴⁹, J.M. Seixas [ID](#)^{84b}, G. Sekhniaidze [ID](#)^{73a}, L. Selem [ID](#)⁶¹, N. Semprini-Cesari [ID](#)^{24b,24a}, A. Semushin [ID](#)^{180,39}, D. Sengupta [ID](#)⁵⁷, V. Senthilkumar [ID](#)¹⁷⁰, L. Serin [ID](#)⁶⁷, M. Sessa [ID](#)^{77a,77b}, H. Severini [ID](#)¹²⁴, F. Sforza [ID](#)^{58b,58a}, A. Sfyrila [ID](#)⁵⁷, Q. Sha [ID](#)¹⁴, E. Shabalina [ID](#)⁵⁶, H. Shaddix [ID](#)¹¹⁹, A.H. Shah [ID](#)³³, R. Shaheen [ID](#)¹⁵¹, J.D. Shahinian [ID](#)¹³², D. Shaked Renous [ID](#)¹⁷⁶, M. Shamim [ID](#)³⁷, L.Y. Shan [ID](#)¹⁴, M. Shapiro [ID](#)^{18a}, A. Sharma [ID](#)³⁷, A.S. Sharma [ID](#)¹⁷¹, P. Sharma [ID](#)³⁰, P.B. Shatalov [ID](#)³⁹, K. Shaw [ID](#)¹⁵³, S.M. Shaw [ID](#)¹⁰⁴, Q. Shen [ID](#)^{145a}, D.J. Sheppard [ID](#)¹⁴⁹, P. Sherwood [ID](#)⁹⁹, L. Shi [ID](#)⁹⁹, X. Shi [ID](#)¹⁴, S. Shimizu [ID](#)⁸⁵, C.O. Shimmin [ID](#)¹⁷⁹, I.P.J. Shipsey [ID](#)^{130,*}, S. Shirabe [ID](#)⁹¹, M. Shiyakova [ID](#)^{40,aa}, M.J. Shochet [ID](#)⁴¹, D.R. Shope [ID](#)¹²⁹, B. Shrestha [ID](#)¹²⁴, S. Shrestha [ID](#)^{123,an}, I. Shreyber [ID](#)⁴⁰, M.J. Shroff [ID](#)¹⁷², P. Sicho [ID](#)¹³⁵, A.M. Sickles [ID](#)¹⁶⁹, E. Sideras Haddad [ID](#)^{34g,167}, A.C. Sidley [ID](#)¹¹⁸, A. Sidoti [ID](#)^{24b}, F. Siegert [ID](#)⁵¹, Dj. Sijacki [ID](#)¹⁶, F. Sili [ID](#)⁹³, J.M. Silva [ID](#)⁵³, I. Silva Ferreira [ID](#)^{84b}, M.V. Silva Oliveira [ID](#)³⁰, S.B. Silverstein [ID](#)^{48a}, S. Simion [ID](#)⁶⁷, R. Simoniello [ID](#)³⁷, E.L. Simpson [ID](#)¹⁰⁴, H. Simpson [ID](#)¹⁵³, L.R. Simpson [ID](#)¹⁰⁹, S. Simsek [ID](#)⁸³, S. Sindhu [ID](#)⁵⁶, P. Sinervo [ID](#)¹⁶², S.N. Singh [ID](#)²⁷, S. Singh [ID](#)³⁰, S. Sinha [ID](#)⁴⁹, S. Sinha [ID](#)¹⁰⁴, M. Sioli [ID](#)^{24b,24a}, K. Sioulas [ID](#)⁹, I. Siral [ID](#)³⁷, E. Sitnikova [ID](#)⁴⁹, J. Sjölin [ID](#)^{48a,48b}, A. Skaf [ID](#)⁵⁶, E. Skorda [ID](#)²¹, P. Skubic [ID](#)¹²⁴, M. Slawinska [ID](#)⁸⁸, I. Slazyk [ID](#)¹⁷, V. Smakhtin [ID](#)¹⁷⁶, B.H. Smart [ID](#)¹³⁸, S.Yu. Smirnov [ID](#)^{141b}, Y. Smirnov [ID](#)⁸³, L.N. Smirnova [ID](#)^{39,a}, O. Smirnova [ID](#)¹⁰¹, A.C. Smith [ID](#)⁴³, D.R. Smith [ID](#)¹⁶⁶, E.A. Smith [ID](#)⁴¹, J.L. Smith [ID](#)¹⁰⁴, M.B. Smith [ID](#)³⁵, R. Smith [ID](#)¹⁵⁰, H. Smitmanns [ID](#)¹⁰³, M. Smizanska [ID](#)⁹⁴, K. Smolek [ID](#)¹³⁶, P. Smolyanskiy [ID](#)¹³⁶, A.A. Snegarev [ID](#)⁴⁰, H.L. Snoek [ID](#)¹¹⁸, S. Snyder [ID](#)³⁰, R. Sobie [ID](#)^{172,ac}, A. Soffer [ID](#)¹⁵⁸, C.A. Solans Sanchez [ID](#)³⁷, E.Yu. Soldatov [ID](#)⁴⁰, U. Soldevila [ID](#)¹⁷⁰, A.A. Solodkov [ID](#)^{34g}, S. Solomon [ID](#)²⁷, A. Soloshenko [ID](#)⁴⁰, K. Solovieva [ID](#)⁵⁵, O.V. Solovyanov [ID](#)⁴², P. Sommer [ID](#)⁵¹, A. Sonay [ID](#)¹³, W.Y. Song [ID](#)^{163b}, A. Sopczak [ID](#)¹³⁶, A.L. Sopio [ID](#)⁵³, F. Sopkova [ID](#)^{29b}, J.D. Sorenson [ID](#)¹¹⁶, I.R. Sotarriva Alvarez [ID](#)¹⁴², V. Sothilingam [ID](#)^{64a}, O.J. Soto Sandoval [ID](#)^{141c,141b}, S. Sottocornola [ID](#)⁶⁹, R. Soualah [ID](#)^{89a}, Z. Soumami [ID](#)^{36e}, D. South [ID](#)⁴⁹, N. Soybelman [ID](#)¹⁷⁶, S. Spagnolo [ID](#)^{71a,71b}, M. Spalla [ID](#)¹¹³, D. Sperlich [ID](#)⁵⁵, B. Spisso [ID](#)^{73a,73b}, D.P. Spiteri [ID](#)⁶⁰, L. Splendori [ID](#)¹⁰⁵, M. Spousta [ID](#)¹³⁷, E.J. Staats [ID](#)³⁵, R. Stamen [ID](#)^{64a}, E. Stanecka [ID](#)⁸⁸, W. Stanek-Maslouska [ID](#)⁴⁹, M.V. Stange [ID](#)⁵¹, B. Stanislaus [ID](#)^{18a}, M.M. Stanitzki [ID](#)⁴⁹, B. Stapf [ID](#)⁴⁹, E.A. Starchenko [ID](#)³⁹, G.H. Stark [ID](#)¹⁴⁰, J. Stark [ID](#)⁹², P. Staroba [ID](#)¹³⁵, P. Starovoitov [ID](#)^{89b}, R. Staszewski [ID](#)⁸⁸, G. Stavropoulos [ID](#)⁴⁷, A. Steff [ID](#)³⁷, P. Steinberg [ID](#)³⁰, B. Stelzer [ID](#)^{149,163a}, H.J. Stelzer [ID](#)¹³³, O. Stelzer-Chilton [ID](#)^{163a}, H. Stenzel [ID](#)⁵⁹, T.J. Stevenson [ID](#)¹⁵³, G.A. Stewart [ID](#)³⁷, J.R. Stewart [ID](#)¹²⁵, M.C. Stockton [ID](#)³⁷, G. Stoicea [ID](#)^{28b}, M. Stolarski [ID](#)^{134a}, S. Stonjek [ID](#)¹¹³, A. Straessner [ID](#)⁵¹,

J. Strandberg ^{[151](#)}, S. Strandberg ^{[48a,48b](#)}, M. Stratmann ^{[178](#)}, M. Strauss ^{[124](#)}, T. Strebler ^{[105](#)}, P. Strizenec ^{[29b](#)}, R. Ströhmer ^{[173](#)}, D.M. Strom ^{[127](#)}, R. Stroynowski ^{[46](#)}, A. Strubig ^{[48a,48b](#)}, S.A. Stucci ^{[30](#)}, B. Stugu ^{[17](#)}, J. Stupak ^{[124](#)}, N.A. Styles ^{[49](#)}, D. Su ^{[150](#)}, S. Su ^{[63](#)}, W. Su ^{[145b](#)}, X. Su ^{[63](#)}, D. Suchy ^{[29a](#)}, K. Sugizaki ^{[132](#)}, V.V. Sulin ^{[39](#)}, M.J. Sullivan ^{[95](#)}, D.M.S. Sultan ^{[130](#)}, L. Sultanaliyeva ^{[39](#)}, S. Sultansoy ^{[3b](#)}, S. Sun ^{[177](#)}, W. Sun ^{[14](#)}, O. Sunneborn Gudnadottir ^{[168](#)}, N. Sur ^{[105](#)}, M.R. Sutton ^{[153](#)}, H. Suzuki ^{[164](#)}, M. Svatos ^{[135](#)}, P.N. Swallow ^{[33](#)}, M. Swiatlowski ^{[163a](#)}, T. Swirski ^{[173](#)}, I. Sykora ^{[29a](#)}, M. Sykora ^{[137](#)}, T. Sykora ^{[137](#)}, D. Ta ^{[103](#)}, K. Tackmann ^{[49,z](#)}, A. Taffard ^{[166](#)}, R. Tafirout ^{[163a](#)}, Y. Takubo ^{[85](#)}, M. Talby ^{[105](#)}, A.A. Talyshiev ^{[39](#)}, K.C. Tam ^{[65b](#)}, N.M. Tamir ^{[158](#)}, A. Tanaka ^{[160](#)}, J. Tanaka ^{[160](#)}, R. Tanaka ^{[67](#)}, M. Tanasini ^{[152](#)}, Z. Tao ^{[171](#)}, S. Tapia Araya ^{[141f](#)}, S. Tapprogge ^{[103](#)}, A. Tarek Abouelfadl Mohamed ^{[110](#)}, S. Tarem ^{[157](#)}, K. Tariq ^{[14](#)}, G. Tarna ^{[28b](#)}, G.F. Tartarelli ^{[72a](#)}, M.J. Tartarin ^{[92](#)}, P. Tas ^{[137](#)}, M. Tasevsky ^{[135](#)}, E. Tassi ^{[45b,45a](#)}, A.C. Tate ^{[169](#)}, G. Tateno ^{[160](#)}, Y. Tayalati ^{[36e,ab](#)}, G.N. Taylor ^{[108](#)}, W. Taylor ^{[163b](#)}, A.S. Tegetmeier ^{[92](#)}, P. Teixeira-Dias ^{[98](#)}, J.J. Teoh ^{[162](#)}, K. Terashi ^{[160](#)}, J. Terron ^{[102](#)}, S. Terzo ^{[13](#)}, M. Testa ^{[54](#)}, R.J. Teuscher ^{[162,ac](#)}, A. Thaler ^{[80](#)}, O. Theiner ^{[57](#)}, T. Theveneaux-Pelzer ^{[105](#)}, O. Thielmann ^{[178](#)}, D.W. Thomas ^{[98](#)}, J.P. Thomas ^{[21](#)}, E.A. Thompson ^{[18a](#)}, P.D. Thompson ^{[21](#)}, E. Thomson ^{[132](#)}, R.E. Thornberry ^{[46](#)}, C. Tian ^{[63](#)}, Y. Tian ^{[57](#)}, V. Tikhomirov ^{[83](#)}, Yu.A. Tikhonov ^{[39](#)}, S. Timoshenko ^{[39](#)}, D. Timoshyn ^{[137](#)}, E.X.L. Ting ^{[1](#)}, P. Tipton ^{[179](#)}, A. Tishelman-Charny ^{[30](#)}, S.H. Tlou ^{[34g](#)}, K. Todome ^{[142](#)}, S. Todorova-Nova ^{[137](#)}, S. Todt ^{[51](#)}, L. Toffolin ^{[70a,70c](#)}, M. Togawa ^{[85](#)}, J. Tojo ^{[91](#)}, S. Tokár ^{[29a](#)}, O. Toldaiev ^{[69](#)}, G. Tolkachev ^{[105](#)}, M. Tomoto ^{[85,114](#)}, L. Tompkins ^{[150,p](#)}, E. Torrence ^{[127](#)}, H. Torres ^{[92](#)}, E. Torró Pastor ^{[170](#)}, M. Toscani ^{[31](#)}, C. Tosciri ^{[41](#)}, M. Tost ^{[11](#)}, D.R. Tovey ^{[146](#)}, T. Trefzger ^{[173](#)}, P.M. Tricarico ^{[13](#)}, A. Tricoli ^{[30](#)}, I.M. Trigger ^{[163a](#)}, S. Trincaz-Duvold ^{[131](#)}, D.A. Trischuk ^{[27](#)}, A. Tropina ^{[40](#)}, L. Truong ^{[34c](#)}, M. Trzebinski ^{[88](#)}, A. Trzupek ^{[88](#)}, F. Tsai ^{[152](#)}, M. Tsai ^{[109](#)}, A. Tsiamis ^{[159](#)}, P.V. Tsiarehka ^{[40](#)}, S. Tsigaridas ^{[163a](#)}, A. Tsirigotis ^{[159,v](#)}, V. Tsiskaridze ^{[162](#)}, E.G. Tskhadadze ^{[156a](#)}, M. Tsopoulou ^{[159](#)}, Y. Tsujikawa ^{[90](#)}, I.I. Tsukerman ^{[39](#)}, V. Tsulaia ^{[18a](#)}, S. Tsuno ^{[85](#)}, K. Tsuru ^{[122](#)}, D. Tsybychev ^{[152](#)}, Y. Tu ^{[65b](#)}, A. Tudorache ^{[28b](#)}, V. Tudorache ^{[28b](#)}, S. Turchikhin ^{[58b,58a](#)}, I. Turk Cakir ^{[3a](#)}, R. Turra ^{[72a](#)}, T. Turtuvshin ^{[40,ad](#)}, P.M. Tuts ^{[43](#)}, S. Tzamarias ^{[159,e](#)}, E. Tzovara ^{[103](#)}, Y. Uematsu ^{[85](#)}, F. Ukegawa ^{[164](#)}, P.A. Ulloa Poblete ^{[141c,141b](#)}, E.N. Umaka ^{[30](#)}, G. Unal ^{[37](#)}, A. Undrus ^{[30](#)}, G. Unel ^{[166](#)}, J. Urban ^{[29b](#)}, P. Urrejola ^{[141a](#)}, G. Usai ^{[8](#)}, R. Ushioda ^{[161](#)}, M. Usman ^{[111](#)}, F. Ustuner ^{[53](#)}, Z. Uysal ^{[83](#)}, V. Vacek ^{[136](#)}, B. Vachon ^{[107](#)}, T. Vafeiadis ^{[37](#)}, A. Vaitkus ^{[99](#)}, C. Valderanis ^{[112](#)}, E. Valdes Santurio ^{[48a,48b](#)}, M. Valente ^{[163a](#)}, S. Valentineti ^{[24b,24a](#)}, A. Valero ^{[170](#)}, E. Valiente Moreno ^{[170](#)}, A. Vallier ^{[92](#)}, J.A. Valls Ferrer ^{[170](#)}, D.R. Van Arneman ^{[118](#)}, T.R. Van Daalen ^{[143](#)}, A. Van Der Graaf ^{[50](#)}, H.Z. Van Der Schyf ^{[34g](#)}, P. Van Gemmeren ^{[6](#)}, M. Van Rijnbach ^{[37](#)}, S. Van Stroud ^{[99](#)}, I. Van Vulpen ^{[118](#)}, P. Vana ^{[137](#)}, M. Vanadia ^{[77a,77b](#)}, U.M. Vande Voorde ^{[151](#)}, W. Vandelli ^{[37](#)}, E.R. Vandewall ^{[125](#)}, D. Vannicola ^{[158](#)}, L. Vannoli ^{[54](#)}, R. Vari ^{[76a](#)}, E.W. Varnes ^{[7](#)}, C. Varni ^{[18b](#)}, D. Varouchas ^{[67](#)}, L. Varriale ^{[170](#)}, K.E. Varvell ^{[154](#)}, M.E. Vasile ^{[28b](#)}, L. Vaslin ^{[85](#)}, M.D. Vassilev ^{[150](#)}, A. Vasyukov ^{[40](#)}, L.M. Vaughan ^{[125](#)}, R. Vavricka ^{[137](#)}, T. Vazquez Schroeder ^{[13](#)}, J. Veatch ^{[32](#)}, V. Vecchio ^{[104](#)}, M.J. Veen ^{[106](#)}, I. Veliscek ^{[30](#)}, L.M. Veloce ^{[162](#)}, F. Veloso ^{[134a,134c](#)}, S. Veneziano ^{[76a](#)}, A. Ventura ^{[71a,71b](#)}, S. Ventura Gonzalez ^{[139](#)}, A. Verbytskyi ^{[113](#)}, M. Verducci ^{[75a,75b](#)}, C. Vergis ^{[97](#)}, M. Verissimo De Araujo ^{[84b](#)}, W. Verkerke ^{[118](#)},

J.C. Vermeulen ^{[118](#)}, C. Vernieri ^{[150](#)}, M. Vessella ^{[166](#)}, M.C. Vetterli ^{[149,aj](#)}, A. Vgenopoulos ^{[103](#)}, N. Viaux Maira ^{[141f](#)}, T. Vickey ^{[146](#)}, O.E. Vickey Boeriu ^{[146](#)}, G.H.A. Viehhauser ^{[130](#)}, L. Vigani ^{[64b](#)}, M. Vigi ^{[113](#)}, M. Villa ^{[24b,24a](#)}, M. Villaplana Perez ^{[170](#)}, E.M. Villhauer ^{[53](#)}, E. Vilucchi ^{[54](#)}, M.G. Vincter ^{[35](#)}, A. Visibile ^{[118](#)}, C. Vittori ^{[37](#)}, I. Vivarelli ^{[24b,24a](#)}, E. Voevodina ^{[113](#)}, F. Vogel ^{[112](#)}, J.C. Voigt ^{[51](#)}, P. Vokac ^{[136](#)}, Yu. Volkotrub ^{[87b](#)}, E. Von Toerne ^{[25](#)}, B. Vormwald ^{[37](#)}, K. Vorobev ^{[39](#)}, M. Vos ^{[170](#)}, K. Voss ^{[148](#)}, M. Vozak ^{[37](#)}, L. Vozdecky ^{[124](#)}, N. Vranjes ^{[16](#)}, M. Vranjes Milosavljevic ^{[16](#)}, M. Vreeswijk ^{[118](#)}, N.K. Vu ^{[145b,145a](#)}, R. Vuillermet ^{[37](#)}, O. Vujinovic ^{[103](#)}, I. Vukotic ^{[41](#)}, I.K. Vyas ^{[35](#)}, J.F. Wack ^{[33](#)}, S. Wada ^{[164](#)}, C. Wagner ^{[150](#)}, J.M. Wagner ^{[18a](#)}, W. Wagner ^{[178](#)}, S. Wahdan ^{[178](#)}, H. Wahlberg ^{[93](#)}, C.H. Waits ^{[124](#)}, J. Walder ^{[138](#)}, R. Walker ^{[112](#)}, W. Walkowiak ^{[148](#)}, A. Wall ^{[132](#)}, E.J. Wallin ^{[101](#)}, T. Wamorkar ^{[18a](#)}, A.Z. Wang ^{[140](#)}, C. Wang ^{[103](#)}, C. Wang ^{[11](#)}, H. Wang ^{[18a](#)}, J. Wang ^{[65c](#)}, P. Wang ^{[104](#)}, P. Wang ^{[99](#)}, R. Wang ^{[62](#)}, R. Wang ^{[6](#)}, S.M. Wang ^{[155](#)}, S. Wang ^{[14](#)}, T. Wang ^{[63](#)}, T. Wang ^{[63](#)}, W.T. Wang ^{[81](#)}, W. Wang ^{[14](#)}, X. Wang ^{[169](#)}, X. Wang ^{[145a](#)}, X. Wang ^{[49](#)}, Y. Wang ^{[115a](#)}, Y. Wang ^{[63](#)}, Z. Wang ^{[109](#)}, Z. Wang ^{[145b,52,145a](#)}, Z. Wang ^{[109](#)}, C. Wanotayaroj ^{[85](#)}, A. Warburton ^{[107](#)}, R.J. Ward ^{[21](#)}, A.L. Warnerbring ^{[148](#)}, N. Warrack ^{[60](#)}, S. Waterhouse ^{[98](#)}, A.T. Watson ^{[21](#)}, H. Watson ^{[53](#)}, M.F. Watson ^{[21](#)}, E. Watton ^{[60](#)}, G. Watts ^{[143](#)}, B.M. Waugh ^{[99](#)}, J.M. Webb ^{[55](#)}, C. Weber ^{[30](#)}, H.A. Weber ^{[19](#)}, M.S. Weber ^{[20](#)}, S.M. Weber ^{[64a](#)}, C. Wei ^{[63](#)}, Y. Wei ^{[55](#)}, A.R. Weidberg ^{[130](#)}, E.J. Weik ^{[121](#)}, J. Weingarten ^{[50](#)}, C. Weiser ^{[55](#)}, C.J. Wells ^{[49](#)}, T. Wenaus ^{[30](#)}, B. Wendland ^{[50](#)}, T. Wengler ^{[37](#)}, N.S. Wenke ^{[113](#)}, N. Wermes ^{[25](#)}, M. Wessels ^{[64a](#)}, A.M. Wharton ^{[94](#)}, A.S. White ^{[62](#)}, A. White ^{[8](#)}, M.J. White ^{[1](#)}, D. Whiteson ^{[166](#)}, L. Wickremasinghe ^{[128](#)}, W. Wiedenmann ^{[177](#)}, M. Wielers ^{[138](#)}, C. Wiglesworth ^{[44](#)}, D.J. Wilbern ^{[124](#)}, H.G. Wilkens ^{[37](#)}, J.J.H. Wilkinson ^{[33](#)}, D.M. Williams ^{[43](#)}, H.H. Williams ^{[132](#)}, S. Williams ^{[33](#)}, S. Willocq ^{[106](#)}, B.J. Wilson ^{[104](#)}, D.J. Wilson ^{[104](#)}, P.J. Windischhofer ^{[41](#)}, F.I. Winkel ^{[31](#)}, F. Winklmeier ^{[127](#)}, B.T. Winter ^{[55](#)}, M. Wittgen ^{[150](#)}, M. Wobisch ^{[100](#)}, T. Wojtkowski ^{[61](#)}, Z. Wolffs ^{[118](#)}, J. Wollrath ^{[37](#)}, M.W. Wolter ^{[88](#)}, H. Wolters ^{[134a,134c](#)}, M.C. Wong ^{[140](#)}, E.L. Woodward ^{[43](#)}, S.D. Worm ^{[49](#)}, B.K. Wosiek ^{[88](#)}, K.W. Woźniak ^{[88](#)}, S. Wozniowski ^{[56](#)}, K. Wraight ^{[60](#)}, C. Wu ^{[21](#)}, M. Wu ^{[115b](#)}, M. Wu ^{[117](#)}, S.L. Wu ^{[177](#)}, S. Wu ^{[14](#)}, X. Wu ^{[57](#)}, X. Wu ^{[63](#)}, Y. Wu ^{[63](#)}, Z. Wu ^{[4](#)}, J. Wuerzinger ^{[113,ah](#)}, T.R. Wyatt ^{[104](#)}, B.M. Wynne ^{[53](#)}, S. Xella ^{[44](#)}, L. Xia ^{[115a](#)}, M. Xia ^{[15](#)}, M. Xie ^{[63](#)}, A. Xiong ^{[127](#)}, J. Xiong ^{[18a](#)}, D. Xu ^{[14](#)}, H. Xu ^{[63](#)}, L. Xu ^{[63](#)}, R. Xu ^{[132](#)}, T. Xu ^{[109](#)}, Y. Xu ^{[143](#)}, Z. Xu ^{[53](#)}, Z. Xu ^{[115a](#)}, B. Yabsley ^{[154](#)}, S. Yacoob ^{[34a](#)}, Y. Yamaguchi ^{[85](#)}, E. Yamashita ^{[160](#)}, H. Yamauchi ^{[164](#)}, T. Yamazaki ^{[18a](#)}, Y. Yamazaki ^{[86](#)}, S. Yan ^{[60](#)}, Z. Yan ^{[106](#)}, H.J. Yang ^{[145a,145b](#)}, H.T. Yang ^{[63](#)}, S. Yang ^{[63](#)}, T. Yang ^{[65c](#)}, X. Yang ^{[37](#)}, X. Yang ^{[14](#)}, Y. Yang ^{[160](#)}, Y. Yang ^{[63](#)}, W-M. Yao ^{[18a](#)}, C.L. Yardley ^{[153](#)}, H. Ye ^{[56](#)}, J. Ye ^{[14](#)}, S. Ye ^{[30](#)}, X. Ye ^{[63](#)}, Y. Yeh ^{[99](#)}, I. Yeletsikh ^{[40](#)}, B. Yeo ^{[18b](#)}, M.R. Yexley ^{[99](#)}, T.P. Yildirim ^{[130](#)}, P. Yin ^{[43](#)}, K. Yorita ^{[175](#)}, S. Younas ^{[28b](#)}, C.J.S. Young ^{[37](#)}, C. Young ^{[150](#)}, N.D. Young ^{[127](#)}, Y. Yu ^{[63](#)}, J. Yuan ^{[14,115c](#)}, M. Yuan ^{[109](#)}, R. Yuan ^{[145b,145a](#)}, L. Yue ^{[99](#)}, M. Zaazoua ^{[63](#)}, B. Zabinski ^{[88](#)}, I. Zahir ^{[36a](#)}, A. Zaio ^{[58b,58a](#)}, Z.K. Zak ^{[88](#)}, T. Zakareishvili ^{[170](#)}, S. Zambito ^{[57](#)}, J.A. Zamora Saa ^{[141d](#)}, J. Zang ^{[160](#)}, D. Zanzi ^{[55](#)}, R. Zanzottera ^{[72a,72b](#)}, O. Zaplatilek ^{[136](#)}, C. Zeitnitz ^{[178](#)}, H. Zeng ^{[14](#)}, J.C. Zeng ^{[169](#)}, D.T. Zenger Jr ^{[27](#)}, O. Zenin ^{[39](#)}, T. Ženiš ^{[29a](#)}, S. Zenz ^{[97](#)}, S. Zerradi ^{[36a](#)}, D. Zerwas ^{[67](#)}, M. Zhai ^{[14,115c](#)}, D.F. Zhang ^{[146](#)}, J. Zhang ^{[144a](#)}, J. Zhang ^{[6](#)}, K. Zhang ^{[14,115c](#)}, L. Zhang ^{[63](#)}, L. Zhang ^{[115a](#)}, P. Zhang ^{[14,115c](#)}, R. Zhang ^{[177](#)}, S. Zhang ^{[92](#)}, T. Zhang ^{[160](#)}, X. Zhang ^{[145a](#)}, Y. Zhang ^{[143](#)}, Y. Zhang ^{[99](#)}, Y. Zhang ^{[63](#)},

Y. Zhang ^{115a}, Z. Zhang ^{18a}, Z. Zhang ^{144a}, Z. Zhang ⁶⁷, H. Zhao ¹⁴³, T. Zhao ^{144a},
Y. Zhao ³⁵, Z. Zhao ⁶³, Z. Zhao ⁶³, A. Zhemchugov ⁴⁰, J. Zheng ^{115a}, K. Zheng ¹⁶⁹,
X. Zheng ⁶³, Z. Zheng ¹⁵⁰, D. Zhong ¹⁶⁹, B. Zhou ¹⁰⁹, H. Zhou ⁷, N. Zhou ^{145a}, Y. Zhou ¹⁵,
Y. Zhou ^{115a}, Y. Zhou⁷, C.G. Zhu ^{144a}, J. Zhu ¹⁰⁹, X. Zhu^{145b}, Y. Zhu ^{145a}, Y. Zhu ⁶³,
X. Zhuang ¹⁴, K. Zhukov ⁶⁹, N.I. Zimine ⁴⁰, J. Zinsser ^{64b}, M. Ziolkowski ¹⁴⁸, L. Živković ¹⁶,
A. Zoccoli ^{24b,24a}, K. Zoch ⁶², T.G. Zorbas ¹⁴⁶, O. Zormpa ⁴⁷, W. Zou ⁴³, L. Zwalinski ³⁷

¹ Department of Physics, University of Adelaide, Adelaide; Australia

² Department of Physics, University of Alberta, Edmonton AB; Canada

³ ^(a) Department of Physics, Ankara University, Ankara; ^(b) Division of Physics, TOBB University of Economics and Technology, Ankara; Türkiye

⁴ LAPP, Université Savoie Mont Blanc, CNRS/IN2P3, Annecy; France

⁵ APC, Université Paris Cité, CNRS/IN2P3, Paris; France

⁶ High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America

⁷ Department of Physics, University of Arizona, Tucson AZ; United States of America

⁸ Department of Physics, University of Texas at Arlington, Arlington TX; United States of America

⁹ Physics Department, National and Kapodistrian University of Athens, Athens; Greece

¹⁰ Physics Department, National Technical University of Athens, Zografou; Greece

¹¹ Department of Physics, University of Texas at Austin, Austin TX; United States of America

¹² Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan

¹³ Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain

¹⁴ Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; China

¹⁵ Physics Department, Tsinghua University, Beijing; China

¹⁶ Institute of Physics, University of Belgrade, Belgrade; Serbia

¹⁷ Department for Physics and Technology, University of Bergen, Bergen; Norway

¹⁸ ^(a) Physics Division, Lawrence Berkeley National Laboratory, Berkeley CA; ^(b) University of California, Berkeley CA; United States of America

¹⁹ Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany

²⁰ Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland

²¹ School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom

²² ^(a) Department of Physics, Bogazici University, Istanbul; ^(b) Department of Physics Engineering, Gaziantep University, Gaziantep; ^(c) Department of Physics, Istanbul University, Istanbul; Türkiye

²³ ^(a) Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogotá; ^(b) Departamento de Física, Universidad Nacional de Colombia, Bogotá; Colombia

²⁴ ^(a) Dipartimento di Fisica e Astronomia A. Righi, Università di Bologna, Bologna; ^(b) INFN Sezione di Bologna; Italy

²⁵ Physikalisches Institut, Universität Bonn, Bonn; Germany

²⁶ Department of Physics, Boston University, Boston MA; United States of America

²⁷ Department of Physics, Brandeis University, Waltham MA; United States of America

²⁸ ^(a) Transilvania University of Brasov, Brasov; ^(b) Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest; ^(c) Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi; ^(d) National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca; ^(e) National University of Science and Technology Politehnica, Bucharest; ^(f) West University in Timisoara, Timisoara; ^(g) Faculty of Physics, University of Bucharest, Bucharest; Romania

²⁹ ^(a) Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava; ^(b) Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic

³⁰ Physics Department, Brookhaven National Laboratory, Upton NY; United States of America

³¹ Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Física, y CONICET, Instituto de Física de Buenos Aires (IFIBA), Buenos Aires; Argentina

- ³² *California State University, CA; United States of America*
- ³³ *Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom*
- ³⁴ ^(a) *Department of Physics, University of Cape Town, Cape Town;* ^(b) *iThemba Labs, Western Cape;* ^(c) *Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg;* ^(d) *National Institute of Physics, University of the Philippines Diliman (Philippines);* ^(e) *University of South Africa, Department of Physics, Pretoria;* ^(f) *University of Zululand, KwaDlangezwa;* ^(g) *School of Physics, University of the Witwatersrand, Johannesburg; South Africa*
- ³⁵ *Department of Physics, Carleton University, Ottawa ON; Canada*
- ³⁶ ^(a) *Faculté des Sciences Ain Chock, Université Hassan II de Casablanca;* ^(b) *Faculté des Sciences, Université Ibn-Tofail, Kénitra;* ^(c) *Faculté des Sciences Semailia, Université Cadi Ayyad, LPHEA-Marrakech;* ^(d) *LPMR, Faculté des Sciences, Université Mohamed Premier, Oujda;* ^(e) *Faculté des sciences, Université Mohammed V, Rabat;* ^(f) *Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco*
- ³⁷ *CERN, Geneva; Switzerland*
- ³⁸ *Affiliated with an institute formerly covered by a cooperation agreement with CERN*
- ³⁹ *Affiliated with an institute covered by a cooperation agreement with CERN*
- ⁴⁰ *Affiliated with an international laboratory covered by a cooperation agreement with CERN*
- ⁴¹ *Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America*
- ⁴² *LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France*
- ⁴³ *Nevis Laboratory, Columbia University, Irvington NY; United States of America*
- ⁴⁴ *Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark*
- ⁴⁵ ^(a) *Dipartimento di Fisica, Università della Calabria, Rende;* ^(b) *INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati; Italy*
- ⁴⁶ *Physics Department, Southern Methodist University, Dallas TX; United States of America*
- ⁴⁷ *National Centre for Scientific Research “Demokritos”, Agia Paraskevi; Greece*
- ⁴⁸ ^(a) *Department of Physics, Stockholm University;* ^(b) *Oskar Klein Centre, Stockholm; Sweden*
- ⁴⁹ *Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany*
- ⁵⁰ *Fakultät Physik, Technische Universität Dortmund, Dortmund; Germany*
- ⁵¹ *Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany*
- ⁵² *Department of Physics, Duke University, Durham NC; United States of America*
- ⁵³ *SUPA — School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom*
- ⁵⁴ *INFN e Laboratori Nazionali di Frascati, Frascati; Italy*
- ⁵⁵ *Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany*
- ⁵⁶ *II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany*
- ⁵⁷ *Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland*
- ⁵⁸ ^(a) *Dipartimento di Fisica, Università di Genova, Genova;* ^(b) *INFN Sezione di Genova; Italy*
- ⁵⁹ *II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany*
- ⁶⁰ *SUPA — School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom*
- ⁶¹ *LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France*
- ⁶² *Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America*
- ⁶³ *Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei; China*
- ⁶⁴ ^(a) *Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg;* ^(b) *Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany*
- ⁶⁵ ^(a) *Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong;* ^(b) *Department of Physics, University of Hong Kong, Hong Kong;* ^(c) *Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong; China*
- ⁶⁶ *Department of Physics, National Tsing Hua University, Hsinchu; Taiwan*
- ⁶⁷ *IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91405, Orsay; France*
- ⁶⁸ *Centro Nacional de Microelectrónica (IMB-CNM-CSIC), Barcelona; Spain*
- ⁶⁹ *Department of Physics, Indiana University, Bloomington IN; United States of America*
- ⁷⁰ ^(a) *INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine;* ^(b) *ICTP, Trieste;* ^(c) *Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine; Italy*

- ⁷¹ ^(a) INFN Sezione di Lecce; ^(b) Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy
- ⁷² ^(a) INFN Sezione di Milano; ^(b) Dipartimento di Fisica, Università di Milano, Milano; Italy
- ⁷³ ^(a) INFN Sezione di Napoli; ^(b) Dipartimento di Fisica, Università di Napoli, Napoli; Italy
- ⁷⁴ ^(a) INFN Sezione di Pavia; ^(b) Dipartimento di Fisica, Università di Pavia, Pavia; Italy
- ⁷⁵ ^(a) INFN Sezione di Pisa; ^(b) Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy
- ⁷⁶ ^(a) INFN Sezione di Roma; ^(b) Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy
- ⁷⁷ ^(a) INFN Sezione di Roma Tor Vergata; ^(b) Dipartimento di Fisica, Università di Roma Tor Vergata, Roma; Italy
- ⁷⁸ ^(a) INFN Sezione di Roma Tre; ^(b) Dipartimento di Matematica e Fisica, Università Roma Tre, Roma; Italy
- ⁷⁹ ^(a) INFN-TIFPA; ^(b) Università degli Studi di Trento, Trento; Italy
- ⁸⁰ Universität Innsbruck, Department of Astro and Particle Physics, Innsbruck; Austria
- ⁸¹ University of Iowa, Iowa City IA; United States of America
- ⁸² Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America
- ⁸³ Istinye University, Sariyer, Istanbul; Türkiye
- ⁸⁴ ^(a) Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora; ^(b) Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; ^(c) Instituto de Física, Universidade de São Paulo, São Paulo; ^(d) Rio de Janeiro State University, Rio de Janeiro; ^(e) Federal University of Bahia, Bahia; Brazil
- ⁸⁵ KEK, High Energy Accelerator Research Organization, Tsukuba; Japan
- ⁸⁶ Graduate School of Science, Kobe University, Kobe; Japan
- ⁸⁷ ^(a) AGH University of Krakow, Faculty of Physics and Applied Computer Science, Krakow; ^(b) Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland
- ⁸⁸ Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland
- ⁸⁹ ^(a) Khalifa University of Science and Technology, Abu Dhabi; ^(b) University of Sharjah, Sharjah; United Arab Emirates
- ⁹⁰ Faculty of Science, Kyoto University, Kyoto; Japan
- ⁹¹ Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka; Japan
- ⁹² L2IT, Université de Toulouse, CNRS/IN2P3, UPS, Toulouse; France
- ⁹³ Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina
- ⁹⁴ Physics Department, Lancaster University, Lancaster; United Kingdom
- ⁹⁵ Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom
- ⁹⁶ Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia
- ⁹⁷ Department of Physics and Astronomy, Queen Mary University of London, London; United Kingdom
- ⁹⁸ Department of Physics, Royal Holloway University of London, Egham; United Kingdom
- ⁹⁹ Department of Physics and Astronomy, University College London, London; United Kingdom
- ¹⁰⁰ Louisiana Tech University, Ruston LA; United States of America
- ¹⁰¹ Fysiska institutionen, Lunds universitet, Lund; Sweden
- ¹⁰² Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain
- ¹⁰³ Institut für Physik, Universität Mainz, Mainz; Germany
- ¹⁰⁴ School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom
- ¹⁰⁵ CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France
- ¹⁰⁶ Department of Physics, University of Massachusetts, Amherst MA; United States of America
- ¹⁰⁷ Department of Physics, McGill University, Montreal QC; Canada
- ¹⁰⁸ School of Physics, University of Melbourne, Victoria; Australia
- ¹⁰⁹ Department of Physics, University of Michigan, Ann Arbor MI; United States of America
- ¹¹⁰ Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America
- ¹¹¹ Group of Particle Physics, University of Montreal, Montreal QC; Canada
- ¹¹² Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany
- ¹¹³ Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München; Germany
- ¹¹⁴ Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan

- ¹¹⁵ ^(a) Department of Physics, Nanjing University, Nanjing; ^(b) School of Science, Shenzhen Campus of Sun Yat-sen University; ^(c) University of Chinese Academy of Science (UCAS), Beijing; China
- ¹¹⁶ Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America
- ¹¹⁷ Institute for Mathematics, Astrophysics and Particle Physics, Radboud University/Nikhef, Nijmegen; Netherlands
- ¹¹⁸ Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands
- ¹¹⁹ Department of Physics, Northern Illinois University, DeKalb IL; United States of America
- ¹²⁰ ^(a) New York University Abu Dhabi, Abu Dhabi; ^(b) United Arab Emirates University, Al Ain; United Arab Emirates
- ¹²¹ Department of Physics, New York University, New York NY; United States of America
- ¹²² Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan
- ¹²³ Ohio State University, Columbus OH; United States of America
- ¹²⁴ Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America
- ¹²⁵ Department of Physics, Oklahoma State University, Stillwater OK; United States of America
- ¹²⁶ Palacký University, Joint Laboratory of Optics, Olomouc; Czech Republic
- ¹²⁷ Institute for Fundamental Science, University of Oregon, Eugene, OR; United States of America
- ¹²⁸ Graduate School of Science, University of Osaka, Osaka; Japan
- ¹²⁹ Department of Physics, University of Oslo, Oslo; Norway
- ¹³⁰ Department of Physics, Oxford University, Oxford; United Kingdom
- ¹³¹ LPNHE, Sorbonne Université, Université Paris Cité, CNRS/IN2P3, Paris; France
- ¹³² Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America
- ¹³³ Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America
- ¹³⁴ ^(a) Laboratório de Instrumentação e Física Experimental de Partículas — LIP, Lisboa; ^(b) Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa; ^(c) Departamento de Física, Universidade de Coimbra, Coimbra; ^(d) Centro de Física Nuclear da Universidade de Lisboa, Lisboa; ^(e) Departamento de Física, Escola de Ciências, Universidade do Minho, Braga; ^(f) Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain); ^(g) Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal
- ¹³⁵ Institute of Physics of the Czech Academy of Sciences, Prague; Czech Republic
- ¹³⁶ Czech Technical University in Prague, Prague; Czech Republic
- ¹³⁷ Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic
- ¹³⁸ Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom
- ¹³⁹ IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France
- ¹⁴⁰ Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America
- ¹⁴¹ ^(a) Departamento de Física, Pontificia Universidad Católica de Chile, Santiago; ^(b) Millennium Institute for Subatomic physics at high energy frontier (SAPHIR), Santiago; ^(c) Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, y Departamento de Física, Universidad de La Serena; ^(d) Universidad Andres Bello, Department of Physics, Santiago; ^(e) Instituto de Alta Investigación, Universidad de Tarapacá, Arica; ^(f) Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso; Chile
- ¹⁴² Department of Physics, Institute of Science, Tokyo; Japan
- ¹⁴³ Department of Physics, University of Washington, Seattle WA; United States of America
- ¹⁴⁴ ^(a) Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao; ^(b) School of Physics, Zhengzhou University; China
- ¹⁴⁵ ^(a) State Key Laboratory of Dark Matter Physics, School of Physics and Astronomy, Shanghai Jiao Tong University, Key Laboratory for Particle Astrophysics and Cosmology (MOE), SKLPPC, Shanghai; ^(b) State Key Laboratory of Dark Matter Physics, Tsung-Dao Lee Institute, Shanghai Jiao Tong University, Shanghai; China
- ¹⁴⁶ Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom
- ¹⁴⁷ Department of Physics, Shinshu University, Nagano; Japan

- ¹⁴⁸ *Department Physik, Universität Siegen, Siegen; Germany*
- ¹⁴⁹ *Department of Physics, Simon Fraser University, Burnaby BC; Canada*
- ¹⁵⁰ *SLAC National Accelerator Laboratory, Stanford CA; United States of America*
- ¹⁵¹ *Department of Physics, Royal Institute of Technology, Stockholm; Sweden*
- ¹⁵² *Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America*
- ¹⁵³ *Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom*
- ¹⁵⁴ *School of Physics, University of Sydney, Sydney; Australia*
- ¹⁵⁵ *Institute of Physics, Academia Sinica, Taipei; Taiwan*
- ¹⁵⁶ ^(a) *E. Andronikashvili Institute of Physics, Iv. Javakhishvili Tbilisi State University, Tbilisi;* ^(b) *High Energy Physics Institute, Tbilisi State University, Tbilisi;* ^(c) *University of Georgia, Tbilisi; Georgia*
- ¹⁵⁷ *Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel*
- ¹⁵⁸ *Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel*
- ¹⁵⁹ *Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece*
- ¹⁶⁰ *International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan*
- ¹⁶¹ *Graduate School of Science and Technology, Tokyo Metropolitan University, Tokyo; Japan*
- ¹⁶² *Department of Physics, University of Toronto, Toronto ON; Canada*
- ¹⁶³ ^(a) *TRIUMF, Vancouver BC;* ^(b) *Department of Physics and Astronomy, York University, Toronto ON; Canada*
- ¹⁶⁴ *Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan*
- ¹⁶⁵ *Department of Physics and Astronomy, Tufts University, Medford MA; United States of America*
- ¹⁶⁶ *Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of America*
- ¹⁶⁷ *University of West Attica, Athens; Greece*
- ¹⁶⁸ *Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden*
- ¹⁶⁹ *Department of Physics, University of Illinois, Urbana IL; United States of America*
- ¹⁷⁰ *Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia — CSIC, Valencia; Spain*
- ¹⁷¹ *Department of Physics, University of British Columbia, Vancouver BC; Canada*
- ¹⁷² *Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada*
- ¹⁷³ *Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany*
- ¹⁷⁴ *Department of Physics, University of Warwick, Coventry; United Kingdom*
- ¹⁷⁵ *Waseda University, Tokyo; Japan*
- ¹⁷⁶ *Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot; Israel*
- ¹⁷⁷ *Department of Physics, University of Wisconsin, Madison WI; United States of America*
- ¹⁷⁸ *Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany*
- ¹⁷⁹ *Department of Physics, Yale University, New Haven CT; United States of America*
- ¹⁸⁰ *Yerevan Physics Institute, Yerevan; Armenia*

^a *Also Affiliated with an institute covered by a cooperation agreement with CERN*

^b *Also at An-Najah National University, Nablus; Palestine*

^c *Also at Borough of Manhattan Community College, City University of New York, New York NY; United States of America*

^d *Also at Center for High Energy Physics, Peking University; China*

^e *Also at Center for Interdisciplinary Research and Innovation (CIRI-AUTH), Thessaloniki; Greece*

^f *Also at CERN, Geneva; Switzerland*

^g *Also at CMD-AC UNEC Research Center, Azerbaijan State University of Economics (UNEC); Azerbaijan*

^h *Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland*

ⁱ *Also at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona; Spain*

^j *Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece*

^k *Also at Department of Mathematical Sciences, University of South Africa, Johannesburg; South Africa*

- ^l Also at Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei; China
- ^m Also at Department of Physics, Bolu Abant Izzet Baysal University, Bolu; Türkiye
- ⁿ Also at Department of Physics, California State University, Sacramento; United States of America
- ^o Also at Department of Physics, King's College London, London; United Kingdom
- ^p Also at Department of Physics, Stanford University, Stanford CA; United States of America
- ^q Also at Department of Physics, Stellenbosch University; South Africa
- ^r Also at Department of Physics, University of Fribourg, Fribourg; Switzerland
- ^s Also at Department of Physics, University of Thessaly; Greece
- ^t Also at Department of Physics, Westmont College, Santa Barbara; United States of America
- ^u Also at Faculty of Physics, Sofia University, 'St. Kliment Ohridski', Sofia; Bulgaria
- ^v Also at Hellenic Open University, Patras; Greece
- ^w Also at Henan University; China
- ^x Also at Imam Mohammad Ibn Saud Islamic University; Saudi Arabia
- ^y Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain
- ^z Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany
- ^{aa} Also at Institute for Nuclear Research and Nuclear Energy (INRNE) of the Bulgarian Academy of Sciences, Sofia; Bulgaria
- ^{ab} Also at Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco
- ^{ac} Also at Institute of Particle Physics (IPP); Canada
- ^{ad} Also at Institute of Physics and Technology, Mongolian Academy of Sciences, Ulaanbaatar; Mongolia
- ^{ae} Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan
- ^{af} Also at Institute of Theoretical Physics, Ilia State University, Tbilisi; Georgia
- ^{ag} Also at National Institute of Physics, University of the Philippines Diliman (Philippines); Philippines
- ^{ah} Also at Technical University of Munich, Munich; Germany
- ^{ai} Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing; China
- ^{aj} Also at TRIUMF, Vancouver BC; Canada
- ^{ak} Also at Università di Napoli Parthenope, Napoli; Italy
- ^{al} Also at University of Colorado Boulder, Department of Physics, Colorado; United States of America
- ^{am} Also at University of the Western Cape; South Africa
- ^{an} Also at Washington College, Chestertown, MD; United States of America
- ^{ao} Also at Yeditepe University, Physics Department, Istanbul; Türkiye
- * Deceased