



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/230740/>

Version: Published Version

Article:

Bavula, V.V. (2022) Localizable sets and the localization of a ring at a localizable set. Journal of Algebra, 610. pp. 38-75. ISSN: 0021-8693

<https://doi.org/10.1016/j.jalgebra.2022.06.034>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Contents lists available at ScienceDirect

Journal of Algebra

www.elsevier.com/locate/jalgebra



Localizable sets and the localization of a ring at a localizable set



V.V. Bavula

Department of Pure Mathematics, University of Sheffield, Hicks Building, Sheffield S3 7RH, UK

ARTICLE INFO

Article history:

Received 7 June 2021
Available online 26 July 2022
Communicated by Louis Rowen

MSC:

16S85
16P50
16P60
16U20

Keywords:

Localizable set
Localization of a ring at a localizable set
Goldie's theorem
The left quotient ring of a ring
The largest left quotient ring of a ring
Maximal localizable set
Maximal left denominator set
The left localization radical of a ring
Maximal left localization of a ring

ABSTRACT

The aim of the paper is to develop the most general theory of one-sided fractions. The concepts of localizable set, localization of a ring and a module at a localizable set are introduced and studied. Localizable sets are generalization of Ore sets and denominator sets, and the localization of a ring/module at a localizable set is a generalization of localization of a ring/module at a denominator set. For a semiprime left Goldie ring, it is proven that the set of maximal left localizable sets that contain all regular elements is equal to the set of maximal left denominator sets (and they are explicitly described). For a semiprime Goldie ring, it is proven that the following five sets coincide: the maximal Ore sets, the maximal denominator sets, the maximal left or right or two-sided localizable sets that contain all regular elements (and they are explicitly described).

© 2022 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

1. Introduction 39

E-mail address: v.bavula@sheffield.ac.uk.

<https://doi.org/10.1016/j.jalgebra.2022.06.034>

0021-8693/© 2022 The Author(s). Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

2. Localizable sets and the localization of a ring at a localizable set	47
3. Localization of a ring at an Ore set	59
4. Localization of a ring at an almost Ore set	62
5. Classification of maximal localizable sets and maximal Ore sets in semiprime Goldie ring . . .	65
6. Localization of a module at a localizable set	68
7. Examples	70
References	75

1. Introduction

In the paper all rings are unital. When we say ring we mean a K -algebra over a commutative ring K that belongs to the centre of the ring.

The goal of the paper is to start to develop the most general theory of *one-sided fractions*. For that the following new concepts are introduced: the almost Ore set, the localizable set and the localizable perfect set. Their relations are given by the chain of inclusions:

$$\{\text{Denominator sets}\} \subseteq \{\text{Ore sets}\} \subseteq \{\text{almost Ore sets}\} \subseteq \{\text{perfect localizable sets}\} \\ \subseteq \{\text{localizable sets}\}.$$

The ring $R\langle S^{-1} \rangle$. Let R be a ring and S be a multiplicative set in R (that is $SS \subseteq S$, $1 \in S$ and $0 \notin S$). Let $R\langle X_S \rangle$ be a ring freely generated by the ring R and a set $X_S = \{x_s \mid s \in S\}$ of free noncommutative indeterminates (indexed by the elements of the set S). Let us consider the factor ring

$$R\langle S^{-1} \rangle := R\langle X_S \rangle / I_S \tag{1}$$

of the ring $R\langle X_S \rangle$ at the ideal I_S generated by the set of elements $\{sx_s - 1, x_s s - 1 \mid s \in S\}$.

The kernel of the ring homomorphism

$$R \rightarrow R\langle S^{-1} \rangle, \quad r \mapsto r + I_S \tag{2}$$

is denoted by $\text{ass}(S) = \text{ass}_R(S)$. The ideal $\text{ass}_R(S)$ of R has a complex structure, its description is given in Proposition 2.12 when S is a left localizable set. Lemma 1.2, Proposition 1.1.(1) and its proof describe a large chunk of the ideal $\text{ass}_R(S)$, which is the ideal $\mathfrak{a}(S)$. The proof of Proposition 1.1 contains an explicit description of the ideal $\mathfrak{a}(S)$. The ideal $\mathfrak{a}(S)$ is the key part in the definition of perfect localizable sets.

Localizable sets.

Definition. A multiplicative set S of a ring R is called a *left localizable set* of R if

$$R\langle S^{-1} \rangle = \{\bar{s}^{-1}\bar{r} \mid \bar{s} \in \bar{S}, \bar{r} \in \bar{R}\} \neq \{0\}$$

where $\overline{R} = R/\mathfrak{a}$, $\mathfrak{a} = \text{ass}_R(S)$ and $\overline{S} = (S + \mathfrak{a})/\mathfrak{a}$, i.e., every element of the ring $R\langle S^{-1} \rangle$ is a left fraction $\overline{s}^{-1}\overline{r}$ for some elements $\overline{s} \in \overline{S}$ and $\overline{r} \in \overline{R}$. Similarly, a multiplicative set S of a ring R is called a *right localizable set* of R if

$$R\langle S^{-1} \rangle = \{\overline{r}\overline{s}^{-1} \mid \overline{s} \in \overline{S}, \overline{r} \in \overline{R}\} \neq \{0\},$$

i.e., every element of the ring $R\langle S^{-1} \rangle$ is a right fraction $\overline{r}\overline{s}^{-1}$ for some elements $\overline{s} \in \overline{S}$ and $\overline{r} \in \overline{R}$. A right and left localizable set of R is called a *localizable set* of R .

The sets of left localizable, right localizable and localizable sets of R are denoted by $\mathbb{L}_l(R)$, $\mathbb{L}_r(R)$ and $\mathbb{L}(R)$, respectively. Clearly, $\mathbb{L}(R) = \mathbb{L}_l(R) \cap \mathbb{L}_r(R)$. In order to study these three sets simultaneously we use the following notation $\mathbb{L}_*(R)$ where $*$ \in $\{l, r, \emptyset\}$ and \emptyset is the empty set ($\mathbb{L}(R) = \mathbb{L}_\emptyset(R)$). Let

$$\text{ass } \mathbb{L}_*(R) = \{\text{ass}_R(S) \mid S \in \mathbb{L}_*(R)\}. \tag{3}$$

For an ideal \mathfrak{a} of R , let $\mathbb{L}_*(R, \mathfrak{a}) = \{S \in \mathbb{L}_*(R) \mid \text{ass}_R(S) = \mathfrak{a}\}$. Then

$$\mathbb{L}_*(R) = \coprod_{\mathfrak{a} \in \text{ass } \mathbb{L}_*(R)} \mathbb{L}_*(R, \mathfrak{a}) \tag{4}$$

is a disjoint union of non-empty sets.

The ideals $\mathfrak{a}(S)$, $\mathfrak{a}'(S)$ and $\mathfrak{a}''(S)$. For each element $r \in R$, let $r \cdot : R \rightarrow R, x \mapsto rx$ and $\cdot r : R \rightarrow R, x \mapsto xr$. The sets $'\mathcal{C}_R := \{r \in R \mid \ker(r \cdot) = 0\}$ and $\mathcal{C}'_R := \{r \in R \mid \ker(\cdot r) = 0\}$ are called the *sets of left and right regular elements* of R , respectively. Their intersection $\mathcal{C}_R = '\mathcal{C}_R \cap \mathcal{C}'_R$ is the *set of regular elements* of R . The rings $Q_{l,cl}(R) := \mathcal{C}_R^{-1}R$ and $Q_{r,cl}(R) := RC_R^{-1}$ are called the *classical left and right quotient rings* of R , respectively. Goldie’s Theorem states that the ring $Q_{l,cl}(R)$ is a semisimple Artinian ring iff the ring R is semiprime, $\text{udim}(R) < \infty$ and the ring R satisfies the a.c.c. on left annihilators (udim stands for the uniform dimension).

Proposition 1.1. *Let R be a ring and S be a non-empty subset of R .*

1. *Suppose that there exists an ideal \mathfrak{b} of R such that $(S + \mathfrak{b})/\mathfrak{b} \subseteq \mathcal{C}_{R/\mathfrak{b}}$. Then there is the least ideal, say $\mathfrak{a} = \mathfrak{a}(S)$, that satisfies this property.*
2. *Suppose that there exists an ideal \mathfrak{b} of R such that $(S + \mathfrak{b})/\mathfrak{b} \subseteq '\mathcal{C}_{R/\mathfrak{b}}$. Then there is the least ideal, say $\mathfrak{a}' = \mathfrak{a}'(S)$, that satisfies this property; and $\mathfrak{a}'(S) \subseteq \mathfrak{a}(S)$.*
3. *Suppose that there exists an ideal \mathfrak{b} of R such that $(S + \mathfrak{b})/\mathfrak{b} \subseteq \mathcal{C}'_{R/\mathfrak{b}}$. Then there is the least ideal, say $\mathfrak{a}'' = \mathfrak{a}''(S)$, that satisfies this property; and $\mathfrak{a}''(S) \subseteq \mathfrak{a}(S)$.*

For a multiplicative set S in a ring R , we fix the following notation (unless it is stated otherwise): $\mathfrak{a} = \mathfrak{a}(S)$ and $\mathfrak{a}' = \mathfrak{a}'(S)$ (see Proposition 1.1),

$${}'R := R/{}'\mathfrak{a}, \quad {}'\pi : R \rightarrow {}'R, \quad r \mapsto {}'r = r + {}'\mathfrak{a}, \quad {}'S = {}'\pi(S), \tag{5}$$

$$R' := R/\mathfrak{a}', \quad \pi' : R \rightarrow R', \quad r \mapsto r' = r + \mathfrak{a}', \quad S' = \pi'(S). \tag{6}$$

The proof of Proposition 1.1 is given in Section 2. The ideals $\mathfrak{a}(S)$, ${}'\mathfrak{a}(S)$ and $\mathfrak{a}'(S)$ are defined in an explicit way, see (10), (11) and (12), respectively. They play an important role in the proofs of many results of this paper.

Lemma 1.2. *Given $S \in \mathbb{L}_*(R)$ where $*$ $\in \{l, r, \emptyset\}$. Then $\text{ass}_R(S) \supseteq \mathfrak{a}(S)$ where $\mathfrak{a}(S)$ is the least ideal of R such that $(S + \mathfrak{a}(S))/\mathfrak{a}(S) \subseteq \mathcal{C}_{R/\mathfrak{a}(S)}$, see Proposition 1.1.(1).*

The structure of the ring $R\langle S^{-1} \rangle$ and its universal property. Let R be a ring. A multiplicative subset S of R is called a *left Ore set* if it satisfies the *left Ore condition*: for each $r \in R$ and $s \in S$,

$$Sr \cap Rs \neq \emptyset.$$

Let $\text{Ore}_l(R)$ be the set of all left Ore sets of R . For $S \in \text{Ore}_l(R)$, $\text{ass}_l(S) := \{r \in R \mid sr = 0 \text{ for some } s \in S\}$ is an ideal of the ring R .

A left Ore set S is called a *left denominator set* of the ring R if $rs = 0$ for some elements $r \in R$ and $s \in S$ implies $tr = 0$ for some element $t \in S$, i.e., $r \in \text{ass}_l(S)$. Let $\text{Den}_l(R)$ be the set of all left denominator sets of R . For $S \in \text{Den}_l(R)$, let

$$S^{-1}R = \{s^{-1}r \mid s \in S, r \in R\}$$

be the *left localization* of the ring R at S (the *left quotient ring* of R at S). Let us stress that in Ore’s method of localization one can localize *precisely* at left denominator sets. In a similar way, right Ore and right denominator sets are defined. Let $\text{Ore}_r(R)$ and $\text{Den}_r(R)$ be the set of all right Ore and right denominator sets of R , respectively. For $S \in \text{Ore}_r(R)$, the set $\text{ass}_r(S) := \{r \in R \mid rs = 0 \text{ for some } s \in S\}$ is an ideal of R . For $S \in \text{Den}_r(R)$,

$$RS^{-1} = \{rs^{-1} \mid s \in S, r \in R\}$$

is the *right localization* of the ring R at S .

Given ring homomorphisms $\nu_A : R \rightarrow A$ and $\nu_B : R \rightarrow B$. A ring homomorphism $f : A \rightarrow B$ is called an *R-homomorphism* if $\nu_B = f\nu_A$. A left and right Ore set is called an *Ore set*. Similarly, a left and right denominator set is called a *denominator set*. Let $\text{Ore}(R)$ and $\text{Den}(R)$ be the set of all Ore and denominator sets of R , respectively. For $S \in \text{Den}(R)$,

$$S^{-1}R \simeq RS^{-1}$$

(an R -isomorphism) is the *localization* of the ring R at S , and $\text{ass}(S) := \text{ass}_l(S) = \text{ass}_r(R)$.

For a ring R and $* \in \{l, r, \emptyset\}$, $\text{Den}_*(R, 0)$ be the set of $*$ denominator sets T of R such that $T \subseteq \mathcal{C}_R$, i.e., the multiplicative set T is a $*$ Ore set of R that consists of regular elements of the ring R . For a ring R , we denote by R^\times its group of units (invertible elements) of the ring R . Theorem 1.3 describes the structure and the universal property of the ring $R\langle S^{-1} \rangle$ and gives a characterization of the ideal $\text{ass}_R(S)$.

Theorem 1.3. *Let $S \in \mathbb{L}_*(R, \mathfrak{a})$ where $* \in \{l, \emptyset\}$, $\overline{R} = R/\mathfrak{a}$, $\pi : R \rightarrow \overline{R}$, $r \mapsto \overline{r} = r + \mathfrak{a}$ and $\overline{S} = \pi(S)$. Then*

1. $\overline{S} \in \text{Den}_*(\overline{R}, 0)$.
2. The ring $R\langle S^{-1} \rangle$ is R -isomorphic to the ring $\overline{S}^{-1}\overline{R}$.
3. Let \mathfrak{b} be an ideal of R and $\pi^\dagger : R \rightarrow R^\dagger = R/\mathfrak{b}$, $r \mapsto r^\dagger = r + \mathfrak{b}$. If $S^\dagger = \pi^\dagger(S) \in \text{Den}_*(R^\dagger, 0)$ then $\mathfrak{a} \subseteq \mathfrak{b}$ and the map

$$\overline{S}^{-1}\overline{R} \rightarrow S^{\dagger^{-1}}R^\dagger, \quad \overline{s}^{-1}\overline{r} \mapsto s^{\dagger^{-1}}r^\dagger$$

is a ring epimorphism with kernel $\overline{S}^{-1}(\mathfrak{b}/\mathfrak{a})$. So, the ideal \mathfrak{a} is the least ideal of the ring R such that $S + \mathfrak{a} \in \text{Den}_*(R/\mathfrak{a}, 0)$.

4. Let $f : R \rightarrow Q$ be a ring homomorphism such that $f(S) \subseteq Q^\times$ and the ring Q is generated by $f(R)$ and the set $\{f(s)^{-1} \mid s \in S\}$. Then
 - (a) $\mathfrak{a} \subseteq \ker(f)$ and the map

$$\overline{S}^{-1}\overline{R} \rightarrow Q, \quad \overline{s}^{-1}\overline{r} \mapsto f(s)^{-1}f(r)$$

is a ring epimorphism with kernel $\overline{S}^{-1}(\ker(f)/\mathfrak{a})$, and $Q = \{f(s)^{-1}f(r) \mid s \in S, r \in R\}$.

- (b) Let $\tilde{R} = R/\ker(f)$ and $\tilde{\pi} : R \rightarrow \tilde{R}$, $r \mapsto \tilde{r} = r + \ker(f)$. Then $\tilde{S} := \tilde{\pi}(S) \in \text{Den}_l(\tilde{R}, 0)$ and $\tilde{S}^{-1}\tilde{R} \simeq Q$, an \tilde{R} -isomorphism.

A similar result holds for $* = r$, i.e., for right localizable sets. Statements 3 and 4 of Theorem 1.3 are the universal property of localization of a ring at a (left or right) localizable set. In the particular case when $S \in \text{Den}_*(R)$, these are precisely the universal property of localization of a ring at a (left or right) denominator set.

In view of Theorem 1.3.(1,2), for $S \in \mathbb{L}_*(R)$ we denote by $S^{-1}R$ the ring $R\langle S^{-1} \rangle$ for $* \in \{l, \emptyset\}$ and by RS^{-1} for $* \in \{r, \emptyset\}$. In particular, for $S \in \mathbb{L}(R)$, $R\langle S^{-1} \rangle = S^{-1}R \simeq RS^{-1}$. Elements of the rings $S^{-1}R$ and RS^{-1} are denoted by $s^{-1}r$ and rs^{-1} , respectively, where $s \in S$ and $r \in R$.

Perfect localizable sets. By Lemma 1.2, $\text{ass}_R(S) \supseteq \mathfrak{a}(S)$ for all $S \in \mathbb{L}_*(R)$ where $* \in \{l, r, \emptyset\}$.

Definition. A localizable set $S \in \mathbb{L}_*(R)$ is called *perfect* if $\text{ass}_R(S) = \mathfrak{a}(S)$, i.e., the ideal $\text{ass}_R(S)$ is ‘the least possible’.

Therefore, localizations at perfect localizable sets are the most free/largest possible. Another feature of perfect localizable sets is that the ideal $\text{ass}_R(S) = \mathfrak{a}(S)$ admits an explicit description that can be computed in many examples (see the proof of Proposition 1.1 and (10)).

Let $\mathbb{L}_*(R) = \{S \in \mathbb{L}_*(R) \mid \text{ass}_R(S) = \mathfrak{a}(S)\}$ and $\text{ass } \mathbb{L}_*(R) = \{\text{ass}_R(S) \mid S \in \mathbb{L}_*(R)\}$. Clearly,

$$\mathbb{L}_*(R) = \coprod_{\mathfrak{a} \in \text{ass } \mathbb{L}_*(R)} \mathbb{L}_*(R, \mathfrak{a}) \tag{7}$$

where $\mathbb{L}_*(R, \mathfrak{a}) = \{S \in \mathbb{L}_*(R) \mid \text{ass}_R(S) = \mathfrak{a}\}$. Clearly, $\mathbb{L}_*(R, \mathfrak{a}) = \mathbb{L}_*(R) \cap \mathbb{L}_*(R, \mathfrak{a})$.

The sets $'\mathbb{L}_l(R)$, $\mathbb{L}'_r(R)$ and $'\mathbb{L}'_{l,r}(R)$. We denote by $\text{Ore}_*(R)$ (where $*$ $\in \{l, r, \emptyset\}$) the set of $*$ Ore sets of R . So, $\text{Ore}_l(R)$ is the set of all left Ore sets of R .

Definition. Let $'\mathbb{L}_l(R)$ (resp., $\mathbb{L}'_r(R)$) be the set of all multiplicative sets S of R such that $'S \in \text{Ore}_l(R)$ (resp., $S' \in \text{Ore}_r(R)$) (see (5) and (6)). Let

$$'\mathbb{L}'_{l,r}(R) := '\mathbb{L}_l(R) \cap \mathbb{L}'_r(R) \text{ and } \mathbb{L}\text{Ore}_*(R) := \mathbb{L}_*(R) \cap \text{Ore}_*(R)$$

where $*$ $\in \{l, r, \emptyset\}$. The elements of the set $\mathbb{L}\text{Ore}_*(R)$ are called the *localizable $*$ Ore sets*.

Localizable left/right Ore sets. The study of localizations at left, right, and left and right Ore sets was started in the paper [2]. In particular, [2, Theorem 4.15] states that *every Ore set is a localizable set*, i.e., $\text{Ore}(R) \subseteq \mathbb{L}(R)$. Therefore,

$$\mathbb{L}\text{Ore}(R) = \text{Ore}(R). \tag{8}$$

This fact also follows from Theorem 1.6.(2). Proposition 1.4 establishes relations between the concepts that are introduced above.

Proposition 1.4.

1. $\mathbb{L}\text{Ore}_l(R) \subseteq '\mathbb{L}_l(R) \subseteq \mathbb{L}'_l(R)$ and $\mathbb{L}\text{Ore}_l(R) = '\mathbb{L}_l(R) \cap \text{Ore}_l(R) = \mathbb{L}'_l(R) \cap \text{Ore}_l(R)$.
2. $\mathbb{L}\text{Ore}_r(R) \subseteq \mathbb{L}'_r(R) \subseteq \mathbb{L}''_r(R)$ and $\mathbb{L}\text{Ore}_r(R) = \mathbb{L}'_r(R) \cap \text{Ore}_r(R) = \mathbb{L}''_r(R) \cap \text{Ore}_r(R)$.
3. $\text{Ore}(R) \subseteq '\mathbb{L}'_{l,r}(R) \subseteq \mathbb{L}^p(R)$.

Criterion for a left Ore set to be a left localizable set. For a ring R and its ideal \mathfrak{a} , let

$$\begin{aligned} ' \text{Den}_l(R, \mathfrak{a}) &:= \{S \in \text{Den}_l(R) \mid \text{ass}_l(S) = \mathfrak{a}, S \subseteq ' \mathcal{C}_R\}, \\ \text{Den}'_r(R, \mathfrak{a}) &:= \{S \in \text{Den}_r(R) \mid \text{ass}_r(S) = \mathfrak{a}, S \subseteq \mathcal{C}'_R\}. \end{aligned}$$

Theorem 1.5.(1) is a criterion for a left Ore set to be a left localizable set and Theorem 1.5.(2) describes the structure of the localization of a ring at a localizable left Ore set.

Theorem 1.5. *Let R be a ring, $S \in \text{Ore}_l(R)$, and $\mathfrak{a} = \text{ass}_R(S)$. Then*

1. $S \in \mathbb{L}_l(R)$ iff $'\mathfrak{a} \neq R$ where the ideal $'\mathfrak{a} = '\mathfrak{a}(S)$ of R is as in Proposition 1.1.(2) and (11).
2. Suppose that $'\mathfrak{a} \neq R$. Let $'\pi : R \rightarrow 'R := R/'\mathfrak{a}$, $r \mapsto 'r = r + '\mathfrak{a}$ and $'S = '\pi(S)$. Then
 - (a) $'S \in '\text{Den}_l('R)$.
 - (b) $\mathfrak{a} = '\pi^{-1}(\text{ass}_l('S))$.
 - (c) $S^{-1}R \simeq 'S^{-1}'R$, an R -isomorphism.

Theorem 2.11 is a criterion for a right Ore set to be a localizable set.

Localization at an Ore set. Theorem 1.6.(1) is the reason why every Ore set is localizable. For an Ore set S of a ring R , Theorem 1.6.(1,2) shows that $\mathfrak{a}(S) = \text{ass}_R(S)$ and gives an explicit description of this ideal.

Theorem 1.6. *Let R be a ring and $S \in \text{Ore}(R)$.*

1. $\mathfrak{a} := \{r \in R \mid srt = 0 \text{ for some elements } s, t \in S\}$ is an ideal of R such that $\mathfrak{a} \neq R$.
2. Let $\pi : R \rightarrow \overline{R} := R/\mathfrak{a}$, $r \mapsto \overline{r} = r + \mathfrak{a}$. Then $\overline{S} := \pi(S) \in \text{Den}(\overline{R}, 0)$, $\mathfrak{a} = \mathfrak{a}(S) = \text{ass}_R(S)$, $S \in \mathbb{L}(R, \mathfrak{a})$, and $S^{-1}R \simeq \overline{S}^{-1}\overline{R}$, an R -isomorphism. In particular, every Ore set is localizable.
3. Let \mathfrak{b} be an ideal of R and $\pi^\dagger : R \rightarrow R^\dagger := R/\mathfrak{b}$, $r \mapsto r^\dagger = r + \mathfrak{b}$. If $S^\dagger := \pi^\dagger(S) \in \text{Den}(R^\dagger, 0)$ then $\mathfrak{a} \subseteq \mathfrak{b}$ and the map

$$\overline{S}^{-1}\overline{R} \rightarrow S^{\dagger^{-1}}R^\dagger, \quad \overline{s}^{-1}\overline{r} \mapsto s^{\dagger^{-1}}r^\dagger$$

is a ring epimorphism.

4. Let $f : R \rightarrow Q$ be a ring homomorphism such that $f(S) \subseteq Q^\times$ and the ring Q is generated by $f(R)$ and $\{f(s)^{-1} \mid s \in S\}$. Then
 - (a) $\mathfrak{a} \subseteq \ker(f)$ and the map

$$\overline{S}^{-1}\overline{R} \rightarrow Q, \quad \overline{s}^{-1}\overline{r} \mapsto f(s)^{-1}f(r)$$

is a ring epimorphism with kernel $\overline{S}^{-1}(\ker(f)/\mathfrak{a})$.

- (b) Let $\tilde{R} = R/\ker(f)$ and $\tilde{\pi} : R \rightarrow \tilde{R}$, $r \mapsto r + \ker(f)$. Then $\tilde{S} := \tilde{\pi}(S) \in \text{Den}(\tilde{R}, 0)$ and $\tilde{S}^{-1}\tilde{R} \simeq Q$, an \tilde{R} -isomorphism.

Theorem 1.6.(1,2) also states that every Ore set S of R is localizable and the localization $S^{-1}R$ of the ring R at the Ore set S is R -isomorphic to the localization $\overline{S}^{-1}\overline{R}$ of the ring \overline{R} at the denominator set \overline{S} of \overline{R} .

Corollary 1.7 provides a view from another angle at the localization of a ring at an Ore set.

Corollary 1.7. *Let R be a ring, $S \in \text{Ore}(R)$ and $\mathfrak{a} = \text{ass}_R(S)$. We keep the notation of Theorem 1.6. Let $\mathfrak{a}_l := \text{ass}_l(S)$ and $\pi_l : R \rightarrow R_l := R/\mathfrak{a}_l, r \mapsto r + \mathfrak{a}_l$; $\mathfrak{a}_r := \text{ass}_r(S)$ and $\pi_r : R \rightarrow R_r := R/\mathfrak{a}_r, r \mapsto r + \mathfrak{a}_r$. Then*

1. $\mathfrak{a}_l + \mathfrak{a}_r \subseteq \mathfrak{a}$.
2. $S_l := \pi_l(S) \in \text{Den}'_r(R_l, \mathfrak{a}/\mathfrak{a}_l)$ and $R_l S_l^{-1} \simeq S^{-1}R \simeq R S^{-1}$, R -isomorphisms.
3. $S_r := \pi_r(S) \in {}'\text{Den}_l(R_r, \mathfrak{a}/\mathfrak{a}_r)$ and $S_r^{-1}R_r \simeq S^{-1}R \simeq R S^{-1} \simeq R_l S_l^{-1}$, R -isomorphisms.
4. ${}'\mathfrak{a}(S) = \mathfrak{a}_r$ and $\mathfrak{a}'(S) = \mathfrak{a}_l$.

Proposition 1.8 explains the origin of the construction of localizable sets. For every Ore set it gives an explicit construction of a *localizable set* with the same localization. For a ring R and its ideal \mathfrak{a} , let $\text{Ore}(R, \mathfrak{a}) := \{S \in \text{Ore}(R) \mid \text{ass}_R(S) = \mathfrak{a}\}$.

Proposition 1.8. *Let R be a ring, $S \in \text{Ore}(R, \mathfrak{a})$, $\pi : R \rightarrow \overline{R} := R/\mathfrak{a}, R \mapsto \bar{r} = r + \mathfrak{a}$ and $\tilde{S} = S + \mathfrak{a}$. Then $S \subseteq \tilde{S}$, $\tilde{S} \in \mathbb{L}(R, \mathfrak{a})$ and $\tilde{S}^{-1}R \simeq S^{-1}R$, an R -isomorphism.*

The set $\max \mathbb{L}_*(R)$ **maximal elements in $\mathbb{L}_*(R)$ where $* \in \{l, r, \emptyset\}$** . For a ring R , the set $\max.\text{Den}_l(R)$ of maximal left denominator sets (w.r.t. \subseteq) is a *non-empty* set, [3, Lemma 3.7.(2)]. Let $\max \mathbb{L}_*(R)$ be the set of maximal elements (w.r.t. \subseteq) of the set $\mathbb{L}_*(R)$.

Theorem 1.9. *Let R be a ring. Then $\max \mathbb{L}_*(R) \neq \emptyset$.*

The key idea of the proof of Theorem 1.9 is to use Lemma 1.10 and Zorn’s Lemma.

Lemma 1.10. *Let R be a ring, $S \in \mathbb{L}_*(R, \mathfrak{a})$ and $T \in \mathbb{L}_*(R, \mathfrak{b})$ such that $S \subseteq T$ where $* \in \{l, r, \emptyset\}$. Then $\mathfrak{a} \subseteq \mathfrak{b}$ and for $* \in \{l, \emptyset\}$ the map $S^{-1}R \rightarrow T^{-1}R, s^{-1}r \mapsto t^{-1}r$ is an R -homomorphism with kernel $S^{-1}(\mathfrak{b}/\mathfrak{a}) = \overline{S}^{-1}(\mathfrak{b}/\mathfrak{a})$ where $\overline{S} = \{s + \mathfrak{a} \mid s \in S\}$. A similar result holds for $* = r$.*

Classification of maximal Ore sets of a semiprime Goldie ring. It was proved that the set $\max.\text{Den}_l(R)$ is a finite set if the classical left quotient ring $Q_{l,cl}(R) := \mathcal{C}_R^{-1}R$ of R is a semisimple Artinian ring, [4], or a left Artinian ring, [5], or a left Noetherian ring, [6]. In each of the three cases an explicit description of the set $\max.\text{Den}_l(R)$ is given. For a ring R , let $\min(R)$ be the set of its minimal prime ideals. For a ring R , the rings

$$Q_{l,cl}(R) := \mathcal{C}_R^{-1}R \text{ and } Q_{r,cl}(R) := RC_R^{-1}$$

are called the *classical left and right quotient rings* provided they exist, respectively. If both rings exist then they are isomorphic and the ring

$$Q_{cl}(R) := Q_{l,cl}(R) \simeq Q_{r,cl}(R)$$

is called the *classical quotient ring* of R . Ideals of a ring are called *incomparable* if none of them is contained in the other.

The next theorem is an explicit description of maximal Ore sets of a semiprime Goldie ring.

Theorem 1.11. *Let R be a semiprime Goldie ring and $\mathcal{N}_* := \{S \in \max \mathbb{L}_*(R) \mid \mathcal{C}_R \subseteq S\}$ where $*$ $\in \{l, r, \emptyset\}$. Then*

1. $\max \text{Ore}(R) = \max \text{Den}(R) = \{\mathcal{C}(\mathfrak{p}) \mid \mathfrak{p} \in \min(R)\} = \mathcal{N}_*$ for all $*$ $\in \{l, r, \emptyset\}$ where $\mathcal{C}(\mathfrak{p}) := \{c \in R \mid c + \mathfrak{p} \in \mathcal{C}_{R/\mathfrak{p}}\}$. So, every maximal Ore set of R is a maximal denominator set, and vice versa.
2. For all $S \in \max \text{Ore}(R)$ the ring $S^{-1}R$ is a simple Artinian ring.
3. $Q_{cl}(R) \simeq \prod_{S \in \max \text{Ore}(R)} S^{-1}R$.
4. $\max \text{ass Ore}(R) = \text{ass max Ore}(R) = \min(R)$. In particular, the ideals in the set $\text{ass max Ore}(R)$ are incomparable.

Classification of maximal left localizable sets of a semiprime left Goldie ring that contain the set of regular elements of the ring. Theorem 1.12 is such a classification.

Theorem 1.12. *Let R be a semiprime left Goldie ring and $\mathcal{N} := \{S \in \max \mathbb{L}_l(R) \mid \mathcal{C}_R \subseteq S\}$. Then $\mathcal{N} = \max \text{Den}_l(R) = \{\mathcal{C}(\mathfrak{p}) \mid \mathfrak{p} \in \min(R)\}$.*

So, every maximal left localizable set of a semiprime left Goldie ring that contains the set of regular elements of the ring is a maximal left denominator set, and vice versa. The proof of Theorem 1.12 is based on Theorem 1.13.

Theorem 1.13. *Let R be a ring and $S_1, \dots, S_n \in \mathbb{L}_*(R)$ where $*$ $\in \{l, r, \emptyset\}$, $\mathfrak{p}_i = \text{ass}_R(S_i)$, $R_i := R/\mathfrak{p}_i$ and Q_i be the localization of R at S_i . Suppose that the rings Q_i are simple Artinian rings, $\bigcap_{i=1}^n \mathfrak{p}_i = 0$ and $\bigcap_{j \neq i} \mathfrak{p}_j \neq 0$ for $i = 1, \dots, n$. Then*

1. The rings R_i are semiprime $*$ Goldie rings.
2. $\min(R) = \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$.
3. $Q_i \simeq Q_{*,cl}(R_i)$ for $i = 1, \dots, n$ (an R -isomorphism).
4. $S_i \subseteq \mathcal{C}(\mathfrak{p}_i)$ for $i = 1, \dots, n$.
5. $Q_{*,cl}(R) \simeq \prod_{i=1}^n Q_i$.
6. For all $i = 1, \dots, n$, $\mathcal{C}(\mathfrak{p}_i) \in \max \mathbb{L}_*(R)$.

In Section 7, examples are considered. In [7], we continue to develop the most general theory of one-sided fractions. The aim of [7] is to introduce 10 types of saturations of a set in a ring and using them to study localizations of a ring at localizable sets, their groups of units and various maximal localizable sets satisfying some natural conditions. The results are obtained for denominator sets (the classical situation), Ore sets and localizable sets.

2. Localizable sets and the localization of a ring at a localizable set

In this section, (left, right) localizable sets of a ring and the construction of localization of a ring at them are introduced and studied. A proof of Theorem 1.3 is given. A criterion for a left (resp., right) Ore set to be a left (resp., right) localizable set is presented, Theorem 1.5.(1) (resp., Theorem 2.11.(1)). A description of the ideal $\text{ass}_R(S)$ of R is given, Proposition 2.12. Proofs of Proposition 1.1, Proposition 1.4, Theorem 1.9, Lemma 1.10, Theorem 1.12 and Theorem 1.13 are given.

The following notation is fixed: R is a ring, S is a multiplicative set of R , $\text{ass}_l(S) := \{r \in R \mid sr = 0 \text{ for some } s \in S\}$ and $\text{ass}_r(S) := \{r \in R \mid rs = 0 \text{ for some } s \in S\}$. We use standard terminology on localizations of a ring at denominator sets, see [8,10,11].

Proof of Proposition 1.1. We keep the notation of Proposition 1.1. Let Γ be the set of ordinals. The ideal \mathfrak{a} (resp., \mathfrak{a}') is the union

$$\mathfrak{a} = \bigcup_{\lambda \in \Gamma} \mathfrak{a}_\lambda \quad (\text{resp.}, \mathfrak{a}' = \bigcup_{\lambda \in \Gamma} \mathfrak{a}'_\lambda, \mathfrak{a}'' = \bigcup_{\lambda \in \Gamma} \mathfrak{a}''_\lambda) \tag{9}$$

of ascending chain of ideals $\{\mathfrak{a}_\lambda\}_{\lambda \in \Gamma}$ (resp., $\{\mathfrak{a}'_\lambda\}_{\lambda \in \Gamma}$, $\{\mathfrak{a}''_\lambda\}_{\lambda \in \Gamma}$), where $\lambda \leq \mu$ in Γ implies $\mathfrak{a}_\lambda \subseteq \mathfrak{a}_\mu$ (resp., $\mathfrak{a}'_\lambda \subseteq \mathfrak{a}'_\mu$, $\mathfrak{a}''_\lambda \subseteq \mathfrak{a}''_\mu$). The ideals \mathfrak{a}_λ (resp., \mathfrak{a}'_λ , \mathfrak{a}''_λ) are defined inductively as follows: the ideal $\mathfrak{a}_0 = \mathfrak{a}(S, R)$ (resp., \mathfrak{a}'_0 , \mathfrak{a}''_0) is generated by the set $\{r \in R \mid sr = 0 \text{ or } rt = 0 \text{ for some elements } s, t \in S\}$ (resp., $\{r \in R \mid rt = 0 \text{ for some element } t \in S\}$, $\{r \in R \mid sr = 0 \text{ for some element } s \in S\}$), and for $\lambda \in \Gamma$ such that $\lambda > 0$ (where below $(\{\dots\})$ means the ideal of R generated by the set $\{\dots\}$),

$$\mathfrak{a}_\lambda = \begin{cases} \bigcup_{\mu < \lambda \in \Gamma} \mathfrak{a}_\mu & \text{if } \lambda \text{ is a limit ordinal,} \\ \left(\{r \in R \mid sr \in \mathfrak{a}_{\lambda-1} \text{ or } rt \in \mathfrak{a}_{\lambda-1} \text{ for some } s, t \in S\} \right) & \text{if } \lambda \text{ is not a limit ordinal,} \end{cases} \tag{10}$$

(resp.,

$$\mathfrak{a}'_\lambda = \begin{cases} \bigcup_{\mu < \lambda \in \Gamma} \mathfrak{a}'_\mu & \text{if } \lambda \text{ is a limit ordinal,} \\ \left(\{r \in R \mid rt \in \mathfrak{a}'_{\lambda-1} \text{ for some } t \in S\} \right) & \text{if } \lambda \text{ is not a limit ordinal,} \end{cases} \tag{11}$$

$$\mathfrak{a}''_\lambda = \begin{cases} \bigcup_{\mu < \lambda \in \Gamma} \mathfrak{a}''_\mu & \text{if } \lambda \text{ is a limit ordinal,} \\ \left(\{r \in R \mid sr \in \mathfrak{a}''_{\lambda-1} \text{ for some } s \in S\} \right) & \text{if } \lambda \text{ is not a limit ordinal,} \end{cases} \tag{12}$$

and the proposition follows (since $\alpha_\lambda \subseteq \alpha_\lambda$ and $\alpha'_\lambda \subseteq \alpha_\lambda$ for all $\lambda \in \Gamma$). \square

The first ordinal such that the ascending chain of ideals in (9) stabilizes is called the Γ -length of the ideal denoted $l_\Gamma(\mathfrak{a})$ (resp., $l_\Gamma(\mathfrak{a}')$, $l_\Gamma(\mathfrak{a}')$).

Lemma 2.1. *Let R be a ring, S be a multiplicative set, $\mathfrak{a} = \text{ass}_R(S)$, \mathfrak{b} be an ideal of the ring R such that $\mathfrak{b} \subseteq \mathfrak{a}$, $\tilde{R} = R/\mathfrak{b}$ and $\tilde{S} = (S + \mathfrak{b})/\mathfrak{b} = \{s \in \mathfrak{b} \mid s \in S\}$. Then*

1. \tilde{S} is a multiplicative set of the ring \tilde{R} , $\tilde{R}\langle\tilde{S}^{-1}\rangle \simeq R\langle S^{-1}\rangle$ and $\text{ass}_{\tilde{R}}(\tilde{S}) = \mathfrak{a}/\mathfrak{b}$.
2. Let $\bar{R} = R/\mathfrak{a}$ and $\bar{S} = (S + \mathfrak{a})/\mathfrak{a}$. Then $R\langle S^{-1}\rangle \simeq \bar{R}\langle\bar{S}^{-1}\rangle$, $\bar{S} \subseteq \mathcal{C}_{\bar{R}}$ and $\text{ass}_{\bar{R}}(\bar{S}) = \mathfrak{a}/\mathfrak{a} = 0$.

Proof. Straightforward. \square

Proof of Lemma 1.2. Let $\mathfrak{a} = \text{ass}_R(S)$ and $\bar{R} = R/\mathfrak{a}$. Then $\bar{S} = (S + \mathfrak{a})/\mathfrak{a} \in \mathcal{C}_{\bar{R}}$. Hence, $\mathfrak{a} \supseteq \mathfrak{a}(S)$, by the minimality of the ideal $\mathfrak{a}(S)$, see Proposition 1.1.(1). \square

For a ring R , its ideal \mathfrak{a} , and $*$ $\in \{l, r, \emptyset\}$, let $\text{Den}_*(R, \mathfrak{a})$ be the set of $*$ denominator sets S of R such that $\text{ass}_*(S) = \mathfrak{a}$.

Lemma 2.2 and Lemma 2.3 show that denominator sets are localizable sets, the localization of a ring at a denominator set is the same as the localization of a ring at the denominator set treated as a localizable set, and $\text{ass}_*(S) = \mathfrak{a}(S)$ for all $S \in \text{Den}_*(R, \mathfrak{a})$. So, the localization at a localizable set is a generalization of the localization at a denominator set.

Lemma 2.2. *Given $S \in \text{Den}_*(R, \mathfrak{a})$ where $*$ $\in \{l, r, \emptyset\}$. Then $\mathfrak{a} = \mathfrak{a}(S)$ where $\mathfrak{a}(S)$ is the least ideal of R such that $(S + \mathfrak{a}(S))/\mathfrak{a}(S) \in \mathcal{C}_{R/\mathfrak{a}(S)}$, see Proposition 1.1.(1).*

Proof. By the very definition of the ideal \mathfrak{a} , $\mathfrak{a} \subseteq \mathfrak{a}(S)$. Since $(S + \mathfrak{a})/\mathfrak{a} \in \mathcal{C}_{R/\mathfrak{a}}$, we have the inverse inclusion $\mathfrak{a} \supseteq \mathfrak{a}(S)$, by the minimality of the ideal $\mathfrak{a}(S)$. Therefore $\mathfrak{a} = \mathfrak{a}(S)$. \square

Lemma 2.3. $\text{Den}_*(R) \subseteq \mathbb{L}_*(R)$ where $*$ $\in \{l, r, \emptyset\}$. For all ideals \mathfrak{a} of R , $\text{Den}_*(R, \mathfrak{a}) \subseteq \mathbb{L}_*(R, \mathfrak{a})$ and for all $S \in \text{Den}_l(R, \mathfrak{a})$, $S^{-1}R \simeq R\langle S^{-1}\rangle$ and $\mathfrak{a} = \mathfrak{a}(S) = \text{ass}_R(S)$.

Proof. Given $S \in \text{Den}_l(R, \mathfrak{a})$. By Lemma 1.2 and Lemma 2.2, $\mathfrak{a} = \mathfrak{a}(S) \subseteq \text{ass}_R(R)$. Let $\bar{R} = R/\mathfrak{a}$ and $\bar{S} = (S + \mathfrak{a})/\mathfrak{a}$. Then

$$\bar{S} \in \text{Den}_l(R, 0) \text{ and } S^{-1}R \simeq \bar{S}^{-1}\bar{R}.$$

The ring $\bar{S}^{-1}\bar{R}$ is generated by the ring \bar{R} and the set $\{\bar{s}^{-1} \mid \bar{s} \in \bar{S}\}$. Therefore, there is natural ring epimorphism $R\langle S^{-1}\rangle \rightarrow \bar{S}^{-1}\bar{R}$ which is an R -homomorphism. Hence, $\text{ass}_R(S) \subseteq \mathfrak{a}$, and so $\mathfrak{a} = \mathfrak{a}(S) = \text{ass}_R(S)$ and $R\langle S^{-1}\rangle \simeq \bar{S}^{-1}\bar{R}$. \square

Proof of Theorem 1.3. To prove the theorem it suffices to consider the case $* = l$.

1 and 2. Let $\mathcal{R} = R\langle S^{-1} \rangle$. By Lemma 2.1.(2),

$$R\langle S^{-1} \rangle \simeq \overline{R}\langle \overline{S}^{-1} \rangle, \quad \overline{S} \subseteq \mathcal{C}_{\overline{R}} \text{ and } \text{ass}_{\overline{R}}(\overline{S}) = 0.$$

Hence, $\overline{R} \subseteq \mathcal{R}$. Then, for all elements $\overline{s} \in \overline{S}$ and $\overline{r} \in \overline{R}$,

$$\overline{r}\overline{s}^{-1} = \overline{s}_1^{-1}\overline{r}_1^{-1}$$

for some elements $\overline{s}_1 \in \overline{S}$ and $\overline{r}_1 \in \overline{R}$. Then $\overline{s}_1\overline{r} = \overline{r}_1\overline{s}$ in \overline{R} . Hence, $\overline{S} \in \text{Den}_l(\overline{R}, 0)$, and so $\overline{R}\langle \overline{S}^{-1} \rangle \simeq \overline{S}^{-1}\overline{R}$, by Lemma 2.3.

3. There is a natural R -epimorphism $R\langle S^{-1} \rangle \rightarrow S^{\dagger -1}R^{\dagger}$, and statement 3 follows from statement 2.

4. The ring homomorphism $r : R \rightarrow Q$ induces the ring epimorphism $f : R\langle S^{-1} \rangle \rightarrow Q$ since the ring Q is generated by $f(R)$ and the set $\{f(s)^{-1} \mid s \in S\}$. Since $R\langle S^{-1} \rangle \simeq \overline{S}^{-1}\overline{R}$ (statement 2), the statement (a) follows. In particular,

$$Q = \{f(s)^{-1}f(r) \mid s \in S, r \in R\}.$$

Furthermore, $Q = \{f(\overline{s})^{-1}f(\overline{r}) \mid s \in S, r \in R\}$ since $\mathfrak{a} \subseteq \ker(f)$. Hence, $Q = \{f(\tilde{s})^{-1}f(\tilde{r}) \mid s \in S, r \in R\}$. Therefore, $\tilde{S} \in \text{Den}_l(\tilde{R}, 0)$ and $\tilde{S}^{-1}\tilde{R} \simeq Q$, via f . \square

Corollary 2.4. *If $S \in \mathbb{L}(R, \mathfrak{a})$ then $S \in \mathbb{L}_l(R, \mathfrak{a})$, $S \in \mathbb{L}_r(R, \mathfrak{a})$ and the localizations of R at S as a localizable set, a left localizable set and a right localizable set are R -isomorphic.*

Proof. The corollary follows from Theorem 1.3.(1,2). \square

Proof of Lemma 1.10. Recall that $\mathfrak{a} = \text{ass}_R(S)$ and $\mathfrak{b} = \text{ass}_R(T)$. Let Q be a subring of $T^{-1}R$ which is generated by the images of the ring R and the set $\{s^{-1} \mid s \in S\}$ in $T^{-1}R$ (recall that $S \subseteq T$). Applying Theorem 1.3.(4a) to the ring homomorphism $R \rightarrow Q \subseteq T^{-1}R, r \mapsto \frac{r}{1}$ we obtain the ring R -homomorphism

$$S^{-1}R \rightarrow T^{-1}R, \quad s^{-1}r \mapsto s^{-1}r.$$

Since $S^{-1}R = \overline{S}^{-1}\overline{R}$ and $T^{-1}R = \overline{T}^{-1}(R/\mathfrak{b})$ where $\overline{T} = \{t + \mathfrak{b} \mid t \in T\}$, the kernel of the R -homomorphism is $\overline{S}^{-1}(\mathfrak{b}/\mathfrak{a})$. \square

The maximal elements in $\mathbb{L}_*(R)$ where $* \in \{l, r, \emptyset\}$.

Lemma 2.5. *Let R be a ring, $* \in \{l, r, \emptyset\}$, and a set $\{S_i\}_{i \in I} \subseteq \mathbb{L}_*(R)$ be such that for any two elements S_i and S_j there is an element S_k such that $S_i \cup S_j \subseteq S_k$. Then $S := \bigcup_{i \in I} S_i \in \mathbb{L}_*(R)$ and the ring $S^{-1}R = \text{injlim } S_i^{-1}R$ is the injective limit of R -homomorphism of rings $\{S_i^{-1}R\}_{i \in I}$ given by the R -homomorphisms $S_i^{-1}R \rightarrow S_j^{-1}R$ in Lemma 1.10 in case $S_i \subseteq S_j$.*

Proof. Straightforward (use Lemma 1.10). \square

Proof of Theorem 1.9. The Theorem 1.9 follows from Zorn’s Lemma and Lemma 2.5. \square

Classification of maximal left localizable sets of a semiprime left Goldie ring that contain the set of regular elements.

Proof of Theorem 1.13. 1. By the assumption $\bigcap_{i=1}^n \mathfrak{p}_i = 0$. So, we have ring monomorphisms

$$R \rightarrow \mathcal{R} := \prod_{i=1}^n R_i \rightarrow Q := \prod_{i=1}^n Q_i, \quad r \mapsto (r + \mathfrak{p}_1, \dots, r + \mathfrak{p}_n), \quad (r_1, \dots, r_n) \mapsto \left(\frac{r_1}{1}, \dots, \frac{r_n}{1} \right).$$

We identify the rings R and \mathcal{R} with their images in the ring Q .

(i) R_i is a prime $*$ Goldie ring with $Q_{*,cl}(R_i) \simeq Q_i$, an R -isomorphism: By Theorem 1.3.(1,2),

$$\overline{S}_i := (S_i + \mathfrak{p}_i) / \mathfrak{p}_i \in \text{Den}_*(R_i, 0) \text{ and } \overline{S}_i^{-1} R_i \simeq Q_i,$$

an R -isomorphism. Since the ring Q_i is a simple Artinian ring, $\overline{S}_i \subseteq \mathcal{C}_{R_i} \subseteq Q_i^\times$. Hence, $\mathcal{C}_{R_i} \in \text{Den}_*(R, 0)$ and $Q_i \simeq Q_{*,cl}(R_i)$. By Goldie’s Theorem, R_i is a prime $*$ Goldie ring.

(ii) $\mathfrak{p}_i \in \text{Spec}(R)$ for $i = 1, \dots, n$: By the statement (i), the ring R_i is prime, and so $\mathfrak{p}_i \in \text{Spec}(R)$.

(iii) *The ring R is a semiprime $*$ Goldie ring:* The ring R is a semiprime ring since $\bigcap_{i=1}^n \mathfrak{p}_i = 0$ and the ideals \mathfrak{p}_i are prime.

Let $\text{udim}_{*,R}$ denote the $*$ uniform dimension of the $*$ R -module ($\text{udim}_{l,R}$ is the left uniform dimension, $\text{udim}_{r,R}$ is the right uniform dimension and udim_R stands for $\text{udim}_{l,R}$ and $\text{udim}_{r,R}$). The ring $R_i = R/\mathfrak{p}_i$ is a prime $*$ Goldie ring. Hence, $\text{udim}_{*,R}(R_i) = \text{udim}_{*,R}(R_i) < \infty$. Since $R \subseteq \prod_{i=1}^n R_i$,

$$\text{udim}_{*,R}(R) \leq \text{udim}_{*,R} \left(\prod_{i=1}^n R_i \right) = \sum_{i=1}^n \text{udim}_{*,R}(R_i) = \sum_{i=1}^n \text{udim}_{*,R_i}(R_i) < \infty.$$

Let X be a non-empty subset of R and $*.\text{ann}_R(X)$ be its $*$ annihilator ($l.\text{ann}_R(X) = \{r \in R \mid rX = 0\}$ is the left annihilator of X in R , etc). Since $R \subseteq \mathcal{R}$, $*.\text{ann}_R(X) = R \cap *.\text{ann}_{\mathcal{R}}(X)$. By the statement (i), the ring $\mathcal{R} = \prod_{i=1}^n R_i$ satisfies the a.c.c. on $*$ annihilators, hence so does the ring R . The proof of the statement (iii) is complete.

2. (i) $\min(R) \subseteq \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$: Given $\mathfrak{p} \in \min(R)$. Then $\bigcap_{i=1}^n \mathfrak{p}_i = \{0\} \subseteq \mathfrak{p}$, hence $\mathfrak{p}_i \subseteq \mathfrak{p}$ for some i , and so $\mathfrak{p}_i = \mathfrak{p}$, by the minimality of \mathfrak{p} .

(ii) $\min(R) = \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$: By the statement (i), the ring R is semiprime, i.e., $\bigcap_{\mathfrak{p} \in \min(R)} \mathfrak{p} = 0$. Since $\min(R) \subseteq \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$ and $\bigcap_{j \neq i} \mathfrak{p}_j \neq 0$ for all $i = 1, \dots, n$, we must have

$$\min(R) = \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}.$$

- 3. Statement 3 has already been proven, see the statement (i) in the proof of statement 1.
- 4. See the proof of the statement (i) in the proof of statement 1.
- 5. The ring R is a semiprime $*$ Goldie ring. By [4, Theorem 4.1],

$$Q_{*,cl}(R) \simeq \prod_{\mathfrak{p} \in \min(R)} Q_{*,cl}(R/\mathfrak{p}).$$

By statement 2, $\min(R) = \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$ and $Q_i \simeq Q_{*,cl}(R_i)$ for $i = 1, \dots, n$ (statement 3). Now, statement 5 follows.

- 6. By [4, Theorem 4.1],

$$\mathcal{C}(\mathfrak{p}_i) \in \max \text{Den}_*(R), \quad \text{ass}_* \mathcal{C}(\mathfrak{p}_i) = \mathfrak{p}_i \quad \text{and} \quad \mathcal{C}(\mathfrak{p}_i)^{-1}R \simeq Q_i,$$

a simple Artinian ring. Suppose that $\mathcal{C}(\mathfrak{p}_i) \subseteq T$ for some $T \in \mathbb{L}_*(R)$. By Lemma 1.10,

$$\mathfrak{p}_i = \text{ass}_R(\mathcal{C}(\mathfrak{p}_i)) \subseteq \mathfrak{a} := \text{ass}_R(T)$$

and there is an R -homomorphism $Q_i = \mathcal{C}(\mathfrak{p}_i)^{-1}R \rightarrow R\langle T^{-1} \rangle$. Since Q_i is a simple Artinian ring, $\mathfrak{p}_i = \mathfrak{a}$. Then, by Theorem 1.3.(1),

$$\overline{T} = (T + \mathfrak{p}_i)/\mathfrak{p}_i \in \text{Den}_*(R_i, 0).$$

Hence, $\overline{T} \subseteq \mathcal{C}_{R_i}$, and so $T \subseteq \mathcal{C}(\mathfrak{p}_i)$, and statement 6 follows. \square

Proof of Theorem 1.12. Let $\mathcal{M} = \max \text{Den}_l(R)$. By [4, Theorem 4.1],

$$\mathcal{M} = \{\mathcal{C}(\mathfrak{p}) \mid \mathfrak{p} \in \min(R)\}$$

and the set \mathcal{M} satisfies the conditions of Theorem 1.13. By Theorem 1.13.(2,6),

$$\mathcal{M} \subseteq \max \mathbb{L}_*(R).$$

Since $\mathcal{C}_R \subseteq \mathcal{C}(\mathfrak{p})$ for all $\mathfrak{p} \in \min(R)$, $\mathcal{M} \subseteq \mathcal{N}$. To finish the proof it remains to show that $\mathcal{N} \subseteq \mathcal{M}$. Given $S \in \mathcal{N}$. Let $\mathfrak{a} = \text{ass}_R(S)$.

(i) $\mathfrak{a} = \bigcap_{\mathfrak{p} \in \min(R), \mathfrak{a} \subseteq \mathfrak{p}} \mathfrak{p}$: Since $S \in \mathcal{N}$, $\mathcal{C}_R \subseteq S$, and so there is an R -homomorphisms $f : Q := \mathcal{C}_R^{-1}R \rightarrow S^{-1}R$. Therefore, $\mathfrak{a} = R \cap \ker(f)$ and $\ker(f) = \mathcal{C}_R^{-1}\mathfrak{a}$. By [4, Theorem 4.1],

$$\mathcal{C}_R^{-1}R \simeq \prod_{\mathfrak{p} \in \min(R)} \mathcal{C}(\mathfrak{p})^{-1}R$$

is a finite direct product of simple Artinian rings $\mathcal{C}(\mathfrak{p})^{-1}R$ with $\text{ass}_R(\mathcal{C}(\mathfrak{p})) = \mathfrak{p}$, and the statement (i) follows.

(ii) $S \subseteq \mathcal{C}(\mathfrak{p})$ for some (unique) $\mathfrak{p} \in \min(R)$ such that $\mathfrak{a} \subseteq \mathfrak{p}$: By Theorem 1.3.(1), $\overline{S} := (S + \mathfrak{a})/\mathfrak{a} \in \text{Den}_l(\overline{R}, 0)$ where $\overline{R} = R/\mathfrak{a}$ is a semiprime ring with

$$\min(\overline{R}) = \{\mathfrak{p}/\mathfrak{a} \mid \mathfrak{p} \in \min(R), \mathfrak{a} \subseteq \mathfrak{p}\},$$

by the statement (i). Since $\mathcal{C}_R \subseteq \mathcal{C}(\mathfrak{p})$ for all $\mathfrak{p} \in \min(R)$, we have that $\overline{\mathcal{C}}_R := (\mathcal{C}_R + \mathfrak{a})/\mathfrak{a} \subseteq \mathcal{C}_{\overline{R}}$ since

$$\overline{R} = R/\mathfrak{a} = R / \bigcap_{\mathfrak{p} \in D} \mathfrak{p} \subseteq \prod_{\mathfrak{p} \in D} R/\mathfrak{p}$$

where $D := \{\mathfrak{p} \in \min(R) \mid \mathfrak{a} \subseteq \mathfrak{p}\}$, see the statement (i). Since $\mathcal{C}_R \in \text{Ore}_l(R)$, $\overline{\mathcal{C}}_R \in \text{Den}_l(\overline{R}, 0)$ and

$$\overline{\mathcal{C}}_R^{-1} \overline{R} \simeq \mathcal{C}_R^{-1}(R/\mathfrak{a}) \simeq \mathcal{C}_R^{-1}R/\mathcal{C}_R^{-1}\mathfrak{a} \simeq \prod_{\mathfrak{p} \in D} \mathcal{C}(\mathfrak{p})^{-1}R.$$

Therefore, \overline{R} is a semiprime left Goldie ring with $\min(\overline{R}) = \{\mathfrak{p}/\mathfrak{a} \mid \mathfrak{p} \in D\}$.

Since $\overline{S} \in \text{Den}_l(\overline{R}, 0)$, the left denominator set \overline{S} is contained in a maximal denominator set \overline{R} , i.e., $\overline{S} \subseteq \mathcal{C}_{\overline{R}}(\mathfrak{p}/\mathfrak{a})$ for some prime ideal $\mathfrak{p} \in D$. Then

$$S \subseteq \mathcal{C}_R(\mathfrak{p})$$

since $\overline{R}/(\mathfrak{p}/\mathfrak{a}) \simeq R/\mathfrak{p}$. \square

The set $T_{*,\mathfrak{a}}(R)$ and the ring $Q_{*,\mathfrak{a}}(R)$. Lemma 2.6.(4) describes the largest element $T_{*,\mathfrak{a}}(R)$ in the set $\mathbb{L}_*(R, \mathfrak{a})$.

Lemma 2.6. *Let R be a ring and $\mathfrak{a} \in \text{ass } \mathbb{L}_*(R)$ where $*$ $\in \{l, r, \emptyset\}$.*

1. *The set $\mathbb{L}_*(R, \mathfrak{a})$ is a commutative multiplicative semigroup where for $S, T \in \mathbb{L}_*(R, \mathfrak{a})$, ST is the submonoid of (R, \cdot) generated by S and T .*
2. *For all $S, T \in \mathbb{L}_*(R, \mathfrak{a})$, $S \cup T \subseteq ST$.*
3. *For all $S_1, S_2, T \in \mathbb{L}_*(R, \mathfrak{a})$ such that $S_1 \subseteq S_2$, $TS_1 \subseteq TS_2$.*
4. *The set $T_{*,\mathfrak{a}} := \bigcup_{S \in \mathbb{L}_*(R, \mathfrak{a})} S$ is the largest element in the set $\mathbb{L}_*(R, \mathfrak{a})$ (w.r.t. \subseteq), and $ST_{*,\mathfrak{a}} = T_{*,\mathfrak{a}}$ for all $S \in \mathbb{L}_*(R, \mathfrak{a})$.*
5. *For all $S, T \in \mathbb{L}_*(R, \mathfrak{a})$ such that $S \subseteq T$, $S^{-1}R \subseteq T^{-1}R \subseteq T_{*,\mathfrak{a}}(R)^{-1}R = \bigcup_{S' \in \mathbb{L}_*(R, \mathfrak{a})} S'^{-1}R$.*

Proof. Let $\pi : R \rightarrow \overline{R} := R/\mathfrak{a}$, $r \mapsto \overline{r} = r + \mathfrak{a}$.

1. By Theorem 1.6.(1), $\overline{S}, \overline{T} \in \text{Den}_*(\overline{R}, 0)$. By [3, Theorem 2.1.(1)], $\overline{ST} \in \text{Den}_*(\overline{R}, 0)$. Hence,

$$\overline{ST} = \overline{S}\overline{T} \in \text{Den}_*(\overline{R}, 0) \text{ and } \overline{ST} \subseteq ((\overline{ST})^{-1}\overline{R})^\times.$$

By Theorem 1.3.(3), $ST \in \mathbb{L}_*(R, \mathfrak{a})$.

2 and 3. Statements 2 and 3 are obvious.

4. Statement 4 follows from statement 2 and Lemma 2.5.

5. Statement 5 follows from Lemma 1.10 and Lemma 2.5. \square

Definition. Let R be a ring and $\mathfrak{a} \in \text{ass } \mathbb{L}_*(R, \mathfrak{a})$ where $*$ \in $\{l, r, \emptyset\}$. The localization of the ring R at $T_{*,\mathfrak{a}}(R)$ (see Lemma 2.6.(5)) is denoted by $\mathbb{Q}_{*,\mathfrak{a}}(R)$.

A left or right Ore set of a ring is called *regular* if it consists of regular elements of the ring. A regular left/right Ore set is automatically a left/right denominator set.

Theorem 2.7. [3, Theorem 2.1] For a ring R there is a largest (w.r.t. inclusion) regular left Ore set $S_l(R)$ in R and the ring $Q_l(R) := S_l(R)^{-1}R$ is called the largest left quotient ring of R .

For a ring R there is a largest regular Ore set $S(R)$ in R and the ring

$$Q(R) := S(R)^{-1}R \simeq RS(R)^{-1}$$

is called the *largest quotient ring* of R , [3, Theorem 4.1.(2)]. The interested reader is referred to [3] for more information about the largest regular left/right Ore sets and the largest left/right quotient rings. Notice that for a ring R , its classical left/right quotient ring does not always exist.

Theorem 2.8 describes the multiplicative sets $T_{*,\mathfrak{a}}(R)$ and the rings $\mathbb{Q}_{*,\mathfrak{a}}(R)$ via the largest regular Ore sets and the largest quotient rings.

Theorem 2.8. Let R be a ring, $\mathfrak{a} \in \text{ass } \mathbb{L}_*(R)$ and $\pi : R \rightarrow \overline{R} := R/\mathfrak{a}$, $r \mapsto \overline{r} = r + \mathfrak{a}$. Then

1. $T_{*,\mathfrak{a}}(R) = \pi^{-1}(S_*(\overline{R}))$ where $S_*(\overline{R})$ is the largest regular $*$ Ore set in \overline{R} .
2. $\mathbb{Q}_{*,\mathfrak{a}}(R) \simeq Q_*(\overline{R})$, an R -isomorphism where $Q_*(\overline{R}) = S_*(\overline{R})^{-1}\overline{R}$ for $*$ \in $\{l, \emptyset\}$ and $Q_r(\overline{R}) = \overline{R}S_r(\overline{R})^{-1}$.
3. $T_{*,\mathfrak{a}}(R) = \sigma^{-1}(Q_{*,\mathfrak{a}}(R)^\times) = \sigma^{-1}(Q_*(\overline{R})^\times)$ where $\sigma : R \rightarrow \mathbb{Q}_{*,\mathfrak{a}}(R) \simeq Q_*(\overline{R})$, $r \mapsto \frac{r}{1}$.

Proof. 1. The set $T := \pi^{-1}(S_*(\overline{R}))$ is a multiplicative set in R since $\pi(T) = S_*(\overline{R})$ is so. Clearly, $T_* := T_{*,\mathfrak{a}}(R) \subseteq T$ since $\pi(T_*) \in \text{Den}_*(\overline{R}, 0)$, by Theorem 1.3.(1). Since

$$\pi(T) = S_*(\overline{R}) \in \text{Den}_*(\overline{R}, 0) \text{ and } \mathfrak{b} := \text{ass}_R(T) \neq R,$$

we have that $T \in \mathbb{L}_*(R, \mathfrak{b})$ and $\mathfrak{b} \subseteq \mathfrak{a}$. Since $T_* \subseteq T$, we have that $\mathfrak{a} \subseteq \mathfrak{b}$, and so $\mathfrak{a} = \mathfrak{b}$. Therefore,

$$T \in \mathbb{L}_*(R, \mathfrak{a}).$$

Since $T_* \subseteq T$, we must have $T_* = T$, by the maximality of T_* .

2. Statement 2 follows from statement 1.
3. The map σ is the composition of the homomorphisms

$$R \xrightarrow{\pi} \overline{R} \xrightarrow{\overline{\sigma}} Q_{*,\mathfrak{a}}(R) \simeq Q_*(\overline{R})$$

where $\overline{\sigma}(r + \mathfrak{a}) = \frac{r}{1}$. By [3, Theorem 2.8.(1)], $S_*(\overline{R}) = \overline{\sigma}^{-1}(Q_*(\overline{R})^\times)$. Now, statement 3 follows from statement 1. \square

The \mathbb{L}_* -radical $\mathbb{L}_*\text{rad}(R)$.

Definition. An element S of the set $\max \mathbb{L}_*(R)$ where $*$ \in $\{l, r, \emptyset\}$ is called the *maximal left/right localizable set* and the *maximal localizable set*, resp., and the rings $S^{-1}R$, RS^{-1} and $S^{-1}R \simeq RS^{-1}$ are called the *maximal left/right localization* and the *maximal localization* of R , respectively.

Definition. The intersection

$$\mathbb{L}_*\text{rad}(R) = \bigcap_{S \in \max \mathbb{L}_*(R)} \text{ass}_R(S) \tag{13}$$

is called the *\mathbb{L}_* -radical* of R .

For a ring R , there is the canonical exact sequence where $*$ \in $\{l, r, \emptyset\}$,

$$0 \rightarrow \mathbb{L}_*\text{rad}(R) \rightarrow R \xrightarrow{\sigma} \prod_{S \in \max \mathbb{L}_*(R)} S^{-1}R, \quad \sigma := \prod_{S \in \max \mathbb{L}_*(R)} \sigma_S, \tag{14}$$

where $\sigma_S : R \rightarrow S^{-1}R, r \mapsto \frac{r}{1}$. A similar sequence exists for $* = r$.

Definition. The sets $\mathcal{L}\mathbb{L}_*(R) := \bigcup_{S \in \mathbb{L}_*(R)} S$ and $\mathcal{N}\mathcal{L}\mathbb{L}_*(R) := R \setminus \mathcal{L}\mathbb{L}_*(R)$ are called the set of *\mathbb{L}_* -localizable* and *\mathbb{L}_* -non-localizable* elements of R , resp., and the intersection

$$\mathcal{C}\mathbb{L}_*(R) = \bigcap_{S \in \max \mathbb{L}_*(R)} S$$

is called the *set of completely \mathbb{L}_* -localizable elements* of the ring R .

By the very definition the sets $\mathcal{L}\mathbb{L}_*(R)$, $\mathcal{N}\mathcal{L}\mathbb{L}_*(R)$ and $\mathcal{C}\mathbb{L}_*(R)$ are invariant under the action of the automorphism group of the ring R , i.e., they are *characteristic sets*.

The sets $'\mathbb{L}_l(R)$, $\mathbb{L}'_r(R)$ and $'\mathbb{L}'_{l,r}(R)$. Recall that for an ideal \mathfrak{a} of a ring R ,

$$'Den_l(R, \mathfrak{a}) = \{S \in Den_l(R, \mathfrak{a}) \mid S \subseteq 'C_R\} \quad \text{and} \quad Den'_r(R, \mathfrak{a}) = \{S \in Den_r(R, \mathfrak{a}) \mid S \subseteq C'_R\},$$

and a localizable set $S \in \mathbb{L}_*(R)$ is called *perfect* if $\text{ass}_R(S) = \mathfrak{a}(S)$. So, the perfect localizable sets S of R have the ‘smallest’ possible ideal $\text{ass}_R(S)$. Proposition 2.9 shows that the sets ${}^l\mathbb{L}_l(R)$, $\mathbb{L}'_r(R)$ and ${}^l\mathbb{L}'_{l,r}(R)$ are perfect localizable sets of the ring R . See the Introduction for their definitions.

Proposition 2.9. *We keep the notation as above.*

1. ${}^l\mathbb{L}_l(R) \subseteq \mathbb{L}^p_l(R)$ and for each $S \in {}^l\mathbb{L}_l(R)$, $\text{ass}_R(S) = \mathfrak{a}(S) = {}^l\pi^{-1}(\text{ass}_l({}^lS))$ and $R\langle S^{-1} \rangle \simeq {}^lS^{-1}R$ where $\text{ass}_l({}^lS) = \{r \in {}^lR \mid {}^lS r = 0 \text{ for some } {}^lS \in {}^lS\}$. Furthermore, ${}^lS \in {}^l\text{Den}_l({}^lR, \mathfrak{a}/{}^l\mathfrak{a})$ where $\mathfrak{a} = \text{ass}_R(S)$.
2. $\mathbb{L}'_r(R) \subseteq \mathbb{L}^p_r(R)$ and for each $S \in \mathbb{L}'_r(R)$, $\text{ass}_R(S) = \mathfrak{a}(S) = \pi'^{-1}(\text{ass}_r(S'))$ and $R\langle S^{-1} \rangle \simeq R'S'^{-1}$ where $\text{ass}_r(S') = \{r' \in R' \mid r's' = 0 \text{ for some } s' \in S'\}$. Furthermore, $S' \in \text{Den}'_r(R', \mathfrak{a}/\mathfrak{a}')$ where $\mathfrak{a} = \text{ass}_R(S)$.
3. ${}^l\mathbb{L}'_{l,r}(R) \subseteq \mathbb{L}^p(R)$ and for each $S \in {}^l\mathbb{L}'_{l,r}(R)$, $\text{ass}_R(S) = \mathfrak{a}(S) = {}^l\pi^{-1}(\text{ass}_l({}^lS)) = \pi'^{-1}(\text{ass}_r(S'))$ and $R\langle S^{-1} \rangle \simeq {}^lS^{-1}R \simeq R'S'^{-1}$. Furthermore, ${}^lS \in {}^l\text{Den}_l({}^lR, \mathfrak{a}/{}^l\mathfrak{a})$ and $S' \in \text{Den}'_r(R', \mathfrak{a}/\mathfrak{a}')$ where $\mathfrak{a} = \text{ass}_R(S)$.

Proof. 1. Recall that $\mathfrak{a}(S)$ is the ideal in Proposition 1.1.(1).

(i) ${}^l\mathfrak{a} \subseteq \mathfrak{a}(S)$: The inclusion follows from Proposition 1.1.(2) (by the minimality of the ideal ${}^l\mathfrak{a}$ since $(S + \mathfrak{a}(S))/\mathfrak{a}(S) \subseteq \mathcal{C}_{R/\mathfrak{a}(S)} \subseteq {}^l\mathcal{C}_{R/\mathfrak{a}(S)}$).

(ii) $\mathfrak{b} := {}^l\pi^{-1}(\text{ass}_l({}^lS)) \subseteq \mathfrak{a}(S)$: The inclusion $\mathfrak{b} \subseteq \mathfrak{a}(S)$ follows from the inclusion ${}^l\mathfrak{a} \subseteq \mathfrak{a}(S)$ and the definition of the ideals \mathfrak{b} and $\mathfrak{a}(S)$.

(iii) $\mathfrak{b} = \text{ass}_R(S)$: By Lemma 1.2, $\mathfrak{a}(S) \subseteq \text{ass}_R(S)$. Now,

$$\mathfrak{b} \subseteq \text{ass}_R(S),$$

by the statement (ii). Since $\tilde{S} := (S + \mathfrak{b})/\mathfrak{b} \in \text{Den}_l(R/\mathfrak{b}, 0)$, we must have

$$\mathfrak{b} \supseteq \text{ass}_R(S),$$

by the minimality of the ideal $\text{ass}_R(S)$ (Theorem 1.3.(3)), and the statement (iii) follows.

By the statements (ii) and (iii),

$$\text{ass}_R(S) = \mathfrak{a}(S) = \mathfrak{b}$$

(since $\mathfrak{b} \subseteq \mathfrak{a}(S) \subseteq \text{ass}_R(S) = \mathfrak{b}$) and $R\langle S^{-1} \rangle \simeq {}^lS^{-1}R$. Clearly, ${}^lS \in \text{Den}_l({}^lR, \mathfrak{a}/{}^l\mathfrak{a})$.

2. Statement 2 can be proven in a similar/dual way as statement 1.

3. Statement 3 follows from statements 1 and 2. \square

For $L \in \{{}^l\mathbb{L}_l(R), \mathbb{L}'_r(R), {}^l\mathbb{L}'_{l,r}(R)\}$, let $\text{ass}(L) := \{\text{ass}_R(S) \mid S \in L\}$. Then

$${}^l\mathbb{L}_l(R) = \coprod_{\mathfrak{a} \in \text{ass } {}^l\mathbb{L}_l(R)} {}^l\mathbb{L}_l(R, \mathfrak{a}) \text{ where } {}^l\mathbb{L}_l(R, \mathfrak{a}) = \{S \in {}^l\mathbb{L}_l(R) \mid \text{ass}_R(S) = \mathfrak{a}\}, \quad (15)$$

$$\mathbb{L}'_r(R) = \coprod_{\mathfrak{a} \in \text{ass } \mathbb{L}'_r(R)} \mathbb{L}'_r(R, \mathfrak{a}) \text{ where } \mathbb{L}'_r(R, \mathfrak{a}) = \{S \in \mathbb{L}'_r(R) \mid \text{ass}_R(S) = \mathfrak{a}\}, \quad (16)$$

$$\begin{aligned} {}'\mathbb{L}'_{l,r}(R) &= \coprod_{\mathfrak{a} \in \text{ass } {}'\mathbb{L}'_{l,r}(R)} {}'\mathbb{L}'_{l,r}(R, \mathfrak{a}) \text{ where } {}'\mathbb{L}'_{l,r}(R, \mathfrak{a}) = \{S \in {}'\mathbb{L}'_{l,r}(R) \mid \text{ass}_R(S) = \mathfrak{a}\}. \\ & \hspace{15em} (17) \end{aligned}$$

Left/right localizable Ore sets. Recall that for a ring R , the sets

$$\begin{aligned} \mathbb{LOre}_l(R) &:= \mathbb{L}_l(R) \cap \text{Ore}_l(R), \quad \mathbb{LOre}_r(R) := \mathbb{L}_r(R) \cap \text{Ore}_r(R), \\ \mathbb{LOre}(R) &:= \mathbb{L}(R) \cap \text{Ore}(R) = \text{Ore}(R) \end{aligned}$$

are called left, right and Ore localizable, respectively. Clearly, $\mathbb{LOre}(R) = \mathbb{LOre}_l(R) \cap \mathbb{LOre}_r(R)$ since

$$\begin{aligned} \mathbb{LOre}(R) &= \mathbb{L}(R) \cap \text{Ore}(R) = (\mathbb{L}_l(R) \cap \mathbb{L}_r(R)) \cap (\text{Ore}_l(R) \cap \text{Ore}_r(R)) \\ &= (\mathbb{L}_l(R) \cap \text{Ore}_l(R)) \cap (\mathbb{L}_r(R) \cap \text{Ore}_r(R)) = \mathbb{LOre}_l(R) \cap \mathbb{LOre}_r(R). \end{aligned}$$

For each element $*$ $\in \{l, r, \emptyset\}$, let $\text{ass } \mathbb{LOre}_*(R) := \{\text{ass}_R(S) \mid S \in \mathbb{LOre}_*(R)\}$. Then

$$\begin{aligned} \mathbb{LOre}_*(R) &= \coprod_{\mathfrak{a} \in \text{ass } \mathbb{LOre}_*(R)} \mathbb{LOre}_*(R, \mathfrak{a}) \\ \text{where } \mathbb{LOre}_*(R, \mathfrak{a}) &:= \{S \in \mathbb{LOre}_*(R) \mid \text{ass}_R(S) = \mathfrak{a}\}. \end{aligned} \quad (18)$$

Clearly, $\mathbb{LOre}_*(R, \mathfrak{a}) = \mathbb{L}_*(R, \mathfrak{a}) \cap \text{Ore}_*(R)$. Since $\text{Ore}(R) \subseteq \mathbb{L}(R)$ see (8), we have that $\mathbb{LOre}(R) = \text{Ore}(R)$. So, the letter ‘ \mathbb{L} ’ is redundant in the definition of the sets ‘ $\mathbb{LOre}(R)$ ’, ‘ $\text{ass } \mathbb{LOre}(R)$ ’ and ‘ $\max \mathbb{LOre}(R, \mathfrak{a})$ ’, and we drop it. So, (18) takes the form

$$\text{Ore}(R) = \coprod_{\mathfrak{a} \in \text{ass } \text{Ore}(R)} \text{Ore}(R, \mathfrak{a}) \text{ where } \text{Ore}(R, \mathfrak{a}) := \{S \in \text{Ore}(R) \mid \text{ass}_R(S) = \mathfrak{a}\}. \quad (19)$$

Proposition 2.10.

1. Given $S \in \mathbb{LOre}_l(R)$. Then ${}'\mathfrak{a} \neq R$ (see Proposition 1.1.(2)) and the ideal ${}'\mathfrak{a}$ is the least ideal \mathfrak{b} of the ring R such that $(S + \mathfrak{b})/\mathfrak{b} \in {}'\text{Den}_l(R/\mathfrak{b})$. In particular, $(S + {}'\mathfrak{a})/{}'\mathfrak{a} \in {}'\text{Den}_l(R/{}'\mathfrak{a})$.
2. Given $S \in \mathbb{LOre}_r(R)$. Then $\mathfrak{a}' \neq R$ (see Proposition 1.1.(3)) and the ideal \mathfrak{a}' is the least ideal \mathfrak{b} of the ring R such that $(S + \mathfrak{b})/\mathfrak{b} \in \text{Den}'_r(R/\mathfrak{b})$. In particular, $(S + \mathfrak{a}')/\mathfrak{a}' \in \text{Den}'_r(R/\mathfrak{a}')$.

Proof. 1. By Proposition 1.1 and Lemma 1.2, ${}'\mathfrak{a} \subseteq \mathfrak{a}(S) \subseteq \text{ass}_R(S) \neq R$. Then

$${}'S := (S + {}'\mathfrak{a})/{}'\mathfrak{a} \in {}'\text{Den}_l(R/{}'\mathfrak{a}).$$

Since the ideal \mathfrak{a} is the least ideal of the ring R such that $(S + \mathfrak{a})/\mathfrak{a} \in {}^{\prime}\mathcal{C}_{R/\mathfrak{a}}$, statement 1 follows.

2. Statement 2 is proven in a dual way to statement 1. \square

Proof of Proposition 1.4. 1. By Proposition 2.10.(1), $\mathbb{L}\text{Ore}_l(R) \subseteq {}^{\prime}\mathbb{L}_l(R)$. By Proposition 2.9.(1), ${}^{\prime}\mathbb{L}_l(R) \subseteq \mathbb{L}_l^p(R)$. Now,

$$\begin{aligned} \mathbb{L}\text{Ore}_l(R) &= {}^{\prime}\mathbb{L}_l(R) \cap \mathbb{L}\text{Ore}_l(R) = {}^{\prime}\mathbb{L}_l(R) \cap \mathbb{L}(R) \cap \text{Ore}_l(R) = {}^{\prime}\mathbb{L}_l(R) \cap \text{Ore}_l(R), \\ \mathbb{L}\text{Ore}_l(R) &= \mathbb{L}_l^p(R) \cap \mathbb{L}\text{Ore}_l(R) = \mathbb{L}_l^p(R) \cap \mathbb{L}(R) \cap \text{Ore}_l(R) = \mathbb{L}_l^p(R) \cap \text{Ore}_l(R). \end{aligned}$$

2. Statement 2 can be proven in a similar way to statement 1.

3. Since $\text{Ore}(R) = \mathbb{L}\text{Ore}_l(R) \cap \mathbb{L}\text{Ore}_r(R)$, statement 3 follows from statements 1 and 2 and the fact that $\text{Ore}(R) \subseteq \mathbb{L}(R)$, see (8). \square

Using some of the above results we obtain criteria for a left/right Ore set to be a left/right localizable set, Theorem 1.5 and Theorem 2.11.

Proof of Theorem 1.5. 1 (\Rightarrow) If $S \in \mathbb{L}_l(R)$ then $\mathfrak{a} \neq R$. Now, the implication follows from the inclusions ${}^{\prime}\mathfrak{a} \subseteq \mathfrak{a}(S) \subseteq \mathfrak{a}$ (Lemma 1.2).

2(a) If ${}^{\prime}\mathfrak{a} \neq R$ then clearly ${}^{\prime}S \in {}^{\prime}\text{Den}({}^{\prime}R)$.

1 (\Leftarrow) Since ${}^{\prime}\mathfrak{a} \subseteq \mathfrak{a}(S) \subseteq \mathfrak{a}$ (Lemma 1.2), the implication follows from the statement 2(a) and Lemma 2.1.(1).

2(b,c) Since ${}^{\prime}\mathfrak{a} \subseteq \mathfrak{a}$ (Lemma 1.2),

$$R\langle S^{-1} \rangle = {}^{\prime}R\langle {}^{\prime}S^{-1} \rangle,$$

by Lemma 2.1.(2). Now, the statements (b) and (c) follow from the inclusion ${}^{\prime}S \in {}^{\prime}\text{Den}_l({}^{\prime}R)$ (the statement (a)). \square

Theorem 2.11.(1) is a criterion for a right Ore set to be a left localizable set and Theorem 2.11.(2) describes the structure of the localization of a ring at a localizable right Ore set.

Theorem 2.11. *Let R be a ring, $S \in \text{Ore}_r(R)$, and $\mathfrak{a} = \text{ass}_R(S)$. Then*

1. $S \in \mathbb{L}_r(R)$ iff $\mathfrak{a}' \neq R$ where the ideal $\mathfrak{a}' = \mathfrak{a}'(S)$ of R is as in Proposition 1.1.(3) and (12).
2. Suppose that $\mathfrak{a}' \neq R$. Let $\pi' : R \rightarrow R'; = R/\mathfrak{a}'$, $r \mapsto r' = r + \mathfrak{a}'$ and $S' = \pi'(S)$. Then
 - (a) $S' \in \text{Den}'_r(R')$,
 - (b) $\mathfrak{a} = (\pi')^{-1}(\text{ass}_r(S'))$,
 - (c) $RS^{-1} \simeq R'S'^{-1}$, an R -isomorphism.

Proof. The proof of the theorem is dual to the proof of Theorem 1.5. \square

Description of the ideal $\text{ass}_R(S)$ for $S \in \mathbb{L}_*(R)$. For each localizable set $S \in \mathbb{L}_*(R, \mathfrak{a})$, Proposition 2.12 describes the ideal \mathfrak{a} .

Proposition 2.12. *Let R be a ring and S be a multiplicative set in R .*

1. *Let $S \in \mathbb{L}_l(R, \mathfrak{a})$. For each pair of elements $s \in S$ and $r \in R$ fix a pair of elements $s_1 \in S$ and $r_1 \in R$ such that $s_1r - r_1s \in \mathfrak{a}$, and let \mathfrak{b}_l be the ideal of R generated by the elements $s_1r - r_1s$, $\pi_l : R \rightarrow R_l := R/\mathfrak{b}_l$, $r \mapsto r + \mathfrak{b}_l$. Then $S_l := \pi_l(S) \in \mathbb{LOre}_l(R_l, \mathfrak{a}/\mathfrak{b}_l)$ and $\mathfrak{a} = \pi_l^{-1}(\mathfrak{a}(S_l))$ where the ideal $\mathfrak{a}(S_l)$ of the ring R_l is defined in Proposition 1.1.(1).*
2. *Let $S \in \mathbb{L}_r(R, \mathfrak{a})$. For each pair of elements $s \in S$ and $r \in R$ fix a pair of elements $s_1 \in S$ and $r_1 \in R$ such that $rs_1 - sr_1 \in \mathfrak{a}$, and let \mathfrak{b}_r be the ideal of R generated by the elements $rs_1 - sr_1$, $\pi_r : R \rightarrow R_r := R/\mathfrak{b}_r$, $r \mapsto r + \mathfrak{b}_r$. Then $S_r := \pi_r(S) \in \mathbb{LOre}_r(R_r, \mathfrak{a}/\mathfrak{b}_r)$ and $\mathfrak{a} = \pi_r^{-1}(\mathfrak{a}(S_r))$.*
3. *Let $S \in \mathbb{L}(R, \mathfrak{a})$, $\mathfrak{b} = \mathfrak{b}_l + \mathfrak{b}_r$ and $\tilde{\pi} : R \rightarrow \tilde{R}$, $r \mapsto r + \mathfrak{b}$. Then $\tilde{S} := \tilde{\pi}(S) \in \text{Ore}(\tilde{R}, \mathfrak{a}/\mathfrak{b})$ and $\mathfrak{a} = \tilde{\pi}^{-1}(\mathfrak{a}(\tilde{S}))$.*

Proof. 1. By the very definition, $\mathfrak{b}_l \subseteq \mathfrak{a}$. By Lemma 2.1.(2),

$$S_l \in \mathbb{L}_l(R_l, \mathfrak{a}/\mathfrak{b}_l).$$

By the definition of \mathfrak{b}_l , $S_l \in \mathbb{LOre}_l(R_l, \mathfrak{a}/\mathfrak{b}_l)$. By Proposition 1.4.(1), $\mathfrak{a}/\mathfrak{b}_l = \mathfrak{a}(S_l)$, and so $\mathfrak{a} = \pi_l^{-1}(\mathfrak{a}(S_l))$.

2 and 3. Statements 2 and 3 can be proven in a similar way. \square

Proposition 2.13 gives a sufficient condition for an epimorphic image of a left localizable set to be a left localizable left Ore set.

Proposition 2.13. *Let R be a ring, $S \in \mathbb{L}_l(R, \mathfrak{a})$, \mathfrak{b} be an ideal of R such that $\mathfrak{a} \subseteq \mathfrak{b}$ and $\bar{R} = R/\mathfrak{b}$. Suppose that the left ideal $S^{-1}R\mathfrak{b}$ is an ideal of the ring $S^{-1}R$ such that $S^{-1}R\mathfrak{b} \neq S^{-1}R$. Then $\bar{S} := (S + \mathfrak{b})/\mathfrak{b} \in \mathbb{LOre}_l(\bar{R}, \bar{\mathfrak{c}})$ for some ideal \mathfrak{c} of R such that $\mathfrak{b} \subseteq \mathfrak{c}$ where $\bar{\mathfrak{c}} = \mathfrak{c}/\mathfrak{b}$.*

Proof. Since $S \in \mathbb{L}_l(R, \mathfrak{a})$ and $\mathfrak{a} \subseteq \mathfrak{b}$, $\bar{S} \in \text{Ore}_l(\bar{R})$. Let \mathfrak{b}_1 be the kernel of the ring homomorphism

$$R \rightarrow S^{-1}R/S^{-1}R\mathfrak{b}, \quad r \mapsto \frac{r}{1} + S^{-1}R\mathfrak{b}.$$

Clearly, $\mathfrak{b} \subseteq \mathfrak{b}_1$, $\tilde{S} = (S + \mathfrak{b}_1)/\mathfrak{b}_1 \in \text{Den}_l(R/\mathfrak{b}_1, 0)$ and $S^{-1}R/S^{-1}R\mathfrak{b} \simeq \tilde{S}^{-1}(R/\mathfrak{b}_1)$. By Theorem 1.3.(3), $\bar{S} \in \mathbb{LOre}_l(\bar{R}, \bar{\mathfrak{c}})$ for some ideal \mathfrak{c} of R such that $\mathfrak{b} \subseteq \mathfrak{c}$. \square

3. Localization of a ring at an Ore set

The localization of a ring at an Ore set was introduced in [3]. The aim of this section is to prove Theorem 1.6 which among other things explains what is the localization of a ring at an Ore set and why it always exists. Proofs of Corollary 1.7 and Proposition 1.8 are given.

Proof of Theorem 1.6. 1. (i) $\mathfrak{a} + \mathfrak{a} \subseteq \mathfrak{a}$: Given elements $a, a' \in \mathfrak{a}$. Then $sat = s'a't' = 0$ for some elements $s, s', t, t' \in S$. Fix elements $s_1, t_1 \in S$ such that $s_1s = \alpha s'$ and $tt_1 = t'\beta$ for some elements $\alpha, \beta \in R$. Then $s_1s, tt_1 \in S$ and

$$s_1s(a + a')t_1 = s_1(sat)t_1 + \alpha(s'a't')\beta = 0 + 0 = 0,$$

and so $a + a' \in \mathfrak{a}$.

(ii) $R\mathfrak{a}R \subseteq \mathfrak{a}$: Given elements $r, r' \in R$ and $a \in \mathfrak{a}$. Then $sas' = 0$ for some elements $s, s' \in S$. Using the left and right Ore conditions, we have the equalities

$$s_1r = r_1s \quad \text{and} \quad r'_1s'_1 = s'r'_1$$

for some elements $s_1, s'_1 \in S$ and $r_1, r'_1 \in R$. Then

$$s_1rar'_1s'_1 = r_1sas'r'_1 = r_10r'_1 = 0,$$

and so $rar' \in \mathfrak{a}$, and the statement (ii) follows.

The statements (i) and (ii) imply that the set \mathfrak{a} is an ideal of the ring R .

(iii) $\mathfrak{a} \neq R$: If $\mathfrak{a} = R$ then $1 \in \mathfrak{a}$, and so $s \cdot 1 \cdot t = 0$ for some elements $s, t \in S$. Then $0 = st \in S$, a contradiction.

2. (i) $S \cap \mathfrak{a} = \emptyset$: If $r \in S \cap \mathfrak{a}$ then $srt = 0$ for some elements $s, t \in S$. This is not possible since $0 = srt \in S$, a contradiction.

(ii) $\overline{S} \in \text{Den}(\overline{R}, 0)$: Clearly, $\overline{S} \in \text{Ore}(\overline{R})$. If $\overline{s\overline{r}} = 0$ or $\overline{r'\overline{s'}} = 0$ for some elements $s, s' \in S$ and $r, r' \in R$. Then $sr \in \mathfrak{a}$ and $r's' \in \mathfrak{a}$. Then $s_1srs'_1 = 0$ or $s_2r's'_1s'_2 = 0$ for some elements $s_1, s_2, s'_1, s'_2 \in S$, and so $r \in \mathfrak{a}$ and $r' \in \mathfrak{a}$, i.e., $\overline{r} = 0$ and $\overline{r'} = 0$, and the statement (ii) follows.

(iii) $S^{-1}R \simeq \overline{S}^{-1}\overline{R}$ and $\mathfrak{a} = \text{ass}_R(S)$: The statement (iii) follows from Theorem 1.3.(2,3).

(iv) $\mathfrak{a} = \mathfrak{a}(S) = \text{ass}_R(S)$: The result follows from the inclusions $\mathfrak{a} \subseteq \mathfrak{a}(S) \subseteq \text{ass}_R(S)$ and the equality $\mathfrak{a} = \text{ass}_R(S)$ (the statement (iii)).

(v) $S \in \mathbb{L}(R, \mathfrak{a})$: The statement (v) follows from statements (iii) and (iv).

3. (i) $\mathfrak{a} \subseteq \mathfrak{b}$: Given an element $a \in \mathfrak{a}$. Then $sat = 0$ for some elements $s, t \in S$, and so $s^\dagger a^\dagger t^\dagger = 0$. Hence $a^\dagger = 0$ in R^\dagger since $s^\dagger, t^\dagger \in \mathcal{C}_{R^\dagger}$, and so $a \in \mathfrak{b}$, and so $\mathfrak{a} \subseteq \mathfrak{b}$.

(ii) The map $\overline{S}^{-1}\overline{R} \rightarrow S^{\dagger^{-1}}R^\dagger, \overline{s}^{-1}\overline{r} \mapsto s^{\dagger^{-1}}r^\dagger$ is a ring epimorphism: The statement (ii) follows from the universal property of localization.

4(a) (i) $Q = \{f(s)^{-1}f(r) \mid s \in S, r \in R\} = \{f(r)f(s)^{-1} \mid s \in S, r \in R\}$: The statement (i) follows from the fact that $S \in \text{Ore}(R)$ and that the ring Q is generated by the sets $f(R)$ and $\{f(s)^{-1} \mid s \in S\}$.

(ii) $\mathfrak{a} \subseteq \ker(f)$: Given an element $a \in \mathfrak{a}$. Then $sat = 0$ for some elements $s, t \in S$, and so

$$0 = f(sat) = f(a)f(s)f(t).$$

Hence, $f(a) = 0$ since $f(s), f(t) \in Q^\times$, and so $a \in \ker(f)$, and the statement (ii) follows.

(iii) *The map $\overline{S}^{-1}\overline{R} \rightarrow Q, \overline{s}^{-1}\overline{r} \mapsto f(s)^{-1}f(r)$ is a ring epimorphism*: Since the ring Q is generated by the set $f(R)$ and $\{f(s)^{-1} \mid s \in S\}$, and $\mathfrak{a} \subseteq \ker(f)$, the statement (iii) follows from the universal property of localization.

(iv) *The kernel of the map in the statement (iii) is $\overline{S}^{-1}(\ker(f)/\mathfrak{a})$* : The statement is obvious.

4(b) The statement (b) follows from the statement (a). \square

Definition ([3]). Let $S \in \text{Ore}(R)$. The ring $\overline{S}^{-1}\overline{R} \simeq \overline{RS}^{-1}$ in Theorem 1.6.(2) is called the *localization of the ring R at the Ore set S* and is denoted by $S^{-1}R = RS^{-1}$. The ideal \mathfrak{a} in Theorem 1.6.(1) is denoted by $\text{ass}(S)$.

Statement 4 (and statement 3) of Theorem 1.6 is the universal property of localization of a ring at Ore set. If the Ore set is a denominator set this is precisely the universal property of localization at a denominator set.

Proof of Corollary 1.7. 1. Statement 1 follows from Theorem 1.6.(1).

2. (i) $S_l \in \text{Ore}_r(R_l)$ with $\text{ass}_l(S_l) = 0$: Since $S \in \text{Ore}(R)$ and $R_l = R/\text{ass}_l(S)$, the statement (i) is obvious.

(ii) $S_l \in \text{Den}'_r(R_l, \mathfrak{a}/\mathfrak{a}_l)$: Since $S \in \text{Ore}(R)$, we have that $S_l \in \text{Den}_r(R_l)$ since $\text{ass}_l(S_l) = 0 \subseteq \text{ass}_r(S_l)$, by the statement (i). By Theorem 1.6.(1),

$$\text{ass}_r(S_l) = \mathfrak{a}/\mathfrak{a}_l.$$

In more detail, $\pi_l(r) \in \text{ass}_r(S_l)$ iff $0 = \pi_l(r)\pi_l(t) = \pi_l(rt)$ for some element $t \in S$ iff $rt \in \mathfrak{a}_l$ iff $srt = 0$ for some $s \in S$ iff $r \in \mathfrak{a}$, by Theorem 1.6.(1).

(iii) $R_l S_l^{-1} \simeq S^{-1}R \simeq RS^{-1}$: The statement (iii) follows from Theorem 1.6.(2).

3. Statement 3 is proven in a similar fashion as statement 2.

4. Let $'\mathfrak{a} = \mathfrak{a}(S)$ and $\mathfrak{a}' = \mathfrak{a}'(S)$. By the very definition of the ideals \mathfrak{a}_l and \mathfrak{a}_r ,

$$'\mathfrak{a} \supseteq \mathfrak{a}_r \text{ and } \mathfrak{a}' \supseteq \mathfrak{a}_l.$$

The reverse inclusions follow from statements 2 and 3 and the ‘minimality of the ideals $'\mathfrak{a}$ and \mathfrak{a}' in the sense of Proposition 1.1.(2,3). \square

Proof of Proposition 1.8. (i) \tilde{S} is a multiplicative set of R such that $S \subseteq \tilde{S}$: The set $\overline{S} = \pi(S)$ is a multiplication set of the ring \overline{R} , hence the set $\tilde{S} = \pi^{-1}(\overline{S})$ is a multiplicative set of R .

(ii) $\tilde{S} \in \mathbb{L}(R)$: It suffices to show that $\tilde{S} \in \mathbb{L}_l(R)$ since then by symmetry we will get $\tilde{S} \in \mathbb{L}_r(R)$. Given elements $s \in \tilde{S}$ and $r \in R$. Since $\pi(\tilde{s}) = \overline{S} \in \text{Den}_l(\overline{R}, 0)$, $\overline{s_1 r} = \overline{r_1 s}$ for some elements $\overline{s_1} \in \overline{S}$ and $\overline{r_1} \in \overline{R}$. Then $s_1 r - r_1 s \in \mathfrak{a}$. By Theorem 1.6.(1), $\mathfrak{a} \subseteq \text{ass}_R(\tilde{S})$, $\tilde{S} \in \mathbb{L}_l(R)$.

(iii) $\tilde{S} \in \mathbb{L}(R, \mathfrak{a})$: Let $\tilde{\mathfrak{a}} = \text{ass}_R(\tilde{S})$. We have to show that $\tilde{\mathfrak{a}} = \mathfrak{a}$. Since $S \subseteq \tilde{S}$, we have the inclusion

$$\mathfrak{a} \subseteq \tilde{\mathfrak{a}},$$

by Lemma 1.10. Since $\pi(\tilde{S}) = \overline{S} \in \text{Den}_l(R, 0)$, we have the reverse inclusion

$$\mathfrak{a} \supseteq \tilde{\mathfrak{a}}$$

(by Theorem 1.6.(2,4)), and so $\tilde{\mathfrak{a}} = \mathfrak{a}$.

(iv) $\tilde{S}^{-1}R \simeq S^{-1}R$ is an R -isomorphism: By the statement (iii), $\tilde{S}^{-1}R \simeq \pi(S)^{-1}\overline{R} = \overline{S}^{-1}\overline{R} = S^{-1}R$. \square

The following obvious lemma is a useful criterion when an epimorphism image of Ore set is an Ore set.

Lemma 3.1. Let R be a ring, $S \in \text{Ore}(R)$ and \mathfrak{b} be an ideal of R . Then $\overline{S} := (S + \mathfrak{b})/\mathfrak{b} \in \text{Ore}(R/\mathfrak{b})$ iff $S \cap \mathfrak{b} = \emptyset$.

Proof. Straightforward. \square

An element a of ring R is called a *normal element* if $aR = Ra$. Suppose that R is a K -algebra over a field K . A K -linear map $* : R \rightarrow R$, $r \mapsto r^*$ is an *involution* if $(rs)^* = s^*r^*$ and $s^{**} = s$ for all elements $s, r \in R$.

We keep the notation of Corollary 1.7. The next example shows that for an Ore set S of R , the ideals \mathfrak{a}_l , \mathfrak{a}_r and \mathfrak{a} are distinct and $\mathfrak{a} = \mathfrak{a}_l + \mathfrak{a}_r$, but, in general, $\mathfrak{a} \not\subseteq \mathfrak{a}_l + \mathfrak{a}_r$ (see the second example).

Example. Let $P = K[x_1, x_2, \dots, y_1, y_2, \dots]$ be a polynomial algebra over a field K in countably many variables $x_1, x_2, \dots, y_1, y_2, \dots$. Let R be a K -algebra generated by P and an element a subject to the defining relations:

$$ax_1 = 0, \quad ax_i = x_{i-1}a \quad (i \geq 2), \quad y_1a = 0, \quad y_ia = ay_{i-1} \quad (i \geq 2).$$

The K -algebra R admits a K -involution $*$ where

$$a^* = a, \quad x_i^* = y_i \quad \text{and} \quad y_i^* = x_i \quad \text{for all } i \geq 1.$$

The element a is a left normal element of R , i.e.,

$$aR \subseteq Ra.$$

In view of the involution $*$ and the fact $a^* = a$, the element a is also a right normal, and so the element a is a normal element of the ring R . Hence,

$$S_a := \{a^i \mid i \in \mathbb{N}\} \in \text{Ore}(R), \quad \text{ass}_l(S_a) = (x_1, x_2, \dots), \quad \text{ass}_r(R) = (y_1, y_2, \dots),$$

$\text{ass}(S_a) = (x_1, \dots, y_1, \dots) = \text{ass}_l(S_a) + \text{ass}_r(S_a)$, and $S_a^{-1}R \simeq RS_a^{-1} \simeq K[a, a^{-1}]$ (Corollary 1.7.(2,3)). Clearly, the ideals $\text{ass}_l(S_a)$, $\text{ass}_r(S_a)$ and $\text{ass}(S_a)$ are distinct and

$$\text{ass}(S_a) = \text{ass}_l(S_a) + \text{ass}_r(S_a).$$

The next example shows that, in general, $\text{ass}(S_a)$ properly contains the sum $\text{ass}_l(S_a) + \text{ass}_r(S_a)$.

Example. Let $P = K[x_1, x_2, \dots, y_1, y_2, \dots]$ be a polynomial algebra over a field K in variables $x_1, x_2, \dots, y_1, y_2, \dots$. Let R be a K -algebra generated by P and an element a subject to the defining relations:

$$ax_1 = 0, \quad ax_i = x_{i-1}a \quad (i \geq 2), \quad y_1a = 0, \quad y_2a = a(y_1 + x_2), \quad y_i a = ay_{i-1} \quad (i \geq 3).$$

Then a is a normal element of R (since $aR \subseteq Ra$ (as $ay_1 = (y_2 - x_1)a$) and $Ra \subseteq aR$), hence $S_a := \{a^i \mid i \in \mathbb{N}\} \in \text{Ore}(R)$, $\text{ass}_l(S_a) = (x_1, x_2, \dots)$, $\text{ass}_r(R) = (y_1)$,

$\text{ass}(S_a) = (x_1, x_2, \dots, y_1, y_2, \dots) \not\supseteq \text{ass}_l(S_a) + \text{ass}_r(S_a)$ and $S_a^{-1}R \simeq RS_a^{-1} \simeq K[a, a^{-1}]$ (Corollary 1.7.(2,3)).

4. Localization of a ring at an almost Ore set

The aim of this section is to introduce almost Ore sets, to give a criterion for an almost Ore set to be a localizable set (Theorem 4.1.(1), Theorem 4.2.(2) and Theorem 4.3.(2)) and for each localizable almost Ore set S to give an explicit description of the ideal $\text{ass}_R(S)$ and of the ring $S^{-1}R$ (Theorem 4.1.(2), Theorem 4.2.(3) and Theorem 4.3.(3)).

Almost Ore sets. Let R be a ring. Recall that a multiplicative set S of R is a *left* (resp., *right*) *Ore set* if the left (resp., right) Ore condition holds: For any elements $s \in S$ and $r \in R$, there are elements $s_1 \in S$ and $r_1 \in R$ such that $s_1r = r_1s$ (resp., $rs_1 = sr_1$). A left and right Ore set of R is called an *Ore set* of R .

Definition. Let R be a ring. A multiplicative set S of R is called an *almost left* (resp., *right*) *Ore set* of R if the *almost left* (resp., *right*) *Ore condition* holds:

(ALO) For any elements $s \in S$ and $r \in R$ there are elements $s_1, s_2 \in S$ and $r_1 \in R$ such that $(s_1r - r_1s)s_2 = 0$.

(ARO) For any elements $s \in S$ and $r \in R$ there are elements $s_1, s_2 \in S$ and $r_1 \in R$ such that $s_2(rs_1 - sr_1) = 0$.

Let $\text{AOre}_l(R)$ and $\text{AOre}_r(R)$ be the sets of almost left and right Ore sets, respectively. Elements of the set $\text{AOre}(R) := \text{AOre}_l(R) \cap \text{AOre}_r(R)$ are called *almost Ore sets*.

Clearly, $\text{Ore}_*(R) \subseteq \text{AOre}_*(R)$ for $*$ in $\{l, r, \emptyset\}$. Let

$$\begin{aligned} \mathbb{L}\text{AO}_l(R) &:= \mathbb{L}_l(R) \cap \text{AOre}_l(R), \\ \mathbb{L}\text{AO}_r(R) &:= \mathbb{L}_r(R) \cap \text{AOre}_r(R), \\ \mathbb{L}\text{AO}(R) &:= \mathbb{L}\text{AO}_l(R) \cap \mathbb{L}\text{AO}_r(R) = \mathbb{L}(R) \cap \text{AOre}(R). \end{aligned}$$

Elements of the sets $\mathbb{L}\text{AO}_l(R)$ and $\mathbb{L}\text{AO}_r(R)$ are called *left localizable almost left Ore sets* and *right localizable almost right Ore sets*, respectively. Elements of the set $\mathbb{L}\text{AO}(R)$ are called *localizable almost Ore sets*.

Let $\text{ass } \mathbb{L}\text{AO}_*(R) := \{\text{ass}_R(S) \mid S \in \mathbb{L}\text{AO}_*(R)\}$. Then

$$\begin{aligned} \mathbb{L}\text{AO}_*(R) &= \coprod_{\mathfrak{a} \in \text{ass } \mathbb{L}\text{AO}_*(R)} \mathbb{L}\text{AO}_*(R, \mathfrak{a}) \\ \text{where } \mathbb{L}\text{AO}_*(R, \mathfrak{a}) &:= \{S \in \mathbb{L}\text{AO}_*(R) \mid \text{ass}_R(S) = \mathfrak{a}\}. \end{aligned} \tag{20}$$

Clearly, $\mathbb{L}\text{AO}_*(R, \mathfrak{a}) = \mathbb{L}\text{AO}_*(R) \cap \mathbb{L}_*(R, \mathfrak{a})$.

Theorem 4.1.(1) is a criterion for an almost Ore set to be a localizable set. It also gives an explicit description of the ideal $\text{ass}_R(S)$ for each $S \in \mathbb{L}\text{AO}(R)$ (Theorem 4.1.(2b)).

Theorem 4.1. *Let R be a ring, $S \in \text{AOre}(R)$, \mathfrak{a}_l and \mathfrak{a}_r be the ideals of R generated by the sets $\text{ass}_l(S) = \{r \in R \mid sr = 0 \text{ for some } s \in S\}$ and $\text{ass}_r(S) = \{r \in R \mid rs = 0 \text{ for some } s \in S\}$, respectively.*

1. *The following statements are equivalent:*
 - (a) $S \in \mathbb{L}(R)$.
 - (b) $\mathfrak{a}_l + \mathfrak{a}_r \neq R$.
2. *Suppose that $\tilde{\mathfrak{a}} = \mathfrak{a}_l + \mathfrak{a}_r \neq R$. Let $\tilde{R} = R/\tilde{\mathfrak{a}}$ and $\tilde{S} = (S + \tilde{\mathfrak{a}})/\tilde{\mathfrak{a}}$. Then*
 - (a) $\tilde{S} \in \text{Ore}(\tilde{R})$.
 - (b) $\text{ass}_R(S) = \tilde{\pi}^{-1}(\mathfrak{a}^\circ)$ where $\tilde{\pi} : R \rightarrow \tilde{R}$, $r \mapsto \tilde{r} = r + \tilde{\mathfrak{a}}$ and \mathfrak{a}° is the ideal in Theorem 1.6.(1) for the Ore set $\tilde{S} \in \text{Ore}_l(\tilde{R})$, that is $\mathfrak{a}^\circ = \{\tilde{r} \in \tilde{R} \mid \tilde{s}\tilde{r}\tilde{t} = 0 \text{ for some elements } \tilde{s}, \tilde{t} \in \tilde{S}\}$.
 - (c) Let $\mathfrak{a} = \text{ass}_R(S)$ and $\pi : R \rightarrow \overline{R} := R/\mathfrak{a}$, $r \mapsto \bar{r} = r + \mathfrak{a}$. Then $\overline{S} := \pi(S) \in \text{Den}(R, 0)$.
 - (d) $S^{-1}R \simeq \overline{S}^{-1}\overline{R}$, an R -isomorphism.

Proof. Recall that $\mathfrak{a} = \text{ass}_R(S)$ and $\mathfrak{a}(S)$ is the least ideal of the ring R such that $(S + \mathfrak{a}(S))/\mathfrak{a}(S) \subseteq \mathcal{C}_{R/\mathfrak{a}(S)}$ (Proposition 1.1.(1)).

(i) $\tilde{\mathfrak{a}} \subseteq \mathfrak{a}(S) \subseteq \mathfrak{a}$: The first inclusion follows from (10) and the second one does from Lemma 1.2.

1 ($a \Rightarrow b$) If $S \in \mathbb{L}(R)$ then $\mathfrak{a} \neq R$, and so $\tilde{\mathfrak{a}} \neq R$, by the statement (i).

1 ($b \Rightarrow a$)

(ii) If $\tilde{\mathfrak{a}} \neq R$ then $\tilde{S} \in \text{Ore}(\tilde{R})$: The statement (ii) follows at once from the conditions (ALO) and (ARO).

Since each Ore set is localizable, the implication ($b \Rightarrow a$) follows from the statement (ii).

2(a) The statement (a) is the same as the statement (ii).

2(b,c,d) The statements (b), (c) and (d) follow from the statement (a) and Theorem 1.6.(1,2). \square

Criterion for an almost left/right Ore set to be a left localizable set. Theorem 4.2 is such a criterion.

Theorem 4.2. *Let R be a ring, $S \in \text{AOre}_l(R)$, $\mathfrak{a} = \text{ass}_R(S)$, \mathfrak{a}_r be the ideal of R generated by the set $\text{ass}_r(S) = \{r \in R \mid rs = 0 \text{ for some } s \in S\}$. Then*

1. $S_r := (S + \mathfrak{a}_r)/\mathfrak{a}_r \in \text{Ore}_l(R/\mathfrak{a}_r)$.
2. $S \in \mathbb{L}_l(R)$ iff $S_r \in \mathbb{L}_l(R/\mathfrak{a}_r)$ iff $'\mathfrak{a}(S_r) \neq R/\mathfrak{a}_r$ iff $'\mathfrak{a}(S) \neq R$ (see Proposition 1.1.(2) and (11)).
3. Suppose that $'\mathfrak{a} := '\mathfrak{a}(S) \neq R$. Let $'\pi : R \rightarrow 'R = R/'\mathfrak{a}$, $r \mapsto 'r = r + '\mathfrak{a}$ and $'S = '\pi(S)$. Then
 - (a) $'S \in '\text{Den}_l('R)$.
 - (b) $\mathfrak{a} = '\pi^{-1}(\text{ass}_l('S))$.
 - (c) $S^{-1}R \simeq 'S^{-1}'R$, an R -isomorphism.

Proof. 1. Statement 1 follows at once from the condition (ALO) and the definition of the ideal \mathfrak{a}_r .

2. Since $\mathfrak{a}_r \subseteq \mathfrak{a}(S) \subseteq \text{ass}_R(S)$ (Lemma 1.2), $S \in \mathbb{L}_l(R)$ iff $S_r \in \mathbb{L}_l(R/\mathfrak{a}_r)$, by Lemma 2.1.(1). Recall that $S_r \in \text{Ore}_l(R/\mathfrak{a}_r)$ (statement 1). Now, by Theorem 1.5.(1),

$$S_r \in \mathbb{L}_l(R/\mathfrak{a}_r) \text{ iff } '\mathfrak{a}(S_r) \neq R/\mathfrak{a}_r.$$

Since $\mathfrak{a}_r \subseteq '\mathfrak{a}(S)$, $'\mathfrak{a}(S)/\mathfrak{a}_r = '\mathfrak{a}(S_r)$. Therefore,

$$'\mathfrak{a}(S_r) \neq R/\mathfrak{a}_r \text{ iff } '\mathfrak{a}(S) \neq R.$$

3. Since $'\mathfrak{a} \neq R$, the set S_r is a localizable left Ore set of the ring R/\mathfrak{a}_r , by statements 1 and 2. Since $\mathfrak{a}_r \subseteq '\mathfrak{a}(S) \subseteq \mathfrak{a}(S) \subseteq \text{ass}_R(S)$, $R\langle S^{-1} \rangle \simeq (R/\mathfrak{a}_r)\langle S_r^{-1} \rangle$ (Lemma 2.1.(1)) and $'\mathfrak{a}(S)/\mathfrak{a}_r = '\mathfrak{a}(S_r)$, statement 3 follows from Theorem 1.5.(2). \square

Theorem 4.3. *Let R be a ring, $S \in \text{AOre}_r(R)$, $\mathfrak{a} = \text{ass}_R(S)$, \mathfrak{a}_l be the ideal of R generated by the set $\text{ass}_l(S) = \{r \in R \mid sr = 0 \text{ for some } s \in S\}$. Then*

1. $S_l := (S + \mathfrak{a}_l)/\mathfrak{a}_l \in \text{Ore}_r(R/\mathfrak{a}_l)$.
2. $S \in \mathbb{L}_r(R)$ iff $S_l \in \mathbb{L}_r(R/\mathfrak{a}_l)$ iff $\mathfrak{a}'(S_l) \neq R/\mathfrak{a}_l$ iff $\mathfrak{a}'(S) \neq R$ (see Proposition 1.1.(3) and (12)).
3. Suppose that $\mathfrak{a}' := \mathfrak{a}'(S) \neq R$. Let $\pi' : R \rightarrow R' = R/\mathfrak{a}'$, $r \mapsto r' = r + \mathfrak{a}'$ and $S' = \pi'(S)$. Then
 - (a) $S' \in \text{Den}'_r(R')$.
 - (b) $\mathfrak{a} = \pi'^{-1}(\text{ass}_r(S'))$.
 - (c) $RS^{-1} \simeq R'S'^{-1}$, an R -isomorphism.

Proof. The proof of the theorem is ‘dual’ to the proof of Theorem 4.2. \square

Localizable almost Ore sets are perfect localizable sets. Corollary below shows that localizable almost Ore sets are perfect localizable sets.

Proposition 4.4.

1. $\mathbb{L}AO_l(R) \subseteq {}^l\mathbb{L}_l(R) \subseteq \mathbb{L}_l^p(R)$ and $\mathbb{L}AO_l(R) = {}^l\mathbb{L}_l(R) \cap \text{AOre}_l(R) = \mathbb{L}_l^p(R) \cap \text{AOre}_l(R)$.
2. $\mathbb{L}AO_r(R) \subseteq \mathbb{L}'_r(R) \subseteq \mathbb{L}_r^p(R)$ and $\mathbb{L}AO_r(R) = \mathbb{L}'_r(R) \cap \text{AOre}_r(R) = \mathbb{L}_r^p(R) \cap \text{AOre}_r(R)$.
3. $\mathbb{L}AO(R) \subseteq {}^l\mathbb{L}'_{l,r}(R) \subseteq \mathbb{L}^p(R)$.

Proof. 1. Statement 1 follows from Theorem 4.2.(3).
 2. Statement 2 follows from Theorem 4.3.(3).
 3. Statement 3 follows from statements 1 and 2. \square

5. Classification of maximal localizable sets and maximal Ore sets in semiprime Goldie ring

The aim of this section is to classify the maximal Ore sets in a semiprime Goldie ring (Theorem 1.11.(1)). One of the key results that is used in the proof of Theorem 1.11 is Theorem 5.2. The concept the maximal left denominator set of a ring was introduced and studied in [3].

Lemma 5.1. *Let R be a ring and $S \in \text{Ore}(R)$. For any elements $s, t \in S$, there is an element $\nu \in S$ such that $\nu = x_1s = x_2t = sy_1 = ty_2$ for some elements $x_i, y_i \in R$ for $i = 1, 2$.*

Proof. Since $S \in \text{Ore}_l(R)$, $s_1s = r_1t$ for some elements $s_1 \in S$ and $r_1 \in R$. Since $S \in \text{Ore}_r(R)$, $ss_2 = tr_2$ for some elements $s_2 \in S$ and $r_2 \in R$. Then $\nu = ss_2s_1s \in S$ and

$$\nu = s \cdot (s_2 s_1 s) = s s_2 s_1 \cdot s = t \cdot r_2 s_1 s = s s_2 r_1 \cdot t,$$

as required. \square

Theorem 5.2. *Let R be a semiprime left Goldie ring. Then $\mathcal{C}_R \cap \text{ass}_R(S) = \emptyset$ for all $S \in \text{Ore}(R)$.*

Proof. The ring R is a semiprime left Goldie ring. By Goldie’s Theorem, its classical left quotient ring $Q = Q_{l,cl}(R) := \mathcal{C}_R^{-1}R$ is a semisimple Artinian ring, $Q = \prod_{i=1}^n Q_i$ where Q_i are simple Artinian rings. The map

$$\sigma : R \rightarrow Q = \prod_{i=1}^n Q_i, \quad r \mapsto \frac{r}{1} = (r_1, \dots, r_n)$$

is a ring monomorphism. We identify the ring R via σ with its image in Q . So, $r = \frac{r}{1} = (r_1, \dots, r_n)$ where $r_i = \frac{r}{1} \in Q_i$.

Suppose that $\mathcal{C}_R \cap \text{ass}_R(S) \neq \emptyset$ for some $S \in \text{Ore}(R)$, we seek a contradiction. Fix an element $c \in \mathcal{C}_R \cap \text{ass}_R(S)$. By Theorem 1.6.(1),

$$sct = 0$$

for some elements $s, t \in S$. By Lemma 5.1, we can assume that $s = t$, i.e., $scs = 0$, i.e., $Q_i \ni s_i c_i s_i = 0$ for all $i = 1, \dots, n$ where $s = (s_1, \dots, s_n)$ and $c = (c_1, \dots, c_n)$. Clearly, $c_i \in Q_i^\times$ for all $i = 1, \dots, n$. Hence,

$$s_i \notin Q_i^\times \text{ for all } i = 1, \dots, n$$

(since $s_i c_i s_i = 0$). The ideal of R , $\mathfrak{a}_l = \text{ass}_l(S)$, is a nonzero ideal (since $s \cdot cs = 0$ and $0 \neq cs \in \mathfrak{a}_l$). The ring Q is a left Noetherian ring. Hence $\mathcal{C}_R^{-1}\mathfrak{a}_l$ is an ideal of Q . If $s_i = 0$ then $\mathcal{C}_R^{-1}\mathfrak{a}_l \cap Q_i = Q_i$. Suppose that $s_i \neq 0$. Then $0 \neq c_i s_i \in \mathcal{C}_R^{-1}\mathfrak{a}_l \cap Q_i$, and so

$$0 \neq \mathcal{C}_R^{-1}\mathfrak{a}_l \cap Q_i = Q_i$$

since Q_i is a simple ring. Therefore, $\mathcal{C}_R^{-1}\mathfrak{a}_l \cap Q_i = Q_i$ for all $i = 1, \dots, n$, and so

$$\mathcal{C}_R^{-1}\mathfrak{a}_l = \prod_{i=1}^n Q_i = Q.$$

Hence, $\mathcal{C}_R \cap \mathfrak{a}_l \neq \emptyset$. Fix an element $a \in \mathcal{C}_R \cap \mathfrak{a}_l$. Then $s'a = 0$ for some $s' \in S$ (since $a \in \mathfrak{a}_l = \text{ass}_l(S)$) but $a \in \mathcal{C}_R$, hence $0 = s' \in S$, a contradiction. \square

Classification of maximal Ore sets of a semiprime Goldie ring.

Proof of Theorem 1.11. 1. By [4, Theorem 4.1],

$$\max \text{Den}(R) = \{C(\mathfrak{p}) \mid \mathfrak{p} \in \min(R)\}.$$

By Theorem 1.12,

$$\{C(\mathfrak{p}) \mid \mathfrak{p} \in \min(R)\} = \mathcal{N}_* \text{ for } * \in \{l, r, \emptyset\}.$$

To finish the proof of statement 1 it suffices to show that every Ore set S of the ring R is contained in a maximal denominator set.

The ring R is a semiprime Goldie ring. The left and right quotient ring of R ,

$$Q_{cl}(R) = C_R^{-1}R \simeq RC_R^{-1} \simeq \prod_{i=1}^n Q_i$$

is a product of simple Artinian ring Q_i . Let $\mathfrak{a} = \text{ass}_R(S)$. By Theorem 5.2,

$$C_R^{-1}\mathfrak{a} \neq Q.$$

Hence, up to order, $C_R^{-1}\mathfrak{a} = \prod_{i=m+1}^n Q_i$ for some m such that $1 \leq m < n$. We have ring homomorphisms

$$\sigma : R \xrightarrow{\pi} R/\mathfrak{a} \xrightarrow{\tau} C_R^{-1}(R/\mathfrak{a}) \simeq \overline{Q} := \prod_{i=1}^m Q_i$$

where $\sigma = \tau\pi$, $\pi(r) = \bar{r} := r + \mathfrak{a}$ and $\tau(\bar{r}) = \frac{\bar{r}}{1}$. The homomorphism π is an epimorphism and the homomorphism τ is a monomorphism. Let $\overline{S} = \pi(S)$. By Theorem 1.6.(2),

$$\overline{S} \in \text{Den}(R/\mathfrak{a}, 0).$$

In particular, $\overline{S} \subseteq \overline{Q}^\times$. Therefore, $\sigma_1(S) \subseteq Q_1^\times$ where $\sigma_1 : R \rightarrow Q \rightarrow Q_1$ where the first map is $r \mapsto \frac{r}{1}$ and the second is the projection onto Q_1 . Therefore,

$$S \subseteq \sigma_1^{-1}(Q_1^\times).$$

By the explicit description of the set $\max \text{Den}(R)$ at the beginning of the proof,

$$\sigma_1^{-1}(Q_1^\times) \in \max \text{Den}(R),$$

as required.

2. In view of the first equality in statement 1, statement 2 is [4, Theorem 4.1.(2d)].
3. In view of the first equality in statement 1, statement 3 is [4, Theorem 4.1].
4. In view of the first equality in statement 1, statement 4 is [4, Theorem 4.1.(2c)]. \square

6. Localization of a module at a localizable set

The aim of the section is to introduce the concept of localization of a module at a localizable set and to consider its basic properties.

Definition. Let R be a ring, $S \in \mathbb{L}_*(R, \mathfrak{a})$ where $*$ \in $\{l, \emptyset\}$ and M be a left R -module. Then the left $S^{-1}R$ -module

$$S^{-1}M := S^{-1}R \otimes_R M$$

is called the (left) localization of M at S . If $S \in \mathbb{L}_*(R, \mathfrak{a})$ where $*$ \in $\{r, \emptyset\}$ and M be a right R -module. Then the right RS^{-1} -module

$$MS^{-1} := M \otimes_R RS^{-1}$$

is called the (right) localization of M at S .

We consider the case when $*$ \in $\{l, \emptyset\}$ and M is a left R -module. By the very definition, $S^{-1}M$ is a left $S^{-1}R$ -module. The elements of the $S^{-1}R$ -module $S^{-1}M$ are denoted by $s^{-1}m$. In particular, $s^{-1}r \otimes m = s^{-1}rm$ for $s \in S$, $r \in R$ and $m \in M$, and $\frac{m}{1} := 1 \otimes m$. The map

$$i_M : M \rightarrow S^{-1}M, \quad m \mapsto 1 \otimes m$$

is an R -homomorphism.

Proposition 6.1. Let R be a ring, $S \in \mathbb{L}_*(R, \mathfrak{a})$ where $*$ \in $\{l, \emptyset\}$, M be an R -module, and $i_M : M \rightarrow S^{-1}M, m \mapsto 1 \otimes m$. Then

1. $S^{-1}M \simeq \overline{S}^{-1}(M/\mathfrak{a}M)$ where $\overline{S} := (S + \mathfrak{a})/\mathfrak{a} \in \text{Den}_l(\overline{R}, 0)$ and $\overline{R} = R/\mathfrak{a}$ (Theorem 1.3.(1)).
2. Let \mathcal{M} be an $S^{-1}R$ -module and $f : M \rightarrow \mathcal{M}$ be an R -homomorphism. Then there is a unique $S^{-1}R$ -homomorphism $S^{-1}f : S^{-1}M \rightarrow \mathcal{M}$ such that $f = S^{-1}f \circ i_M$.
3. $\mathfrak{t}_S(M) := \ker(i_M) = \{m \in M \mid sm \in \mathfrak{a}M \text{ for some } s \in S\}$.

Proof. 1. Let $\overline{M} = M/\mathfrak{a}M$. Then $S^{-1}(\mathfrak{a}M) = S^{-1}R \otimes_R \mathfrak{a}M = S^{-1}R\mathfrak{a} \otimes_R M = 0$, and so

$$S^{-1}M = S^{-1}\overline{M} = S^{-1}R \otimes_R \overline{M} \simeq \overline{S}^{-1}\overline{R} \otimes_{\overline{R}} \overline{M} = \overline{S}^{-1}\overline{M}.$$

2. The R -homomorphism $f : M \rightarrow \mathcal{M}$ determines a ring homomorphism

$$R \rightarrow \text{End}_{\mathbb{Z}}(\mathcal{M}), \quad r \mapsto (m \mapsto rm).$$

The images of the elements of the set S in $\text{End}_{\mathbb{Z}}(\mathcal{M})$ are units. Now, statement 2 follows from Theorem 1.3.(4).

3. Statement 3 follows from statement 1. \square

For a ring R , let $R\text{-mod}$ be the category of left R -modules. By Proposition 6.1.(1), the localization at S functor,

$$S^{-1} : R\text{-mod} \rightarrow S^{-1}R\text{-mod}, \quad M \mapsto S^{-1}M,$$

is the composition of two right exact functors

$$S^{-1} = \overline{S}^{-1} \circ (R/\mathfrak{a} \otimes_R -). \tag{21}$$

Therefore, the functor S^{-1} is also a *right exact functor* for all $S \in \mathbb{L}_*(R)$: Given a short exact sequence of R -modules $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$, then the sequence of $S^{-1}R$ -modules

$$0 \rightarrow \overline{S}^{-1}(M_1 \cap \mathfrak{a}M_2/\mathfrak{a}M_1) \rightarrow S^{-1}M_1 \rightarrow S^{-1}M_2 \rightarrow S^{-1}M_3 \rightarrow 0 \tag{22}$$

is exact. Notice that

$$M_1 \cap \mathfrak{a}M_2/\mathfrak{a}M_1 \simeq \overline{M}_1 \cap (\mathfrak{a}M_2/\mathfrak{a}M_1).$$

An R -module M is called *S-torsion* (resp., *S-torsionfree*) if $S^{-1}M = 0$ (resp., $\mathfrak{t}_S(M) = 0$, i.e., the map $i_M : M \rightarrow S^{-1}M$, $m \mapsto 1 \otimes m$ is an R -module monomorphism). Let $\mathfrak{f}_S(M) = \text{im}(i_M)$, the image of the map i_M , and we have a short exact sequence of R -modules

$$0 \rightarrow \mathfrak{t}_S(M) \rightarrow M \rightarrow \mathfrak{f}_S(M) \rightarrow 0. \tag{23}$$

Clearly, $\mathfrak{a}M \subseteq \mathfrak{t}_S(M)$,

$$\mathfrak{t}_S(M)/\mathfrak{a}M = \text{tor}_{\overline{S}}(\overline{M}) := \{\overline{m} \in \overline{M} \mid \overline{s}m = 0 \text{ for some element } \overline{s} \in \overline{S}\}$$

where $\overline{M} = M/\mathfrak{a}M$ and

$$\mathfrak{f}_S(M) = M/\mathfrak{t}_S(M) \simeq (M/\mathfrak{a}M)/(\mathfrak{t}_S(M)/\mathfrak{a}M) \simeq \overline{M}/\text{tor}_{\overline{S}}(\overline{M}).$$

By taking the short exact sequence (23) modulo $\mathfrak{a}M$, we obtain a short exact sequence of \overline{R} -modules

$$0 \rightarrow \text{tor}_{\overline{S}}(\overline{M}) \rightarrow \overline{M} \rightarrow \overline{M}/\text{tor}_{\overline{S}}(\overline{M}) \rightarrow 0. \tag{24}$$

Lemma 6.2. *Let R be a ring, $S \in \mathbb{L}_*(R, \mathfrak{a})$ where $*$ $\in \{l, \emptyset\}$ and M be an R -module. Then*

1. $\mathfrak{t}_S \mathfrak{f}_S(M) = 0$ and so the R -module $\mathfrak{f}_S(M)$ is S -torsionfree.
2. $\mathfrak{f}_S \mathfrak{f}_S(M) = \mathfrak{f}_S(M)$.

Proof. 1. $\mathfrak{t}_S \mathfrak{f}_S(M) = \mathfrak{t}_S(\overline{M}/\text{tor}_{\overline{S}}(\overline{M})) = \text{tor}_{\overline{S}}(\overline{M}/\text{tor}_{\overline{S}}(\overline{M})) = 0$.

2. $\mathfrak{f}_S \mathfrak{f}_S(M) \simeq \mathfrak{f}_S(M)/\mathfrak{t}_S \mathfrak{f}_S(M) = \mathfrak{f}_S(M)$ since $\mathfrak{t}_S \mathfrak{f}_S(M) = 0$, by statement 1. \square

Theorem 6.3 is a criterion for the functor $S^{-1} : M \rightarrow S^{-1}M$ to be exact.

Theorem 6.3. *Let R be a ring, $S \in \mathbb{L}_*(R, \mathfrak{a})$ where $*$ $\in \{l, \emptyset\}$, $\overline{R} = R/\mathfrak{a}$ and $\overline{S} := (S + \mathfrak{a})/\mathfrak{a}$. The functor S^{-1} is exact iff for all R -modules M_1 and M_2 such that $M_1 \subseteq M_2$, the \overline{R} -modules $M_1 \cap \mathfrak{a}M_2/\mathfrak{a}M_1$ is \overline{S} -torsion.*

Proof. The theorem follows from the exact sequence (22). \square

Corollary 6.4. *Let R be a ring, $S \in \mathbb{L}_*(R, \mathfrak{a})$ where $*$ $\in \{l, \emptyset\}$, $\overline{R} = R/\mathfrak{a}$ and $\overline{S} := (S + \mathfrak{a})/\mathfrak{a}$. If the functor S^{-1} is exact then the \overline{R} -module $\mathfrak{a}/\mathfrak{a}^2$ is \overline{S} -torsion (recall that $\overline{S} \in \text{Den}_l(\overline{R}, 0)$, by Theorem 1.3.(1)).*

Proof. Applying Theorem 6.3 to the pair of R -modules $M_1 = \mathfrak{a} \subseteq M_2 = R$, we conclude that the \overline{R} -module $(M_1 \cap \mathfrak{a}M_2)/\mathfrak{a}M_1 = \mathfrak{a}/\mathfrak{a}^2$ is \overline{S} -torsion. \square

For an R -module M and $S \in \mathbb{L}_*(R, \mathfrak{a})$, we have a descending chain of R -modules

$$\mathfrak{t}_S(M) \supseteq \mathfrak{t}_S^2(M) \supseteq \dots \supseteq \mathfrak{t}_S^n(M) \supseteq \dots$$

where $\mathfrak{t}_S^n(M) = \mathfrak{t}_S \mathfrak{t}_S \dots \mathfrak{t}_S(M)$, n times.

7. Examples

In this section, we consider several examples and present explicitly some of the concepts that are introduced in the paper.

The algebras \mathbb{S}_n , $n \geq 1$, of one-sided inverses. Let K be a field and K^\times be its group of units, and $P_n := K[x_1, \dots, x_n]$ be a polynomial algebra over K .

Definition ([1]). The algebra \mathbb{S}_n of one-sided inverses of P_n is an algebra generated over a field K by $2n$ elements $x_1, \dots, x_n, y_1, \dots, y_n$ that satisfy the defining relations:

$$y_1 x_1 = \dots = y_n x_n = 1, \quad [x_i, y_j] = [x_i, x_j] = [y_i, y_j] = 0 \quad \text{for all } i \neq j,$$

where $[a, b] := ab - ba$, the commutator of elements a and b .

Let us recall some results of [1] that are used in proofs below. By the very definition, the algebra \mathbb{S}_n is obtained from the polynomial algebra P_n by adding commuting, left

(or right) inverses of its canonical generators. The algebra S_1 is a well-known primitive algebra [9], p. 35, Example 2. Over the field \mathbb{C} of complex numbers, the completion of the algebra S_1 is the *Toeplitz algebra* which is the C^* -algebra generated by a unilateral shift on the Hilbert space $l^2(\mathbb{N})$ (note that $y_1 = x_1^*$). The Toeplitz algebra is the universal C^* -algebra generated by a proper isometry.

Clearly, $S_n = S_1^{\otimes n}$ and $S_1 = K\langle x, y \mid yx = 1 \rangle = \bigoplus_{i,j \geq 0} Kx^i y^j$. For each natural number $d \geq 1$, let $M_d(K) := \bigoplus_{i,j=0}^{d-1} KE_{ij}$ be the algebra of d -dimensional matrices where $\{E_{ij}\}$ are the matrix units, and $M_\infty(K) := \varinjlim M_d(K) = \bigoplus_{i,j \in \mathbb{N}} KE_{ij}$ be the algebra (without 1) of infinite dimensional matrices. The algebra S_1 contains the ideal $F := \bigoplus_{i,j \in \mathbb{N}} KE_{ij}$, where

$$E_{ij} := x^i y^j - x^{i+1} y^{j+1}, \quad i, j \geq 0. \tag{25}$$

For all natural numbers i, j, k , and l , $E_{ij} E_{kl} = \delta_{jk} E_{il}$ where δ_{jk} is the Kronecker delta function. The ideal F is an algebra (without 1) isomorphic to the algebra $M_\infty(K)$ via $E_{ij} \mapsto E_{ij}$. For all $i, j \geq 0$,

$$xE_{ij} = E_{i+1,j}, \quad yE_{ij} = E_{i-1,j} \quad (E_{-1,j} := 0), \tag{26}$$

$$E_{ij}x = E_{i,j-1}, \quad E_{ij}y = E_{i,j+1} \quad (E_{i,-1} := 0). \tag{27}$$

$$S_1 = K \oplus xK[x] \oplus yK[y] \oplus F, \tag{28}$$

the direct sum of vector spaces. Then

$$S_1/F \simeq K[x, x^{-1}] =: L_1, \quad x \mapsto x, \quad y \mapsto x^{-1}, \tag{29}$$

since $yx = 1, xy = 1 - E_{00}$ and $E_{00} \in F$.

Lemma 7.1 is used in the proof of Proposition 7.2.

Lemma 7.1. *Let R be a ring, \mathfrak{a} be an ideal of R , and $\pi : R \rightarrow \overline{R} := R/\mathfrak{a}, r \mapsto r + \mathfrak{a}$. Suppose that S is a multiplicative set in R such that $\overline{S} := \pi(S) \in \text{Den}_*(\overline{R}, \overline{\mathfrak{b}})$ and $\mathfrak{a} \subseteq \text{ass}_*(S)$ where $*$ $\in \{l, r, \emptyset\}$. Then $S \in \text{Den}_*(R, \mathfrak{b})$ where $\mathfrak{b} = \pi^{-1}(\overline{\mathfrak{b}})$.*

Proof. We prove the lemma for $*$ = l . The other two cases can be proven in a similar way. For each element $r \in R$, let $\overline{r} = \pi(r)$.

(i) $S \in \text{Ore}_l(R)$: Given elements $s \in S$ and $r \in R$. Then $\overline{s} \in \overline{S}$ and $\overline{r} \in \overline{R}$. Since \overline{S} is a left Ore set in \overline{R} , $\overline{s}_1 \overline{r} = \overline{r}_1 \overline{s}$ for some elements $s_1 \in S$ and $r_1 \in R$. Hence,

$$a := s_1 r - r_1 s \in \mathfrak{a}.$$

Since $\mathfrak{a} \subseteq \text{ass}_l(S)$, we can choose an element, say $s_2 \in S$, such that $0 = s_2 a = s_2 s_1 r - s_2 r_1 s$, and the statement (i) follows.

(ii) $\text{ass}_l(S) = \mathfrak{b}$: Given an element $b \in \mathfrak{b}$. Then $\bar{b} \in \bar{\mathfrak{b}}$, and so $\overline{sb} = 0$ for some element $s \in S$ (since $\bar{S} \in \text{Den}_l(\bar{R}, \bar{\mathfrak{b}})$). Hence, $sb \in \mathfrak{a}$, and so $tsb = 0$ for some element $t \in S$ (since $\mathfrak{a} \subseteq \text{ass}_l(S)$). Therefore, $b \in \text{ass}_l(S)$ and $\mathfrak{b} \subseteq \text{ass}_l(S)$.

Conversely, given an element $a \in \text{ass}_l(S)$. Then $sa = 0$ for some element $s \in S$. Then $\overline{sa} = 0$, and so $\bar{a} \in \bar{\mathfrak{b}}$ and $a \in \mathfrak{b}$. Therefore, $\mathfrak{b} \supseteq \text{ass}_l(S)$, and the statement (ii) follows.

(iii) $S \in \text{Den}_l(R, \mathfrak{b})$: In view of the statements (i) and (ii), we have to show that if $as = 0$ for some elements $a \in R$ and $s \in S$ then $a \in \mathfrak{b}$. Clearly, $\overline{as} = 0$, and so $\bar{a} \in \bar{\mathfrak{b}}$. Hence, $a \in \pi^{-1}(\bar{\mathfrak{b}}) = \mathfrak{b}$, as required. \square

The algebra \mathbb{S}_n admits the *involution*

$$\eta : \mathbb{S}_n \rightarrow \mathbb{S}_n, \quad x_i \mapsto y_i, \quad y_i \mapsto x_i, \quad i = 1, \dots, n,$$

i.e., it is a K -algebra anti-isomorphism ($\eta(ab) = \eta(b)\eta(a)$ for all $a, b \in \mathbb{S}_n$) such that $\eta^2 = \text{id}_{\mathbb{S}_n}$, the identity map on \mathbb{S}_n . So, the algebra \mathbb{S}_n is *self-dual* (i.e., it is isomorphic to its opposite algebra, $\eta : \mathbb{S}_n \simeq \mathbb{S}_n^{\text{op}}$). This means that left and right algebraic properties of the algebra \mathbb{S}_n are the same.

Let $\mathfrak{a}_n := (x_1y_1 - 1, \dots, x_ny_n - 1)$, an ideal of \mathbb{S}_n . By [1, Eq. (19)], the factor algebra

$$\mathbb{S}_n/\mathfrak{a}_n = L_n = K[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$$

is the Laurent polynomial algebra. Clearly, $L_n^\times = \{\lambda x^\alpha \mid \lambda \in K^\times, \alpha \in \mathbb{Z}^n\}$ where $x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$. Let

$$\sigma : \mathbb{S}_n \rightarrow L_n, \quad a \mapsto a + \mathfrak{a}_n.$$

Proposition 7.2. *Let $X = \langle x_1, \dots, x_n \rangle$ and $Y = \langle y_1, \dots, y_n \rangle$ be multiplicative submonoids of (\mathbb{S}_n, \cdot) that are generated by the elements in the brackets. Then*

1. $Y \in \text{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ and $Y^{-1}\mathbb{S}_n = L_n$.
2. $X \in \text{Den}_r(\mathbb{S}_n, \mathfrak{a}_n)$ and $\mathbb{S}_nX^{-1} = L_n$.

Proof. Recall that $\mathbb{S}_n = \mathbb{S}_1^{\otimes n}$. By [1, Eq. (19)], $\mathfrak{a}_n = \mathfrak{p}_1 + \cdots + \mathfrak{p}_i + \cdots + \mathfrak{p}_n$ where

$$\mathfrak{p}_1 = F \otimes \mathbb{S}_{n-1}, \dots, \mathfrak{p}_i = \mathbb{S}_{i-1} \otimes F \otimes \mathbb{S}_{n-i}, \dots, \mathfrak{p}_n = \mathbb{S}_{n-1} \otimes F.$$

By (26), $\mathfrak{p}_i \subseteq \text{ass}_l(S_i)$ where $S_i = \{y_i^j \mid j \geq 0\} \subseteq Y$. Hence, $\mathfrak{a}_n \subseteq \text{ass}_l(Y)$. Notice that $Y \in \text{Den}_l(L_n, 0)$. By Lemma 7.1, $Y \in \text{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$. Now,

$$Y^{-1}\mathbb{S}_n \simeq Y^{-1}(\mathbb{S}_n/\mathfrak{a}_n) = Y^{-1}L_n = L_n.$$

2. By applying the involution η of the algebra \mathbb{S}_n to statement 1 we obtain statement 2 (since $\eta(\mathfrak{a}_n) = \mathfrak{a}_n$, $\eta(Y) = X$ and $\eta(X) = Y$). \square

For the algebra \mathbb{S}_n and its multiplicative set Y , Lemma 7.3 presents explicitly all the ingredients of Proposition 1.1 and Theorem 1.3.

Lemma 7.3.

1. $Y \in \text{Ore}(\mathbb{S}_n)$ and $Y \notin \text{Den}_r(\mathbb{S}_n)$, $Y \subseteq {}'C_{\mathbb{S}_n}$, $\text{ass}_l(Y) = \mathfrak{a}_n$, and $\text{ass}_r(Y) = 0$.
2. The ideals $\mathfrak{a}(Y) = \mathfrak{a}(Y)' = \mathfrak{a}_n$ and ${}'\mathfrak{a}(Y) = 0$ (see Proposition 1.1).
3. We keep the notation of Theorem 1.3. Then for all $* \in \{l, r, \emptyset\}$,
 - (a) $Y \in \mathbb{L}_*(\mathbb{S}_n, \mathfrak{a})$, $\mathfrak{a} = \mathfrak{a}_n$, and $Y^{-1}\mathbb{S}_n \simeq \mathbb{S}_n Y^{-1} \simeq L_n$,
 - (b) $\overline{\mathbb{S}}_n := \mathbb{S}_n/\mathfrak{a} = \mathbb{S}_n/\mathfrak{a}_n = L_n$,
 - (c) $\overline{Y} = \tilde{Y} \in \text{Den}_*(\overline{\mathbb{S}}_n, 0)$.

Proof. 1. The equalities $y_i x_i = 1$, $i = 1, \dots, n$, imply that $y_i \in {}'C_{\mathbb{S}_n}$, and so $Y \subseteq {}'C_{\mathbb{S}_n}$. Hence, $\text{ass}_r(Y) = 0$. By Proposition 7.2.(1), $\text{ass}_l(Y) = \mathfrak{a}_n$. Hence, $Y \notin \text{Den}_r(\mathbb{S}_n)$ (since $0 \neq \mathfrak{a}_n = \text{ass}_l(Y) \not\subseteq \text{ass}_r(Y) = 0$).

By Proposition 7.2.(1), $Y \in \text{Ore}_l(\mathbb{S}_n)$. To finish the proof of statement 1, it remains to show that $Y \in \text{Ore}_r(\mathbb{S}_n)$. Since $\mathbb{S}_n = \mathbb{S}_1^{\otimes n}$, it suffices to prove the statement for $n = 1$, that is $Y = \{y^i \mid i \geq 0\}$, we drop the subscript ‘1’. The algebra \mathbb{S}_1 is generated by the elements x and y , and $Y = \{y^i \mid i \geq 0\}$. So, it suffices to check that the right Ore condition holds for the elements $x \in \mathbb{S}_1$ and $y \in Y$, i.e. to prove that there are elements $a \in \mathbb{S}_1$ and y^i such that $xy^i = ya$. It suffices to take $i = 2$ and $a = 1 - E_{11}$:

$$xy^2 = (1 - (1 - xy))y = (1 - E_{00})y = y - E_{01} = y(1 - E_{11}).$$

2. By statement 1, $Y \subseteq {}'C_{\mathbb{S}_n}$, and so ${}'\mathfrak{a}(Y) = 0$. By Proposition 7.2.(1), $Y \in \text{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$. Hence, $\mathfrak{a}(Y) = \mathfrak{a}_n$. On the one hand, $\mathfrak{a}(Y)' \subseteq \mathfrak{a}(Y) = \mathfrak{a}_n$, by Proposition 1.1.(3). On the other hand, $\mathfrak{a}_n \subseteq \mathfrak{a}(Y)'$, by (26). Therefore, $\mathfrak{a}(Y)' = \mathfrak{a}_n$.

3. The case $* = l$ follows from the fact that $Y \in \text{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ (Proposition 7.2.(1)). It suffices to consider the case where $* = r$. By statement 1, $\text{ass}_l(Y) = \mathfrak{a}_n$. Clearly, $\text{ass}_l(Y) \subseteq \text{ass}_R(Y)$. Since $\mathbb{S}_n/\text{ass}_l(Y) = \mathbb{S}_n/\mathfrak{a}_n = L_n$ and the elements of the set Y are units in the Laurent polynomial ring L_n , we have that $\text{ass}_R(Y) = \mathfrak{a}_n$, $Y \in \mathbb{L}_r(\mathbb{S}_n, \mathfrak{a}_n)$ and $\mathbb{S}_n Y^{-1} \simeq L_n$. Now statements (b) and (c) follow. \square

For the algebra \mathbb{S}_n and its multiplicative set X , Lemma 7.4 presents explicitly all the ingredients of Proposition 1.1 and Theorem 1.3.

Lemma 7.4.

1. $X \in \text{Ore}(\mathbb{S}_n)$ and $X \notin \text{Den}_l(\mathbb{S}_n)$, $X \subseteq C'_{\mathbb{S}_n}$, $\text{ass}_r(X) = \mathfrak{a}_n$, and $\text{ass}_l(X) = 0$.
2. The ideals $\mathfrak{a}(X) = {}'\mathfrak{a}(X) = \mathfrak{a}_n$ and $\mathfrak{a}(X)' = 0$ (see Proposition 1.1).
3. We keep the notation of Theorem 1.3. Then for all $* \in \{l, r, \emptyset\}$,
 - (a) $X \in \mathbb{L}_*(\mathbb{S}_n, \mathfrak{a})$, $\mathfrak{a} = \mathfrak{a}_n$, and $X^{-1}\mathbb{S}_n \simeq \mathbb{S}_n X^{-1} \simeq L_n$,

- (b) $\overline{S}_n := S_n/\mathfrak{a} = S_n/\mathfrak{a}_n = L_n,$
- (c) $\overline{X} = \tilde{X} \in \text{Den}_*(\overline{S}_n, 0).$

Proof. Since $\eta(Y) = X$ and $\eta(\mathfrak{a}_n) = \mathfrak{a}_n,$ the lemma follows from Lemma 7.3. \square

Localization of a ring at the powers of a normal non-nilpotent element. In commutative algebra and algebraic geometry a localization of a commutative ring at the powers of a non-nilpotent element plays a prominent role in many proofs. An analogue of this situation for a noncommutative ring is a localization of a ring at the powers of a normal non-nilpotent element, see Lemma 7.5 for details.

Let R be a ring and $x \in R$ be a normal non-nilpotent element ($Rx = xR$ and $x^i \neq 0$ for all $i \geq 1$). Then $S_x := \{x^i \mid i \in \mathbb{N}\}$ is an Ore set.

Lemma 7.5. *We keep the notation of Theorem 1.6. Let R be a ring, $x \in R$ be a normal non-nilpotent element, and $S_x := \{x^i \mid i \in \mathbb{N}\}.$*

1. $\mathfrak{a} = \mathfrak{a}(S_x) = \text{ass}_R(S_x) = \{r \in R \mid x^i r x^j = 0 \text{ for some } i, j \geq 0\}.$
2. Let $\pi : R \rightarrow \overline{R} := R/\mathfrak{a}, r \mapsto \overline{r} = r + \mathfrak{a}.$ Then $\overline{S}_x := \pi(S_x) \in \text{Den}(\overline{R}, 0), S_x \in \mathbb{L}(R, \mathfrak{a}),$ and $S_x^{-1}R \simeq \overline{S}_x^{-1}\overline{R},$ an R -isomorphism.
3. Let \mathfrak{b} be an ideal of R and $\pi^\dagger : R \rightarrow R^\dagger := R/\mathfrak{b}, r \mapsto r^\dagger = r + \mathfrak{b}.$ If $x^\dagger \in \mathcal{C}_{R^\dagger}$ then $\mathfrak{a} \subseteq \mathfrak{b}$ and the map

$$\overline{S}_x^{-1}\overline{R} \rightarrow S_x^{-1}R^\dagger, \overline{x}^{-i}\overline{r} \mapsto x^{\dagger^{-i}}r^\dagger$$

is a ring epimorphism.

4. Let $f : R \rightarrow Q$ be a ring homomorphism such that $f(x) \in Q^\times$ and the ring Q is generated by $f(R)$ and $f(x)^{-1}.$ Then
 - (a) $\mathfrak{a} \subseteq \ker(f)$ and the map

$$\overline{S}_x^{-1}\overline{R} \rightarrow Q, \overline{x}^{-i}\overline{r} \mapsto f(x)^{-i}f(r)$$

is a ring epimorphism with kernel $\overline{S}_x^{-1}(\ker(f)/\mathfrak{a}).$

- (b) Let $\tilde{R} = R/\ker(f)$ and $\tilde{\pi} : R \rightarrow \tilde{R}, r \mapsto r + \ker(f).$ Then $\tilde{S}_x := \tilde{\pi}(S_x) \in \text{Den}(\tilde{R}, 0)$ and $\tilde{S}_x^{-1}\tilde{R} \simeq Q,$ an \tilde{R} -isomorphism.

Proof. 1. Statement 1 follows from Theorem 1.6.(1,2) where $S = S_x.$

2–4. Statements 2–4 follow from Theorem 1.6.(2)–Theorem 1.6.(4), respectively. \square

Localization of a ring at the powers of a normal nonzero idempotent element. If, in addition, the above element x is an idempotent we can find ingredients of Lemma 7.5 explicitly, see the corollary below.

Corollary 7.6. *We keep the notation of Lemma 7.5. Let R be a ring, $x \in R \setminus \{0\}$ be a normal idempotent, and $S_x := \{x^i \mid i \in \mathbb{N}\} = \{1, x\}$.*

1. $R = xRx \times (1-x)R(1-x)$ is a direct product of rings and $\mathfrak{a} = \mathfrak{a}(S_x) = \text{ass}_R(S_x) = (1-x)R(1-x)$.
2. $S_x^{-1}R \simeq R/(1-x)R(1-x) \simeq xRx$.

Proof. 1. The elements of R , $e_1 = x$ and $e_2 = 1 - x$, are orthogonal idempotents ($e_1e_2 = e_2e_1 = 0$) such that $1 = e_1 + e_2$. Hence, the ring R is canonically isomorphic to the matrix ring

$$\begin{pmatrix} R_{11} & R_{12} \\ R_{21} & R_{22} \end{pmatrix} \text{ where } R_{ij} = e_i R e_j \text{ for } i, j = 1, 2,$$

and each element $r \in R$ is identified with the matrix

$$\begin{pmatrix} r_{11} & r_{12} \\ r_{21} & r_{22} \end{pmatrix} \text{ where } r_{ij} = e_i r e_j \text{ for } i, j = 1, 2.$$

Since the element x is a normal idempotent of R , $R_{12} = xR(1-x) = Rx(1-x) = 0$ and $R_{21} = (1-x)Rx = (1-x)xR = 0$. Therefore, the ring $R = xRx \times (1-x)R(1-x)$ is a direct product of rings. Statement 1 follows from Lemma 7.5.(1).

2. Statement 2 follows from statement 1 and Lemma 7.5.(2). \square

References

- [1] V.V. Bavula, The algebra of one-sided inverses of a polynomial algebra, J. Pure Appl. Algebra 214 (2010) 1874–1897, arXiv:0903.0641 [math.RA].
- [2] V.V. Bavula, The algebra of integro-differential operators on an affine line and its modules, J. Pure Appl. Algebra 217 (2013) 495–529, arXiv:1011.2997 [math.RA].
- [3] V.V. Bavula, The largest left quotient ring of a ring, Commun. Algebra 44 (8) (2016) 3219–3261, arXiv:1101.5107 [math.RA].
- [4] V.V. Bavula, New criteria for a ring to have a semisimple left quotient ring, J. Algebra Appl. 14 (6) (2015) 1550090.
- [5] V.V. Bavula, Left localizations of left Artinian rings, J. Algebra Appl. 15 (9) (2016) 1650165, arXiv:1405.0214 [math.RA].
- [6] V.V. Bavula, Criteria for a ring to have a left Noetherian largest left quotient ring, Algebr. Represent. Theory 21 (2) (2018) 359–373.
- [7] V.V. Bavula, Localizations of a ring at localizable sets, their groups of units and saturations, Math. Comput. Sci. 16 (1) (2022), Paper No. 10, 15 pp.
- [8] A.V. Jategaonkar, Localization in Noetherian Rings, London Math. Soc. Lect. Note Ser., vol. 98, Cambridge Univ. Press, 1986.
- [9] N. Jacobson, Structure of Rings, rev. ed., Am. Math. Soc. Colloq., vol. XXXVII, Am. Math. Soc., Providence, 1968.
- [10] J.C. McConnell, J.C. Robson, Noncommutative Noetherian Rings. With the Cooperation of L.W. Small, revised edition, Graduate Studies in Mathematics, vol. 30, American Mathematical Society, Providence, RI, 2001, 636 pp.
- [11] B. Stenström, Rings of Quotients, Springer-Verlag, Berlin, Heidelberg, New York, 1975.