

This is a repository copy of Patterns of Self-Harm Functions among Clinic-Referred Youth: Associations with Treatment Outcomes.

White Rose Research Online URL for this paper: https://eprints.whiterose.ac.uk/id/eprint/230974/

Version: Supplemental Material

Article:

Carrier, C., Herbert, M., Cottrell, D. orcid.org/0000-0001-8674-0955 et al. (2 more authors) (2025) Patterns of Self-Harm Functions among Clinic-Referred Youth: Associations with Treatment Outcomes. Behaviour Research and Therapy, 193. 104843. ISSN: 0005-7967

https://doi.org/10.1016/j.brat.2025.104843

This is an author produced version of an article published in Behaviour Research and Therapy, made available under the terms of the Creative Commons Attribution License (CC-BY), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Supplementary Materials:

Patterns of Self-Harm Functions among Clinic-Referred Youth: Associations with Treatment Outcomes

Christina Carrier, Monique Herbert, David Cottrell, Alexandra Wright-Hughes, & Madison Aitken

Table S1Descriptive Statistics for Baseline Variables by Latent Class

- Variable	M(SD) or %						
	Anger/ Desperation	Anxious/ Depressed	Mixed Intra/ Interpersonal	Intrapersonal/ Internalizing	Few functions		
Age	14.24 (1.54)	14.52 (1.35)	14.31 (1.34)	14.54 (1.33)	14.18 (1.29)		
Sex (Female)	92	92	88	90	87		
Medication use	5	8	6	6	3		
Physical abuse	30	23	33	22	22		
Sexual abuse	20	18	19	15	16		
Suicidal ideation severity	9.99 (9.00)	11.42 (9.58)	11.24 (8.63)	12.14 (9.18)	9.17 (9.04)		

Table S2Descriptives Statistics for Variables at 12-month Follow-Up by Latent Class

	M(SD) or %							
Variable	Anger/ Desperation	Anxious/ Depressed	Mixed Intra/ Interpersonal	Intrapersonal/ Internalizing	Few functions			
Suicidal ideation severity	4.55 (6.98)	6.62 (8.74)	5.08 (6.68)	6.42 (8.19)	3.49 (6.31)			
Self-harm presence	71	80	60	72	62			