



This is a repository copy of *How to get physiologically relevant data with students using Lumbriculus variegatus*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/231151/>

Version: Published Version

Article:

Seeley, A. orcid.org/0000-0001-7085-4296, Corns, L.F. orcid.org/0000-0003-1139-3527, Rouse, J.L. orcid.org/0000-0001-8457-4623 et al. (1 more author) (2025) How to get physiologically relevant data with students using *Lumbriculus variegatus*. *Advances in Physiology Education*. ISSN: 1043-4046

<https://doi.org/10.1152/advan.00084.2025>

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial (CC BY-NC) licence. This licence allows you to remix, tweak, and build upon this work non-commercially, and any new works must also acknowledge the authors and be non-commercial. You don't have to license any derivative works on the same terms. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

1 SOURCEBOOK OF LABORATORY ACTIVITIES IN PHYSIOLOGY

2 RUNNING HEAD: Laboratory practical classes utilising *L. variegatus*

3 How to get physiologically relevant data with students 4 using *Lumbriculus variegatus*

5 Aidan Seeley,¹ Laura F Corns,² James L Rouse,^{3,4} Nicholas S Freestone⁵

6 ¹ Swansea University Medical School, Swansea University, Swansea, United Kingdom

7 ² School of Biosciences, University of Sheffield, Sheffield, United Kingdom

8 ³ School of Biology, University of Leeds, Leeds, United Kingdom

9 ⁴ Applied Insect Sciences, Ripon, United Kingdom

10 ⁵ School of Life Sciences, Pharmacy and Chemistry, Kingston University, London, United

11 Kingdom

12 Correspondence: Aidan Seeley (Aidan.Seeley@swansea.ac.uk)

13

14

ABSTRACT

The decline of *in vivo* teaching in higher education has resulted in graduates lacking essential experimental skills. To address this gap, we present an easy and cost-effective practical class using the emerging invertebrate model organism, *Lumbriculus variegatus*, as an additional *in vivo* model for education. This practical class enables students to observe the effects of pharmacologically active compounds on the stereotypical behaviours of body reversal and helical swimming in *L. variegatus* through tactile stimulation. During this class, students will conduct drug dilution calculations, administer test compounds, and conduct an *in vivo* behavioural experiment. Results from this class demonstrate drug effects *in vivo* and enable students to observe reversible or irreversible behavioural effects, depending on the compound tested. This class demonstrates *L. variegatus* as a model for hands-on *in vivo* teaching, providing students with critical laboratory experience without the need for vertebrate or higher-order mammal models. Furthermore, the approach outlined here is scalable and an adaptable teaching methodology that enhances student engagement with *in vivo* teaching without costly equipment or complex animal husbandry.

NEW & NOTEWORTHY

There are increasing societal expectations that the extent of experimentation using animals is reduced in both research and educational fields. This leaves educators and researchers with a problem, how to foster the development of skills and advances in knowledge in future generations? This paper seeks to address this problem by providing clear examples of an experimental model that can be used to study fundamental physiological and pharmacological processes using the invertebrate organism, *Lumbriculus variegatus*.

Keywords: 3Rs, *Lumbriculus variegatus*, *L. variegatus*, partial replacement, undergraduate experiments

INTRODUCTION

Animal use in education has seen a decline, with an 88% decrease in animals used for education and training since 2001 in the United Kingdom alone (1, 2), which has resulted in students completing their degree without being taught *in vivo* skills (3). Learned societies have highlighted the importance of animal use in education, such as the British Pharmacological Society (4), the Human Anatomy & Physiology Society (5) and the American Physiological Society (6), and experiences of using living systems are impactful learning experiences for students (7, 8). Furthermore, this decline in *in vivo* teaching has produced an *in vivo* skills gap in graduates (9, 10), and so additional teaching methodologies for delivering *in vivo* teaching are required to address this.

Invertebrate models, such as *Drosophila melanogaster* and *Caenorhabditis elegans*, while extensively used in research, have extensive applications in education (11). The ease of rearing invertebrates, the low costs of acquisition and maintenance, and their similarities, or differences, to vertebrate counterparts all contribute to their value as experimental subjects for use in education (12).

The freshwater annelid, *Lumbriculus variegatus*, was first proposed as an inexpensive and accessible organism for education by Charles Drewes in 1996 (13). As invertebrates, *L. variegatus* are not covered under the Animal (Scientific Procedures) Act 1986 in the United Kingdom, or the Animal Welfare Act Regulations or the Public Health Service Policy on Humane Care and Use of Laboratory Animals in the United States, with similar legislation in many other countries allowing widespread adoption of the techniques outlined in this paper. This organism, therefore, offers the opportunity for utilisation within education settings. The use of invertebrate models for practical teaching, which traditionally utilises mammalian species, can serve as a partial replacement in line with the principles of Replacement,

Reduction, & Refinement (3Rs), while still offering students valuable hands-on experience for behavioural studies.

Objectives and Overview

The use of *L. variegatus* within biomedical education has recently had a small resurgence owing to their low cost, ease of use, lack of regulatory restrictions, and utilisation of commonly used equipment and resources (14, 15).

The practical class described here provides the opportunity to observe the effects of drug compounds in the invertebrate model, *L. variegatus*, using methodology initially described by Seeley *et al.* (14). The selection of test compounds can easily be aligned to a range of physiological applications and students performing the outlined practical class will observe the effects of diverse compounds on the behaviours of *L. variegatus*' ability to perform stereotyped behaviours before, during and after exposure to compounds.

The objectives of the activities described in this exercise are (i) conduct drug dilution calculations and prepare drug solutions, (ii) complete a behavioural study examining the effects of drug administration on stereotypical movements of body reversal and helical swimming by *L. variegatus* through collection of quantitative data and (iii) evaluate and analyse data relating to changes in stereotypical movements of *L. variegatus* following the addition and removal of drug solutions using basic statistical techniques.

Background

Lumbriculus variegatus, more commonly known as the blackworm or Californian blackworm, is an asexually reproducing, regenerative detritivore species that inhabits shallow freshwater ponds, lakes and marshes (14, 16). Initially described by Drewes in 1999, *L. variegatus* display easily quantifiable

stereotyped locomotor behaviours of body reversal and helical swimming following tactile stimulation (16) whereby tactile stimulation of the anterior segments of *L. variegatus* evokes body reversal through bending movements that reverse the head and tail positions (Figure 1A and Supplemental Video 1), while stimulation of the posterior region evokes helical swimming movements characterised by rapid helical body bends (Figure 1B and Supplemental Video 2). These behaviours are analogous to the behaviours observed in the extensively characterised *C. elegans* model, which displays quantifiable body bending and locomotor activity (17), but the larger size of *L. variegatus* (50 – 80 mm) compared with *C. elegans* (≈ 1 mm) makes them easier to view as an individual without microscopy techniques.

A 2021 study by Seeley *et al.* (14) described methods for quantifying the effects of exposure to compounds on inhibition of *L. variegatus* stereotypical movements in response to tactile stimulation. These stereotypical movement assays can be utilised to give students hands-on *in vivo* training, using a simple three-point scale (14), enabling students to measure the effects of compounds on reducing tactile stimulation to elicit two different behaviours (body reversal and helical swimming). It should be noted that this assay can only measure a decreased response to tactile stimulation, so it allows students to distinguish *L. variegatus* that perform body reversal or helical swimming movements in response to tactile stimulation (Figure 1, Supplemental Video 1 – 2) from *L. variegatus* that do not (Supplemental Video 3). The relative ease of the tactile stimulation application and the simplicity of the three-point scale minimise the risk of misinterpretation of the movements and limit variation between students conducting the assay.

L. variegatus have been studied extensively for the impact of pollutants (18–23) and are increasingly being used to study diverse compounds including; the selective-serotonin reuptake inhibitor, fluoxetine (24), non-steroid anti-inflammatory drugs such as diclofenac (25), the channel-blockers dantrolene,

quinine and lidocaine (14), anti-histamines such as mepyramine and loratadine (15), alcohol (26), nicotine (27) and cannabidiol (CBD) (28). These studies have examined behavioural effects such as reproduction, feeding, tactile stimulation to elicit stereotyped behaviours of body reversal and helical swimming and unstimulated locomotor activity as experimental endpoints. Additional studies have described the effects of biogenic amines in *L. variegatus* (29), demonstrated the presence of glutaminergic neurons (30) and proposed the presence of a cholinergic system within this organism through quantification of acetylcholine levels and cholinesterase activity (27). These findings demonstrate the applicability of *L. variegatus* for use in physiology education while avoiding the costly and complex issues associated with vertebrate teaching models.

Here, we provide a comprehensive guide for administering the stereotypical movement assay (14) as an *in vivo* behavioural practical class using *L. variegatus* within a teaching environment.

Learning Objectives

After completing this practical class, students should be able to do the following:

1. Prepare, dilute and administer drug compounds to *L. variegatus*
2. Observe the effects of compounds on the body reversal and helical swimming stereotypical behaviours of *L. variegatus* using the three-point scale from Seeley *et al.* (14)
3. Measure and plot relationships between drug concentrations and stereotypical behaviour responses following tactile stimulation.
4. Apply basic statistical analyses to determine the effects of drugs of interest on the ability of *L. variegatus* to perform stereotypical behaviours during and after drug exposure.

Activity Level

The practical class presented here has been used with all levels of undergraduate students and Masters students in the general biomedical science disciplines. This method provides an early opportunity for students to engage with *in vivo* techniques, learn new skills, and gain knowledge of a research environment while generating potentially meaningful data. Furthermore, these techniques can, and have been, easily adapted for use in outreach activities with the general public.

Prerequisite Student Knowledge or Skills

Before students perform this practical class, they should have a good understanding of animal models in research and should be able to use a micropipette and calculate drug dilutions before the class commences, or additional instruction provided before conducting the class. Students would benefit from an understanding of the principles of Replacement, Reduction & Refinement (the 3Rs), and students may also benefit from pre-reading materials on the use of *L. variegatus*, such as the previous studies on using *L. variegatus* in education settings (14, 15).

Time Required

The class described here can be completed within a three-hour period, which includes a brief lecture covering the use of animals in research, the 3Rs and the *L. variegatus* model and its stereotyped behaviours.

MATERIALS AND METHODS

Equipment and Supplies

Where suppliers have been cited below, alternative supplies are routinely available in many countries. Where a supplier has not been listed, these are available from standard scientific supply companies or directly from commercial sources.

- *L. variegatus* can be purchased from Carolina Biological Supply (North Carolina, United States, Cat No. 141720), Blades Biological Ltd (Kent, United Kingdom), or purchased from local aquarium stores before the class.
- Artificial pond water constitutions; sodium chloride (Melford Laboratories Ltd, Ipswich, United Kingdom, Cat No. S23020), potassium chloride (Melford Laboratories Ltd, Ipswich, United Kingdom, Cat No. P41000), calcium nitrate tetrahydrate (Duchefa Biochemie, Haarlem, Netherlands, Cat No. C0505), magnesium sulfate heptahydrate (Duchefa Biochemie, Haarlem, Netherlands, Cat No. M0513) and HEPES (Sigma-Aldrich, Dorset, United Kingdom, Cat No. 54457), deionised water.
- Equipment for short-term culturing (up to four weeks) of *L. variegatus*: small commercial aquaria or buckets, and brown paper towels.
- (Optional) Equipment for longer-term culturing (longer than four weeks) of *L. variegatus*: Commercial aquaria, water filters, air stones, a timed light source, sandy or fine sediment and spirulina powder.
- TetraMin® flakes.
- Micropipettes (10, 20, 200, 1000 µL volumes).
- Standard laboratory plasticware – six-well plates, Pasteur pipettes, pipette tips, 15 mL Falcon centrifuge tubes.
- Marker pens.
- Liquid waste containers.
- Timers.
- Drug compounds for testing; exemplary compounds are shown in Table 1.
- Dimethyl sulfoxide (DMSO) (Sigma-Aldrich, Dorset, United Kingdom, Cat No. 34869) if a drug vehicle is required, as indicated in Table 1.

- 70% ethanol solution.

Animal Subjects

For short-term cultures, which are sustainable for up to four weeks before the practical class, *L. variegatus* cultures should be maintained in artificial pond water composed of 1 mM sodium chloride; 13 μ M potassium chloride; 4 μ M calcium nitrate tetrahydrate; 17 μ M magnesium sulfate heptahydrate; 71 μ M HEPES buffer in UV-treated deionised water (14, 20) at room temperature (18–21°C) in small aquaria or buckets with strips of brown paper towels (31), which mimic leaf litter of natural habitats (32), and fed TetraMin® flakes weekly.

Instructors may, optionally, establish longer-term cultures, which can be maintained for periods of years. Instructors may wish to establish longer-term cultures if there are multiple iterations of the practical class throughout the academic year or to reduce the reordering of cultures ahead of the class. If electing to culture *L. variegatus* in the longer term, cultures can be laboratory-reared in artificial pond water, as above, and should be maintained at room temperature and subject to a 16:8-h light-dark cycle in aquaria with sandy or other fine sediment added to the aquaria. *L. variegatus* cultures should be fed TetraMin® flakes and 10 mg/L spirulina weekly. Continuous water aeration and filtration can be achieved by the use of commercial air stones and aquarium filters, respectively. As per previous studies, it is recommended that populations be maintained for a minimum of 3 months before use to allow for population expansion by asexual reproduction (14, 15, 26–28).

Instructors of this class are responsible for following the regulations for animal research within their home institution.

Instructions

The following are specific instructions for setting up the practical class.

Preparations before student arrival

1. Prepare artificial pond water, as outlined above. This can be prepared as a 100X solution and diluted to 1X ahead of the practical class. Artificial pond water can be maintained at room temperature for up to six months.
2. The instructor should separate *L. variegatus* into individual wells of a six-well plate 18 – 24 hours before the practical class begins to allow for *L. variegatus* acclimatisation, as per previous studies (14, 15, 26–28). Each well should contain approximately 4 mL of artificial pond water. Individual *L. variegatus* should be randomly selected, lacking any obvious morphological defects, and range from 2 - 8 cm in length as per previous studies (14, 15, 26–28). The number of plates required will depend on class size, but we recommend one six-well plate for each pair of students. This can easily be modified depending on class size and student level.
3. Prepare drug solutions to be used during the practical class by dissolving the appropriate mass in artificial pond water. Example drug compounds and their tested concentration ranges, as well as effective concentrations, are shown in Table 1. Drug compounds may require a vehicle, and we have observed that 0.5 % (v/v) DMSO does not affect *L. variegatus* stereotypical behaviours (14, 15, 28). Furthermore, 0.58 % (v/v) ethanol has been shown to have no significant effect on *L. variegatus* (26) and so, depending on the drug selected, ethanol may be used as a vehicle control at ≤ 0.5 % (v/v). If a vehicle is required, then instructors should utilise a suitable vehicle control, such as 0.5 % (v/v) DMSO or ethanol in artificial pond water. If the drug compound is water-soluble and does not require a vehicle, then artificial pond water only should be used as an experimental control.

Instructions for students

1. Following the 18 – 24 hour acclimatisation period, students should remove all artificial pond water from each well of the six-well plate, using a Pasteur pipette, to remove any debris. This should be immediately replaced with 4 mL of fresh artificial pond water.
2. Before administering drug compounds, the baseline ability of the worm to perform stereotypical behaviours should be tested and recorded. This is done by alternately stimulating the anterior or posterior ends of the body with a clean plastic pipette tip or Pasteur pipette to stimulate body reversal (Figure 1A and Supplemental Video 1) and helical swimming (Figure 1B and Supplemental Video 2). Tactile stimulation should be done by gently swiping the tip across the plate to touch the worm. Tactile stimulation should not be done in a stabbing motion, as this can induce autotomy: the sudden separation of *L. variegatus* into two or more segments due to compression (33).
3. Students should administer tactile stimulation to each *L. variegatus* five times on the anterior and posterior end, with a 5 – 10 second time interval between tactile stimuli. Students should score these movements using the three-point scale (14) whereby:
A score of 3 corresponds to full stereotypical movements, as demonstrated in Figure 1 and Supplemental Video 1 – 2.
A score of 2 corresponds to partial stereotypical movement, whereby movement is less than Figure 1 and Supplemental Video 1 – 2, but more movement than Supplemental Video 3.
A score of 1 for no stereotypical movement following tactile stimulation, as shown in Supplemental Video 3.
Scores should be recorded for each stimulation at both the anterior and posterior tactile stimulation of *L. variegatus* and recorded in Supplemental Table 1.

4. Once students have completed their baseline scores, the artificial pond water should then be removed and immediately replaced with 4 mL of a vehicle or experimental control or test drug solutions.
5. After a 10-minute incubation with the drug solution and vehicle or experimental control, *L. variegatus* can be tested again using the same procedure to evaluate the effects of drug exposure on stereotypical movements.

Students should administer tactile stimuli beginning with the vehicle or experimental control and then progressing from the lowest to the highest concentrations. After which, students should discard the pipette tip before starting the next round of tactile stimuli to prevent cross-contamination with higher concentrations of the test compound.

The class can be terminated after evaluating the effects of drug exposure on *L. variegatus*' response to tactile stimuli should instructors wish to reduce the overall time of the practical class, or students can evaluate the effects following the removal of the drug.
6. To evaluate the effects following the removal of the drug compounds, the drug solutions and vehicle or experimental control are aspirated from the well and disposed of in suitable liquid waste containers.
7. To remove any latent drug and vehicle or experimental control residue, 4 mL of fresh pond water should be added and immediately aspirated and then replaced with 4 mL of fresh artificial pond water.
8. The ability of *L. variegatus* to perform stereotypical movements can then be retested after 10 minutes to measure reversibility of any observed effects.
9. Depending on time constraints, *L. variegatus* can be retested 24 hours after exposure to examine the effects on stereotypical movements and to measure toxicity or lethality. Toxicity in

L. variegatus is defined as tissue pallor or partial tissue degeneration, with lethality resulting in complete tissue degeneration.

10. Upon completion of the practical class, *L. variegatus* should not be returned to cultures due to the unknown long-term effects of exposure to test compounds. At experimental endpoints, *L. variegatus* should be euthanised by rapid submersion in 70% ethanol.

Quantification and Statistical Analyses

Students should be encouraged to plot the effects of the drug compound on the ability of tactile stimulation to elicit stereotypical movements. Students should calculate the average score for each *L. variegatus* at the different drug concentrations and time points tested for both movements, and data can be normalised to baseline ability by using the calculation:

$$\text{Ability to Complete Movement (Relative to Baseline)} = \frac{\text{Average movement score of } L. \text{ variegatus at the tested concentration or time point}}{\text{Average movement score of } L. \text{ variegatus at baseline.}}$$

Using this calculation, students should have a baseline score equal to 1 and if the drug has affected the ability of tactile stimulation to elicit stereotypical movements in *L. variegatus* then a score of <1 will be calculated. Student data can be visualised for each student or collated to show the impact of repeated experiments. Plotting the collated data will emphasise to students the requirement for technical and experimental replicates.

The type of statistical analyses applied will vary depending on student level and access to statistical software. Students could compare stereotypical movements of *L. variegatus* during drug exposure conditions with stereotypical movements taken at baseline using paired nonparametric two-tailed *t*-

tests, in line with previous publications (14, 15, 26–28). For analysis of stereotypical movements after the removal of the drug compound, students may analyse these using a two-way ANOVA with Dunnett's post-test to compare baseline stereotypical movements with stereotypical movements at both the 10-minute and 24-hour recovery time points, as in previous studies utilising this method (14, 15, 26–28).

Troubleshooting

As *L. variegatus* undergo asexual reproduction they may separate during the acclimatisation period, with the newly separated *L. variegatus* having reduced tactile responses. Additionally, a small number of *L. variegatus* may expire during the acclimatisation process. It is recommended that instructors have a small number of additional *L. variegatus* for the class which have been acclimatised to replace any *L. variegatus* which have undergone asexual separation into two or more *L. variegatus* or those that may have expired during the acclimatisation stage.

L. variegatus will show reduced stereotypical movements if continuously stimulated and may stop responding if repeatedly stimulated without an interval. As outlined above, each stimulus should be administered after a 5 – 10 second time lapse between tactile stimuli. Instructors should suggest that students stimulate each worm individually at the anterior, then stimulate each worm individually at the posterior, and repeat for five stimulation events at the anterior and posterior to prevent a diminished response of *L. variegatus*.

Additionally, students may report that their worm has suddenly split into two worms. While mechanical pressure has been shown to induce autotomy of *L. variegatus* (33), the sudden separation of the worm in these classes is normally due to excessive force during stimulation. The method for tactile stimulation of the stereotypical behaviours is shown in Supplemental Video 1 - 3.

325

326 New test compounds can be selected to align with course curricula, but some compounds may not be
327 soluble in artificial pond water nor produce an effect in *L. variegatus*. Instructors are recommended to
328 check the solubility of test compounds before the practical class using either DMSO or ethanol as a
329 vehicle, not exceeding 0.5% (v/v). Furthermore, the effects of some compounds, such as mecamylamine,
330 may not appear after a 10-minute exposure to the compound but effects may be observed at recovery
331 time points (10 minutes or 24 hours after exposure and incubation in artificial pond water only) and so it
332 is recommended that recovery timepoints be completed to observe any delayed effects of drug
333 exposure. Instructors may wish to include a compound which has been shown to inhibit stereotypical
334 movement in response to tactile stimulation of *L. variegatus* as a positive control. Compounds which
335 produce a known effect are shown in Table 1.

336

337 Safety Considerations

338 When preparing the artificial pond water, Instructors should wear appropriate personal protection
339 equipment, including gloves, goggles, and a mask, as calcium nitrate tetrahydrate is harmful if
340 swallowed and causes serious eye damage.

341 Instructors should check with their institutional review boards regarding the use of proposed test
342 compounds with students to ensure compliance with regional and institutional regulations. Each test
343 compound will have its specific safety considerations, but at a minimum, standard laboratory personal
344 protective equipment should be used throughout the practical class. Instructors are advised to consult
345 the manufacturer's safety data sheet on the safe disposal of test compounds. Furthermore, instructors
346 are responsible for ensuring that appropriate health and safety documentation is completed for any
347 drug compound selected for investigation and whether their institute has any ethical requirements for
348 the use of *L. variegatus*.

L. variegatus should not be released into the environment to prevent environmental and ecological contamination and should be euthanised in 70% ethanol and disposed of by appropriate disposal routes.

RESULTS AND DISCUSSION

Expected Results

Depending on the test compound selected, students will observe a dose-dependent inhibition of body reversal and/or helical swimming stereotypical movements. Tested compounds with known effects on the response to tactile stimulation to elicit stereotypical movements in *L. variegatus* are shown in Table 1. Exemplary data from using this methodology is shown in Figure 2, which demonstrates the effects of acute exposure on the ability of tactile stimulation to elicit body reversal (Figure 2A) and helical swimming (Figure 2B). The ability of *L. variegatus* to perform these movements 10 minutes and 24 hours after exposure is also shown (Figure 2C-D), demonstrating the irreversible effects of this test compound. Compounds which affect stereotypical movements reversibly include lidocaine, mepyramine, loratadine and ethanol (14, 15, 26), while irreversible inhibition occurs when exposed to quinine, histamine or nicotine (14, 15, 27).

If instructors are selecting a novel drug compound to test in *L. variegatus*, exposure may result in (i) no observable effect on *L. variegatus*' ability to perform stereotypical movements following tactile stimulation, (ii) partial inhibition of stereotypical movements whereby *L. variegatus* responds to tactile stimulation but movements may be slowed or incomplete, (iii) complete inhibition of stereotypical movements whereby *L. variegatus* has no response to tactile stimulation or (iv) rapid toxicity characterised by rapid writhing of *L. variegatus* followed by a period of stasis and potential decomposition observed up to 24 hours after exposure.

Misconceptions

A common point of confusion is that the inhibition of stereotypical movements in *L. variegatus* following tactile stimulation is interpreted as lethality following drug exposure or recovery time points. However, the toxicity of drug compounds is confirmed by visible partial tissue degeneration and/or pallor and lethality as complete tissue degeneration or whole-organism tissue pallor, which is observable 24 hours post-drug exposure. The inability of *L. variegatus* to respond to tactile stimulation at any time point is not indicative of organism expiry, as this may be a long-term inhibitory effect-of drug exposure on stereotypical movement.

Evaluation of Student Work

Experiments described here using the *L. variegatus* model have been used from the first year of undergraduate studies through to more advanced levels of study. Students may present their results in several different ways, such as a laboratory report, or we have had students use their findings to produce a scientific poster and present their poster with a two-minute summary. In this format, students gain valuable experience in presenting their findings clearly and concisely, while gaining skills in science communication. Anecdotally, students have reported that presenting their own data provides them with increased confidence in their presentation skills. For students who prefer not to work with living animals, data sets from the class can be shared, or exemplary data can be obtained from the corresponding author.

Inquiry Applications

Commonly, students will enquire what it is they *should* see rather than reporting what they see. To conduct these experiments ethically and ensure we are continually adding to the knowledge base on *L. variegatus* responses, we often test new compounds rather than ones with known effects. In this manner, there is no desired result, and students are encouraged to report on what they see, not what

they think they *should* see. To remove this confusion, students are blinded to compounds to prevent any bias and relieve student pressure on aiming to get the 'required' result.

Whilst the methods described above most obviously lend themselves to a Facilitated Inquiry approach, they have also more recently been used as a vehicle for co-creation of the curriculum (15) whereby students at final year undergraduate level, as well as Master's level, are involved in the selection of compounds, the laboratory set up, class delivery, and analysis of collated data, with this data directly feeding into their projects. This aligns with an Open Inquiry approach where students themselves generate hypotheses and then design the experiments to test the hypotheses. This ensures that the data generated from these organisms has research applications and is a more ethical use of animals for education and training purposes. Moreover, the methodologies outlined here are well-suited for independent student projects and can easily be tailored to systems/drugs of interest, highlighting their application in physiology education.

Wider Applications

This practical class can be used to explore the behavioural effects of diverse pharmacologically active compounds following short-term(10 minutes) exposure, as outlined here, the effects of repeated exposures, or longer-term exposures, up to 24 hours, as found within the literature (27, 28).

As a problem-based learning exercise, students could be provided with compounds known to produce effects, such as GABA (Figure 2), and drugs which do not produce an effect, such as the GABA agonist, baclofen (Table 1), and asked to identify the compounds. Alternatively, students could be tasked with identifying a parental compound and inactive metabolites, which have diminished effects on behaviour (28). Further, students could investigate agonist and antagonist pairings and compare the effects in

isolation, and then in combination, which has previously been utilised for co-created practical classes using *L. variegatus*, examining the effects of histamine and the antihistamine drugs loratadine and mepyramine (15).

Additional experiments using *L. variegatus* could run across multiple practical classes, or as student projects, to enable students to investigate single compounds in depth and produce a more comprehensive report of their findings. Indeed, multiple studies have quantified unstimulated locomotor activity of *L. variegatus* using low-cost equipment and free-to-use software (14, 26–28) or investigated effects of compounds on dorsal blood vessel pulse rates using a variety of techniques (23, 28, 29, 31, 34). These manipulations or additions open up a range of new experimental possibilities for the utilisation of *L. variegatus* within practical classes.

ADDITIONAL RESOURCES

For additional information demonstrating the effect of pharmacologically active compounds on *L. variegatus*, please see refs 14, 15, 26–28. Additional information and support for implementing this resource can be found at www.SWIRLswansea.com.

SUPPLEMENTAL MATERIAL

Supplemental Video 1. DOI: <https://doi.org/10.6084/m9.figshare.28727045.v3>

Supplemental Video 2. DOI: <https://doi.org/10.6084/m9.figshare.28727051.v2>

Supplemental Video 3. DOI: <https://doi.org/10.6084/m9.figshare.28727042.v3>

Supplemental Table 1. DOI: <https://doi.org/10.6084/m9.figshare.28727048.v2>

ACKNOWLEDGMENTS

Thank you to all students who have participated in laboratory activities coordinated by the Swansea Worm Integrative Research Laboratory (SWIRL) and contributed to the development of this class and our knowledge of *L. variegatus* responses to drug compounds.

GRANTS

This work is supported by the British Pharmacological Society's Education Grant 2019 and 2021.

DISCLOSURES

The authors declare that they have no conflict of interest.

REFERENCES

1. Statistics of Scientific Procedures on Living Animals Great Britain 2001 [Online]. *GOV.UK*: 2002.
<https://www.gov.uk/government/statistics/statistics-of-scientific-procedures-on-living-animals-great-britain-2001> [29 Jan. 2025].
2. Statistics of scientific procedures on living animals, Great Britain: 2023 [Online]. *GOV.UK*: 2024.
<https://www.gov.uk/government/statistics/statistics-of-scientific-procedures-on-living-animals-great-britain-2023> [29 Jan. 2025].
3. **British Pharmacological Society**. A survey of integrative physiology/pharmacology teaching undertaken by the BPS and the Physiological Society. *pA2 Online* 3: 10–11, 2004.
4. **British Pharmacological Society**. Curriculum for the use of research animals [Online]. 2019.
<https://www.bps.ac.uk/education-engagement/research-animals/curriculum-for-the-use-of-research-animals> [29 May 2020].
5. **Human Anatomy and Physiology Society**. Animal Use Position Statement [Online]. 2016.
<https://www.hapsweb.org/about-us/position-statements/> [18 July 2025].
6. **The American Physiological Society**. APS Position Statement on the Use of Animals in Teaching. *The Physiologist* 48: 206–208, 2005.
7. **Moore WA, Noonan AC**. Using live tissue laboratories to promote clinical reasoning in doctor of physical therapy students. *Advances in Physiology Education* 34: 54–58, 2010. doi: 10.1152/advan.00059.2009.
8. **Ra'anan AW**. The evolving role of animal laboratories in physiology instruction. *Advances in Physiology Education* 29: 144–150, 2005. doi: 10.1152/advan.00017.2005.

- 478 9. **The Association of the British Pharmaceutical Industry.** Skills Needs for Biomedical Research
479 [Online]. <https://www.abpi.org.uk/publications/skills-biomedical-research/>.
- 480 10. **The Association of the British Pharmaceutical Industry.** Bridging the skills gap in the
481 biopharmaceutical industry [Online]. [https://www.abpi.org.uk/publications/bridging-the-skills-](https://www.abpi.org.uk/publications/bridging-the-skills-gap-in-the-biopharmaceutical-industry/)
482 [gap-in-the-biopharmaceutical-industry/](https://www.abpi.org.uk/publications/bridging-the-skills-gap-in-the-biopharmaceutical-industry/).
- 483 11. **Wilson-Sanders SE.** Invertebrate models for biomedical research, testing, and education. *ILAR J* 52:
484 126–152, 2011. doi: 10.1093/ilar.52.2.126.
- 485 12. **Deyrup-Olsen I, Linder TM.** Use of invertebrate animals to teach physiological principles. *Am J*
486 *Physiol* 260: S22–24, 1991. doi: 10.1152/advances.1991.260.6.S22.
- 487 13. **Drewes CD.** Heads or Tails? Patterns of segmental regeneration in a freshwater oligochaete
488 [Online]. In: *Tested studies for laboratory teaching*, p. 23–35.
489 <https://www.ableweb.org/biologylabs/wp-content/uploads/volumes/vol-17/2-drewes.pdf>.
- 490 14. **Seeley A, Bellamy C, Davies NA, Wallace MJ.** *Lumbriculus variegatus*: A novel organism for *in vivo*
491 pharmacology education. *Pharmacol Res Perspect* 9: e00853, 2021. doi: 10.1002/prp2.853.
- 492 15. **Carriere JJ, Davies NA, Cunningham MR, Wallace MJ, Seeley A.** Co-created *in vivo* pharmacology
493 practical classes using the novel organism *Lumbriculus variegatus*. *Pharmacol Res Perspect* 11:
494 e01158, 2023. doi: 10.1002/prp2.1158.
- 495 16. **Drewes CD.** Helical swimming and body reversal behaviors in *Lumbriculus variegatus* (Annelida:
496 Clitellata: Lumbriculidae). *Hydrobiologia* 406: 263–269, 1999. doi: 10.1023/A:1003784100638.

- 497 17. **Parida L.** The locomotory characteristics of *Caenorhabditis elegans* in various external
498 environments: A review. *Applied Animal Behaviour Science* 255: 105741, 2022. doi:
499 10.1016/j.applanim.2022.105741.
- 500 18. **Aikins DM, Mehler WT, Veilleux HD, Zhang Y, Goss GG.** The Acute and Chronic Effects of a
501 Sediment-Bound Synthetic Musk, Galaxolide, on *Hyaella azteca*, *Chironomus dilutus*, and
502 *Lumbriculus variegatus*. *Arch Environ Contam Toxicol* 84: 227–236, 2023. doi: 10.1007/s00244-023-
503 00978-3.
- 504 19. **Colombo V, Pettigrove VJ, Hoffmann AA, Golding LA.** Effects of *Lumbriculus variegatus* (Annelida,
505 Oligochaete) bioturbation on zinc sediment chemistry and toxicity to the epi-benthic invertebrate
506 *Chironomus tepperi* (Diptera: Chironomidae). *Environ Pollut* 216: 198–207, 2016. doi:
507 10.1016/j.envpol.2016.05.063.
- 508 20. **O’Gara BA, Bohannon VK, Teague MW, Smeaton MB.** Copper-induced changes in locomotor
509 behaviors and neuronal physiology of the freshwater oligochaete, *Lumbriculus variegatus*. *Aquat*
510 *Toxicol* 69: 51–66, 2004. doi: 10.1016/j.aquatox.2004.04.006.
- 511 21. **Sardo AM, Soares AMVM.** Assessment of the effects of the pesticide imidacloprid on the
512 behaviour of the aquatic oligochaete *Lumbriculus variegatus*. *Arch Environ Contam Toxicol* 58:
513 648–656, 2010. doi: 10.1007/s00244-010-9470-0.
- 514 22. **Silva CJ, Patrício Silva AL, Campos D, Soares AM, Pestana JL, Gravato C.** *Lumbriculus variegatus*
515 (oligochaeta) exposed to polyethylene microplastics: biochemical, physiological and reproductive
516 responses. *Ecotoxicol Environ Saf* 207: 111375, 2021. doi: 10.1016/j.ecoenv.2020.111375.

- 517 23. **Vought V, Wang H-S.** Impact of common environmental chemicals bisphenol A and bisphenol S on
518 the physiology of *Lumbriculus variegatus*. *Environ Toxicol Pharmacol* 60: 225–229, 2018. doi:
519 10.1016/j.etap.2018.05.003.
- 520 24. **Nentwig G.** Effects of pharmaceuticals on aquatic invertebrates. Part II: the antidepressant drug
521 fluoxetine. *Arch Environ Contam Toxicol* 52: 163–170, 2007. doi: 10.1007/s00244-005-7190-7.
- 522 25. **Karlsson MV, Marshall S, Gouin T, Boxall ABA.** Routes of uptake of diclofenac, fluoxetine, and
523 triclosan into sediment-dwelling worms. *Environ Toxicol Chem* 35: 836–842, 2016. doi:
524 10.1002/etc.3020.
- 525 26. **Seeley A, Mahmood R, Bellamy C, Roome EG, Williams BS, Davies NA, Wallace MJ.**
526 Concentration- and time-dependent behavioural effects of ethanol on *Lumbriculus variegatus*.
527 *Genes Brain Behav* 23: e70006, 2024. doi: 10.1111/gbb.70006.
- 528 27. **Davies NA, Carriere JJ, Gopal A, Rajan A, Wallace MJ, Seeley A.** The inhibitory effect of nicotine on
529 *Lumbriculus variegatus* stereotypical movements and locomotor activity. *Pharmacol Biochem*
530 *Behav* 247: 173953, 2025. doi: 10.1016/j.pbb.2024.173953.
- 531 28. **Williams BS, Jomy G, Flanagan M, Carriere JJ, Labdon GE, Hawkes GS, McRobbie-Aston J, Wallace**
532 **MJ, Price CL, Davies NA, Seeley A.** The behavioral, physiological, and biochemical responses of
533 *Lumbriculus variegatus* exposed to cannabidiol and its metabolites. *Environ Toxicol Chem* 44:
534 1297–1309, 2025. doi: 10.1093/etjnl/vgaf048
- 535 29. **Crisp KM, Grupe RE, Lobsang TT, Yang X.** Biogenic amines modulate pulse rate in the dorsal blood
536 vessel of *Lumbriculus variegatus*. *Comp Biochem Physiol C Toxicol Pharmacol* 151: 467–472, 2010.
537 doi: 10.1016/j.cbpc.2010.02.003.

- 538 30. **Lybrand ZR, Martinez-Acosta VG, Zoran MJ.** Coupled sensory interneurons mediate escape neural
539 circuit processing in an aquatic annelid worm, *Lumbriculus variegatus*. *J Comp Neurol* 528: 468–
540 480, 2020. doi: 10.1002/cne.24769.
- 541 31. **Bohrer KE.** Effects of Drugs on Pulsation Rate of *Lumbriculus variegatus* (Blackworms). *Association*
542 *for Biology Laboratory Education (ABLE) 2005 Proceedings* 27: 127–146, 2006.
- 543 32. **Martinez Acosta VG, Arellano-Carbajal F, Gillen K, Tweeten KA, Zattara EE.** It Cuts Both Ways: An
544 Annelid Model System for the Study of Regeneration in the Laboratory and in the Classroom. *Front*
545 *cell dev biol* 9:780422, 2021. doi: 10.3389/fcell.2021.780422.
- 546 33. **Lesiuk NM, Drewes CD.** Autotomy reflex in a freshwater oligochaete, *Lumbriculus variegatus*
547 (Clitellata: Lumbriculidae). *Hydrobiologia* 406: 253–261, 1999. doi: 10.1023/A:1003756722019.
- 548 34. **Lesiuk NM, Drewes CD.** Blackworms, Blood Vessel Pulsations & Drug Effects. *Am Biol Teach* 61:
549 48–53, 1999. doi: 10.2307/4450609.

550

551

FIGURE LEGENDS

Figure 1. Stereotypical movements of *L. variegatus* following tactile stimulation. Images show the stereotypical movement (A) body reversal, whereby bending movements reverse the head and tail positions, and (B) helical swimming, characterised by rapid helical body bends, of *L. variegatus* following tactile stimulation with a pipette tip on the anterior and posterior regions, respectively. Images show the stereotypical movements over a 2 – 3 second period following stimulation.

Figure 2. The effects of gamma-aminobutyric acid (GABA) on *Lumbriculus variegatus* behaviour. *L. variegatus* were exposed to GABA (0 – 100 mM) and tested for the ability of tactile stimulation to elicit (A) body reversal or (B) helical swimming. GABA was then removed, and the ability of *L. variegatus* to perform (C) body reversal or (D) helical swimming was tested 10 minutes after removal (Recovery [10 mins]) and 24 hours after removal (Recovery [24h]). Data are expressed as a ratio of the movement score after exposure relative to the movement score at baseline. Error bars represent the standard error of the mean, $n=8$ with a single *L. variegatus* exposed to each concentration per experimental repeat * $p<.05$, ** $p<.01$, where * refers to statistical significance between baseline and GABA exposure or statistical significance between baseline and Recovery (10 min), no statistical significance between baseline and Recovery (24 h) was observed.

568 **TABLES**

569 **Table 1. Test compounds known to affect *L. variegatus*' response to tactile stimulation.** Suggested
570 stock and working concentrations are provided, along with the known mammalian molecular target and
571 known effect in *L. variegatus*. Test compounds are soluble in artificial pond water up to the Stock
572 Concentration, unless denoted by #, in which the compound requires to be initially dissolved in 100%
573 DMSO and then diluted to 0.5% (v/v) DMSO in artificial pond water before use. * is shown in Figure 2. *
574 See Ref 14 ** See Ref 15, *** See Ref 26, **** See Ref 27, ***** See Ref 28.

575
576


Test compound	Known target in mammals	Stock concentration	Working concentration	Concentration that decreases stereotypical movements after 10 minutes
Acetylcholine	Non-selective acetylcholine receptor agonist	100 mM	100 mM	≥25 mM
Baclofen	GABA _B agonist	20 mM	20 mM	No effect
Bicuculline #	GABA _A antagonist	50 mM	250 µM	≥50 µM
Cannabidiol (CBD) #	Various	4 mM	20 µM	≥5 µM *****
Dantrolene #	Ryanodine receptor (RyR1) antagonist	10 mM	50 µM	≥25 µM *
Donepezil	Acetylcholinesterase inhibitor	100 µM	100 µM	≥50 µM
Dopamine	Non-selective dopamine receptor	100 mM	100 mM	≥10 mM

	agonist			
Ethanol	Various	500 mM	500 mM	≥ 250 mM ***
GABA ⁺	Non-selective GABA receptor agonist	100 mM	100 mM	≥ 10 mM
Glycine	Non-selective glycine receptor agonist	500 mM	500 mM	500 mM
Histamine	Non-selective histamine receptor agonist	10 mM	10 mM	≥ 1 mM **
Haloperidol [#]	Dopamine receptor (D ₂) antagonist	20 mM	100 μM	≥ 50 μM
Lidocaine	Voltage-gated sodium channel inhibitor	1 mM	1 mM	≥ 0.5 mM *
Loratadine [#]	Histamine receptor 1 (H1) antagonist	12 mM	60 μM	≥ 6 μM **
Mecamylamine	Non-selective of the nicotinic acetylcholine receptor antagonist	100 μM	100 μM	≥ 50 μM
Mepyramine	Histamine receptor 1 (H1) antagonist	10 mM	10 mM	≥ 5mM **
Naproxen	COX Inhibitor	25 mM	125 μM	≥ 50 μM
Nicotine	Nicotinic acetylcholine receptor agonist	1 mM	1 mM	≥ 0.1 mM *****

Quinine #	A nonselective sodium and potassium channel blocker	200 mM	1 mM	$\geq 0.1 \text{ mM}^*$
Serotonin	Non-selective 5-HT receptor agonist	2 mM	2 mM	$\geq 0.25 \text{ mM}$
Tubocurarine	Non-selective of the nicotinic acetylcholine receptor antagonist	100 μM	100 μM	$\geq 25 \text{ } \mu\text{M}$

577

A



B



