



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/231268/>

Version: Published Version

Article:

Ruxton, C.H.S., Bateson, M., Garner, N. et al. (2025) Inequity and food insecurity and effects on obesity and associated co-morbidities: meeting report and research priorities. *British Journal of Nutrition*, 134 (2). pp. 169-176. ISSN: 0007-1145

<https://doi.org/10.1017/s0007114525103930>

© The Author(s), 2025. Published by Cambridge University Press on behalf of The Nutrition Society. Reproduced in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Workshop Report

Cite this article: Ruxton CHS, Bateson M, Garner N, Harvey K, Lambie-Mumford H, Levy L, McDonald A, Parretti HM, Walker AW, Thies F, Minihane AM, Fairweather-Tait SJ, Woodside JV, Lovegrove JA, and Corfe BM (2025). Inequity and food insecurity and effects on obesity and associated co-morbidities: meeting report and research priorities. *British Journal of Nutrition*, page 1 of 8. doi: [10.1017/S0007114525103930](https://doi.org/10.1017/S0007114525103930)

Received: 12 December 2024

Revised: 3 July 2025

Accepted: 8 July 2025

Keywords:

Obesity; Food insecurity; Metabolism; Research recommendations

Abbreviations:

FIS, food insecurity; HFSS, foods/drinks high in fat sugar and salt

Corresponding author:

Bernard Corfe; Email: bernard.corfe@newcastle.ac.uk

Inequity and food insecurity and effects on obesity and associated co-morbidities: meeting report and research priorities

Carrie H. S. Ruxton¹ , Melissa Bateson², Nikki Garner³, Kate Harvey⁴, Hannah Lambie-Mumford⁵, Louis Levy⁶, Alana McDonald⁷, H. M. Parretti³, Alan W. Walker⁸, Frank Thies⁸, Anne Marie Minihane³ , Susan J. Fairweather-Tait³ , Jayne V. Woodside⁹ , Julie A. Lovegrove¹⁰ and Bernard M. Corfe¹¹ 

¹Nutrition Communications, Cupar, UK; ²Biosciences Institute/Centre for Behaviour and Evolution, Newcastle University, Newcastle upon Tyne, UK; ³Norwich Medical School and Norwich Institute of Healthy Ageing (NIHA), University of East Anglia, Norwich, UK; ⁴Department of Psychology, The University of Reading, PO Box 217, Reading, UK; ⁵Department of Politics, University of Sheffield, Sheffield, UK; ⁶Office of Health Improvement and disparities, Department of Health and Social Care, UK; Faculty of Health, Medicine and Society, University of Chester, Chester, UK; ⁷Food Standards Scotland, Aberdeen, UK; ⁸Rowett Institute, University of Aberdeen, Aberdeen, UK; ⁹Centre for Public Health, Queen's University Belfast, Belfast, UK; ¹⁰Hugh Sinclair Unit of Human Nutrition and the Institute for Cardiovascular and Metabolic Research, Department of Food and Nutritional Sciences, University of Reading, Reading, UK and ¹¹Human Nutrition and Exercise Research Centre, Population and Health Science Institute, Newcastle University, Newcastle upon Tyne, UK

Abstract

A Rank Forum was convened to discuss the evidence around food insecurity (FIS), its impact on health, and interventions which could make a difference both at individual and societal level, with a focus on the UK. This paper summarises the proceedings and recommendations. Speakers highlighted the growing issue of FIS due to current economic and social pressures. The health implications of FIS vary geographically since food insecure women in higher income regions tend to be living with overweight or obesity, in contrast to those living in low-to-middle-income countries. This paradox could be due to stress and/or metabolic or behavioural responses to an unpredictable food supply. The gut microbiota may play a role given the negative effects of low fibre diets on bacterial diversity. Solutions to FIS involve individual behavioural change, targeted services and societal/policy change. Obesity-related services are currently difficult to access. Whilst poverty is the root cause of FIS, it cannot be solved solely by making healthy food cheaper due to ingrained beliefs, attitudes and behaviours in target groups. Person-centred models, such as Capability-Opportunity-Motivation Behavioural Change Techniques and *Elicit-Provide-Elicit* communication techniques, are recommended. Societal change or improved resilience through psychological support may be more equitable ways to address FIS. They can combine with fiscal or food environment policies to shift purchasing towards healthier foods. Policy implementation can be slow to enact due to the need for strong evidence, consultation and political will. Eradicating FIS must involve co-creation of interventions and policies to ensure a consensus on solutions.

Anderson⁽¹⁾ defines food insecurity (FIS) as 'not having access by all people at all times to enough food for an active, healthy life' and proposes this involves both the *availability* of nutritionally adequate and safe foods as well as socially acceptable means to *access* these, which precludes stealing, use of food banks or scavenging. Whilst FIS is intimately intertwined with poverty, low income is not the sole factor determining FIS. Other drivers and barriers include, for example, the psychological impact of awareness of supply issues including environmental or geopolitical stressors, which include impacts such as the war in Ukraine, Brexit and the cost-of-living crisis following the COVID-19 pandemic of 2020–2022. Since FIS is an important risk factor for obesity and certain non-communicable diseases, this RANK symposium sought to draw together experts from a range of backgrounds to explore factors driving FIS and the impact of food poverty, from basic molecular metabolic mechanisms through to impacts on populations and healthcare provision. The speaker and topic list for the symposium is shown in [Table 1](#).

Food inequity: understanding the national landscape

Household FIS is a concept to describe 'uncertainty about future food availability and access, insufficiency in the amount and kind of food required for a healthy lifestyle, or the need to use socially unacceptable ways to acquire food'⁽¹⁾. The FAO defines four dimensions of food

Table 1. Speakers and talk titles at the rank symposium on ‘inequity and food insecurity: effects on obesity and associated co-morbidities’

Inequity and food insecurity (setting the scene)	
Hannah Lambie-Mumford (Sheffield)	Food inequity: understanding the national landscape.
Metabolic impacts of food insecurity	
Melissa Bateson (Newcastle)	The metabolic impacts of food insecurity.
Alan Walker (Aberdeen)	Food insecurity and the microbiome.
Design considerations and behaviour change	
Kate Harvey (Reading)	Achieving behaviour change through consumer acceptance.
Nikki Garner (UEA)	Supporting behaviour change in practice.
Helen Parretti (UEA)	Obesity and weight management in primary care.
Achieving policy change	
Alana McDonald (Food Standards Scotland)	Improving health through diet: the Scottish perspective.
Louis Levy (Independent Expert)	Shifting nutrition policy to improve health: the evidence needs of policy makers.

security: physical availability of food, economic or physical access to food, food utilisation by people and the stability of the first three dimensions⁽²⁾.

The study of household FIS in the UK has evolved considerably over the last two decades, from relatively little work before the rise of food banks in the 2010s to a vibrant multidisciplinary field⁽³⁾. Government monitoring has also evolved over this time, and now central and devolved governments measure household food security routinely. Regular surveys are conducted by different organisations. For example, the Food Standards Agency⁽⁴⁾ tracks experiences over the past 12 months while the Department of Work and Pensions tracks over a 30-day recall period (Bates, 2017). Questions around FIS and related indicators are included in the Scottish Government’s Scottish Health Survey⁽⁵⁾ and the Welsh Government’s National Survey for Wales⁽⁶⁾. The most recent results from the Food Standards Agency’s Food and You survey⁽⁷⁾ found that 30 % of consumers reported having marginal, low or very low food security over a 12-month period. The Department of Work and Pensions’ Family Resources Survey⁽⁸⁾, using a survey window of 30 d, reported a lower proportion of consumers in this situation (12 %). The Food Foundation’s FIS tracker publishes data multiple times a year using a 30-day recall period⁽⁹⁾. Focussing on the June 2023 tracker, the proportion of households reporting moderate or severe FIS was 17 % compared with the pre-pandemic baseline of 16 %. Households more likely to report FIS are those with children (24 %), receiving benefits (49 %) or including an adult with disability (42 %). Other research has identified further characteristics of those at higher risk of FIS⁽¹⁰⁾ (Table 2).

There are now a range of interventions in place that are designed to promote better access to food from the state, charities and local communities. National government interventions include Healthy Start vouchers⁽¹¹⁾ (run through the NHS in England, Northern Ireland and Wales), Best Start Grant and Best Start Foods (run by the Scottish Government⁽¹²⁾), free school meal provision and Universal Free School Meals. Local Welfare Assistance payments (provided through local authorities in

Table 2. Characteristics most associated with self-reported FIS at different severity

Any food insecurity	Severe food insecurity
Younger age (16–34 years)	Unemployment
Non-white identity	Disability or life-limiting health problems
Children in the household	Income in the lowest quartile
Lower educational level	
Unemployment	
Disability or life-limiting health problems	
Income in the lowest quartile	

FIS, food insecurity.

England), Crisis Grants through the Scottish Welfare Fund (Scottish Government), Discretionary Support (Northern Ireland) and Emergency Assistance Payments (Welsh Government) are short-term financial payments designed to support people in financial crisis meet basic needs. Charitable and community provision has grown significantly in recent decades including food banks, social supermarkets and pantries, sometimes with the support for local and national businesses⁽³⁾. Increasingly, research is considering ‘what works’ to address FIS including projects funded through the UKRI Transforming UK Food Systems Strategic Priorities Fund⁽¹³⁾ and the Salient Food Systems trials⁽¹⁴⁾. Evaluative research from these teams and others will be vital to inform future policy and practice across the UK Policy.

The metabolic and biological impacts of food insecurity

A study, albeit from the USA, found that women experiencing FIS were characterised by greater variability in temporal patterns of eating both within and between days, lower consumption of fibre and protein and higher consumption of carbohydrates compared with food secure women⁽¹⁵⁾. Amongst these patterns, the one most strongly associated with FIS was a variable time gap between eating occasions, such as when breakfast is typically eaten. In a comparison of low-income UK adults (income below the UK median of £30 800 and not in full-time education), mean daily energy intake and inter-day variations in energy intake were similar regardless of whether individuals were classified as food secure or insecure which may suggest differences in eating patterns between the UK and USA⁽¹⁶⁾. People who are food insecure have increased odds of living with overweight or obesity, with the most marked differences seen in adult women from high-income countries who have 50 % increased odds of obesity compared with women who are food secure⁽¹⁷⁾. This is sometimes called the ‘*food insecurity-obesity paradox*’, but the phenomenon is less pronounced in men and children generally, or in women from low/middle-income countries⁽¹⁷⁾. There is also evidence from some countries, such as Canada, that individuals experiencing FIS eat more highly processed foods⁽¹⁸⁾.

Even after correcting for income and socio-economic status, people living with FIS have 40–70 % increased odds of obesity and an increased chance of developing metabolic syndrome⁽¹⁹⁾ as well as some non-communicable diseases⁽²⁰⁾. This suggests that these enhanced risks are unlikely to be linked solely to poverty.

Furthermore, FIS is associated with accelerated biological ageing estimated using telomere length⁽²¹⁾. As expected, given these findings, a greater all-cause mortality is observed in populations with FIS beginning at an OR of 1.28 (CI = 1.08, 1.52) in those who are marginally food insecure rising to an OR of 2.60 (CI = 2.17, 3.12) in those with severe FIS⁽²²⁾. Data from animal models (such as starlings⁽²³⁾ or rats⁽²⁴⁾) and human randomised controlled trials^(25–27) strongly suggest that health and obesity associations with FIS are causal. Potential reasons may include irregular eating patterns and anticipated food scarcity leading to increases in body weight or disrupted energy balance, and preferences for energy dense foods favouring weight gain⁽²⁸⁾. Research in starlings subjected to FIS found shorter telomere length and inhibited feather regrowth, indicating reduced investment in somatic maintenance and repair⁽²³⁾. Variable or unpredictable food intake is sufficient to cause weight gain in the absence of changes to diet quality or total energy intake⁽²³⁾. Therefore, it is hypothesised that FIS directly triggers reductions in energy expenditure or more efficient energy bioavailability from food. The biological mechanisms underlying this effect are unclear but may include a stress response to irregular food supply which impacts hunger, appetite and preference for energy dense foods; an adverse impact of fasting on dietary thermogenesis or voluntary energy expenditure; or an inability to develop the anticipatory hormonal responses to meals requisite for homeostatic control of blood glucose and fats, for example normal insulin release. It is also plausible that changes to the gut microbiota may be implicated in the altered metabolic response to FIS. The gut microbiome supports the extraction of energy and nutrients from foods, produces multiple metabolites with relevance to health, supporting normal immunity and protection of the host against pathogens⁽²⁹⁾. However, the gut microbiota is dependent upon a steady supply of substrate for its maintenance, with non-digestible carbohydrates a major substrate⁽³⁰⁾.

Although each person has a unique microbiome signature, it is known that the composition and activities of the microbiota can change in response to diet, medication, stress and illness⁽³⁰⁾. The microbiome can therefore be altered in response to irregular eating patterns and poor-quality diets, which may, in turn, increase the risk of chronic disease, for example bowel cancer, inflammatory bowel disease, diabetes or allergy⁽³¹⁾. In contrast, higher fibre diets promote more beneficial species (particularly lactic acid bacteria) and their metabolites (particularly short-chain fatty acids) within the gut, which may lower disease and allergy risks⁽³²⁾. However, whilst certain disease states are associated with reduced bacterial diversity and altered metabolic profiles in observational studies, there is a paucity of cause-and-effect data from well-conducted longitudinal studies and clinical trials. A barrier to this is the innate variation in the gut microbiome that makes identifying specific and reproducible disease signatures challenging⁽³³⁾.

Literature on the relationship between FIS and the microbiome remains sparse, although differences in faecal metabolite profiles between food secure and insecure USA students have been noted⁽³⁴⁾. Underlying mechanisms to explain these could include the impact of lower fibre, nutrient-poor diets on the gut microbiota, as well as stress and environmental factors. Emerging research from animal studies^(35–37) suggests negative effects of some of these components on the gut microbiota, glucose tolerance, inflammation and opportunistic pathogen expansion, although findings still need to be verified in humans.

As mentioned previously, lower fibre intakes are more common in populations experiencing FIS⁽¹⁵⁾. A lack of fibre may have long-term health impacts for individuals since the

breakdown of fibre by gut bacteria can promote gut mucosa integrity through various actions including (a) the utilisation of bacterial metabolites such as short-chain fatty acids; (b) more rapid gut transit which reduces colonic exposure to putrefactive microbiota fermentation products; (c) release of polyphenol compounds from plant fibres; and (d) microbial synthesis of nutrients including vitamin K, folate and riboflavin.

FIS, behaviour change and the care pathway

Obesity affects more than 60 % of the UK adult population but is more common in populations experiencing FIS⁽¹⁹⁾. Prevention strategies do not appear to have been successful given the UK's static and high incidence of obesity. In the NHS in England, adult weight management is currently delivered via a tiered system. Individuals move between tiers depending on the intensity of intervention required, often needing to complete a lower tier to access a higher Tier (e.g. Tier 2 for referral to Tier 3) (Figure 1).

Tier 1 interventions include public health campaigns, adverts, websites and brief interventions in primary care. Brief interventions in primary care can be effective as demonstrated in the BWeL trial⁽³⁸⁾.

Tier 2 is aimed at individuals with a BMI ≥ 30 kg/m². Interventions include group interventions and the prescription of Orlistat, a lipase inhibitor.

Tier 3 involves more specialist one-to-one interventions, using multidisciplinary teams and typically in a hospital setting. It is aimed at individuals with severe or complex obesity (BMI ≥ 40 kg/m² or ≥ 35 kg/m² with obesity related co-morbidities). Within this tier, options include very low-calorie diets or the prescription of glucagon-like peptide-1 receptor agonists, such as Semaglutide or Liraglutide⁽³⁹⁾.

Tier 4 includes bariatric (metabolic) surgery, but individuals are required to receive input from tier 3 multidisciplinary teams before being referred for this (typically 12 months)⁽⁴⁰⁾.

Although the recent approvals by NICE for new pharmacotherapies, such as Liraglutide⁽⁴¹⁾ and Semaglutide⁽⁴²⁾, may improve options for the treatment of obesity, other barriers exist to effective clinical management of obesity. These include a patchy provision of tier 3 service⁽⁴³⁾ and a lack of GP confidence and training to initiate conversations about patients' weight⁽⁴⁴⁾. From the patient perspective, misunderstanding or fear of stigmatisation can hamper engagement with health services⁽⁴⁵⁾. One study reported that people take, on average, nine years to approach their GP about their weight⁽⁴⁶⁾. Therefore, people living with FIS who may be at a high risk of obesity may face difficulty seeking and receiving help to manage their obesity.

Understanding and supporting behaviour change: Lower socio-economic status is associated with a higher BMI⁽⁴⁷⁾. This relationship appears to be mediated by health motives, i.e. how much participants 'bought into' healthy eating and concerns about body weight, but not by executive functioning (i.e. an individual's capacity for planning ahead and monitoring their own behaviour). In the UK, FIS in adults was indirectly associated with higher BMI via greater psychological distress and 'eating to cope'⁽⁴⁸⁾. Whilst the Theory of Planned Behaviour is a useful framework to study how beliefs, attitudes, perceived identity control and subjective norms influence behaviour⁽⁴⁹⁾, it is more useful to change health behaviour rather than simply observe it.

The Behaviour Change Wheel provides a conceptual framework⁽⁵⁰⁾ to analyse behavioural drivers with three layers of drivers likely to influence behaviour. These are (i) individual sources of




Figure 1. Weight management tiers in adult clinical settings (NICE, 2025; ref⁽³⁹⁾).

behaviour e.g., social, physical, psychological (ii) interventions such as education, coercion, incentivisation and (iii) policy options including guidelines, fiscal measures and service provision. As developed by the Behavioural Change Wheel, successful interventions focus on the COM-B dimensions, which are Capability, Opportunity and Motivation leading to Behavioural change⁽⁵⁰⁾. In the context of healthier diet, ‘capability’ reflects an individual’s psychological or physical ability to choose healthier diets; ‘opportunity’ refers to the food environment including enablers and barriers within this and ‘motivation’ encompasses reflective and automatic mechanisms which either activate or inhibit behaviour.

Qualitative research has examined current barriers to healthier diets in adults at risk of FIS. Poor mental health, financial stress and high food prices were barriers to change, while current food choice was influenced by social factors (e.g., child food preferences) and environmental factors (e.g., temptations in local food shops)⁽⁵¹⁾. Parents living with FIS commented that they bought treat foods or takeaways because this was affordable and made their children happy. Often healthy eating knowledge is overridden by desire or pressure for convenience, poor cooking facilities, time pressure, stress and desire to avoid conflict at mealtimes. Population and policy interventions could help to address some of these barriers to healthier eating and may involve planning, legislation, regulation and fiscal measures (Table 3).

Behavioural change techniques tested in weight management interventions suggest goal-setting approaches are effective. For example, a meta-analysis of thirty-five interventions found that the most effective interventions were based at home or in community settings⁽⁶¹⁾. As noted in this study, healthy eating, self-monitoring, delivery through personal contact and targeting multiple behaviours further increased effectiveness. Intervention designs are increasingly using co-production, where target communities, such as people experiencing FIS, are actively involved in the development of behavioural interventions. Effective co-production, according to authors who developed a children’s health strategy, is guided by nine principles built on the values of equality, reciprocity and agency⁽⁶²⁾.

Co-production is underpinning a new study – the CHarMING project – to develop interventions to support multiple behaviour change⁽⁶³⁾. So far, early insight has been gained into the

Table 3. Policy opportunities to influence food choice in populations with FIS

Policy area	Examples
Environmental/ Social Planning	Restricting hot food takeaways near schools ⁽⁵²⁾
Communication/ Marketing	Health messaging ⁽⁵³⁾ ; marketing ⁽⁵⁴⁾ ; calorie reduction ⁽⁵⁵⁾
Legislation	Restrictions on placement and promotion of HFSS products ⁽⁵⁶⁾
Service Provision	Weight management services ⁽⁵⁷⁾ ; interventions including bariatric surgery and pharmacotherapy
Regulation	Calorie labelling on menus ⁽⁵⁸⁾
Fiscal Measures	Soft drinks industry levy ⁽⁵⁹⁾
Guidelines	Obesity-related guidance, e.g. NICE, FSA ⁽⁶⁰⁾

FIS, food insecurity; FSA, Food Standards Agency.

misalignment between target community and intervention developers’ goals, resulting in improved training and supportive conversations about behaviour change. Figure 2 summarises three steps that could be considered when working with groups experiencing FIS to instigate more supportive, client-centred conversations about health⁽⁶³⁾.

Achieving policy change

It is said that ‘what surrounds us, shapes us’. Within Western food environments, there is an abundance of food and drink that is high in energy, saturated fat, free sugars and salt. Foods high in saturated fat, salt and free sugars are heavily advertised, marketed and promoted, encouraging consumers to buy more, or to make unplanned purchases⁽⁶⁴⁾. In practice, this has a disproportionately negative impact on lower income families, compounding inequalities in health and household budgets. Regulating the food environment is an equitable way of improving health. Evidence from Scotland suggests that discretionary foods account for 20 % of total daily energy and half of daily sugar intake⁽⁶⁵⁾. Discretionary items (confectionery, cakes, biscuits, pastries, savoury snacks, sugar-sweetened beverages and alcoholic drinks) tend to be high in calories, low in nutritional value and are unnecessary for health. Mean daily intakes of discretionary foods do not appear to differ by deprivation score, but there are significant differences in the consumption of sugar-sweetened beverages (113 g in most deprived v. 59 g per day in least deprived) providing 9 g v. 3 g in daily intakes of free sugars, respectively⁽⁶⁶⁾. Cost is just one of the factors influencing food choice in lower income groups with transportation, energy costs and cooking/storage facilities also being important. FIS is greater in single parent and single adult households,⁽⁵⁾ which are more likely to be impacted by food costs, particularly those in lower socio-economic groups or with disabilities and long-term health conditions⁽⁶⁷⁾. Furthermore, 34 % of people blamed the cost of living situation for their diets being less healthy, while 45 % said they were deprioritising healthy eating.

Addressing barriers within the food environment is a more equitable way of improving population diets than relying on individual change: examples include reformulation⁽⁶⁸⁾, restricting promotion of foods/drinks high in fat sugar and salt⁽⁶⁹⁾, nutrition guidance for the out-of-home sector and a code of practice for children’s menus⁽⁷⁰⁾. Food Standards Scotland is supporting and




Figure 2. Key steps to delivering client-centred conversations about health.




Figure 3. How nutrition research fits into the policy cycle.

building on this work, as outlined in its new public health nutrition strategy⁽⁷¹⁾ and the Eat Well Your Way online resource⁽⁷²⁾. Future policy work will need to support sustainability goals, such as meat and dairy reduction⁽⁷³⁾, while considering the cost barriers experienced by lower income consumers.

Policy sits in a context of political, economic, sociological, technological, legal and environmental considerations that influence strategic organisations. Civil servants follow UK Government's Green book⁽⁷⁴⁾ that outlines the ROAMEF policy development cycle: this takes decision-makers through the Rationale, Objectives, Appraisal (of different options), Monitoring (of implementation), Evaluation (of outcomes) and,

finally, the Feedback⁽⁷⁴⁾. Good policymaking sets benefit against potential detriment, impact on inequalities, unintended consequences, cost/benefit, return for investment and opportunity costs. Successful policy depends on evidence, politics and delivery coming together. UK nutrition policy – with the Scientific Advisory Committee on Nutrition at its core – follows the key steps of identifying a need to evaluate and review risks (Figure 3). The process of evidence gathering and consultation leading to policy change is not usually a quick outcome. One example is the folic acid fortification policy,⁽⁷⁵⁾ which began in 1991 with the identification of a need (prevention of neural tube defects in infants), moved onto appraisal of policy options in 2001 but has,

since then, seen ongoing cycles of requests for evidence, scrutiny and consultation. Indeed, despite agreement for mandatory fortification of non-wholemeal flour with folic acid in September 2021, at time of submission of this paper (summer 2024) policy still had not been implemented.

Regarding FIS, good policy depends on clear articulation of the issue with firm definitions; for example, 'sufficient food', 'nutritious food', 'normal growth' and 'food preferences' remain difficult to quantify and open to interpretation. If FIS is defined as lacking 'nutritious food', as proposed by several organisations, then most people in the UK are in this category since very few meet current dietary recommendations. Another challenge is the lack of consistency in dietary messaging especially the proliferation of 'healthy, sustainable diets' endorsed by expert bodies, such as Eat-Lancet, One Blue Dot, Mediterranean diet, which could create confusion since they add complexity to the basic principles defined by the Eatwell Guide. It is often assumed that eating a healthy diet is more expensive than the current average diet, but this is not necessarily the case⁽⁷⁶⁾. A whole systems approach to dietary recommendations with cross-government co-operation could be beneficial as this would help resolve current tensions between sustainability, nutrition and affordability.

Discussion

It is clear that FIS has a causal impact on metabolic health and body weight control which, over time, could have a detrimental effect on disease risk. Altered temporal patterns of eating may have significant effects on body weight and disease risk. However, it remains unclear why some forms of fasting (e.g. intermittent fasting) can benefit weight management while short periods of fasting in the context of FIS in Westernised countries is linked with excess weight gain⁽⁷⁷⁾. Analysis of the impact of FIS on the microbiome is in its infancy, but there is emerging evidence that the health impacts of FIS may, in part, be mediated via the microbiome.

Future nutrition policy to reduce FIS needs to be effective, avoid widening existing inequalities and guard against unintended consequences. Supporting evidence will need to come from collaboration across different scientific disciplines and effective interventions will need to be based on participatory research. Solving FIS should be viewed within the context of the whole food system with the goal of making sustainable diets affordable and accessible to all.

Research gaps

Forum participants identified key research gaps in understanding the impacts of FIS and determining which interventions could make sustainable diets affordable, accessible and acceptable to all. These were:

Improved experimental methods and models that can more reliably and precisely capture the different components of FIS, e.g. food and nutrient intakes, unpredictability of food supply, anxiety, emotional distress and chaotic eating, so that these can be examined separately for their impact on health. Both human and animal models should be considered, and mechanistic work is important.

Evidence is needed on what makes people more susceptible or resilient to the health effects of FIS. For example, why are women experiencing FIS in higher income regions at a greater risk of obesity than women experiencing FIS in low-to-middle-income

countries? What psychological interventions could help people to become more resilient? Evaluation of the long-term health effects of FIS, including health effects other than obesity, is merited, and any potential transgenerational (epigenetic) impacts of FIS should be tracked.

Research should capture the views of those directly affected by FIS and living with obesity. Greater use of co-creation in research and policy development is recommended to meet the needs, priorities and health aspirations of communities affected by FIS. Exploration of how technologies, such as artificial intelligence is merited, to support data collection, identify novel interventions and drive behavioural change.

We need more robust evidence to support future policy options, which can include high-quality interventions to test the impact of recommended diets or trade policies on health and climate, and how advice lands in real world settings especially in groups experiencing FIS. There is also a need for more evaluation of existing policy implementation and how this relates to dietary and health inequalities. Exploration of behaviourally relevant alternatives to food banks would be useful.

Given the emerging data on the microbiota, dietary components such as fibre could be investigated as a useful intervention to ameliorate some of the health impacts of FIS.

Finally, the ultimate goal of research and policy development should be the eradication of FIS without widening existing health inequalities. Evaluation of policy actions taking place in similarly sized countries with less inequity would be helpful.

Acknowledgements. The authors thank the Department of Food and Nutritional Sciences, University of Reading, for hosting the forum on 9th June 2023.

The forum, including a fee to CHSR for report writing, was fully supported by an award from The Rank Prize Funds.

All authors reviewed the manuscript and agreed its content; C. H. S. R. attended and contributed to the forum and wrote the first draft of the manuscript; M. B., N. G., K. H., H. L.-M., L. L., A. McD., H. P. and A. W. W. contributed presentations (as indicated in Table 1) and to the discussion, L. L.'s contribution was made in an independent capacity; K. H. attended the forum and contributed to the discussion; F. T., A. M. M., S. F.-T., J. V. W. co-designed the meeting and contributed to the discussion; J. A. L. and B. M. C. co-designed the meeting, contributed to the discussion and edited the final version of the manuscript.

There are no conflicts of interest.

References

1. Anderson SA (1990) Core indicators of nutritional state for difficult-to-sample populations. *J Nutr* **120**, 1559–1600.
2. Food and Agriculture Organization (FAO) (2008) An Introduction to the Basic Concepts of Food Security. Rome, EC-FAO Food Security Programme. <https://www.fao.org/docrep/013/al936e/al936e00.pdf> (accessed 04 January 2024).
3. Loopstra R & Lambie-Mumford H (2023) Food banks: understanding their role in the food insecure population in the UK. *Proc Nutr Soc* **82**, 253–263.
4. Food Standards Agency (2023) Consumer Insights Tracker. <https://www.food.gov.uk/research/household-food-insecurity-in-the-uk-6-results> (accessed 04 January 2024).
5. Scottish Government (2022) The Scottish Health Survey. <https://www.gov.scot/collections/scottish-health-survey/> (accessed 04 January 2024).
6. Welsh Government (2023) National Survey for Wales. <https://www.gov.wales/national-survey-wales> (accessed 04 January 2024).
7. Food Standards Agency (2022) Food and You 2: Wave 4. <https://www.food.gov.uk/research/food-and-you-2/food-and-you-2-wave-4> (accessed 25 July 2023).

8. Department for Work and Pensions (2023) Family Resources Survey: Financial Year 2021 to 2022. <https://www.gov.uk/government/statistics/family-resources-survey-financial-year-2021-to-2022> (accessed 25 July 2023).
9. Food Foundation (2023) Food Insecurity Tracking. <https://foodfoundation.org.uk/initiatives/food-insecurity-tracking#tabs/Round-13> (accessed 04 January 2024).
10. Loopstra R, Reeves A & Tarasuk V (2019) The rise of hunger among low-income households: an analysis of the risks of food insecurity between 2004 and 2016 in a population-based study of UK adults. *J Epidemiol Community Health* **73**, 668–673.
11. UK Government (2006) Healthy Start. <https://www.gov.uk/healthy-start> (accessed 04 January 2024).
12. Scottish Government (2022) Best Start Grant and Best Start Foods. <https://www.mygov.scot/best-start-grant-best-start-foods> (accessed 04 January 2024).
13. Strategic Priorities Fund (2024) Transforming UK Food Systems Programme. <https://ukfoodsystems.ukri.org/> (accessed 04 January 2024).
14. SALIENT Salient Consortium. <https://www.salientfoodtrials.uk/> (accessed 04 January 2024).
15. Nettle D & Bateson M (2019) Food-insecure women eat a less diverse diet in a more temporally variable way: evidence from the US National Health and Nutrition Examination Survey, 2013–2014. *J Obes* **2019**, 7174058.
16. Shinwell J, Bateson M, Nettle D, *et al.* (2022) Food insecurity and patterns of dietary intake in a sample of UK adults. *Br J Nutr* **128**, 770–777.
17. Nettle D, Andrews C & Bateson M (2017) Food insecurity as a driver of obesity in humans: the insurance hypothesis. *Behav Brain Sci* **40**, E105.
18. Dicken SJ, Qamar S & Batterham RL (2023) Who consumes ultra-processed food? A systematic review of sociodemographic determinants of ultra-processed food consumption from nationally representative samples. *Nutr Res Rev* **31**, 1–41.
19. Park SH & Strauss SM (2020) Food insecurity as a predictor of metabolic syndrome in U.S. female adults. *Public Health Nurs* **37**, 663–670.
20. Salinas-Roca B, Rubió-Piqué L, Carrillo-Álvarez E, *et al.* (2022) Impact of health and social factors on the cardiometabolic risk in people with food insecurity: a systematic review. *Int J Environ Res Public Health* **19**, 14447.
21. Hastings WJ, Shalev I & Belsky DW (2019) Comparability of biological aging measures in the National Health and Nutrition Examination Study, 1999–2002. *Psychoneuroendocrinology* **106**, 171–178.
22. Gundersen C, Tarasuk V, Cheng J, *et al.* (2018) Food insecurity status and mortality among adults in Ontario, Canada. *PLoS One* **13**, e0202642.
23. Andrews C, Zuidersma E, Verhulst S, *et al.* (2021) Exposure to food insecurity increases energy storage and reduces somatic maintenance in European starlings (*Sturnus vulgaris*). *R Soc Open Sci* **8**, 211099.
24. Myers KP, Majewski M, Schaefer D, *et al.* (2022) Chronic experience with unpredictable food availability promotes food reward, overeating, and weight gain in a novel animal model of food insecurity. *Appetite* **176**, 106120.
25. Alhussain MH, Macdonald IA & Taylor MA (2022) Impact of isoenergetic intake of irregular meal patterns on thermogenesis, glucose metabolism, and appetite: a randomized controlled trial. *Am J Clin Nutr* **115**, 284–297.
26. Alhussain MH, Macdonald IA & Taylor MA (2016) Irregular meal-pattern effects on energy expenditure, metabolism, and appetite regulation: a randomized controlled trial in healthy normal-weight women. *Am J Clin Nutr* **104**, 21–32.
27. Farshchi HR, Taylor MA & Macdonald IA (2004) Decreased thermic effect of food after an irregular compared with a regular meal pattern in healthy lean women. *Int J Obes Relat Metab Disord* **28**, 653–660.
28. Folwarczyn M, Christensen JD, Li NP, *et al.* (2021) Crisis communication, anticipated food scarcity, and food preferences: preregistered evidence of the insurance hypothesis. *Food Qual Prefer* **91**, 104213.
29. Lawley TD & Walker AW (2013) Intestinal colonization resistance. *Immunol* **138**, 1–11.
30. Ruxton CHS, Kajita C, Rocca P, *et al.* (2023) Microbiota and probiotics: chances and challenges – a symposium report. *Gut Microbiome* **4**, e6.
31. Round JL & Mazmanian SK (2009) The gut microbiota shapes intestinal immune responses during health and disease. *Nat Rev Immunol* **9**, 313–323.
32. Duncan SH, Conti E, Ricci L, *et al.* (2023) Links between diet, intestinal anaerobes, microbial metabolites and health. *Biomedicine* **11**, 1338.
33. Walker AW & Hoyle L (2023) Human microbiome myths and misconceptions. *Nat Microbiol* **8**, 1392–1396.
34. Mohr AE, Jasbi P, Vander Wyst KB, *et al.* (2022) Association of food insecurity on gut microbiome and metabolome profiles in a diverse college-based sample. *Sci Rep* **12**, 14358.
35. Devkota S, Wang Y, Musch MW, *et al.* (2012) Dietary-fat-induced taurocholic acid promotes pathobiont expansion and colitis in IL10^{-/-} mice. *Nature* **487**, 104–108.
36. Suez J, Korem T, Zeevi D, *et al.* (2014) Artificial sweeteners induce glucose intolerance by altering the gut microbiota. *Nature* **514**, 181–186.
37. Chassaing B, Koren O, Goodrich JK, *et al.* (2015) Dietary emulsifiers impact the mouse gut microbiota promoting colitis and metabolic syndrome. *Nature* **519**, 92–96.
38. Aveyard P, Lewis A, Tearne S, *et al.* (2016) Screening and brief intervention for obesity in primary care: a parallel, two-arm, randomised trial. *Lancet* **388**, 2492–2500.
39. National Institute for Health and Care Excellence (2025) Overweight and Obesity Management. Clinical Guideline NG246. <https://www.nice.org.uk/guidance/ng246> (accessed 30 June 2025).
40. Hazlehurst JM, Logue J, Parretti HM, *et al.* (2020) Developing integrated clinical pathways for the management of clinically severe adult obesity: a critique of NHS England policy. *Curr Obes Rep* **9**, 530–543.
41. National Institute for Health and Care Excellence (2020) Liraglutide for Managing Overweight and Obesity. <https://www.nice.org.uk/guidance/ta664> (accessed 04 January 2024).
42. National Institute for Health and Care Excellence (2023) Semaglutide for Managing Overweight and Obesity. <https://www.nice.org.uk/guidance/ta875> (accessed 04 January 2024).
43. Public Health England (2015) National Mapping of Weight Management Services. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/484115/Final_Weight_Management_Mapping_Report.pdf (accessed 04 January 2024).
44. Tremblett M, Webb H, Ziebland S, *et al.* (2022) Talking delicately: providing opportunistic weight loss advice to people living with obesity. *SSM Qual Res Health* **2**, 100162.
45. All-Party Parliamentary Group on Obesity (2018) The Current Landscape of Obesity Services. <https://obesityappg.com/inquiries> (accessed 01 August 2023).
46. Hughes CA, Ahern AL, Kasetty H, *et al.* (2021) Changing the narrative around obesity in the UK: a survey of people with obesity and healthcare professionals from the ACTION-IO study. *BMJ Open* **11**, e045616.
47. Robinson E, Jones A & Marty L (2022) The role of health-based food choice motives in explaining the relationship between lower socioeconomic position and higher BMI in UK and US adults. *Int J Obes* **46**, 1818–1824.
48. Keenan GS, Christiansen P & Hardman CA (2021) Household food insecurity, diet quality, and obesity: an explanatory model. *Obesity* **29**, 143–149.
49. Schifter DE & Ajzen I (1985) Intention, perceived control, and weight loss: an application of the theory of planned behavior. *J Pers Soc Psychol* **49**, 843–851.
50. Michie S, van Stralen MM & West R (2011) The behaviour change wheel: a new method for characterising and designing behaviour change interventions. *Implement Sci* **6**, 42.
51. van der Velde LA, Schuilenburg LA, Thiruvikraman JK, *et al.* (2019) Needs and perceptions regarding healthy eating among people at risk of food insecurity: a qualitative analysis. *Int J Equity Health* **18**, 184.
52. Turbutt C, Richardson J & Pettinger C (2019) The impact of hot food takeaways near schools in the UK on childhood obesity: a systematic review of the evidence. *J Pub Health* **41**, 231–239.
53. Schnepfer R, Richard A, Wilhelm FH, *et al.* (2019) A combined mindfulness–prolonged chewing intervention reduces body weight, food craving, and emotional eating. *J Consult Clin Psychol* **87**, 106–111.
54. Mediano Stoltze F, Reyes M, Smith TL, *et al.* (2019) Prevalence of child-directed marketing on breakfast cereal packages before and after Chile's food marketing law: a pre- and post-quantitative content analysis. *Int J Environ Res Public Health* **16**, 4501.
55. Public Health England (2018) Calorie Reduction: The Scope and Ambition for Action. <https://www.gov.uk/government/publications/calorie-reduction-the-scope-and-ambition-for-action> (accessed 06 January 2024).

56. Muir S, Dhuria P, Roe E, *et al.* (2023) UK government's new placement legislation is a 'good first step': a rapid qualitative analysis of consumer, business, enforcement and health stakeholder perspectives. *BMC Med* **21**, 33.
57. Hartmann-Boyce J, Johns DJ, Jebb SA, *et al.* (2014) Ingredients of weight loss programmes. *Obes Rev* **15**, 598–609.
58. Sinclair SE, Cooper M & Mansfield ED (2014) The influence of menu labeling on calories selected or consumed: a systematic review and meta-analysis. *J Acad Nutr Diet* **114**, 1375–1388.
59. Rogers NT, Cummins S, Forde H, *et al.* (2023) Associations between trajectories of obesity prevalence in English primary school children and the UK soft drinks industry levy: an interrupted time series analysis of surveillance data. *PLoS Med* **20**, e1004160.
60. Poobalan AS, Aucott LS, Ahmed S, *et al.* (2010) Analysis of the UK recommendations on obesity based on a proposed implementation framework. *BMC Pub Health* **10**, 17.
61. Bull ER, McCleary N, Li X, *et al.* (2018) Interventions to promote healthy eating, physical activity and smoking in low-income groups: a systematic review with meta-analysis of behavior change techniques and delivery/context. *Int J Behav Med* **25**, 605–616.
62. Albert A, Islam S, Haklay M, *et al.* (2023) Nothing about us without us: a co-production strategy for communities, researchers and stakeholders to identify ways of improving health and reducing inequalities. *Health Expect* **26**, 836–846.
63. Norwich Institute of Healthy Ageing (2023) The CHarMING Project. <https://healthyageingnorwich.com/associated-projects/the-charming-project/> (accessed 31 July 2023).
64. The Food Foundation (2023) From Purse to Plate: Implications of the Cost-Of-Living Crisis on Health. <https://foodfoundation.org.uk/publication/purse-plate-implications-cost-living-crisis-health> (accessed 06 January 2024).
65. Food Standards Scotland (2020) The Scottish Diet: It Needs to Change. 2020 Update. <https://www.foodstandards.gov.scot/publications-and-research/publications/the-scottish-diet-it-needs-to-change-2020-update> (accessed 31 July 2023).
66. Food Standards Scotland (2023) Consumption of Discretionary Foods and Drinks and Other Categories of Dietary Concern in Adults (16+ Years). <https://www.foodstandards.gov.scot/publications-and-research/publications/consumption-of-discretionary-foods-and-drinks-and-other-categories-of-dietary-concern-in-adults-16-yr> (accessed 06 January 2024).
67. Food Standards Scotland (2023) Consumer Attitudes Towards the Diet and Food Environment in Scotland Research Report. <https://www.foodstandards.gov.scot/publications-and-research/publications/consumer-attitudes-towards-the-diet-and-food-environment-in-scotland-research-report> (accessed 31 July 2023).
68. Food and Drink Federation (2019) Reformulation for Health Programme. <https://www.fdfscotland.org.uk/fdf/what-we-do/diet-and-health/reformulation-and-portion-size/reformulation-support-for-scotland/> (accessed 06 January 2024).
69. Department of Health and Social Care (2023) Restricting Promotions of Products High in Fat, Sugar or Salt by Location and by Volume Price: Implementation Guidance. <https://www.gov.uk/government/publications/restricting-promotions-of-products-high-in-fat-sugar-or-salt-by-location-and-by-volume-price/restricting-promotions-of-products-high-in-fat-sugar-or-salt-by-location-and-by-volume-price-implementation-guidance> (accessed 06 January 2024).
70. Food Standards Scotland & Public Health Scotland (2023) Eating Out, Eating Well Framework and Code of Practice for Children's Menus. <https://eatingouteatingwell.publichealthscotland.scot/> (accessed 06 January 2024).
71. Food Standards Scotland (2023) Public Health Nutrition Strategy. <https://www.foodstandards.gov.scot/publications-and-research/publications/public-health-nutrition-strategy> (accessed 31 July 2023).
72. Food Standards Scotland (2023) Eat Well Your Way. <https://www.eatwellyourway.scot/> (accessed 31 July 2023).
73. Climate Change Committee (2022) Is Scotland Climate Ready? 2022 Report to Scottish Parliament Recommendations. <https://www.theccc.org.uk/publication/is-scotland-climate-ready-2022-report-to-scottish-parliament/> (accessed 31 July 2023).
74. HM Treasury (2022) The Green Book. <https://www.gov.uk/government/publications/the-green-book-appraisal-and-evaluation-in-central-government/the-green-book-2020> (accessed 31 July 2023).
75. UK Government (2022) Press release: Folic Acid Added to Flour to Prevent Brain and Spinal Conditions in Foetuses. <https://www.gov.uk/government/news/folic-acid-added-to-flour-to-prevent-brain-and-spinal-conditions-in-foetuses> (accessed 06 January 2024).
76. Scarborough P, Kaur A, Cobiaci L, *et al.* (2016) Eatwell Guide: modelling the dietary and cost implications of incorporating new sugar and fibre guidelines. *BMJ Open* **6**, e013182.
77. Allaf M, Elghazaly H, Mohamed OG, *et al.* (2021) Intermittent fasting for the prevention of cardiovascular disease. *Cochrane Database Syst Rev* **1**, CD013496.