

This is a repository copy of *Observation of a new isomer in 185Au*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/232251/>

Version: Accepted Version

Article:

Fu, X. Y., Liu, Z., Andreyev, A. N. orcid.org/0000-0003-2828-0262 et al. (40 more authors) (2025) Observation of a new isomer in 185Au. Chinese Physics C. 084004. ISSN: 0256-307X

<https://doi.org/10.1088/1674-1137/add70c>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Observation of a new isomer in ^{185}Au

X. Y. Fu,^{1,2} Z. Liu,^{1,2,*} A. N. Andreyev,^{3,4,†} C. F. Jiao,^{5,‡} R. Y. Dong,⁵ W. Q. Zhang,^{1,2} H. Huang,^{1,2} W. Sun,⁶ Q. B. Chen,⁷ S. Q. Zhang,⁸ J. G. Li,^{1,2} X. H. Zhou,^{1,2} X. H. Yu,^{1,2} Q. B. Zeng,^{1,2} D. Seweryniak,⁹ B. Andel,^{10,11} S. Antalic,¹⁰ A. E. Barzakh,¹² M. Block,^{13,14,15} A. Bronis,¹⁰ M. P. Carpenter,⁹ P. Copp,⁹ J. G. Cubiss,^{3,16} D. T. Doherty,¹⁷ P. Van Duppen,¹¹ Z. Favier,¹⁸ F. Giaccoppo,^{13,14} B. Kindler,¹⁴ F. G. Kondev,⁹ T. Lauritsen,⁹ B. Lommel,¹⁴ M. Al Monthery,³ P. Mosat,¹⁰ C. Raison,³ W. Reviol,⁹ G. Savard,⁹ S. Stolze,⁹ G. L. Wilson,¹⁹ B. Ding,^{1,2} S. Guo,^{1,2} C. Y. Guo,⁸ Y. B. Yu,^{1,2} and Y. F. Wu^{1,2}

¹*Institute of Modern Physics, Chinese Academy of Sciences, Lanzhou 730000, China*

²*University of Chinese Academy of Sciences, Beijing 100049, China*

³*School of Physics, Engineering and Technology,*

University of York, York, YO10 5DD, United Kingdom

⁴*Advanced Science Research Center (ASRC), Japan Atomic Energy Agency, Tokai-mura, Japan*

⁵*School of Physics and Astronomy, SUN YAT-SEN University, Zhuhai 519082, P.R. China*

⁶*School of Physics, SUN YAT-SEN University, Guangzhou 510275, China*

⁷*Department of Physics, East China Normal University, Shanghai 200241, China*

⁸*State Key Laboratory of Nuclear Physics and Technology,*

School of Physics, Peking University, Beijing 100871, China

⁹*Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA*

¹⁰*Department of Nuclear Physics and Biophysics,*

Comenius University in Bratislava, 84248 Bratislava, Slovakia

¹¹*KU Leuven, Instituut voor Kern-en Stralingsfysica, 3001 Leuven, Belgium*

¹²*Petersburg Nuclear Physics Institute, NRC Kurchatov Institute, Gatchina 188300, Russia*

¹³*Helmholtz-Institut Mainz, 55128 Mainz, Germany*

¹⁴*GSI Helmholtzzentrum für Schwerionenforschung Darmstadt, 64291 Darmstadt, Germany*

¹⁵*Johannes-Gutenberg Universität, 55099 Mainz, Germany*

¹⁶*School of Physics and Astronomy, University of Edinburgh, Edinburgh, EH9 3FD, United Kingdom*

¹⁷*Department of Physics, University of Surrey, Guildford GU2 7XH, United Kingdom*

¹⁸*Irfu, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France*

¹⁹*Department of Physics and Astronomy, Louisiana State University, Baton Rouge, Louisiana 70803, USA*

(Dated: May 6, 2025)

Delayed γ -ray spectroscopy of ^{185}Au was studied at the Argonne Gas-Filled Analyzer. A New isomer at an excitation energy of 1504.2(4) keV with a half-life of 630(80) ns was identified in γ - γ coincidence analysis, decaying via a 294.8(3) keV transition. Based on Weisskopf estimates, the multipolarity of the 295 keV transition is assigned to be $E1$, $M1$, $E2$ or $M2$. Possible configurations for this new isomer are discussed based on configuration-constrained potential energy surface calculations.

I. INTRODUCTION

In the lead region ($Z = 82$) near the neutron mid-shell ($N = 104$), shape coexistence of spherical, oblate, and prolate shapes is well established. A striking example of shape coexistence is ^{186}Pb , where a triplet of low-lying 0^+ states was found [1, 2]. In the odd- A ^{187}Pb , three bands with different shapes were also reported [3, 4]. In addition, different shapes were also observed in $^{188,189}\text{Po}$ and $^{185,188,189,190}\text{Pb}$ [5–10].

Another interesting phenomenon in this region is the presence of high- K isomers. They are present in well-deformed nuclei and even in transitional nuclei with mass near 190 [11, 12]. High- K isomers arise from the breaking of one or more nucleon pairs to

form multi-quasiparticle (multi-qp) configurations. The K -hindrance and other properties of these states depend strongly on the proton and neutron states involved [13]. In ^{178}Hf , apart from the remarkable 31-year $K^\pi = 16^+$ four-quasineutron isomer [14], a two-quasineutron $K^\pi = 8^-$ isomer arises from the coupling between nucleons in high- j orbitals ($\nu 9/2^+[624] \otimes \nu 7/2^-[514]$). Similar two-quasineutron isomers have been observed systematically in even-mass $N = 106$ isotones from ^{174}Er to ^{188}Pb [11, 15].

Furthermore, in the $N = 106$ odd-mass isotones from ^{175}Tm to ^{187}Tl (except the ^{183}Ir) [16–21], 3-qp high- K isomers, originated from the coupling between the odd proton and the above mentioned $K^\pi = 8^-$ isomer in even-mass cores, have been identified.

The $N = 106$ nucleus, ^{185}Au , was previously studied in decay and in-beam γ -ray spectroscopy [22, 23]. A ground-state quadrupole deformation of $\beta_2 = 0.24$ was determined in a laser spectroscopy experiment [24].

In the present work, we report on the discovery of a new isomeric state in ^{185}Au at an excitation

* liuzhong@impcas.ac.cn

† andrei.andreyev@york.ac.uk

‡ jiaochf@mail.sysu.edu.cn




FIG. 1. Main panel: The γ -ray energy spectrum recorded by the X-array within a $2\ \mu\text{s}$ window after an EVR implantation in the DSSD. A background spectrum taken from the $2\text{--}4\ \mu\text{s}$ after an EVR implantation is subtracted. The three γ -rays highlighted in red were experimentally established as a cascade in ^{185}Au [22]. Inset: Expanded view of prominent γ -ray peaks from 260 to 400 keV.

energy of $1504.2(4)\ \text{keV}$, with a half-life of $T_{1/2} = 630(80)\ \text{ns}$, decaying to the $(17/2^-)$ level previously assigned to the $\pi 1h_{11/2}$ band [22]. The possible spin-parities of this isomer are tentatively assigned based on Weisskopf estimates. The possible configuration of this isomer is discussed within the framework of configuration-constrained potential energy surface (cc-PES) calculations [25].

II. EXPERIMENT

The experiment was conducted at Argonne National Laboratory (ANL), and this study extends prior results reported in earlier experiments from the same experimental program, which including ^{187}Pb , ^{188}Bi , ^{188}Po , ^{183}Hg , and ^{187}Tl [3, 6, 21, 26]. In the present work, the ^{185}Au nucleus was produced in the $^{50}\text{Cr} + ^{142}\text{Nd} \rightarrow ^{192}\text{Po}^* \rightarrow ^{185}\text{Au} + 3p\alpha$ fusion-evaporation reaction. The ^{50}Cr ions were accelerated to an energy of 255 MeV with a typical intensity of 7 pA by the superconducting linear accelerator at ANL. The target material, enriched to 99.8% in $^{142}\text{NdF}_3$, had a thickness of $700\ \mu\text{g}/\text{cm}^2$. To minimize target heating, four target sectors were mounted on a rotating wheel, and the beam was wobbled horizontally by $\pm 2.5\ \text{mm}$ across the target using a magnetic steerer.

Evaporation residues (EVRs) were separated from the primary beam by Argonne Gas-Filled Analyzer (AGFA) [27] which was filled with 0.65 mbar of helium gas. The flight time of EVRs in the AGFA separator is estimated to be several hundred nanoseconds (approximately 500 ns flight time for $v = 0.02c$ EVRs). The energy loss and arrival time of the EVRs were measured by a parallel grid avalanche counter (PGAC). The EVRs were then implanted into a double-sided silicon strip detector (DSSD) located at the focal plane, which recorded the deposition energy of the recoil and subsequent charged particle decay.

The present work was performed with the delayed γ correlation method, using the DSSD and X-array detector in the focal plane. The X-array detector setup consists of four Clover HPGe detectors and was used to detect X rays and γ rays emitted from the EVRs [28]. The typical energy resolution for the X-array was around 4 keV (FWHM) for γ rays of around 300 keV.

III. EXPERIMENTAL RESULTS

The delayed γ -ray spectrum recorded by the X-array detectors within $2\ \mu\text{s}$ after implantation in the DSSD, with the background in the time window of $2\text{--}4\ \mu\text{s}$ subtracted, is shown in Fig. 1. Most of the prominent




FIG. 2. γ -ray energy spectrum gated on the 527-, 462-, 212- and 295-keV transitions (a) - (d) and the sum of these spectra (e).

γ -ray peaks are from known isomers produced in other reaction channels, including the strong γ ray from microsecond isomers in ^{187}Tl and $^{187,188}\text{Pb}$.

The 212-, 462- and 527-keV γ rays shown in Fig. 1, were initially reported in the in-beam experiment by Larabee *et al.* [22] as a prompt 527-462-212 keV transition cascade. Considering the transport time through AGFA is more than hundreds of nanoseconds,




FIG. 3. Time distribution of the summed 212 and 527 keV γ rays after EVR implantation, with a background subtraction made by gating on nearby Compton background. The data are fitted with an exponential function with a constant background.

these transitions are considerably delayed, indicating the existence of an isomer de-exciting via this prompt cascade.

A previously unobserved 294.8(3)-keV transition was identified in the γ -gated spectra of 527-, 462- and 212-keV γ transitions, as shown in Fig. 2(a-c). The spectrum gated on the 295-keV transition and the summed spectrum gated on the four transitions are shown in Fig. 2 (d,e), respectively. This 295-keV γ ray is attributed to the deexcitation of the new isomer in ^{185}Au .

The time distribution of $\Delta T(\text{EVR}-\gamma(212 \text{ keV and } 527 \text{ keV}))$ is shown in Fig. 3. Based on a least-squares fit of this time distribution, with an exponential function plus a constant background, the half-life of the isomer was determined to be 630(80) ns.

Figure 4 illustrates the proposed decay scheme of the isomer along with a partial level scheme of ^{185}Au , and it shows that the new 1504.2(4)-keV isomer decays to the 1209.4(3)-keV ($17/2^-$) level. The spin and parity assignments for established levels were adopted from previous studies [22, 23], while the corresponding energy values were derived from the latest ENDF Library evaluations of ^{185}Au [29].

We noted that no decay to other low-lying levels from this isomer was observed after a thorough analysis of the singles and coincidence spectra. We attempted to perform an intensity balance analysis of the 295-527-462-212 keV γ -ray cascade in the recoil-delayed γ -ray spectrum Fig. 1. However, as shown in the inset of Fig. 1, there is only a slight indication of a peak at 295 keV, and hence no definitive conclusions can be drawn on its multipolarity.




FIG. 4. Partial level scheme of ^{185}Au , spin and parity are all taken from Ref. [22, 23] except the new isomer, the spin parity of the new isomer is from Weisskopf estimates, and states are labeled with their energies in keV.

IV. Discussions and calculations

The Weisskopf single-particle transition probabilities for possible multiplicities ($E1$, $M1$, $E2$, $M2$, $E3$) of the 295-keV γ -ray, corrected for internal conversion, are presented in Table I. The $E3$ and higher multiplicities can be excluded based on the recommended upper limits [30]. The possible multiplicities constrain the spin of the 1504 keV isomer, decaying to $(17/2^-)$, to a range from $13/2$ to $21/2$. The Weisskopf hindrance factors (F_w , which is the reciprocal of transition probability) of the $E1$, $M1$, and $E2$ multiplicities are large, indicating significant hindrance.

TABLE I. The Weisskopf estimates of single-particle transition probabilities for different multiplicities of the 295-keV γ ray.

Multiplicity assumed	Weisskopf half-life estimates (s)	Transition probability (W.u)
$E1$	$8.13(3) \times 10^{-15}$	$1.3(2) \times 10^{-4}$ ^a
$M1$	$8.59(3) \times 10^{-13}$	$1.4(2) \times 10^{-6}$
$E2$	$4.06(2) \times 10^{-9}$	$6.5(8) \times 10^{-3}$
$M2$	$4.29(2) \times 10^{-7}$	$6.9(9) \times 10^{-1}$
$E3$	$3.08(2) \times 10^{-3}$	$4.9(7) \times 10^3$

^a The transition probability of K -allowed $E1$ transitions is multiplied by 10^4 as suggested in Ref. [31, 32].

To assign the possible configuration to this new isomer




FIG. 5. PES of low-lying proton states in ^{185}Au , left is for the $h_{9/2}$ ground-state band and the right is the $h_{11/2}$ band. Calculations using cc-PES calculations, the minima are indicated by black circles, and labeled with their deformation parameters.

in ^{185}Au , we have performed cc-PES calculations [25] based on the non-axial deformed Woods-Saxon (WS) potential [33] with universal parameters [34]. The total energy is composed of a macroscopic part with the standard liquid-drop energy [35]. To avoid the possible pairing collapse in multi-qp states, the approximate particle-number conservation uses the Lipkin-Nogami (LN) method [36]. For a multi-qp state, the microscopic energy incorporates the contribution from the unpaired particles that occupy the single-particle orbits associated with the given configuration. The blocking effects from

TABLE II. Possible 3-qp configurations for the isomeric states in ^{185}Au . Excitation energies and deformation parameters are calculated using cc-PES calculations.

K^π	Configuration	Energy (keV)	β_2	β_4	γ
17/2 ⁻	$\pi 1/2^- [541] \otimes \nu 9/2^- [505] \otimes \nu 7/2^- [503]$	1041	0.153	-0.010	22.907
21/2 ⁺	$\pi 11/2^- [505] \otimes \nu 9/2^- [624] \otimes \nu 1/2^- [521]$	1289	0.168	-0.041	-25.477
17/2 ⁺	$\pi 1/2^- [541] \otimes \nu 9/2^+ [624] \otimes \nu 7/2^- [514]$	1444	0.230	-0.033	1.036
19/2 ⁺	$\pi 3/2^- [532] \otimes \nu 9/2^+ [624] \otimes \nu 7/2^- [514]$	1458	0.246	-0.031	1.144
17/2 ⁺	$\pi 1/2^- [541] \otimes \nu 9/2^+ [624] \otimes \nu 7/2^- [503]$	1582	0.165	-0.037	25.680
17/2 ⁺	$\pi 1/2^- [541] \otimes \nu 9/2^- [505] \otimes \nu 7/2^+ [633]$	1632	0.159	-0.024	24.605
17/2 ⁺	$\pi 1/2^- [530] \otimes \nu 9/2^+ [624] \otimes \nu 7/2^- [514]$	1633	0.252	-0.030	0.210
13/2 ⁺	$\pi 3/2^- [532] \otimes \nu 9/2^+ [624] \otimes \nu 1/2^- [521]$	1652	0.223	-0.038	20.593
15/2 ⁺	$\pi 1/2^- [541] \otimes \nu 7/2^+ [633] \otimes \nu 7/2^- [503]$	1667	0.169	-0.032	25.437
19/2 ⁻	$\pi 3/2^- [532] \otimes \nu 9/2^+ [624] \otimes \nu 7/2^+ [633]$	1733	0.227	-0.037	-14.634
21/2 ⁻	$\pi 5/2^+ [402] \otimes \nu 9/2^+ [624] \otimes \nu 7/2^- [514]$	1757	0.266	-0.028	0.299
17/2 ⁻	$\pi 1/2^- [550] \otimes \nu 9/2^+ [624] \otimes \nu 7/2^+ [633]$	1778	0.153	-0.026	-26.763
19/2 ⁺	$\pi 3/2^- [532] \otimes \nu 9/2^+ [624] \otimes \nu 7/2^- [503]$	1875	0.200	-0.044	-26.624
15/2 ⁻	$\pi 1/2^- [541] \otimes \nu 7/2^- [503] \otimes \nu 7/2^- [514]$	1969	0.214	-0.031	21.749

the unpaired nucleons are considered by removing the singly occupied orbitals from the LN calculation. The cc-PES calculations are carried out in the lattice of deformation space $(\beta_2, \beta_4, \gamma)$. The resultant PES's can self-consistently provide information on the deformation, excitation energy, and pairing interaction properties of multi-qp states. The calculations have successfully reproduced the experimental results for the 3-qp configurations in $N = 106$ odd- A isotones, with excitation energy deviations ranging between 1 and 174 keV (^{181}Re 21/2⁻ isomeric state [19] and ^{175}Tm 23/2⁺ isomeric state [16]). Details of the calculations will be presented in a forthcoming paper Ref. [37].

In the previous experimental studies, the ground-state band (C) was proposed to originate from the strongly mixed $\pi 3/2^- [532]$ and $\pi 1/2^- [541]$ configuration [38, 39], while Band (B) was assigned to an $h_{11/2}$ single-quasiproton configuration [22]. PES calculations for all possible low-lying proton configurations near the Fermi surface are also performed, and the results for the 3/2⁻ [532] and 1/2⁻ [550] configurations are shown in Fig. 5. The results for 3/2⁻ [532] configuration reproduce the $\beta_2 = 0.24$ ground-state deformation from the laser spectroscopy measurements [40]. Band (B) corresponds to the 1/2⁻ [550] orbital, as shown in Fig. 5 (right). Notably, the PES calculations show remarkable triaxial deformations for the two low-lying configurations, similar to those observed in calculations for neighboring $^{183,187}\text{Au}$ [41, 42].

The calculated excitation energies and deformations for possible 3-qp configurations in ^{185m}Au are presented in Table II. These high- K states are all predicted to lie closely to the measured excitation energy of 1504 keV. However, it should be mentioned that the residual interactions between unpaired nucleons are not accounted for in the present calculations. These residual interactions can lead to energy shifts of up to a few

hundred keV, depending on the relative orientation of intrinsic spins of nucleons involved [43–45]. Therefore, in the context of the excitation energy and decay patterns, we are unable to make a firm assignment for the configuration of the observed isomer from the calculated 3-qp configurations. The calculated results incorporating the neutron configuration $\nu 9/2^+ [624] \otimes \nu 7/2^- [514]$ are consistent with systematics of 3-qp configurations in odd-mass nuclei with $N = 106$. However, further experimental data are required to unambiguously assign the spin-parity and configuration of the 1504 keV isomer.

V. Summary

In summary, a new isomer at 1504 keV in ^{185}Au was identified through its deexcitation to a previously known level, by a 295-keV transition in coincidence with the known 212-462-527 keV cascade. Possible multiplicities of $E1$, $M1$, $E2$, and $M2$ are proposed for this transition in comparison to predictions from Weisskopf estimates. By using the cc-PES calculations which described well the systematics of known three-quasiparticle isomers in odd-mass $N = 106$ isotones [37], possible spin-parity and configuration candidates for the new isomer at 1504 keV in ^{185}Au are given.

VI. Acknowledgments

This work was supported by the National Natural Science Foundation of China (Grants No. 12405140, No. 12135004, No. 11961141004, No. U2032138 and No. 12275369) and the Natural Science Foundation of Gansu Province, China (Grant No. 24JRRA033). UK personnel are grateful for financial support from the STFC and

A.N. Andreyev is funded by the Chinese Academy of Sciences President's International Fellowship Initiative (Grant No. 2020VMA0017). This work was supported by the U.S. Department of Energy, Office of Nuclear Physics, under Contract No. DE-AC02-06CH11357 and Grants No. DE-FG02-94ER41041 (UNC) and DE-FG02-97ER41033 (TUNL). This work was also supported by the Slovak Research and Development Agency (contract No. APVV-22-0282) and Slovak Grant Agency VEGA (Contract No. 1/0019/25). This research

used resources of ANL's ATLAS facility, which is a DOE Office of Science User Facility. We thank the Target Labor team of GSI for preparing the targets for the experiment.

VII. References

-
- [1] A. N. Andreyev, M. Huyse, P. Van Duppen, L. Weissman, D. Ackermann, J. Gerl, F. P. Hessberger, S. Hofmann, A. Kleinböhl, G. Münzenberg, S. Reshitko, C. Schlegel, H. Schaffner, P. Cagarda, M. Matos, S. Saro, A. Keenan, C. Moore, C. D. O'Leary, R. D. Page, M. Taylor, H. Kettunen, M. Leino, A. Lavrentiev, R. Wyss, and K. Heyde, *Nature* **405**, 430 (2000).
 - [2] J. Ojala, J. Pakarinen, P. Papadakis, J. Sorri, M. Sandzelius, D. M. Cox, K. Auranen, H. Badran, P. J. Davies, T. Grahn, P. T. Greenlees, J. Henderson, A. Herzán, R.-D. Herzberg, J. Hilton, U. Jakobsson, D. G. Jenkins, D. T. Joss, R. Julin, S. Juutinen, T. Kibédi, J. Konki, G. J. Lane, M. Leino, J. Liimatainen, C. G. McPeake, O. Neuvonen, R. D. Page, E. Parr, J. Partanen, P. Peura, P. Rakhila, J. Revill, P. Ruotsalainen, J. Sarén, C. Scholey, S. Stolze, J. Uusitalo, A. Ward, and R. Wadsworth, *Communications Physics* **5**, 213 (2022).
 - [3] W. Zhang, A. Andreyev, Z. Liu, D. Seweryniak, H. Huang, Z. Li, J. Li, C. Guo, D. Doherty, A. Barzakh, P. Van Duppen, J. Cubiss, B. Andel, S. Antalic, M. Block, A. Bronis, M. Carpenter, P. Copp, B. Ding, Z. Favier, F. Giacoppo, T. Huang, X. Yu, B. Kindler, F. Kondev, T. Lauritsen, G. Li, B. Lommel, H. Lu, M. Al Monthery, P. Mošaf, Y. Niu, C. Raison, W. Reviol, G. Savard, S. Stolze, G. Wilson, H. Wu, Z. Wang, F. Xu, Q. Zeng, and X. Zhou, *Physics Letters B* **829**, 137129 (2022).
 - [4] A. N. Andreyev, M. Huyse, K. Van de Vel, P. Van Duppen, O. Dorvaux, P. Greenlees, K. Helariutta, P. Jones, R. Julin, S. Juutinen, H. Kettunen, P. Kuusiniemi, M. Leino, M. Muikku, P. Nieminen, P. Rakhila, J. Uusitalo, R. Wyss, K. Hauschild, and Y. Le Coz, *Phys. Rev. C* **66**, 014313 (2002).
 - [5] K. Van de Vel, A. N. Andreyev, D. Ackermann, H. J. Boardman, P. Cagarda, J. Gerl, F. P. Heßberger, S. Hofmann, M. Huyse, D. Karlgren, I. Kojouharov, M. Leino, B. Lommel, G. Münzenberg, C. Moore, R. D. Page, S. Saro, P. Van Duppen, and R. Wyss, *Phys. Rev. C* **68**, 054311 (2003).
 - [6] W. Q. Zhang, A. N. Andreyev, Z. Liu, D. Seweryniak, H. Huang, Z. H. Li, J. G. Li, C. Y. Guo, A. E. Barzakh, P. Van Duppen, M. Al Monthery, B. Andel, S. Antalic, M. Block, A. Bronis, M. P. Carpenter, P. Copp, J. G. Cubiss, B. Ding, D. T. Doherty, Z. Favier, F. Giacoppo, T. H. Huang, B. Kindler, F. G. Kondev, T. Lauritsen, G. S. Li, B. Lommel, H. Y. Lu, P. Mošaf, Y. F. Niu, C. Raison, W. Reviol, G. Savard, S. Stolze, G. L. Wilson, H. Y. Wu, Z. H. Wang, F. R. Xu, X. H. Yu, Q. B. Zeng, and X. H. Zhou, *Phys. Rev. C* **106**, 024317 (2022).
 - [7] P. Papadakis, J. Pakarinen, A. Briscoe, D. Cox, R. Julin, K. Auranen, T. Grahn, P. Greenlees, K. Hadyńska-Klek, A. Herzán, R.-D. Herzberg, U. Jakobsson, S. Juutinen, J. Konki, M. Leino, A. Mistry, D. O'Donnell, P. Peura, P. Rakhila, P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, S. Stoltze, J. Sorri, J. Uusitalo, and K. Wrzosek-Lipska, *Physics Letters B* **858**, 139048 (2024).
 - [8] A. M. Baxter, A. P. Byrne, G. D. Dracoulis, P. M. Davidson, T. Kibédi, R. V. F. Janssens, M. P. Carpenter, C. N. Davids, T. L. Khoo, and T. Lauritsen, *Phys. Rev. C* **71**, 054302 (2005).
 - [9] K. Van de Vel, A. N. Andreyev, D. Ackermann, H. J. Boardman, P. Cagarda, J. Gerl, F. P. Heßberger, S. Hofmann, M. Huyse, D. Karlgren, I. Kojouharov, M. Leino, B. Lommel, G. Münzenberg, C. Moore, R. D. Page, S. Saro, P. Van Duppen, and R. Wyss, *The European Physical Journal A - Hadrons and Nuclei* **24**, 57 (2005).
 - [10] A. M. Plaza, J. Pakarinen, P. Papadakis, R.-D. Herzberg, R. Julin, T. R. Rodríguez, A. D. Briscoe, A. Illana, J. Ojala, P. Ruotsalainen, E. Uusikylä, B. Alayed, A. Alharbi, O. Alonso-Sañudo, K. Auranen, V. Bogdanoff, J. Chadderton, A. Esmaylzadeh, C. Fransen, T. Grahn, P. T. Greenlees, J. Jolie, H. Joukainen, H. Jutila, C.-D. Lakenbrink, M. Leino, J. Louko, M. Luoma, A. McCarter, B. S. N. Singh, P. Rakhila, A. Raggio, J. Romero, J. Sarén, M.-M. Satrazani, M. Stryczyk, C. M. Sullivan, Álvaro Tolosa-Delgado, J. Uusitalo, F. von Spee, J. Warbinek, and G. L. Zimba, *Communications Physics* **8**, 8 (2025).
 - [11] G. D. Dracoulis, P. M. Walker, and F. G. Kondev, *Reports on Progress in Physics* **79**, 076301 (2016).
 - [12] F. R. Xu, P. M. Walker, and R. Wyss, *Phys. Rev. C* **59**, 731 (1999).
 - [13] K. Jain, P. Walker, and N. Rowley, *Physics Letters B* **322**, 27 (1994).
 - [14] J. Van Klinken, W. Venema, R. Janssens, and G. Emery, *Nuclear Physics A* **339**, 189 (1980).
 - [15] G. D. Dracoulis, G. J. Lane, F. G. Kondev, H. Watanabe, D. Seweryniak, S. Zhu, M. P. Carpenter, C. J. Chiara, R. V. F. Janssens, T. Lauritsen, C. J. Lister, E. A. McCutchan, and I. Stefanescu, *Phys. Rev. C* **79**, 061303 (2009).
 - [16] R. O. Hughes, G. J. Lane, G. D. Dracoulis, A. P. Byrne, P. H. Nieminen, H. Watanabe, M. P. Carpenter, P. Chowdhury, R. V. F. Janssens, F. G. Kondev, T. Lauritsen, D. Seweryniak, and S. Zhu, *Phys. Rev. C*

- 86**, 054314 (2012).
- [17] G. Dracoulis, F. Kondev, G. Lane, A. Byrne, T. Kibédi, I. Ahmad, M. Carpenter, S. Freeman, R. Janssens, N. Hammond, T. Lauritsen, C. Lister, G. Mukherjee, D. Seweryniak, P. Chowdhury, S. Tandel, and R. Gramer, *Physics Letters B* **584**, 22 (2004).
 - [18] D. Barnéoud, S. André, and C. Foin, *Nuclear Physics A* **379**, 205 (1982).
 - [19] C. Pearson, P. Walker, C. Purry, G. Dracoulis, S. Bayer, A. Byrne, T. Kibédi, and F. Kondev, *Nuclear Physics A* **674**, 301 (2000).
 - [20] A. Byrne, A. Baxter, G. Dracoulis, S. Mullins, T. Kibédi, T. McGoram, K. Helariutta, J. Cocks, P. Jones, R. Julin, S. Juutinen, H. Kankaanpää, H. Kettunen, P. Kuusiniemi, M. Leino, M. Muikku, P. Nieminen, P. Rakhila, and A. Savelius, *The European Physical Journal A - Hadrons and Nuclei* **7**, 41 (2000).
 - [21] C. Y. Guo, W. Q. Zhang, H. Huang, S. Q. Zhang, Z. H. Li, Z. Liu, H. Hua, D. W. Luo, H. Y. Wu, A. N. Andreyev, D. Seweryniak, B. Andel, S. Antalic, A. E. Barzakh, M. Block, A. Bronis, M. P. Carpenter, P. Copp, J. G. Cubiss, D. T. Doherty, P. Van Duppen, Z. Favier, F. Giacoppo, B. Kindler, F. G. Kondev, T. Lauritsen, X. Q. Li, B. Lommel, M. Al Monthery, P. Mošat, L. Ni, C. Raison, W. Reviol, G. Savard, S. Stolze, G. L. Wilson, C. Xu, J. Z. Zhang, S. Y. Zhang, and Z. X. Zhou, *The European Physical Journal A* **60**, 165 (2024).
 - [22] A. Larabee, M. Carpenter, L. Riedinger, L. Courtney, J. Waddington, V. Janzen, W. Nazarewicz, J.-Y. Zhang, R. Bengtsson, and G. Lèander, *Physics Letters B* **169**, 21 (1986).
 - [23] M. Kortelahti, E. F. Zganjar, H. K. Carter, C. Papanicolopoulos, M. A. Grimm, and J. L. Wood, *Journal of Physics G: Nuclear Physics* **14**, 1361 (1988).
 - [24] K. Wallmeroth, G. Bollen, A. Dohn, P. Egelhof, J. Grüner, F. Lindenlauf, U. Krönert, J. Campos, A. Rodriguez Yunta, M. J. G. Borge, A. Venugopalan, J. L. Wood, R. B. Moore, and H. J. Kluge, *Phys. Rev. Lett.* **58**, 1516 (1987).
 - [25] F. Xu, P. Walker, J. Sheikh, and R. Wyss, *Physics Letters B* **435**, 257 (1998).
 - [26] H. Huang, W. Zhang, A. Andreyev, Z. Liu, D. Seweryniak, Z. Li, C. Guo, A. Barzakh, P. Van Duppen, B. Andel, S. Antalic, M. Block, A. Bronis, M. Carpenter, P. Copp, J. Cubiss, B. Ding, D. Doherty, Z. Favier, F. Giacoppo, T. Huang, B. Kindler, F. Kondev, T. Lauritsen, J. Li, G. Li, B. Lommel, H. Lu, M. Al Monthery, P. Mošat, Y. Niu, C. Raison, W. Reviol, G. Savard, S. Stolze, G. Wilson, H. Wu, Z. Wang, F. Xu, Q. Zeng, X. Yu, F. Zeng, and X. Zhou, *Physics Letters B* **833**, 137345 (2022).
 - [27] B. Back, in *EPJ Web Conf.*, Vol. 163 (E3S Web of Conferences, 2017) p. 00003.
 - [28] A. Mitchell, P. Bertone, B. DiGiovine, C. Lister, M. Carpenter, P. Chowdhury, J. Clark, N. DOLympia, A. Deo, F. Kondev, E. McCutchan, J. Rohrer, G. Savard, D. Seweryniak, and S. Zhu, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **763**, 232 (2014).
 - [29] D. Brown, M. Chadwick, R. Capote, *et al.*, *Nuclear Data Sheets* **148**, 1 (2018), special Issue on Nuclear Reaction Data.
 - [30] R. Firestone, V. Shirley, C. Baglin, S. Chu, and J. Zipkin, *Title of the 8th edition of the Table of Isotopes* (Springer, Hungary, 1996) proceedings of the 9th International Symposium on Capture gamma-ray spectroscopy and related topics. V. 2.
 - [31] Z. P. et al., *Physics Letters B* **753**, 182 (2016).
 - [32] P. M. Walker and F. G. Kondev, *The European Physical Journal Special Topics* **233**, 983 (2024).
 - [33] W. Nazarewicz and P. Olanders, *Nuclear Physics A* **441**, 420 (1985).
 - [34] J. Dudek, Z. Szymański, and T. Werner, *Phys. Rev. C* **23**, 920 (1981).
 - [35] W. D. Myers and W. J. Swiatecki, *Nuclear Physics* **81**, 1 (1966).
 - [36] H. Pradhan, Y. Nogami, and J. Law, *Nuclear Physics A* **201**, 357 (1973).
 - [37] R. Y. Dong and C. F. Jiao (2025), in preparation.
 - [38] M. Porquet, C. Bourgeois, P. Kilcher, and J. Sauvage-Letessier, *Nuclear Physics A* **411**, 65 (1983).
 - [39] B. Roussière, F. Ibrahim, J. Libert, J. Sauvage, D. Verney, L. Cabaret, J. E. Crawford, J. Genevey, D. Hojman, G. Huber, A. Knipper, F. Le Blanc, J. K. P. Lee, G. Marguier, J. Oms, J. Pinard, and A. Wojtasiewicz, *Physics of Atomic Nuclei* **64**, 1048 (2001).
 - [40] K. Wallmeroth, G. Bollen, A. Dohn, P. Egelhof, U. Krönert, M. Borge, J. Campos, A. Yunta, K. Heyde, C. De Coster, J. Wood, and H.-J. Kluge, *Nuclear Physics A* **493**, 224 (1989).
 - [41] P. Möller, A. J. Sierk, R. Bengtsson, H. Sagawa, and T. Ichikawa, *Phys. Rev. Lett.* **103**, 212501 (2009).
 - [42] S. Guo, X. Zhou, C. Petrache, E. Lawrie, S. Mthembu, Y. Fang, H. Wu, H. Wang, H. Meng, G. Li, Y. Qiang, J. Wang, M. Liu, Y. Zheng, B. Ding, W. Zhang, A. Rohilla, K. Muhki, Y. Yang, H. Ong, J. Ma, S. Xu, Z. Bai, H. Fan, J. Huang, J. Li, J. Xu, B. Lv, W. Hua, Z. Gan, and Y. Zhang, *Physics Letters B* **828**, 137010 (2022).
 - [43] K. Jain and A. K. Jain, *Phys. Rev. C* **45**, 3013 (1992).
 - [44] J. Boisson, R. Piepenbring, and W. Ogle, *Physics Reports* **26**, 99 (1976).
 - [45] K. Jain, O. Burglin, G. Dracoulis, B. Fabricius, N. Rowley, and P. Walker, *Nuclear Physics A* **591**, 61 (1995).