

Title: Age-specificity in territory quality and spatial structure in a wild bird population

Short title: Space and age in a wild bird population

Key words: Age structure; breeding density; *Parus major*; spatially-variable demography; territory quality; territory choice

Article type: Major article

Wordcount: 7115 (excluding abstract, literature cited, and figure legends)

List of contents

ABSTRACT	2
(1) Introduction	3
(2) Methods	9
(i) Study system and data collection	9
(ii) Data selection	10
(iii) Statistical analyses	11
(3) Results	15
(4) Discussion	19
(5) Conclusions	27
References	29

ABSTRACT

Age influences behaviour, survival, and reproduction; hence variation in population age structure can affect population-level processes. The extent of spatial age structure may be important in driving spatially-variable demography, particularly when space-use is linked to reproduction, yet it is not well understood. We use long-term data from a wild bird population to quantify covariance between territory quality and age and examine spatial age structure. We find associations between age and aspects of territory quality, but little evidence for spatial age structure compared to the spatial structure of territory quality and reproductive output. We also report little between-year repeatability of spatial age structure compared to structure in reproductive output. We suggest that high breeding site fidelity among individuals that survive between years, yet frequent territory turnover driven by high mortality and immigration rates, limits the association between age and territory quality and weakens overall spatial age structure. Greater spatial structure and repeatability in reproductive output compared to age suggests that habitat quality may be more important in driving spatially-variable demography than age in this system. We suggest that the framework developed here can be used in other taxa to assess spatial age structure, particularly in longer-lived species where we predict from our findings there may be greater structure.

(1) Introduction

Age affects many aspects of life, as resources are allocated into processes and traits at different points throughout lifespan to maximise fitness (Stearns 1992) and individuals gain in experience as they age. Further, variation in population age structure, i.e. the proportion of a population in given age cohorts, may affect population dynamics, via effects on reproductive output and social organisation (Coulson et al. 2005; Gamelon et al. 2019; Siracusa et al. 2023; Woodman et al. in press). There is widespread evidence of temporal variation in age structure in wild populations (Coulson et al. 2005; Gamelon et al. 2016; Hoy et al. 2020), but the way in which age structure varies across space seems to be less well understood. Some work identifies spatial variation in age structure in fish populations, often linked to human harvesting of different age-cohorts in given areas (Fowler et al. 2000; McIlwain et al. 2005); and in ungulate and bird populations, associated with age-specificity in habitat use and social behaviour (Ferrer and Bisson 2003; Albery et al. 2022). Given that age is often important for individuals' fitness, spatial age structure is particularly interesting as it might drive local age-related population dynamics. For example, if there is covariance between habitat quality and the age of individuals occupying sites, both could lead to spatially-variable demography, thus potentially biasing estimates of the effects of age on reproduction or, conversely, estimates of environmental effects on trait variation.

One social system where covariance between age and habitat might lead to local age structure is where individuals defend breeding territories. Here, age structure might be particularly important in driving spatial age-related dynamics due to the tight association between breeding territories and reproductive output (Kerbiriou et al. 2006; Jones et al. 2014; Groenendijk et al. 2015). In many animal taxa, individuals undergo reproductive attempts while defending a territory in which resources are used for breeding (Reynolds 1996; Davies et al. 2012). This is particularly prevalent in socially monogamous bird species (Hinde 1956; Greenwood 1980). In such cases, spatial age structure could develop as

non-randomness arises when individuals of either similar or dissimilar age breed in closer proximity to each other than expected from chance.

Several mechanisms might affect spatial age structure in territorial animals; we outline four mechanisms here. (1) First, temporal variation in population age structure might influence how age is arranged in space. Temporal variation in wild population age structure is common, particularly in short-lived iteroparous species, where seasonal fluctuations in the proportion of breeding populations consisting of the youngest age cohort may vary greatly depending on survival and recruitment between breeding events (Gamelon et al. 2016; Woodman et al. 2022). Such variation might affect how age is arranged in space through passive mechanisms whereby clusters of territories occupied by same-age individuals will be more likely to arise when the age distribution is skewed towards this age-cohort. For example, recent work demonstrates how fluctuations in population age structure determine how age is structured within breeding pairs, where there is greater age-assortative pairing when the proportion of yearlings is higher (Woodman et al. 2022). Thus, between-year age distribution is likely to passively affect spatial age structure as clusters of territories occupied by same-age individuals are more likely to arise when much of the population exists in a single age-cohort (figure 1A).

However, age-specific biases in settlement may drive spatial age structure over-and-above that expected from temporal variation in age distribution alone. This may occur if there is spatial structure in territory quality (which may be generated for example through spatial clustering of resources), and there is covariance between age and territory quality, which may arise through two mechanisms. (2a) In many species, older individuals are dominant (Piper 1997; Wilson 2000), thus potentially leading to age-specificity in territory use where older individuals outcompete younger ones to acquire higher quality territories (Ferrer and Bisson 2003; Garneau et al. 2008; Brewer et al. 2009; Stepanuk et al. 2021; figure 1B). (2b) Alternatively, covariance between age and territory quality might arise if higher quality territories elevate survival, and individuals therefore persist for longer in such territories over

time (Kokko et al. 2006). Additionally, individuals might be more likely to remain on higher quality territories, which will generate a similar association between length of tenure and hence age (figure 1B). Therefore, if older individuals are more likely to occupy higher quality territories through either of these mechanisms, spatial structure in territory quality might drive spatial age structure and repeatability in this over time. Conversely, in species where total population size exceeds the number of available breeding territories, competition for breeding sites might be high, thus retention of previous territories regardless of quality might be under stronger selection than occupation of new and higher quality sites. In this case, high breeding site fidelity might break down the relationship between age and territory quality if individuals retain territories as they age regardless of its quality.

(3) Age-specific territory occupation might also be generated through mechanisms that do not directly link to how ageing affects territory acquisition. For example, if there is individual-specificity in territory acquisition mediated by states or phenotypes other than age, but there is covariance between these and age, then association may arise between age and the type of territory (despite no causal direct link between the two). For example, territory-specific occupation depending on prior dispersal is relatively common across taxa (Piper 2011). Specifically, in fragmented landscapes, sites have environmental 'edges', such as forest edges which provide an interface between internal woodland and the external environment (Saunders et al. 1991). In edge environments, there is often a higher incidence of individuals that disperse into the site (Krauss et al. 2003; Wilkin et al. 2007a). Dispersing between birth and first breeding, known as natal dispersal, generally involves greater distances than dispersal between two breeding attempts (Greenwood and Harvey 1982; Johst and Brandl 1999). Thus, individuals moving into a new environment are often younger (Greenwood 1980; Verhulst et al. 1997; Slagsvold and Wiebe 2018). In such cases, if edge territories are more commonly occupied by younger immigrant individuals, then this might provide a mechanism through which spatial age structure develops (figure 1C). Edge territories provide a particularly interesting mechanism by which spatial age structure is generated, not only because of immigrant-specific occupation, but also because they are often of poorer quality (Wilkin et al. 2007a; Purcell et al. 2012; Loveridge et al. 2017),

108 thus they might also directly generate spatial age structure if there is covariance between age and
109 territory quality (mechanisms 2a and 2b).

110




Figure 1 – Overview of mechanisms that might generate spatial age structure across territories referred to in the introduction. The theoretical populations consist of 200 individuals, each of which are

associated with a territory, illustrated across Wytham Woods, Oxford, as an example. Yellow and green territories are occupied by 'young' and 'old' individuals respectively. In (a), individuals occupy territories randomly, but at time t the population consists of a 1:1 ratio of old/young individuals; and at time $t + 1$ a 1:3 ratio of old/young individuals. In (b), the population consists of a 1:1 ratio of old/young individuals, but older individuals occupy the territories of highest quality. In (c), territories which do not border the edge are occupied by a 3:1 ratio of old/young individuals, whereas territories with an edge boundary are occupied by a ratio 1:3 of old/young individuals.

These mechanisms lead to *a priori* expectations that spatial age structure might develop across breeding territories; however, empirical evidence of this is currently sparse. Additionally, although these mechanisms might act concurrently on individuals in time, the areas in space that bias age-specific settlement may differ from each other. For example, if different aspects of territory quality are independent then the relative strength of these resource distributions, if any, in influencing age-specific settlement is unknown. Further, how such mechanisms bias age-specific settlement might be influenced by sex given the different roles that males and females will often play in territory acquisition (Brown 1964; Greenwood 1980). In short, given the role that age structure might play in spatially-structured demography, it is important that we advance our understanding on the evidence for spatial age structure and the mechanisms that drive it.

Here, we outline processes and their consequences for generating spatial age structure on long-term breeding data from a natural population of great tits *Parus major* in Wytham Woods, Oxford. We first assess whether there is covariance between territory quality and the age of individuals in such territories. Second, we evaluate whether there is non-random spatial arrangement of age within breeding seasons. Finally, we assess evidence for age-biases in space that persist across time compared to spatial structure in territory quality and reproductive output. We discuss the relative roles of spatial age structure versus structure in territory quality for producing spatially-variable demography,

and suggest that the framework presented here can be used across populations when considering spatial age structure.

(2) Methods

(i) Study system and data collection

The great tit *Parus major* is a socially monogamous passerine bird found in woodlands across Europe, with pairs defending territories during annual breeding seasons (Hinde 1952). The species has breeding ages ranging 1–9, averaging 1.8 years (Perrins 1979; Gosler 1993; Bouwhuis et al. 2009). Although there are some continuous physiological changes with age (Bouwhuis et al. 2009), the main age effects on individual-level traits are captured by two age-classes: first-years (hereafter yearlings) and older (hereafter adults, Greenwood et al. 1979; Perrins 1979; Sandell and Smith 1991; Gosler 1993; Farine et al. 2015). Specifically, behaviour related to dispersal and territory acquisition predominantly varies between yearlings and adults. Yearlings disperse relatively great distances from their natal origin to breeding sites (median distance 558m for males and 879m for females, Greenwood et al. 1979), and necessarily have no previous breeding site familiarity. In contrast, most adults that survive between breeding seasons retain their previous territories, with instances of breeding dispersal generally associated with previous unsuccessful breeding or divorce of a mate (Harvey et al. 1979a; Könczey et al. 1997; Culina et al. 2015). When breeding dispersal is undertaken, movements occur across smaller spatial scales compared to natal dispersal (median distance 50–143m, Harvey et al. 1979a), with no evidence that the spatial scale of dispersal changes continuously within the adult age class (Harvey et al. 1979a).

Data used here are from a long-term study in Wytham Woods, Oxford (51°46'N, 1°20'W), a 385ha mixed deciduous woodland surrounded by farmland (Savill et al. 2010). The tit population has been monitored since 1947, where breeding adults and their chicks have been marked with unique BTO

(British Trust for Ornithology) rings since the 1960s; and standard reproductive metrics are collected (Perrins 1965). Individuals breed almost exclusively in the 1026 nest-boxes which have remained in fixed positions with known GPS coordinates for the duration of the long-term study (Krebs 1971; Wilkin et al. 2006). The number of birds breeding in natural cavities is estimated to be very low, as the proportion of nest-boxes at which adult females are successfully identified (0.833) is very similar to the overall recapture probability using mark–recapture models (0.83 ± 0.07 ; Bouwhuis et al. 2012; Kidd et al. 2015). Additionally, periodic checks show that very few great tits breed in natural cavities, with no evidence that this is related to nest-box density, the number of which is in excess of the total breeding population (Greenwood et al. 1979; Perrins 1979; East and Perrins 1986). All chicks are ringed at 14-days of age, while adults are trapped at nest-boxes and identified by ring number, or marked with a new ring if they are immigrants. Age is based on year of hatching for local birds, or plumage characteristics for immigrants (Svensson 1992). Although immigration rates are high (46%), most are first caught as yearlings (78%) and can therefore be aged accurately.

(ii) Data selection

We used a dataset that assigned the year of hatching to all individuals between 1950–2022, across which exact age was calculated for 88.8% of 46062 identified breeding individuals (Woodman et al. 2022). In the present study, we included birds in analyses that attempted to breed between 1978–2022, for which data were more complete compared to earlier dates (figure S1). Individuals that were first caught post-fledging are assumed to be immigrants, as locally-hatched tits are marked as nestlings in nest-boxes and the proportion of birds hatched in natural cavities is very low (Kidd et al. 2015). Immigrants that entered the population with adult plumage were assigned a minimum age of 2, and subsequent age estimates were based on this (6.7% and 10.0% of breeding females and males). Age was therefore assigned for 68.7% of all breeding individuals where at least one egg was laid (due

to a combination of nests failing prior to adult trapping and unsuccessful trapping attempts, there are cases where the identity and therefore the age of parents is unknown).

(iii) Statistical analyses

Determining breeding territories

We defined annual breeding territories through a Dirichlet tessellation technique that forms Thiessen polygons (Rhynsburger 1973; Tanemura and Hasegawa 1980) around each occupied nest-box. The polygon includes all space within the habitat that is closer to the focal box than any other (with a boundary also imposed by the woodland edge). This metric of territory has been shown to be biologically meaningful in terms of territory size and territorial neighbours in tit species and is strongly related to other methods of calculating territories (Adams 2001; Wilkin et al. 2006; Schlicht et al. 2014; Firth and Sheldon 2016; Gokcekus et al. 2023). However, a limitation is that unrealistically large polygons are formed in areas where nest-boxes are placed at great distances from each other. We therefore capped territories at 2ha, which is a more realistic maximum scale at which individuals use space, as supported in previous analytical and field studies (Krebs 1971; Both and Visser 2000; Wilkin et al. 2006, 2007a).

Age and territory quality

We first assessed covariance between the age of individuals and their territories' quality. We measured territory quality through four measures: the number of oak trees *Quercus* spp. within 75m of the nest-box; average territory density; the edge distance index; and the long-term nest-box popularity index. Each of these is justified below.

Great tits predominantly provision offspring with caterpillars collected close to their nests (Gosler 1993), thus variation in caterpillar availability is directly linked to reproductive success (Perrins 1991; Keller and van Noordwijk 1994; Rytönen and Krams 2003). Caterpillars are found most abundantly on oak trees (Perrins 1991; Gosler 1993), therefore oak proximity, health and abundance is important for breeding success (Wilkin et al. 2006, 2009; Cole et al. 2021; Gokcekus et al. 2023). A radius of 75m was chosen as the abundance of oaks within this distance has been shown to be particularly important for breeding (Wilkin et al. 2007b; Hinks et al. 2015).

The density of conspecifics breeding in proximity may influence resource availability if foraging ranges overlap, and therefore territory density may also represent an aspect of territory quality. Additionally, territory density may affect site quality through social mechanisms, such as increased competition and emergent need for territory defence leading to reduced foraging (Ydenberg and Krebs 1987), or conversely mutual benefits between familiar neighbours (Grabowska-Zhang et al. 2012b, 2012a; Gokcekus et al. 2023). We calculated average territory density directly from the Thiessen polygon area produced from tessellation by taking the reciprocal of the mean polygon area.

Territories at woodland edges are associated with lower reproductive success in great tits (Wilkin et al. 2007a). Following Wilkin et al. (2007a), we defined the edge distance index (EDI) for each nest-box by multiplying the distance to forest edge by the proportion of woodland habitat within a 75m radius of the box. Thus, boxes within 75m of the edge have an EDI value in proportion to the amount of woodland habitat within this radius, therefore considering not only the distance to edge, but also the number and geometric arrangement of edges relative to each nest-box.

Finally, the frequency with which a territory is occupied in the long-term may provide a measure of quality, as individuals may choose sites that confer reproductive benefits more often, as evidenced in

other species (Sergio and Newton 2003; Smith and Moore 2005; Johnson 2007; Steenhof and Newton 2007; Potti et al. 2018). There is evidence of this in Wytham, where the number of times a nest-box has been occupied positively correlates with the average number of offspring that fledge per breeding attempt (figure S2). We therefore calculated the frequency of occupancy of each nest-box independent of individual breeding site fidelity by calculating the number of times a nest-box has been occupied since 1965 by a new breeding individual (i.e. attempts where either the female or male had previously used the same box were removed). However, nest-box occupation frequency is related to nest-box density (figure S3), because in areas of high density there are multiple unoccupied boxes which would likely be associated with the same territorial range if they were occupied (thus, in regions of high box density, birds may re-occupy the same territory over multiple years, but not necessarily the exact same box). To correct for this, we ran a linear model between the number of boxes within 30m of a focal nest-box (supporting information) and the number of times said box has been occupied by a new breeding individual, and took the residuals as the long-term nest-box popularity index.

We constructed a generalised linear mixed-effects model assuming a binomial error distribution to analyse the association between these four measures of territory quality and the age of the breeding individual. We modelled age (yearling/adult) as the response variable, with the territory quality measures as explanatory variables, which were z-transformed to compare their relative effects in predicting age (the explanatory variables were not highly correlated). Additionally, we modelled age as a continuous trait and assessed non-linear relationships between territory quality and age (supporting information). Individual ID, nest-box ID, and breeding year were included as random effects (table S1). We ran three sets of these models: one with all individuals; one with only females; and one with only males, allowing us to assess potential sex-specific differences in associations between aspects of territory quality and age. These models were also run to test for covariance between territory quality and residency status (supporting information). All models were conducted in R statistical software (R Core Team 2021) using the *lme4* package (Bates et al. 2015).

263

264 **Spatial age structure**

265 For each year's breeding population, we constructed an individual-by-individual matrix denoting
266 breeding neighbours i.e. a network of breeding territories, where nodes represent individuals and
267 edges represent the spatial connectivity of territories. Specifically, edges connected individuals if their
268 territories share a boundary from the tessellation technique and were weighted relative to the distance
269 between nest-boxes of neighbouring territories.

270

271 We created three networks per year: one with all individuals (but removing edges within breeding pairs
272 that occupy the same territory); one with only females; and one with only males. Edges connecting
273 individuals of unknown identity were removed. Across these networks, we calculated the assortativity
274 coefficient of age (yearling/adult), which measures the correlation between individuals' age and that
275 of their territorial neighbours accounting for edge weight (proximity of neighbouring nest-boxes) and
276 the relative proportion of the two age classes across the network. Through this technique, the age-
277 composition of neighbourhoods of birds that are likely to interact during territorial and foraging
278 behaviour contribute to the emergent quantitative signal of spatial age structure to a greater degree
279 than they would if spatial autocorrelation were calculated through pairwise distance of all individuals
280 across the system. Thus, this method allows us to assess evidence for spatial age structure at a
281 biologically relevant scale (analyses were also run treating age as a discrete and continuous trait, as
282 well as calculating assortment of residency status; supporting information). We also ran analyses to
283 compare spatial age structure with spatial structuring of territory quality and reproductive output. To
284 calculate territory quality structure, we assigned the value from the previously described four measures
285 of quality to each node (an occupied nest-box) and calculated the territory quality assortativity
286 coefficient across the network. For spatial reproductive output structure, we ran parallel analyses, but
287 calculated the assortativity in clutch size, chick number, fledgling number, and binary success (where

0 is no fledglings and 1 is at least one fledgling) associated with each nest-box. These analyses were run using the *assortnet* package (Farine 2014).

Temporal repeatability of spatial age structure

Finally, we tested whether spatial regions show temporal repeatability in age-composition, territory quality, and reproductive output. To assess this, within each year, we defined a radius around each occupied nest-box that corresponded to an area of 25, 50, 100, 150 and 200ha, representing neighbourhoods of breeding individuals of variable population sizes. Within each radius we calculated: the proportion of individuals that were adults; the mean number of oaks within 75m of the focal boxes; mean territory density; mean edge distance index; mean nest-box popularity index; mean clutch size; mean chick number; mean fledgling number; and proportion of boxes with binary success. We then calculated the same metrics for all nest-boxes outside of the focal radius. From this, we calculated a ratio index of within versus outside radius for each calculated measure, where a value of one represents the same annual average measure within and outside the radius. We then tested the repeatability of the ratio index associated with each spatial scale across the 45 years of data by calculating the intraclass correlation coefficient (ICC) derived from a linear mixed-effects model, where the response variable is the ratio index, and year and radius area are fitted as categorical grouping variables.

(3) Results

Age and territory quality

Across 20121 breeding individuals, representing 11167 breeding attempts, we found that higher density territories weakly predicted occupation by adults compared to yearlings (odds ratio = 1.094, 95% confidence intervals = 1.051–1.139, $p < 0.001$), with little evidence that the long-term popularity index, number of oaks within 75m or EDI were linked to age (figure 2; table S2). When assessing the

influence of territory quality on the age of individuals separately for both sexes, similar patterns were found, except territories that have a greater long-term popularity index predicted occupation by adult males (OR = 1.055, 95% CIs = 1.008–1.104, $p = 0.021$), but were not associated with female age (OR = 0.982, 95% CIs = 0.941–1.025, $p = 0.405$). Similar associations were found when assessing age as a continuous trait, with little evidence for non-linear relationships (supporting information; table S4; figures S6–8).




Figure 2 – Point estimates of odds ratios obtained from generalised linear mixed-effects models. The analysis examines the association between territory quality on the odds of the breeding individual being an adult. Each point corresponds to a specific level of the fixed effects (on the y-axis), and error bars denote 95% confidence intervals. Green points are from analysis assessing all individuals, purple are only females, and blue are only males.

Spatial age structure

Generally, across the entire sample, we found weak positive age assortment across breeding territories, whether considering the whole population (median r age assortativity coefficient = 0.014, 51% of annual standard errors overlap zero; figure 3), females (median r = 0.002; 58% SEs overlap zero), or males (median r = -0.004; 38% SEs overlap zero), although there was temporal variation with many individual years having little evidence for spatial age structure and some having significant disassortment (i.e. birds of similar age less likely to be associated). In contrast, there was much greater spatial structure in territory quality (oak abundance median r = 0.782; territory density median r = 0.645; EDI median r = 0.817; and long-term popularity index median r = 0.258, no annual SEs overlap zero). There was greater evidence for spatial reproductive output structure compared to age, but the signal of this structure is still relatively weak (clutch size median r = 0.059, 22% SEs overlap zero; chick number median r = 0.047, 29% SEs overlap zero; fledgling number median r = 0.075, 22% SEs overlap zero; binary success median r = 0.057, 24% SEs overlap zero; figure 3; table S5 for all results).




Figure 3 – Annual age and fledgling number assortativity coefficients (r) across territories. Large purple points represent age assortativity values where the standard error does not overlap zero, and large

green points are fledgling number assortativity values where standard error does not overlap zero. The dotted purple line denotes the median age assortativity value, and the green line denotes the median fledgling number assortativity value.

Temporal repeatability of spatial age structure

There was low temporal repeatability in the age-composition of spatial regions at all assessed scales 25–200ha (ICC range: 0.036–0.057; figure 4). In contrast, as would be expected, there is very high temporal repeatability in all assessed territory quality measures (figure 4). Compared to age-composition, there is higher temporal repeatability in average reproductive measures of neighbourhoods, with mean clutch size (ICC range: 0.234–0.447) displaying the highest repeatability (figure 4; table S6 for all results).




Figure 4 – Temporal repeatability of age-composition, territory quality and reproductive output of neighbourhoods at spatial scales of 25, 50, 100, 150 and 200ha. Repeatability was derived from the ICC from a linear mixed-effects model, where the ratio index of within versus outside the defined

spatial scale of the average measure was the response variable, and year and spatial scale area categorical grouping variables. Circles are sized and coloured on a gradient from red to blue as repeatability increases.

(4) Discussion

We quantified the extent of spatial age structure in a large long-term study population of territorial birds, and tested several mechanisms that could generate assortment by age in space. We show that, in general, spatial age structure is weak, despite the occurrence of spatial structure in habitat features, some of which covary with age and could plausibly generate spatial age structure. We also find that there is little temporal repeatability of spatial regions which bias certain age structures, despite stronger evidence for repeatability in spatial structure of reproductive output and territory quality. Here, we discuss these findings, suggesting that spatial structure in territory quality might be a greater contributor to spatially-variable demography and why there is not greater spatial age structure in this system.

Age and territory quality

Across all breeding individuals, we show that older individuals are more likely to acquire territories at higher densities, while there is a lack of evidence for a relationship between age and the abundance of oak trees or distance to edge. Considering females and males separately, we find similar associations between age and territory quality, except that the popularity index of a nest-box predicts occupation by adult males, but not adult females.

Our finding that territories at higher average densities are more likely to be occupied by adults than yearlings is notable, considering that this may result in reduced territory size, increased chance of

encounter with conspecifics during chick-rearing behaviour, and has previously been linked to lower reproductive success (Wilkin et al. 2006). This may relate to evidence that adults have a weaker response to density dependence in great tit populations, where yearling cohorts have a greater effect in reducing recruitment and survival, and more negatively respond to density dependence (Gamelon et al. 2016). Together with our results, this may reflect improved breeding competence of adults compared to yearlings leading to higher territorial densities. Specifically, older individuals may have increased foraging efficiency (Perrins and McCleery 1985), thus negating the need for large territories (indeed, individuals with smaller foraging ranges can be more efficient with higher reproductive success, Cole et al. 2012), therefore generating higher breeding densities among adults and a weaker response to density dependence. Additionally, increased familiarity between adult territorial neighbours may reduce competition associated with foraging (Grabowska-Zhang et al. 2012b; Gokcekus et al. 2023), or even promote joint benefits such as nest defence (Grabowska-Zhang et al. 2012a), thus favouring closer breeding between adults. Recent advances in tracking technology of passerine foraging behaviour (Baldan and van Loon 2022) might advance insight into whether competition is reduced among older familiar neighbours by assessing evidence for increased coordination during foraging compared to unfamiliar neighbours. Further, given the relationship between average territorial density and age, and the fact that overall breeding population size varies markedly between years (annual range of pairs = 115–482), it would be interesting to take our findings further to assess demographic feedbacks when considering changes in population-level breeding density and age structure. While this is not within the scope of the current study, future work could disentangle the relative roles of variation in overall population size, age structure and age-related breeding behaviour in producing interannual changes in population-level territorial breeding density.

Territories that were more popular in the long-term were more likely to be occupied by adult males than yearlings, but this was not the case for females or when considering all individuals together. The use of frequency of occupancy as a measure of territory quality is well-established in avian systems

(Sergio and Newton 2003; Smith and Moore 2005; Johnson 2007; Steenhof and Newton 2007; Potti et al. 2018), and may capture information on cues of quality which individuals respond to in the long-term which cannot be directly measured. The association between more popular territories and adults in males but not females may be explained by sex differences, where male great tits are more important in initial acquisition and defence of territories (Krebs 1971, 1977; Perrins 1979; Björklund et al. 1989), which is widespread in other avian (Brown 1964; Stutchbury 1991; Beletsky 1992; Eikenaar et al. 2009) and taxonomic groups (Clutton-Brock 1989; López and Martín 2002; Milner et al. 2010). Thus, this could suggest that older males with greater competitive ability (Sandell and Smith 1991) outcompete younger individuals to occupy higher quality sites. This may be driven either by novel acquisition of high-quality territories, or by persistence and defence of a territory between years (Greenwood et al. 1979; Harvey et al. 1979a). In the latter case, males may occupy a high-quality territory in their first breeding attempt, which supports successful breeding, and then retain this territory in future breeding. Thus, if survival is high between years, particularly among individuals in high-quality territories, then covariance between male age and territory quality may arise as younger first-year breeders are forced into low-quality territories.

There is little evidence that the abundance of oaks close to the nest predicts age. Oak abundance at this scale has been linked to great tit breeding success (Wilkin et al. 2006, 2007b, 2009; Hinks et al. 2015; Cole et al. 2021; Gokcekus et al. 2023), thus increased experience and competitive ability of adults (Sandell and Smith 1991) might lead to the assumption that older individuals will acquire territories with more oaks. However, although higher oak abundance might provide better quality territories when considering entire populations, little is known about how this interacts with phenological synchrony at the individual-level (Hinks et al. 2015) and how the degree of phenological matching depends on age. Specifically, older great tits typically breed earlier (Harvey et al. 1979b; McCleery and Perrins 1998), possibly due to improved breeding competence leading to greater synchrony between chick-rearing and peak caterpillar abundance. Further, the rate at which

caterpillars are provisioned to chicks strongly correlates with the caterpillar biomass available (Naef-Daenzer et al. 2000). Thus, if a greater proportion of yearlings mistime their breeding compared to adults, then they may be unable to monopolise caterpillar resources even if in oak-rich territories due to temporal mismatch. In addition to greater phenological synchrony, older birds may improve their foraging ability in a broader sense through greater site familiarity from previous occupation (Perrins and McCleery 1985; Piper 2011; Slagsvold and Wiebe 2018), utilisation of a wider variety of food resources (Gosler 1993), and improved foraging coordination with pair-bonded partners (Mariette and Griffith 2015; Griffith 2019). In this sense, the overall abundance of oaks may not be a crucial cue when acquiring territories, rather there is stronger selection on phenological matching to the local environment (Hinks et al. 2015; Cole et al. 2021) and general foraging ability. Additionally, there is little evidence that the edge distance index is associated with age, which has been linked to reduced reproductive output (Wilkin et al. 2007a), and thus there is no apparent age-specificity in how edge effects mediate territory acquisition.

The lack of an effect of oak abundance and edge distance on age-specific territory occupation, as well as the general weak association with other territory attributes, may suggest that territory acquisition in adults is predominantly driven by retaining previous territories in which they successfully bred, regardless of quality. This may be driven by a relative lack of available territories compared to overall population size, thus providing strong selection for the retention of a previously occupied territory (Clark 1994; Harts et al. 2016). High breeding area fidelity is evident in great tits (Harvey et al. 1979a; Könczey et al. 1997; Firth et al. 2018). In this case, the cue which individuals respond to when acquiring a territory may be experience of previous breeding success, as opposed to current environmental quality cues. Consequently, any covariance between older age and high territory quality might break down through two mechanisms: either previous breeding occurred in a low-quality territory, but breeding was successful and thus the territory is retained as an adult; or previous breeding occurred in a high-quality territory, but the attempt was unsuccessful, and thus individuals

move to new poorer quality territories in future years. Breeding success in poor quality territories can regularly occur, for example if intraspecific competition is low, or when individual or partner quality is high. In such cases, older individuals might utilise their higher competitive ability to defend territories in which they have had success regardless of quality (Harvey et al. 1979a; Yasukawa 1979; de Vos 1983). Conversely, unsuccessful breeding may happen in areas of high-quality, through stochastic environmental effects (such as extreme weather or predation), or low individual or partner quality. In these cases, despite being in high-quality territories, pairs may divorce in an attempt to improve future breeding (Culina et al. 2015) and disperse to new territories (Harvey et al. 1979a). This also relates to a broader mechanism which might mask covariation between age and territory quality, whereby there are interannual short-term changes in site quality not captured within this study. For example, shifts in predator space-use across years can lead to high brood mortality in specific regions (Perrins 1965; Dunn 1977). This might induce dispersal away from such areas by adult individuals between breeding attempts, leading to covariation between age and short-term, but not necessarily long-term, territory quality, thus leaving the unoccupied territories for yearlings to acquire.

Given our findings, future research should aim to disentangle the relative effects of territory quality and previous reproductive success on site fidelity in territorial species, as well as assessing behavioural differences in the degree of territory defence in individuals that re-occupy sites of varying quality. Further, examining age-specific behaviour following divorce and acquisition of a new territory might shed light on how age relates to territory quality when removing the effect of site fidelity. For instance, there may be variation in territory quality across divorced individuals that newly settle depending on their own age and that of their new partner. Finally, more work should assess age-specificity in territory acquisition while considering non-breeding individuals that fail to acquire a territory. For example, in great tit populations, there are often more individuals than available territories (Perrins 1979), leaving non-breeders in the population during the spring. In avian species, non-breeders tend to be younger (Stutchbury 1991; Sergio et al. 2009) and the ability to defend a territory against intrusions from non-

breeders increases with age (Arcese 1987). Thus, experiments to identify non-breeders during territorial activity (Firth et al. 2018) and accounting for this cohort might reveal further age-specificity in territory quality.

Spatial age structure

Our study reveals more frequent positive annual spatial age structure, but this structure is weak and often absent, with no clear sex differences. In contrast, there is strong evidence for spatial territory quality structure, with particularly strong assortment in oak abundance, territory density and edge distance (the latter of which is spatially necessitated), and moderate assortment in the long-term popularity index of a nest-box. There is greater evidence for spatial structure in reproductive output compared to age, with fledgling number displaying the greatest structure.

It is interesting that there is not greater spatial age structure not only due to our evidence of covariance between some attributes of territory quality and age, and spatial structure in such territory quality, but also due to social mechanisms that might generate spatial age structure. Winter social associations in great tits are positively age-assorted, albeit fairly weakly (Farine et al. 2015), and social connections carryover into breeding as individuals establish territories closer to previous social associates (Firth and Sheldon 2016). This is likely because having familiar neighbours can reduce the need for territory defence, increase cooperation, and enhance reproductive success (Grabowska-Zhang et al. 2012b, 2012a; Gokcekus et al. 2023). Thus, because older individuals are more likely to be familiar with their neighbours than younger ones (Gokcekus et al. 2023), this might be expected to generate spatial age structure through clusters of older individuals breeding in proximity. However, although this mechanism provides direct benefits at the dyadic-level for closer breeding between adults, there is little overall spatial age structure. This may be partly explained by the species' life-history, where there is high annual mortality (~52%, Bouwhuis et al. 2009). Together with annual influxes of immigrant

birds, this means that the breeding population is often composed of a large proportion of individuals that are first-time breeders (46–83%, Woodman et al. 2022), thus only making it possible for a small proportion of the population to have familiar neighbours. In short, despite benefits for older breeders to retain previous neighbours, high mortality rates lead to a high chance that at least some previous territorial associates will have died, and high competition for breeding sites mean these will be acquired by new and likely yearling breeders (Krebs 1977; Perrins 1979; Björklund et al. 1989), thus reducing overall spatial age structure. This hypothesis should be tested through comparative analyses using the framework developed here by assessing whether there is greater spatial age structure in territorial species with lower mortality rates and thus a greater opportunity to retain previous neighbours, particularly considering familiarity between neighbours is adaptive in many taxonomic groups (Temeles 1994; Frostman and Sherman 2004; Siracusa et al. 2021).

Additionally, although there are benefits of retaining familiar neighbours, this is not necessarily underpinned by an active process where older individuals select territories based on closer breeding to previous associates. Instead, it may be underpinned by a passive process where individuals which survive between years occupy the same locality (Harvey et al. 1979a), leading to the incidental retention of familiar neighbours that also survive and have high site fidelity. Such a passive mechanism is consistent with recent findings that age-assortative mating within pairs of great tits, despite being adaptive (Harvey et al. 1979b), is underpinned by passive processes of pair retention and fluctuations in overall population age structure as opposed to active age-related selection (Woodman et al. 2022). Thus, even if there are direct advantages of retaining familiar neighbours, if it is not underpinned by active processes, then this might weaken its prevalence and therefore overall spatial age structure. Further, closer breeding between non-familiar individuals of different ages may be adaptive in some cases, thus decreasing positive age assortment across territories. For example, although great tits are socially monogamous, approximately 50% of broods contain at least one extra-pair young (Patrick et al. 2012; Firth et al. 2015). There is some evidence that extra-pair copulations are age-related in great

tits, where yearlings are more likely to be cuckolded and have a higher proportion of extra-pair young in their nests than older males (Lubjuhn et al. 2007), potentially related to lower social dominance (Sandell and Smith 1991). Thus, this might provide a mechanism where older male great tits opt to breed closer to subdominant yearlings to gain extra-pair copulations, which are highly structured in space (Roth et al. 2019).

Finally, although there is temporal overlap in when territories are used, there is variation between pairs which will affect how neighbours interact depending on their breeding timing synchrony, thus affecting spatial age structure at any single point in time. Firstly, there is spatial variation in the timing of food availability at a scale relevant to individual territories, underpinned by variation in individual tree phenology (Cole and Sheldon 2017), and this local variation predicts breeding timing in individual pairs (Hinks et al. 2015). Thus, pairs with neighbouring territories may not interact much during chick-rearing behaviour if neighbouring phenology is highly asynchronous (although there is generally spatial structure in breeding phenology across Wytham, Wilkin et al. 2007b). Second, there might be even greater temporal mismatch in occupation of neighbouring territories when considering initial acquisition and defence behaviour, which begins in January or even earlier, months before chick-rearing (Krebs 1977, 1978, 1982). Thus, interactions between individuals and competition associated with territory defence will be variable depending on the occupation status of neighbouring territories at the time of focal territory acquisition. For example, there may be extended periods of territorial activity pre-breeding where neighbouring pairs interact, but then a new pair may acquire a territory which bisects such breeding sites, thus breaking down the territorial boundary between the two original neighbouring pairs. Previous research in this system has assessed age-specificity in nest-site visitation patterns prior to breeding in terms of the number and spatial clustering of nest-boxes visited (Firth et al. 2018). Thus, future work could incorporate such techniques with advances in the tracking of movement (Levin et al. 2015; Baldan and van Loon 2022) and other defence behaviour (Merino Recalde et al. 2023) to advance our understanding on the temporal scale at which individuals interact during territorial

behaviour before breeding, how this is affected as territories are gradually occupied prior to the spring, and age-specific variation in this.

Temporal repeatability of spatial age structure

We find low repeatability in the age-composition of spatial regions between years compared to repeatability in spatial territory quality and reproductive output structure. High repeatability in spatial territory quality structure is expected, as annual variation in average quality measures across spatial regions is changed only depending on which nest-boxes are occupied (i.e. if the same boxes were occupied every year, there would be 100% repeatability). However, there is variation in which boxes are used each year. Thus, in conjunction with our previous finding of high within-year spatial territory quality structure, it is interesting then that we report a higher repeatability in the average reproductive output compared to the age-composition of spatial regions. This suggests that territory quality in this system may be more important in driving spatially-variable demography than local age structure. Additionally, the finding of low repeatability in spatial age-composition is consistent with our suggestion that adult occupation is likely largely driven by persistence in previous territories of varying quality as opposed to active acquisition of high-quality territories (the latter would lead to higher repeatability as older birds would frequently use higher quality sites independent of year). Thus, future studies could use a similar framework presented here to examine the relative importance of spatial age structure and spatial autocorrelation in territory quality in producing repeatable spatially-variable demography. Specifically, we may expect that for longer-lived species breeding in more homogenous environments, spatial age structure may be more repeatable across years and contribute more to spatial structure in reproductive output than territory quality.

(5) Conclusions

Our study highlights association between age and aspects of territory quality in a territorial passerine species, specifically showing that older individuals breed at higher densities and, depending on sex, in sites which are more popular in the long-term. However, we show limited evidence of spatial age structure. We suggest that age-specific occupation of territories is likely driven by persistence in territories across lifespan (largely irrespective of quality), leading to passive retention of some older neighbours which also survive between years. We emphasise the importance of exploring the operation of these mechanisms that might lead to spatial age structure in species with different life-histories. Specifically, it might be expected that in species with lower mortality rates, greater spatial age structure would be generated through passive mechanisms of territorial neighbour retention as more individuals survive between years. The observed patterns of higher spatial territory quality and reproductive output structure (compared to spatial age structure), in addition to greater temporal repeatability in these, contributes valuable insights into the drivers of spatially-variable demography. Namely, given these results, we suggest that spatial clustering of territory quality is likely to be more important in determining repeatable spatial structure in reproductive output compared to spatial age structure.

References

- Adams, E. S. 2001. Approaches to the study of territory size and shape. *Annual Review of Ecology and Systematics* 32:277–303.
- Albery, G. F., T. H. Clutton-Brock, A. Morris, S. Morris, J. M. Pemberton, D. H. Nussey, and J. A. Firth. 2022. Ageing red deer alter their spatial behaviour and become less social. *Nature Ecology and Evolution* 6:1231–1238.
- Arcese, P. 1987. Age, intrusion pressure and defence against floaters by territorial male song sparrows. *Animal Behaviour* 35:773–784.
- Baldan, D., and E. E. van Loon. 2022. Songbird parents coordinate offspring provisioning at fine spatio-temporal scales. *Journal of Animal Ecology* 1316–1326.
- Bates, D., M. Mächler, B. M. Bolker, and S. C. Walker. 2015. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67.
- Beletsky, L. D. 1992. Social Stability and Territory Acquisition in Birds. *Behaviour* 123:290–313.
- Björklund, M., B. Westman, and K. Allander. 1989. Song in Swedish Great Tits: Intra-or Intersexual Communication? *Behaviour* 111:257–269.
- Both, C., and M. E. Visser. 2000. Breeding territory size affects fitness: An experimental study on competition at the individual level. *Journal of Animal Ecology* 69:1021–1030.
- Bouwhuis, S., R. Choquet, B. C. Sheldon, and S. Verhulst. 2012. The forms and fitness cost of senescence: Age-specific recapture, survival, reproduction, and reproductive value in a wild bird population. *American Naturalist* 179.
- Bouwhuis, S., B. C. Sheldon, S. Verhulst, and A. Charmantier. 2009. Great tits growing old: Selective disappearance and the partitioning of senescence to stages within the breeding cycle. *Proceedings of the Royal Society B: Biological Sciences* 276:2769–2777.

629 Brewer, S. K., R. J. DiStefano, and C. F. Rabeni. 2009. The influence of age-specific habitat
630 selection by a stream crayfish community (*Orconectes* spp.) on secondary production. *Hydrobiologia*
631 619:1–10.

632 Brown, J. L. 1964. The Evolution of Diversity in Avian Territorial Systems. *The Wilson Bulletin*
633 76:160–169.

634 Clark, C. W. 1994. Antipredator behavior and the asset-protection principle. *Behavioral Ecology*
635 5:159–170.

636 Clutton-Brock, T. H. 1989. Mammalian Mating Systems. *Proceedings of the Royal Society B:*
637 *Biological Sciences* 236:339–372.

638 Cole, E. F., J. Morand-Ferron, A. E. Hinks, and J. L. Quinn. 2012. Cognitive ability influences
639 reproductive life history variation in the wild. *Current Biology* 22:1808–1812.

640 Cole, E. F., C. E. Regan, and B. C. Sheldon. 2021. Spatial variation in avian phenological response
641 to climate change linked to tree health. *Nature Climate Change* 11:872–878.

642 Cole, E. F., and B. C. Sheldon. 2017. The shifting phenological landscape: Within- and between-
643 species variation in leaf emergence in a mixed-deciduous woodland. *Ecology and Evolution* 7:1135–
644 1147.

645 Coulson, T., J. M. Gaillard, and M. Festa-Bianchet. 2005. Decomposing the variation in population
646 growth into contributions from multiple demographic rates. *Journal of Animal Ecology* 74:789–801.

647 Culina, A., R. Radersma, and B. C. Sheldon. 2015. Trading up: The fitness consequences of divorce
648 in monogamous birds. *Biological Reviews* 90:1015–1034.

649 Davies, N. B., J. R. Krebs, and S. A. West. 2012. *An introduction to behavioural ecology*. John Wiley
650 & Sons.

651 de Vos, G. J. 1983. Social behaviour of black grouse. *Ardea* 71:1–103.

652 Dunn, E. 1977. Predation by Weasels (*Mustela nivalis*) on Breeding Tits (*Parus* Spp.) in Relation to
653 the Density of Tits. *Journal of Animal Ecology* 46:633–652.

654 East, M. L. and C. M. Perrins. 1986. The effect of nestboxes on breeding populations of birds in
655 broadleaved temperate woodlands. *Ibis* 130: 393–401.

656 Eikenaar, C., D. S. Richardson, L. Brouwer, R. Bristol, and J. Komdeur. 2009. Experimental
657 evaluation of sex differences in territory acquisition in a cooperatively breeding bird. *Behavioral*
658 *Ecology* 20:207–214.

659 Farine, D. R. 2014. Measuring phenotypic assortment in animal social networks: Weighted
660 associations are more robust than binary edges. *Animal Behaviour* 89:141–153.

661 Farine, D. R., J. A. Firth, L. M. Aplin, R. A. Crates, A. Culina, C. J. Garroway, C. A. Hinde, et al.
662 2015. The role of social and ecological processes in structuring animal populations: A case study
663 from automated tracking of wild birds. *Royal Society Open Science* 2.

664 Ferrer, M., and I. Bisson. 2003. Age and territory-quality effects on fecundity in the Spanish Imperial
665 Eagle (*Aquila adalberti*). *Auk* 120:180–186.

666 Firth, J. A., J. D. Hadfield, A. W. Santure, J. Slate, and B. C. Sheldon. 2015. The influence of
667 nonrandom extra-pair paternity on heritability estimates derived from wild pedigrees. *Evolution*
668 69:1336–1344.

669 Firth, J. A., and B. C. Sheldon. 2016. Social carry-over effects underpin trans-seasonally linked
670 structure in a wild bird population. *Ecology Letters* 19:1324–1332.

671 Firth, J. A., B. L. Verhelst, R. A. Crates, C. J. Garroway, and B. C. Sheldon. 2018. Spatial, temporal
672 and individual-based differences in nest-site visits and subsequent reproductive success in wild
673 great tits. *Journal of Avian Biology* 49:1–11.

674 Fowler, A. J., L. McLeay, and D. A. Short. 2000. Spatial variation in size and age structures and
675 reproductive characteristics of the King George whiting (*Percoidei* : *Sillaginidae*) in South Australian
676 waters. *Marine and Freshwater Research* 51:11–22.

677 Frostman, P., and P. T. Sherman. 2004. Behavioral response to familiar and unfamiliar neighbors in
678 a territorial cichlid, *Neolamprologus pulcher*. *Ichthyological Research* 51:283–285.

679 Gamelon, M., V. Grøtan, S. Engen, E. Bjørkvoll, M. E. Visser, and B. E. Sæther. 2016. Density
680 dependence in an age-structured population of great tits: Identifying the critical age classes. *Ecology*
681 97:2479–2490.

682 Gamelon, M., S. J. G. Vriend, S. Engen, F. Adriaensen, A. A. Dhondt, S. R. Evans, E. Matthysen, et
683 al. 2019. Accounting for interspecific competition and age structure in demographic analyses of
684 density dependence improves predictions of fluctuations in population size. *Ecology Letters* 22:797–
685 806.

686 Garneau, D. E., T. Boudreau, M. Keech, and E. Post. 2008. Habitat use by black bears in relation to
687 conspecifics and competitors. *Mammalian Biology* 73:48–57.

688 Gokcekus, S., J. A. Firth, C. Regan, E. F. Cole, B. C. Sheldon, and G. F. Albery. 2023. Social
689 Familiarity and Spatially Variable Environments Independently Determine Reproductive Fitness in a
690 Wild Bird. *American Naturalist* 201:813–824.

691 Gosler, Andrew. 1993. The great tit . Hamlyn species guides. Hamlyn, London.

692 Grabowska-Zhang, A. M., B. C. Sheldon, and C. A. Hinde. 2012a. Long-term familiarity promotes
693 joining in neighbour nest defence. *Biology Letters* 8:544–546.

694 Grabowska-Zhang, A. M., T. A. Wilkin, and B. C. Sheldon. 2012b. Effects of neighbor familiarity on
695 reproductive success in the great tit (*Parus major*). *Behavioral Ecology* 23:322–333.

696 Greenwood, P. J. 1980. Mating systems, philopatry and dispersal in birds and mammals. *Animal*
697 *Behaviour* 28:1140–1162.

698 Greenwood, P. J., and P. H. Harvey. 1982. The natal and breeding dispersal of birds. Annual
699 Review of Ecology and Systematics 13:1–21.

700 Greenwood, P. J., P. H. Harvey, and C. M. Perrins. 1979. The role of dispersal in the great tit (*Parus*
701 major): The causes, consequences and heritability of natal dispersal. *Journal of Animal Ecology*
702 48:123–142.

703 Griffith, S. C. 2019. Cooperation and Coordination in Socially Monogamous Birds: Moving Away
704 From a Focus on Sexual Conflict. *Frontiers in Ecology and Evolution* 7.

705 Groenendijk, J., F. Hajek, C. Schenck, E. Staib, P. J. Johnson, and D. W. Macdonald. 2015. Effects
706 of territory size on the reproductive success and social system of the giant otter, south-eastern Peru.
707 *Journal of Zoology* 296:153–160.

708 Harts, A. M. F., K. Jaatinen, and H. Kokko. 2016. Evolution of natal and breeding dispersal: When is
709 a territory an asset worth protecting? *Behavioral Ecology* 27:287–294.

710 Harvey, P. H., P. J. Greenwood, and C. M. Perrins. 1979a. Breeding area fidelity of great tits (*Parus*
711 major). *Journal of Animal Ecology* 48:305–313.

712 Harvey, P. H., P. J. Greenwood, C. M. Perrins, and A. R. Martin. 1979b. Breeding success of great
713 tits *Parus major* in relation to age of male and female parent. *Ibis* 121:216–219.

714 Hinde, R. A. 1952. The Behaviour of the Great Tit (*Parus major*) and Some Other Related Species.
715 *Behaviour* 2:1–201.

716 ———. 1956. Biological significance of the territories of birds. *Ibis* 98.

717 Hinks, A. E., E. F. Cole, K. J. Daniels, T. A. Wilkin, S. Nakagawa, and B. C. Sheldon. 2015. Scale-
718 dependent phenological synchrony between songbirds and their caterpillar food source. *American*
719 *Naturalist* 186:84–97.

720 Hoy, S. R., D. R. MacNulty, D. W. Smith, D. R. Stahler, X. Lambin, R. O. Peterson, J. S. Ruprecht, et
 721 al. 2020. Fluctuations in age structure and their variable influence on population growth. *Functional*
 722 *Ecology* 34:203–216.

723 Johnson, M. D. 2007. Measuring habitat quality: A review. *Condor* 109:489–504.

724 Johst, K., and R. Brandl. 1999. Natal versus breeding dispersal: Evolution in a model system.
 725 *Evolutionary Ecology Research* 1:911–921.

726 Jones, J. A., M. R. Harris, and L. Siefferman. 2014. Physical habitat quality and interspecific
 727 competition interact to influence territory settlement and reproductive success in a cavity nesting
 728 bird. *Frontiers in Ecology and Evolution* 2:1–8.

729 Keller, L. F., and A. J. van Noordwijk. 1994. Effects of local environmental conditions on nestling
 730 growth in the great tit *Parus major* L. *Ardea* 82:349–362.

731 Kerbiriou, C., F. Gourmelon, F. Jiguet, I. Le Viol, F. Bioret, and R. Julliard. 2006. Linking territory
 732 quality and reproductive success in the Red-billed Chough *Pyrrhocorax pyrrhorax*: implications for
 733 conservation management of an endangered population. *Ibis* 148:352–364.

734 Kidd, L. R., B. C. Sheldon, E. G. Simmonds, and E. F. Cole. 2015. Who escapes detection?
 735 Quantifying the causes and consequences of sampling biases in a long-term field study. *Journal of*
 736 *Animal Ecology* 84:1520–1529.

737 Kokko, H., A. López-Sepulcre, and L. J. Morrell. 2006. From hawks and doves to self-consistent
 738 games of territorial behavior. *American Naturalist* 167:901–912.

739 Könczey, R., L. Tóth, and J. Török. 1997. Site fidelity of great and blue tits in the Pilis-Visegrád
 740 mountains. *Opuscula Zoologica*.

741 Krauss, J., I. Steffan-Dewenter, and T. Tschamtkke. 2003. Local species immigration, extinction, and
 742 turnover of butterflies in relation to habitat area and habitat isolation. *Oecologia* 137:591–602.

743 Krebs, J. R. 1971. Territory and Breeding Density in the Great Tit, *Parus Major* L. *Ecology* 52:2–22.

744 ———. 1977. Song and territory in the great tit *Parus major*. in B. Stonehouse and C. M. Perrins,
745 eds. *Evolutionary Ecology*. MacMillan, London.

746 ———. 1978. Song repertoires and territory defence in the great tit. *Nature* 271:539–542.

747 ———. 1982. Territorial Defence in the Great Tit (*Parus major*): Do Residents Always Win?
748 *Behavioral Ecology and Sociobiology* 11:185–194.

749 Levin, I. I., D. M. Zonana, J. M. Burt, and R. J. Safran. 2015. Performance of encounter net tags:
750 Field tests of miniaturized proximity loggers for use on small birds. *PLoS ONE* 10.

751 López, P., and J. Martín. 2002. Effects of Female Presence on Intrasexual Aggression in Male
752 Lizards, *Podarcis hispanicus*. *Aggressive Behavior: Official Journal of the International Society for*
753 *Research on Aggression* 28:491–498.

754 Loveridge, A. J., M. Valeix, N. B. Elliot, and D. W. Macdonald. 2017. The landscape of
755 anthropogenic mortality: how African lions respond to spatial variation in risk. *Journal of Applied*
756 *Ecology* 54:815–825.

757 Lubjuhn, T., T. Gerken, J. Brün, and T. Schmoll. 2007. Yearling male great tits, *Parus major*, suffer
758 more strongly from cuckoldry than older males. *Zoology* 110:387–397.

759 Mariette, M. M., and S. C. Griffith. 2015. The adaptive significance of provisioning and foraging
760 coordination between breeding partners. *The American Naturalist* 185.

761 McCleery, R. H., and C. M. Perrins. 1998. ...temperature and egg-laying trends. *Nature* 391:30–31.

762 McIlwain, J. L., M. R. Claereboudt, H. S. Al-Oufi, S. Zaki, and J. S. Goddard. 2005. Spatial variation
763 in age and growth of the kingfish (*Scomberomorus commerson*) in the coastal waters of the
764 Sultanate of Oman. *Fisheries Research* 73:283–298.

765 Merino Recalde, N. 2023. pykanto: A python library to accelerate research on wild bird song.
766 *Methods in Ecology and Evolution* 14:1994–2002.

767 Milner, R. N. C., I. Booksmythe, M. D. Jennions, and P. R. Y. Backwell. 2010. The battle of the
 768 sexes? Territory acquisition and defence in male and female fiddler crabs. *Animal Behaviour*
 769 79:735–738.

770 Naef-Daenzer, L., B. Naef-Daenzer, and R. G. Nager. 2000. Prey selection and foraging
 771 performance of breeding Great Tits *Parus major* in relation to food availability. *Journal of Avian*
 772 *Biology* 31:206–214.

773 Patrick, S. C., J. R. Chapman, H. L. Dugdale, J. L. Quinn, and B. C. Sheldon. 2012. Promiscuity,
 774 paternity and personality in the great tit. *Proceedings of the Royal Society B: Biological Sciences*
 775 279:1724–1730.

776 Perrins, C. M. 1965. Population fluctuations and clutch-size in the great tit, *Parus major* L. *Journal of*
 777 *Animal Ecology* 34:601–647.

778 ———. 1979. *British tits* (1st ed.). Collins, London.

779 Perrins, C. M. 1991. Tits and their caterpillar food supply. *Ibis* 133:49–54.

780 Perrins, C. M., and R. H. McCleery. 1985. The effect of age and pair bond on the breeding success
 781 of Great Tits *Parus major*. *Ibis* 127:306–315.

782 Piper, W. H. 1997. Social Dominance in Birds. Pages 125–187 in V. Nolan, E. D. Ketterson, and C.
 783 F. Thompson, eds. *Current Ornithology*. Springer US, Boston, MA.

784 Piper, W. H. 2011. Making habitat selection more “familiar”: A review. *Behavioral Ecology and*
 785 *Sociobiology* 65:1329–1351.

786 Potti, J., C. Camacho, D. Canal, and J. Martinez-Padilla. 2018. Long-term occupancy of nest boxes
 787 as a measure of territory quality for Pied Flycatchers. *Journal of Field Ornithology* 89:337–347.

788 Purcell, K. L., C. M. Thompson, and D. A. Drynan. 2012. Response of Brown Creepers to elevation
 789 and forest edges in the southern Sierra Nevada, California. *Condor* 114:185–196.

790 R Core Team. 2021. R: a language and environment for statistical computing. Vienna: R Foundation
 791 for Statistical Computing.

792 Reynolds, J. D. 1996. Animal breeding systems. *Trends in Ecology and Evolution* 11:68–72.

793 Rhynsburger, D. 1973. Analytic Delineation of Thiessen Polygons. *Geographical Analysis* 5:133–
 794 144.

795 Roth, A. M., J. A. Firth, S. C. Patrick, E. F. Cole, and B. C. Sheldon. 2019. Partner's age, not social
 796 environment, predicts extrapair paternity in wild great tits (*Parus major*). *Behavioral Ecology*
 797 30:1782–1793.

798 Rytönen, S., and I. Krams. 2003. Does foraging behaviour explain the poor breeding success of
 799 great tits *Parus major* in northern Europe? *Journal of Avian Biology* 34:288–297.

800 Sandell, M., and H. G. Smith. 1991. Dominance, prior occupancy, and winter residency in the great
 801 tit (*Parus major*). *Behavioral Ecology and Sociobiology* 29:147–152.

802 Saunders, D. A., R. J. Hobbs, and C. R. Margules. 1991. Biological Consequences of Ecosystem
 803 Fragmentation: A Review. *Conservation Biology* 5:18–32.

804 Savill, P., C. Perrins, K. Kirby, and N. Fisher. 2010. Wytham Woods: Oxford's ecological laboratory.
 805 OUP Oxford.

806 Schlicht, L., M. Valcu, and B. Kempenaers. 2014. Thiessen polygons as a model for animal territory
 807 estimation. *Ibis* 156:215–219.

808 Sergio, F., J. Blas, and F. Hiraldo. 2009. Predictors of floater status in a long-lived bird: A cross-
 809 sectional and longitudinal test of hypotheses. *Journal of Animal Ecology* 78:109–118.

810 Sergio, F., and I. Newton. 2003. Occupancy as a Measure of Territory Quality. *Journal of Animal*
 811 *Ecology* 72:857–865.

812 Siracusa, E. R., S. Boutin, B. Dantzer, J. E. Lane, D. W. Coltman, and A. G. McAdam. 2021.
813 Familiar Neighbors, but Not Relatives, Enhance Fitness in a Territorial Mammal. *Current Biology*
814 31:438-445.e3.

815 Siracusa, E. R., A. S. Pereira, J. B. Brask, J. E. Negron-Del Valle, D. Phillips, M. L. Platt, J. P.
816 Higham, et al. 2023. Ageing in a collective: The impact of ageing individuals on social network
817 structure. *Philosophical Transactions of the Royal Society B: Biological Sciences* 378.

818 Slagsvold, T., and K. L. Wiebe. 2018. Immigrants and locally recruited birds differ in prey delivered
819 to their offspring in blue tits and great tits. *Animal Behaviour* 139:127–135.

820 Smith, R. J., and F. R. Moore. 2005. Arrival timing and seasonal reproductive performance in a long-
821 distance migratory landbird. *Behavioral Ecology and Sociobiology* 57:231–239.

822 Stearns, S. C. 1992. *The evolution of life histories*. Oxford University Press, Oxford.

823 Steenhof, K., and I. Newton. 2007. Assessing nesting success and productivity. *Raptor research and*
824 *management techniques*.

825 Stepanuk, J., E. Heywood, J. Lopez, R. DiGiovanni Jr, and L. Thorne. 2021. Age-specific behavior
826 and habitat use in humpback whales: implications for vessel strike. *Marine Ecology Progress Series*
827 663:209–222.

828 Stutchbury, B. J. 1991. Floater behaviour and territory acquisition in male purple martins. *Animal*
829 *Behaviour* 42:435–443.

830 Svensson, L. 1992. *Identification guide to European passerines* (4th, revd. ed.). Lars Svensson ;
831 British Trust for Ornithology, Stockholm : Thetford.

832 Tanemura, M., and M. Hasegawa. 1980. Geometrical models of territory I. Models for synchronous
833 and asynchronous settlement of territories. *Journal of Theoretical Biology* 82:477–496.

834 Temeles, E. J. 1994. The role of neighbours in territorial systems: when are they “dear enemies”?
835 *Animal Behaviour* 47:339–350.

836 Verhulst, S., C. M. Perrins, and R. Riddington. 1997. Natal dispersal of Great Tits in a patchy
837 environment. *Ecology* 78:864–872.

838 Wilkin, T. A., D. Garant, A. G. Gosler, and B. C. Sheldon. 2006. Density effects on life-history traits
839 in a wild population of the great tit *Parus major*: Analyses of long-term data with GIS techniques.
840 *Journal of Animal Ecology* 75:604–615.

841 ———. 2007a. Edge effects in the great tit: Analyses of long-term data with GIS techniques.
842 *Conservation Biology* 21:1207–1217.

843 Wilkin, T. A., L. E. King, and B. C. Sheldon. 2009. Habitat quality, nestling diet, and provisioning
844 behaviour in great tits *Parus major*. *Journal of Avian Biology* 40:135–145.

845 Wilkin, T. A., C. M. Perrins, and B. C. Sheldon. 2007b. The use of GIS in estimating spatial variation
846 in habitat quality: A case study of lay-date in the Great Tit *Parus major*. *Ibis* 149:110–118.

847 Wilson, E. O. 2000. *Sociobiology: the new synthesis (25th anniv.)*. Belknap Press of Harvard
848 University Press, Cambridge, Mass. ; London.

849 Woodman, J. P., E. F. Cole, J. A. Firth, C. M. Perrins, and B. C. Sheldon. 2022. Disentangling the
850 causes of age-assortative mating in bird populations with contrasting life-history strategies. *Journal*
851 *of Animal Ecology*.

852 Woodman, J.P., Gokcekus, S., Beck, K.B., Green, J.P., Nussey, D.H. & Firth, J.A. (2024). The
853 ecology of ageing in wild societies: linking age structure and social behaviour. *Philosophical*
854 *Transactions of the Royal Society B: Biological Sciences*, in press (accepted).

855 Yasukawa, K. 1979. Territory Establishment in Red-Winged Blackbirds: Importance of Aggressive
856 Behavior and Experience. *The Condor* 81:258–264.

857 Ydenberg, R. C., and J. R. Krebs. 1987. The Tradeoff Between Territorial Defense and Foraging in
858 the Great Tit (*Parus major*) 1. *American Zoologist* 27:337–346.

859