

# Effect of bulb type on moth trap catch and composition in UK gardens

Reuben O'Connell-Booth<sup>1,2</sup>  | Christopher Hassall<sup>1</sup>  | William E. Kunin<sup>1</sup>

<sup>1</sup>School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds, UK

<sup>2</sup>Department of Zoology, University of Cambridge, Cambridge, UK

## Correspondence

Reuben O'Connell-Booth, School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT; Department of Zoology, University of Cambridge, Cambridge, CB2 1TN, UK.

Email: [reo29@cam.ac.uk](mailto:reo29@cam.ac.uk)

## Funding information

Natural Environment Research Council, Grant/Award Number: UKR11346

Editor: Alan John Stewart and Associate Editor: Nick Littlewood

## Abstract

1. Light traps are a key method for monitoring moth populations. A wide variety of light sources are employed for this purpose, differing in brightness and spectrum. Relatively little is known about how this affects the resulting sample.
2. We analyse seven moth trap bulb types using 10 years of records from the Garden Moth Scheme (GMS) to provide the largest and most comprehensive comparison of moth trap bulb types to date.
3. The total abundance of moths caught by a trap is strongly linked to bulb wattage and brightness. We estimate that Heath traps catch fewer moths than Robinson or Skinner-stye traps.
4. Ordination models indicated that Mercury Vapour bulbs collected a distinct fauna to Actinic bulbs. Species composition also varied between Actinic bulbs which differ in brightness, with brighter Actinic traps tending to collect a larger proportion of large-winged species.
5. We develop hypotheses whereby large-winged, strong-flying moths are more strongly affected by artificial light (arising from moth traps or other sources). We explore how this hypothesised mechanism may result in negative fitness effects for larger, mobile moth species. Our findings have significant consequences for survey design, citizen science projects and understanding the impact of ALAN on the moth community.

## KEYWORDS

ALAN, artificial light at night, ecological trap, insect declines, insect monitoring, Lepidoptera

## INTRODUCTION

As the largest single radiation of plant-feeding insects (Mitter et al., 2017), the order Lepidoptera comprises one tenth of all described species on Earth, the vast majority of which are moths (van Nieukerken, 2011; Wagner et al., 2021). Butterflies are frequently employed as an indicator of the health of insect communities and of biodiversity more generally (Fleishman & Murphy, 2009). Moths share many of the features which make butterflies attractive in this regard: Lepidoptera tend to respond rapidly to environmental change

(New, 1997) and are distributed globally (Fleishman & Murphy, 2009). Relative to other insect groups, moth species are readily identified owing to the presence of distinctive wing patterns (Thomas, 2005) and the availability of exceptional identification resources (e.g. Sterling et al., 2023; Waring & Townsend, 2018). However, moths can serve as particularly effective indicators of insect diversity due to their substantially greater species richness compared to diurnal Lepidoptera (Wagner et al., 2021) and the ease with which they can be collected in large numbers using light traps—even in highly anthropogenically modified environments such as gardens (Bates et al., 2013).

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *Insect Conservation and Diversity* published by John Wiley & Sons Ltd on behalf of Royal Entomological Society.

Standardised recording of moths in the United Kingdom is conducted through the long-running Rothamsted Insect Survey, which uses a 112-node network of 200 W clear tungsten filament moth traps to measure changes in abundance throughout the country (Bell et al., 2020; Conrad et al., 2006). In addition to standardised monitoring, as many as 2000 volunteers regularly run moth traps in the United Kingdom, many of which submit their records to biodiversity databases or other citizen science organisations (Fox et al., 2011). However, standardised Rothamsted traps are rarely used by citizen scientists or in field experiments due to their large size, mains power requirements and relatively inefficient 200 W tungsten bulb. Instead, a great variety of alternative designs and light sources are employed, hampering standardisation and therefore wider use of community science datasets to monitor moth abundance. In contrast to butterflies, for which standardised 'Pollard walks' (Pollard & Yates, 1993) facilitate straightforward inclusion of volunteer records in large-scale abundance monitoring, volunteer moth records have only recently been included in Butterfly Conservation's annual survey detailing *The State of Britain's Larger Moths* (Fox, 2021). This has been made possible through the application of occupancy models (Dennis et al., 2017). These methods control for variation in moth trap type by treating all records as 'opportunistically gathered', such that citizen science records convey information only on the occurrence or non-occurrence of a species in a location. In this process, all information on species' relative abundance is obscured. Given the central role played by the abundance of common species in ecosystem service delivery (Winfree et al., 2015; Winfree et al., 2025) and recent interest in monitoring insect abundance trends (e.g. Duchenne & Fontaine, 2024; Müller et al., 2024; Seibold et al., 2019), methods capable of estimating moth abundance trends from semi-standardised citizen science records could unlock a novel data source of wide interest and relevance to understanding the status of insects in the United Kingdom and more widely. The first step to broader use of citizen science data for monitoring the abundance of moths is to develop a more complete understanding of the various methods used to collect moths in these surveys.

Many smaller studies have sought to compare one or several moth trap bulb types and receptacle designs to one another in the field. For example, Williams (1951) found that compared to a 125 W Mercury Vapour (MV) Robinson-style traps, Rothamsted-style traps using the same bulb collected 63.82% fewer moths, while Rothamsted-style traps using 200 W 'ordinary' (presumably tungsten filament) bulbs caught 91.44% fewer moths. Williams et al. (1955) found that 200 W 'ordinary' bulbs collected 37.5% fewer moths than 125 W MV, while use of Rothamsted-style trap designs reduced catch by 25% compared to Robinson designs. Heath (1966) found that 15 W blacklight bulbs collect 3.3% more moths than 15 W bulbs and Waring (1980) found that a 6 W Actinic heath trap caught 35.30% of the macrolepidoptera collected by a 125 W Robinson MV. More recent work has shown that various trap parameters can impact catch size, including receptacle design (e.g. Robinson, Heath, Skinner, Rothamsted; Bates et al., 2013), period of the night for which the trap is running (Axmacher & Fiedler, 2004; Nowinszky, 2007), time at which the trap is emptied (discussed in Fry & Waring, 2020), presence

and colour of vanes (Singh et al., 2022), height of the trap (Intachat & Woiwod, 1999), presence and position of packing material (usually egg boxes) inside the trap (discussed in Fry & Waring, 2020), manual versus automatic collection of samples (Axmacher & Fiedler, 2004) and, perhaps most importantly, the brightness and emission spectra of the light source used (Bates et al., 2013; Donners et al., 2018; Fayle et al., 2007; Somers-Yeates et al., 2013; van Grunsven et al., 2014; van Langevelde et al., 2011).

Further, as moth species are thought to differ in their attraction to light (Merckx & Slade, 2014), mediated by variation in mobility (van de Schoot et al., 2024) and possibly spectral sensitivity (Somers-Yeates et al., 2013), it is likely that the species composition, as well as total abundance, of trap catches will be affected by the bulb type used (Donners et al., 2018; Somers-Yeates et al., 2013; van Langevelde et al., 2011). Anecdotal evidence from naturalists has long suggested that traps using different bulbs will produce different samples (Altermatt & Ebert, 2016; Frank, 1988). At the very least, some species are rarely collected at light but appear more abundant using other collection methods (see Sterling & Parsons, 2012). However, as Brehm (2017) notes, field comparisons regularly find little compositional differences between bulbs with fundamentally different light spectra. For example, Infusino et al. (2017) found localised habitat conditions to be a stronger determinant of moth trap composition than light source, and the geometrid assemblages attracted to MV in comparison to incandescent lamps have been found to be surprisingly similar (Intachat & Woiwod, 1999; Jonason et al., 2014). In the more controlled environment of an indoor choice experiment, Brehm et al. (2021) found that moths are preferentially attracted to lamps that emit a high proportion of short-wave radiation. However, in a similar field-based comparison, Niermann and Brehm (2022) found that different LED lamps collected similar numbers and assemblages of moths, independent of whether strong or weak lights were used or if the lamps were of a mixed radiation or UV type, despite the traps being placed close together (26.4 m–46.9 m). It therefore appears that small variations in microhabitat can obscure differences between bulb types, especially in short-term experiments. Additionally, the effective range of moth traps appears to vary between families (Merckx & Slade, 2014) and traps placed close together will suffer from light competition effects (Brehm et al., 2021). These difficulties make designing a field experiment capable of detecting abundance and compositional differences between moth trap light sources a significant logistical challenge, which is yet to be overcome in a comprehensive comparison of commonly used bulb types.

Describing catch size and compositional differences in insect communities is important for monitoring programmes and the design of ecological experiments, and it is important that any biases in widely used sampling methods are fully understood. However, variation in the composition of moth trap samples across trapping methods can also be used to make inferences about the impact of artificial light at night (ALAN) on the moth community. This is because near-identical mechanisms are responsible for both the functioning of moth traps and the disruptive effects of ALAN. As such, while Actinic and MV bulbs are not commonly employed as street lighting, various traps of different spectra are frequently used to explore the effect of ALAN

on moths in choice experiments (e.g. Altermatt & Ebert, 2016; Brehm et al., 2021; Somers-Yeates et al., 2013; van Langevelde et al., 2011).


Here, 10 years of records collected through the Garden Moth Scheme (GMS) in a single land-use type (gardens in the United Kingdom) from seven bulb types and three trap designs are used to explore the impact of spectrum and brightness on the attraction of moths to artificial light. We use weather, bulb-type and trap-type variables to examine how moth abundance changes night-to-night and collate garden habitat variables with information about surrounding land uses to partition garden-to-garden variation attributable to bulb type, urbanisation and distance to street lighting. As such, we provide the largest comparison of moth trap types to date, producing estimates of how moth trap and bulb types compare to one another in terms of catch size, with the aim of assisting in the development of standardised monitoring programmes and the selection of moth trap types for field studies.

Exploring the species composition of moth trap catches, we also test two hypotheses regarding the impact of ALAN on moths. Firstly, we hypothesise that moths display species-specific responses to artificial light of different spectra. This leads to the prediction that moth traps of different spectra will sample different sections of the moth fauna and collect different species. We test this hypothesis by comparing species composition from traps using Actinic to those using MV bulbs (Figure 1). Secondly, we explore the “reduced mobility hypothesis”, described by van de Schoot et al. (2024). Altermatt and Ebert (2016) find that spindle ermine moths (*Yponomeuta cagnagella*) exhibit reduced wing size in areas impacted by ALAN compared to pristine populations. Investigating the mechanism behind this reduced flight-to-light response, van de Schoot et al. (2024) find that urban individuals of this species had on average smaller wings than those emerging from dark, pristine environments. This “reduced mobility hypothesis” states that ALAN exerts selective pressure against mobility in moths, mediated by evolutionary changes in wing size. Here, we develop this hypothesis further. We expect that larger, more mobile moths are more strongly attracted to moth traps, but that this effect will be stronger for brighter light sources. We hypothesise that brighter traps will be able to be perceived by moths from further distances, but that only highly mobile moths will be able to travel to the trap and become caught. We therefore predict that dimmer bulbs will collect a higher proportion of smaller, less mobile species, while brighter bulb types will collect a higher proportion of larger, more mobile species. We test this mobility hypothesis by comparing the size distribution of samples from different bulb types and different levels of ALAN exposure while controlling for other habitat, landscape and geographic variables in ordination models.

## METHODS

### The garden moth scheme dataset

The GMS is a semi-standardised citizen science surveying project (2003 – present), which asks participants to sample moths with a light trap in



**FIGURE 1** (a) Spectral power distribution of a 125 W Mercury Vapour lamp manufactured by Philips, normalised to an irradiance of 1000 lux in 5-nm wavelength bands. (b) Generic spectral power distribution of Actinic lamps across wattages produced by Philips in 5-nm wavelength bands. Irradiance is expressed as a portion of the total irradiance produced, which varies between lamps. The most frequently used lamp type used in the GMS, 125 W MV emits light at a much broader spectrum than the Actinic lamp (data sourced from Philips Lighting, 2025, 2024).

their garden each Friday between March and November. Participants are encouraged to report all records (including trap nights where no moths were recorded), to sample regardless of weather conditions (barring extreme or dangerous weather) and are given a prescribed list of readily identifiable species on which to focus. The scheme has engaged hundreds of participants, producing a large collection of c.36 36-week time series of moth trap catches from a variety of trap types, distributed across Great Britain, Ireland and the Channel Islands. Participants also report information about their garden: the distance to green space, farmland, street lighting, water, woodland and to the coast. Participants provide a list of their garden microhabitats: presence of a bird table, Buddleia, compost heap, >25-m<sup>2</sup> lawn, log pile, long grass, pussy willow, trees over 10 m, oak trees over 10 m, wild honeysuckle, wild ivy, wild-flower meadow are recorded.

### Statistics

All analyses were conducted in R version 4.4.3 (R Core Team, 2025). The percentage landcover in a 100-m radius surrounding each site was

extracted from the UK CEH Land Cover Map 2022 (25-m rasterised pixels; Marston et al., 2024) dataset for each of the 21 landcover classes. Daily 1-km resolution rainfall volume and minimum temperature variables were extracted from the CEDA Archive HadUK Gridded Climate Observations v1.2.0 (Hollis et al., 2019). Moon phase was gathered for each sample night using the package 'moonlit' (Śmielak, 2023), which calculates the percentage of full moonlight intensity based on moon phase, elevation, latitude and longitude (compared to the average full moon value of 0.32 lx), not accounting for cloud cover.

To estimate differences in catch between a range of trap types across each covariate (Model 1) a generalised linear mixed model (GLMM) was fit to time series from each site meeting a set of filtering criteria. Firstly, we separate all GMS records by site (participants' gardens) and year. We filter to include only site-years in Great Britain (excluding records from Ireland and Northern Ireland) in which at least 34 weeks of sampling were conducted. We remove any additional samples taken in winter outside of the GMS sampling window. We exclude timeseries which employed multiple trap or bulb types. Using participants' reported collection method, we identify seven frequently used bulb types (125 W MV, 80 W MV, 60 W Actinic, 40 W Actinic, 20 W Actinic, 15 W Actinic, 6 W Actinic) and three frequently used trap types (Heath, Robinson, Skinner), excluding all records that fall outside these categories. This excludes a small number of participants who used home-made or unconventional traps and bulbs. This produces a dataset of 34–36-week timeseries from seven bulb and three trap categories in Great Britain. Individual gardens may include between 1 and 10 years of records. Graphical description of the spread of trap and bulb categories and covariates in the unfiltered dataset is provided in Supplementary Information Figures S1–S5.

To these data we fitted a GLMM using package *glmmTMB* (Brooks et al., 2017; McGillicuddy et al., 2025) including variables describing bulb type (125 W MV, 80 W MV, 60 W Actinic, 40 W Actinic, 20 W Actinic, 15 W Actinic, 6 W Actinic), trap type (Robinson, Heath, Skinner), week of the year, maximum daily temperature, rainfall volume (in log cubic millimetres) and percentage full moonlight intensity. We model crossed random effects for 'site' (garden) and 'year' (2013–2022) to account for repeated measures and interannual variation in insect abundance.

We undertook extensive residual simulations throughout the modelling process using the R package *DHARMA* (Hartig, 2024) to refine model specification. Data consisted of counts (total number of moths caught in a single trap over one night of sampling) which in preliminary Poisson models indicated overdispersion. As such, a type I negative binomial error term with a log link function was selected (Ver Hoef & Boveng, 2007), coupled with a dispersion model including the variables bulb type, maximum daily temperature, site ID and week of the year. To assess any potential residual non-linearity, we plotted residuals against each covariate individually. This procedure indicated non-linearity between maximum daily temperature and moth abundance. We also found complex multimodal non-linearity between moth abundance and week of the year, corresponding to seasonal variation in moth abundance. We captured this non-linearity using

natural splines with 20 degrees of freedom for week of the year and 5 degrees of freedom for maximum daily temperature.

We tested for multicollinearity using variance inflation factors implemented in R package *performance* (Lüdtke et al., 2021). All variables took values less than 3. Bulb type, trap type, rainfall and moon phase took values between 1.08 and 1.66, while natural spline terms week of the year and maximum daily temperature took values of 2.75 and 2.82 respectively.

The variance explained by fixed effects (marginal  $R^2$ ) was calculated using a likelihood-ratio test implemented in package *MuMIn* (Bartoń, 2025). This test compares the fitted model to a null model including only an intercept, the random effects (site and year) and a dispersion model to estimate the variance explained by fixed effects only. Predictions were made using package *marginalEffects* (Arel-Bundock et al., 2024) and converted to a percentage of the mean 125 W MV prediction for comparability. Post-hoc pairwise comparisons between estimated marginal mean values for bulb and trap-type categories were conducted using package *emmeans* (Lenth, 2024).

Ordination was conducted on samples pooled by site and year to examine species composition variation attributable to bulb type. Only sites with complete GMS microhabitat information were used ( $n = 34$ –36-week timeseries). Categorical distance to feature variables were converted to numerical by taking an average of the distances in the range category. The furthest option in the survey response, ">2 km", was converted to equal 2 km. To determine the ordination method, *Decorana* (DCA) was performed on the log-transformed species matrix as a measure of heterogeneity. A DCA axis 1 value of 2.54 indicated beta-diversity was suitably homogeneous for constrained linear ordination Redundancy Analysis (RDA) rather than Canonical Correspondence Analysis (CCA). Using the function 'ordiR2step' with 49,999 permutations (Oksanen, 2022), a global model with all available variables was compared to an intercept-only model using forward selection to identify and remove extraneous variables. In this procedure variable selection stops if a new variable is either non-significant or if the adjusted  $R^2$  of the model including the new variable exceeds that of the global model. The function provides a test of the significance of each variable to the selected model with an 'ANOVA-like' Monte Carlo test using 999 permutations (Oksanen, 2022), which was then adjusted for multiple comparisons using Holm's correction. Multicollinearity was assessed with variance inflation factors. Percentage suburban landcover had a variance inflation factor of 12.43 and was removed, lowering the variance inflation factors of all variables to below 2.4. All variables excluding suburban landcover were therefore selected. Ten variables with the highest contribution to model adjusted  $R^2$  are provided in Table 1. A full list of selected variables is provided in Supplementary Table S1.

After initial model selection, partial ordination was used to identify the effect of bulb type on species composition. In a partial ordination, conditioning variables (selected CEH landcover variables, latitude, longitude, year, the GMS microhabitat variables and GMS distance to feature variables) are 'partialled out' before analysis by comparing models with and without the variables, leaving only the contribution of the selected constraining variable(s) (Borcard et al., 2018). Following this procedure,

**TABLE 1** GLMM model 1 bulb categories with total number of trap nights and predicted values for each bulb category.

Bulb	Trap nights	Estimate	Std.error	Statistic	p value	CLD	Percent 125 W MV catch
Intercept (125 W MV)	47,565	-2.72	0.14	-19.83	0.00	a	100.00
MV 80	1437	-0.07	0.05	-1.36	0.17	a	93.25
Actinic 60	8884	-0.33	0.03	-11.19	0.00	b	71.85
Actinic 40	8193	-0.51	0.04	-13.92	0.00	c	60.30
Actinic 15	12,622	-0.60	0.04	-16.62	0.00	d	54.76
Actinic 20	4336	-0.60	0.04	-14.56	0.00	d	54.73
Actinic 6	2071	-1.09	0.06	-19.04	0.00	e	33.46

Note: Population-level predictions are provided (setting random effects to 0) using all covariates at their mean values. Compact letter display (CLD) indicates pairwise comparisons in which estimated marginal means were significantly different from one another. Bulbs that share a CLD were not significantly different from one another. Extended version including all covariates is provided in Supplementary Table S1.

the 'mobility hypothesis' (van de Schoot et al., 2024) was tested by regressing species' typical wing size on to RDA axes one and two using the vegan function 'envfit'. This function is suitable for calculating the regression of supplementary variables that pertain to species by setting the 'display' argument to 'species' (Oksanen, 2022). Typical wing size was calculated by averaging minimum and maximum forewing length gathered from Cook et al. (2021) with additional data from Waring and Townsend (2018) and Sterling and Parsons (2012). These data are provided in the Supplementary Information Table S3. Significance of these trait variables was assessed by Monte Carlo simulation (999 permutations). We assess differences in RDA scores across moth families using a Kruskal-Wallis test. Post hoc pairwise comparisons were conducted using Dunn's test with Holm correction to control the family-wise error rate.

The inertia explained by the selected model was evaluated by calculating an unconstrained Principle Component Analysis (PCA) model with no explanatory variables and extracting the first two axes (Zelený, 2022). PCA axes were then used as variables in an RDA to determine the maximum inertia that could be explained by the theoretical best orthogonal explanatory variables. The model  $R^2$  value was then calculated as a proportion of this maximum explicable variation. The significance of the  $R^2$  value was evaluated by comparison to the distribution of 999  $R^2$  values generated by Monte Carlo simulation using randomised environmental variables, implemented in the vegan function 'anova' (Oksanen, 2022).

## RESULTS

### Abundance analysis

In total, 3,801,663 individual moths were included in the abundance analysis across 86,062 trap nights and 575 individual gardens. The mean number of years surveyed per garden was 4.33.

The GLMM Model 1 fixed effects' (daily rainfall volume, maximum daily temperature, trap type, bulb type, week of the year) marginal  $R^2$  was 56.65%. Percentage 125 W MV catch is provided in Table 1 to facilitate comparison of abundance between traps. All bulb categories showed

significant ( $p < 0.0001$ ) effects relative to the 125 W MV reference category, except for 80 W MV, which was not statistically significant ( $p = 0.175$ ). Predicted values show catch increased with wattage (Figure 2a; Table 1). Marginal mean predictions were not significantly different in pairwise comparisons between 80 W MV and 125 W MV ( $p = 0.7171$ ) and between 15 W Actinic and 20 W Actinic bulbs ( $p = 0.9999$ ). All other pairwise comparisons were significant ( $p < 0.0025$ ).


Following an early low peak in March, catch was highest in mid-summer, generally decreased with rainfall, increased with minimum temperature and decreased with moonlight (Figure 2c-f; SI Table 1.) Total abundance of moths across all GMS sites fluctuated between 2013 and 2022, lacking a clear trend (Figure 2g).

Trap type had a significant effect on moth trap catch ( $p < 0.001$ ). Pairwise marginal mean comparisons indicated that Heath traps caught the fewest moths. Heath traps were predicted to catch 11.05% fewer moths than Robinson traps ( $p = 0.0004$ ) and 9.03% fewer moths than Skinner traps ( $p = 0.0022$ ). There was no significant difference between Robinson and Skinner traps ( $p = 0.3770$ ).

### Species composition

The overall unconditioned RDA model was highly significant ( $p < 0.001$ , adj- $R^2$  was 0.29). In this model, all included variables were selected and highly significant ( $p < 0.006$ ). Based on adjusted  $R^2$  (Table 2.), bulb category made the single largest contribution to compositional differences observed in the GMS (adj- $R^2 = 0.0619$ ), followed by latitude (adj- $R^2 = 0.0589$ ) and distance to farmland (adj- $R^2 = 0.0473$ ).

Model 2 (Figure 3) was generated by conditioning variance attributed to all variables aside from bulb category, which was constrained. Model 2 was highly significant ( $p < 0.001$ ). 26.57% of the inertia was conditioned and 4.40% was constrained. The adjusted  $R^2$  was 0.044, representing 29.71% of the maximum variation explicable by two axes. For constrained axes, Eigenvalues were 8.78 for RDA1, 0.515 for RDA2 and 0.233 for RDA3.



**FIGURE 2** Predicted number of moths caught in a single trap night by bulb type (a), trap type (b), percentage full moon (c) log-transformed rainfall volume (d), daily maximum temperature (e), week of the year (f) and year (g).


Variation between MV and Actinic traps aligned with RDA1 (which contained the majority of the total model variance), while centroids of Actinic traps were similar in RDA1, but showed variation in RDA2 (Figure 3). The higher wattage Actinic traps took lower RDA2 scores, but broadly similar RDA1 scores.

The average of each species' reported minimum and maximum forewing length (size) was then regressed onto the ordination axes. Species' position on the ordination was significantly related to their size ( $p > 0.023$ ,  $r = 0.065$ ). The regression aligned with RDA2, the axis partitioning Actinic bulbs by brightness. Size was aligned with the

**TABLE 2** Source, significance, adjusted  $R^2$ , AIC, F values of the 10 variables with the largest impact on moth community composition, measured by adjusted  $R^2$ .


Variable	Cumulative adj- $R^2$	Adj $R^2$	Df	AIC	F	Pr(>F)	Source
Bulb category	0.0619	0.0619	6	15268.43	32.47	0.00112	GMS
Latitude	0.1208	0.0589	1	15083.9	192.20	0.00112	
Distance to farmland	0.1681	0.0473	1	14926.7	163.19	0.00112	GMS
Longitude	0.2054	0.0373	1	14796.34	135.00	0.00112	
Year	0.2210	0.0156	1	14740.7	58.00	0.00112	
Garden size	0.2302	0.0092	3	14709.73	12.34	0.00112	GMS
Distance to woodland	0.2380	0.0079	1	14681.37	30.36	0.00112	GMS
% Suburban landcover	0.2443	0.0063	1	14658.7	24.64	0.00112	CEH
Distance to coastline	0.2499	0.0057	1	14638.19	22.46	0.00112	GMS
Soil Type	0.2554	0.0054	2	14619.41	11.36	0.00112	GMS

Note: Variables were evaluated using OrdR2step with 49,999 permutations.  $R^2$  was adjusted using Holm's correction (adj- $R^2$ ). Degrees of Freedom (Df), AIC (Akaike Information Criterion), F-statistics (F) and F-statistic  $p$ -values (Pr(>F)) are provided. GMS variables originate from the Garden Moth Scheme survey dataset; CEH variables were extracted from the 2021 CEH Landcover map. A full list of evaluated variables is available in Supplementary Information Table S2.

**FIGURE 3** Model 2 partial log-transformed RDA ordination with regressed arrow for size. The effect of all sample variables besides bulb category was 'partialled out' before plotting. Arrow 'size' shows the effect of species' forewing length regressed on to the ordination result.

ordination arrow for 60 W Actinic bulbs and positioned opposite to lower Actinic bulbs and 80 W MV bulbs. Extracting species' RDA scores (Figure 4), a Kruskal–Wallis test indicated a significant difference in RDA1 scores across moth families,  $H(3) = 9.08$ ,  $p = 0.028$ . Post hoc Dunn's test with Holm's correction indicated no significant pairwise differences in RDA1 scores between families (all  $p > 0.06$ ). A

Kruskal–Wallis test indicated a significant difference in RDA2 scores across families,  $H(3) = 14.94$ ,  $p = 0.0018$ . Post hoc Dunn's test using Holm's correction indicated a small significant difference in RDA2 between Noctuidae and Erebidae ( $p = 0.0018$ ; Erebidae median = 0.00441, Noctuidae median =  $-0.00649$ ), but no significant differences between any other pairwise comparison of families.




**FIGURE 4** Comparison of RDA scores between moth families. Species of Geometridae had significantly higher RDA2 scores than those of Noctuidae. RDA scores are extracted from Model 2. Higher RDA1 scores were associated with MV bulb types, while lower RDA2 scores were associated with brighter Actinic bulbs.

Model 3 (Figure 5) constrained the variables 'distance to street lighting' and 'percentage urban landcover', with all other variables conditioned. The resulting ordination was highly significant ( $p < 0.001$ ). The adjusted  $R^2$  was 0.0033, representing 29.71% of the maximum variance explicable by two axes. Overall, 29.75% of the variance was conditioned; 0.37% of the variance was constrained. Eigenvalues for the constrained axes were:  $RDA1 = 0.4052$ ;  $RDA2 = 0.3397$ . The regressed vector for species' size was significantly related to the ordination ( $p = 0.0019$ ,  $R^2 = 0.0288$ ) and aligned with distance to street lighting. The regressed arrows approached orthogonality to percentage urban landcover.

## DISCUSSION

We find that brighter moth trap bulbs generally collect larger numbers of moths. We find that Robinson and Skinner traps tend to collect more moths than Heath-style traps. We find some differences in species composition between bulb types. Ordination model results lead us to hypothesise that MV traps collect a unique assemblage and that brighter, higher wattage Actinic bulbs collect a larger proportion of large-winged and mobile moth species.

Model 1 indicates that higher wattage moth traps generally collect larger numbers of moths. One bulb type, 15 W Actinic, falls out of this pattern and is estimated to collect marginally more (+0.032) moths than 20 W Actinic bulbs. Notably, 15 W Actinic traps can be powered easily by battery, meaning that they may be more freely placed away from competing light sources within the garden. This flexibility may be conducive to higher total catches than otherwise expected by wattage alone. We estimate that Heath-style traps are likely to collect fewer moths than Robinson or Skinner style traps. However, we find no significant difference between Robinson and Skinner designs. We find that catch is negatively affected by moonlight, rainfall and low temperature, in line with previous studies (Holyoak et al., 1997; Jonason et al., 2014; Puskas et al., 2006). However, due to the observational



**FIGURE 5** Model 3 RDA ordination. All variables besides percentage urban landcover and distance to streetlighting were 'partialled out' before plotting. Arrow 'size' shows the effect of species' size regressed on to the ordination result.

nature of our study, a major unmeasured source of confounding is non-random assignment of moth trap type to participants' gardens (selection effects). For instance, where competing light pollution is not an issue, participants may select a dimmer trap which can catch a manageable number of moths or will ensure a larger proportion land inside (rather than surrounding) the trap, more experienced trappers may use more powerful bulb types and participants with the means to buy more expensive traps may also have more diverse gardens (the 'luxury effect', Leong et al., 2018). We are unable to control for this confounding. Our estimates are, however, based on substantially larger sample sizes than previous field-based comparisons (e.g. Blomberg et al., 1976; Brehm, 2017; Brehm et al., 2021; Fayle et al., 2007; Infusino et al., 2017; Merckx & Slade, 2014; Niermann & Brehm, 2022;

Somers-Yeates et al., 2013). We also provide improved estimates over previous analyses of GMS data (Bates et al., 2013) by parameterising the important determinants of moth abundance over 10 years.


Model 2 indicates that bulb types collect somewhat different assemblages of species. Previous reports conflict over compositional differences between moth traps. Between traps of different spectra, several field experiments have found little or no difference in assemblage (Blomberg et al., 1976; Fayle et al., 2007; Merckx & Slade, 2014; Somers-Yeates et al., 2013; van Grunsven et al., 2014), while others note differences, but conflict over underlying causes (cf. Brehm, 2017; Brehm et al., 2021; Infusino et al., 2017; Niermann & Brehm, 2022; see Introduction). Model 2 indicates that there are relatively large differences in species composition between different bulb types compared to other variables examined. Based on adjusted  $R^2$ , compositional differences attributable to bulb type were larger than any other geographic, landcover or garden microhabitat variable tested (Table 2).

What underlies this compositional variation? As RDA models explained a relatively small portion of the total variation in species composition, it is not possible to draw firm conclusions on this question. We find that noctuid species took significantly lower values in RDA2 when compared to erebid species, but there was no significant difference compared to any other family tested. Regressing species' forewing length on to Model 2 indicated that species' forewing length was significantly related to their position on the ordination axes. Larger species had significantly larger negative values in RDA2, while

smaller species took larger positive values in RDA2. RDA2 partitioned Actinic traps by wattage, indicating that species associated with brighter, higher wattage Actinic traps tended to have larger wings than those associated with dimmer, lower wattage Actinic bulbs.

Building on the “reduced mobility hypothesis” described by van de Schoot et al. (2024), we hypothesise that a mechanism contributing to this association is species' mobility. As trap brightness is increased, we expect that its light will be perceptible to moths from further distances, but that many of the moths which perceive the light will not reach the trap and become caught. It seems likely that a contributing factor to whether an individual moth reaches the trap and becomes caught is its size and mobility, as more mobile moths are likely to be capable of travelling further to reach the trap. If a bulb's brightness is increased one would therefore expect to sample the local (mobile and relatively immobile) fauna, plus some portion of the more distant individuals that are mobile enough to reach the trap. The hypothesis is illustrated diagrammatically in Figure 6. This hypothesis predicts that large moths should be over-represented with respect to weak-flying species in bright traps, a prediction tentatively supported by our ordination results.

While also explaining a relatively small portion of the total variation in species composition, Model 3 provides some support for the above hypotheses. In Model 3, which constrained percentage urban landcover together with distance to street lighting, species' position on the ordination was also significantly related to their size. The



**FIGURE 6** Diagram illustrating the mobility hypothesis, building on the mobility mechanism described in Altermatt and Ebert (2016) and tested in van de Schoot et al. (2024). The central, yellow-coloured box represents a moth trap or other light source. Black circles, filled grey, represent the distance from which the trap can be seen; we expect brighter moth traps can be perceived by moths from further away. Silhouettes represent moths in proximity to the trap. Moths 1 and 2 represent small, relatively immobile species, whereas moth 3 represents a more mobile species; coloured circles represent the distance a particular moth is capable of travelling. For the dimmer trap, only moth 1 can see the trap. Moth 1 is also capable of travelling to the trap, so may become caught. For the brighter trap, both moth 2 and moth 3 can see the trap and may become attracted to it. However, moth 2, which is relatively immobile, is not capable of reaching the trap to become caught. In this simple graphical model, increasing the brightness of a trap will change the species assemblage collected, increasing the proportion of large species relative to small species.

regressed arrow aligned with that of distance to street lighting and was orthogonal to that of urban landcover. This association indicates that traps within gardens further from street lighting tended to collect a larger proportion of large-winged moth species, whereas gardens closer to street lighting tended to collect more small-winged species. Orthogonality between the regressed size arrow and percentage urban landcover indicates that this relationship was unrelated to urbanity, a potential confounder of the effect of ALAN on species composition. It follows from the mechanism postulated above that strong-flying moth species are more likely to be caught under streetlights as well as within moth traps. Ordination Model 3 therefore leads us to tentatively hypothesise that there may be selective pressure against mobility in moths due to ALAN. Previous work also supports these hypotheses. Altermatt and Ebert (2016) established that in the spindle ermine moth (*Yponomeuta cagnagella*), ALAN exposure precipitated behavioural and morphological changes causing urban-collected individuals to display 30% reduced flight-to-light response compared to those collected in dark rural areas. van de Schoot et al. (2024) measured the wing dimensions (length, width and area) of these moths, finding that those collected from ALAN-affected areas showed a significant reduction in forewing length. This suggests selective pressure against mobility mediated by changes in forewing length. Findings presented here, that larger-winged moths form a larger proportion of catch within brighter traps, provide some support to this hypothesis. Together, this has led us to hypothesise that ALAN can work as an ecological trap, altering moth community composition by exerting negative fitness effects on large moth species (Figure 6), a mechanism potentially contributing to the faster decline of large species in the United Kingdom (Coulthard et al., 2019).

Another mechanism that could explain the observed patterns is allometry: all else equal, smaller eyes are less sensitive to light as they can collect and focus fewer photons (Warrant, 2017; Warrant & Somanathan, 2022). As eye size appears to scale with wing size allometrically in moths (Stöckl et al., 2022), we are unable to distinguish between these two mechanisms—both may play a role. However, previous work supports the hypothesis that mobility plays a substantial role in the flight-to-light response in moths (Altermatt & Ebert, 2016; van de Schoot et al., 2024).

Our results also suggest that MV traps collect a distinct assemblage to actinic traps (Model 2). However, factors underlying compositional differences between MV and Actinic traps are less clear than those between Actinic traps of different wattages. MV bulbs emit a larger proportion of non-UV and longer-wavelength light than Actinic (Figure 1). Brehm et al. (2021), van Langevelde et al. (2011) and Somers-Yeates et al. (2013) find that light sources of longer-wavelength collect fewer species, suggesting the UV portion of the spectrum is most attractive to moths. Our findings support this conclusion. However, not tested is the combination of UV and longer-wavelength light. Here, Model 2 showed that 80 W MV bulbs were positioned closest to 125 W MV bulbs along RDA1, despite lower and more variable catch size produced by 80 W MV traps (Model 1). This indicates that a portion of the variation between these bulb types is attributable to emission spectra rather than

simply sample size. One study, comparing blacklight-coated (excluding all but UV radiation) MV bulbs to non-coated MV bulbs (Fayle et al., 2007) found that excluding the human-visible portion of the spectrum decreased catch size and diversity. Together with ordination Model 2 presented here, this supports the hypothesis that UV in concert with longer-wavelength light produces a characteristic MV species assemblage, possibly including species which are more sensitive to longer-wavelength light. Further work is required to evaluate traits underlying this variation.

In 2015, MV bulbs were banned in the EU for general purpose use and are no longer widely manufactured (Bates et al., 2013). Here it is shown that the trap type producing the next largest samples is 60 W Actinic, but this comes with the caveat that MV and Actinic bulbs appear to sample different sections of the moth fauna. An alternative not evaluated are LED traps (White et al., 2016; Brehm, 2017; Infusino et al., 2017). These traps can be designed to output a great variety of spectra by combining diodes, are energy efficient and typically light-weight. Several designs are currently available, such as the LepiLED (Brehm, 2017), but the ideal spectral profile of LED lamps is yet to be determined (but cf. Brehm, 2017; Niermann & Brehm, 2022). Results presented here support the idea that inclusion of longer-wavelength emitting diodes are likely to broaden the fauna which LED traps sample (Brehm, 2017). Also not evaluated here are 200 W tungsten bulbs used by the Rothamsted Insect Survey light trap network. These bulbs emit substantially less UV and a broader range of LW radiation than MV traps and are therefore likely to sample a distinct fauna.

## CONCLUSIONS

We found that higher wattage bulbs generally collected larger numbers of moths and that different bulb types collected somewhat different sets of species. We found that Heath-style traps collected fewer moths than Robinson or Skinner style traps but found no significant difference between Robinson and Skinner trap designs.

We also used ordination to explore compositional differences between traps. We hypothesise that: (1) large-winged, mobile species can travel further to light sources and are consequently more likely to be caught by moth traps or under street lighting; (2) this mechanism is a cause of the (relatively weak) patterning observed in ordination models, which indicated brighter moth traps are characterised by a larger proportion of large-winged species; (3) this mechanism implies that the negative fitness effects of ALAN are more pronounced for large-winged moth species; (4) compositional differences between MV and Actinic traps indicate that there may be an 'MV-unique' moth assemblage, potentially caused by the broader spectral emission profile of MV bulbs in comparison to Actinic bulbs. Further work is required to explore these hypotheses in an experimental setting.

While standardised moth collection methods exist in the form of Rothamsted-style traps, these traps are rarely used by citizen scientists. Given that the most widely used MV bulbs have now been phased out of production, there is an opportunity to develop

a new standardised protocol for monitoring moths akin to Pollard walks used to monitor butterflies in citizen science surveys. It is imperative that any candidate ‘standard’ bulb stay in production long into the future. It is also desirable that a candidate bulb is inexpensive, widely available, non-toxic, energy efficient and long-lasting. While LED moth traps are not evaluated here, potential candidate bulbs assessed include widely used 60 W Actinic bulbs for sites where mains power is available, and 15 W Actinic bulbs for sites where battery power is required.

## AUTHOR CONTRIBUTIONS

**Reuben O’Connell-Booth:** Conceptualization; investigation; methodology; writing – original draft; writing – review and editing; visualization; formal analysis; data curation; validation; software. **Christopher Hassall:** Conceptualization; writing – review and editing; supervision; project administration; methodology. **William E. Kunin:** Supervision; writing – review and editing; conceptualization; project administration; methodology.

## ACKNOWLEDGEMENTS

We are very grateful to all the participants and organisers of the Garden Moth Scheme for providing the data on which this paper is based. We thank two reviewers for their helpful comments which have improved the manuscript. This work was supported by the Natural Environment Research Council grant number UKRI1346.

## CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

## DATA AVAILABILITY STATEMENT

Species occurrence data in this study were used under licence from a citizen science recording scheme and are available from the Garden Moth Scheme (<https://gardenmothscheme.org.uk>), subject to a data-sharing agreement. Code used to produce this manuscript are available via FigShare, <https://doi.org/10.6084/m9.figshare.27727659.v2> (O’Connell-Booth et al., 2024).

## ORCID

Reuben O’Connell-Booth  <https://orcid.org/0009-0007-9929-895X>

Christopher Hassall  <https://orcid.org/0000-0002-3510-0728>

## REFERENCES

- Altermatt, F. & Ebert, D. (2016) Reduced flight-to-light behaviour of moth populations exposed to long-term urban light pollution. *Biology Letters*, 12(4), 20160111. Available from: <https://doi.org/10.1098/rsbl.2016.0111>
- Arel-Bundock, V., Greifer, N. & Heiss, A. (2024) How to interpret statistical models using marginaffects for R and python. *Journal of Statistical Software*, 111(9), 1–32. Available from: <https://doi.org/10.18637/jss.v111.i09>
- Axmacher, J.C. & Fiedler, K. (2004) Manual versus automatic moth sampling at equal light sources – a comparison of catches from Mt. Kilimanjaro. *Journal of the Lepidopterists’ Society*, 58(4), 196–202.

- Bartoń, K. (2025) *MuMIn: Multi-Model Inference*. <https://CRAN.R-project.org/package=MuMIn>
- Bates, A.J., Sadler, J.P., Everett, G., Grundy, D., Lowe, N., Davis, G. et al. (2013) Assessing the value of the garden moth scheme citizen science dataset: how does light trap type affect catch? *Entomologia Experimentalis et Applicata*, 146(3), 386–397. Available from: <https://doi.org/10.1111/eea.12038>
- Bell, J.R., Blumgart, D. & Shortall, C.R. (2020) Are insects declining and at what rate? An analysis of standardised, systematic catches of aphid and moth abundances across Great Britain. *Insect Conservation and Diversity*, 13(2), 115–126. Available from: <https://doi.org/10.1111/icad.12412>
- Blomberg, O., Itämiä, J. & Kuusela, K. (1976) Insect catches in a blended and a black light-trap in northern Finland. *Oikos*, 27(1), 57–63. Available from: <https://doi.org/10.2307/3543432>
- Borcard, D., Gillet, F. & Legendre, P. (2018) *Numerical Ecology with R*. Cham: Springer International Publishing (Use R!). Available from: <https://doi.org/10.1007/978-3-319-71404-2>
- Brehm, G. (2017) A new LED lamp for the collection of nocturnal Lepidoptera and a spectral comparison of light-trapping lamps. *Nota Lepidopterologica*, 40(1), 87–108. Available from: <https://doi.org/10.3897/nl.40.11887>
- Brehm, G., Niermann, J., Jaimes Nino, L.M., Enseling, D., Jüstel, T., Axmacher, J.C. et al. (2021) Moths are strongly attracted to ultraviolet and blue radiation. *Insect Conservation and Diversity*, 14(2), 188–198. Available from: <https://doi.org/10.1111/icad.12476>
- Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A. et al. (2017) glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R Journal*, 9(2), 378–400. Available from: <https://doi.org/10.32614/RJ-2017-066>
- Conrad, K.F., Warren, M.S., Fox, R., Parsons, M.S. & Woiwod, I.P. (2006) Rapid declines of common, widespread British moths provide evidence of an insect biodiversity crisis. *Biological Conservation*, 132(3), 279–291. Available from: <https://doi.org/10.1016/j.biocon.2006.04.020>
- Cook, P.M., Tordoff, G.M., Davis, T.M., Parsons, M.S., Dennis, E.B., Fox, R. et al. (2021) Traits data for the butterflies and macro-moths of Great Britain and Ireland. NERC EDS Environmental Information Data Centre, 103, e3670. Available from: <https://doi.org/10.5285/5B5A13B6-2304-47E3-9C9D-35237D1232C6>
- Coulthard, E., Norrey, J., Shortall, C. & Harris, W.E. (2019) Ecological traits predict population changes in moths. *Biological Conservation*, 233, 213–219. Available from: <https://doi.org/10.1016/j.biocon.2019.02.023>
- Dennis, E.B., Morgan, B.J.T., Brereton, T.M., Roy, D.B. & Fox, R. (2017) Using citizen science butterfly counts to predict species population trends. *Conservation Biology: The Journal of the Society for Conservation Biology*, 31(6), 1350–1361. Available from: <https://doi.org/10.1111/cobi.12956>
- Donners, M., van Grunsven, R.H.A., Groenendijk, D., van Langevelde, F., Bikker, J.W., Longcore, T. et al. (2018) Colors of attraction: modeling insect flight to light behavior. *Journal of Experimental Zoology Part A: Ecological and Integrative Physiology*, 329(8–9), 434–440. Available from: <https://doi.org/10.1002/jez.2188>
- Duchenne, F. & Fontaine, C. (2024) Weather explains inter-annual variability, but not the temporal decline, in insect biomass. *Insect Conservation and Diversity*, 17(6), 932–937. Available from: <https://doi.org/10.1111/icad.12769>
- Fayle, T.M., Sharp, R.E. & Majerus, M.E.N. (2007) The effect of moth trap type on catch size and composition of British Lepidoptera. *British Journal of Entomology & Natural History*, 20(4), 221–232.
- Fleishman, E. & Murphy, D.D. (2009) A realistic assessment of the indicator potential of butterflies and other charismatic taxonomic groups. *Conservation Biology*, 23(5), 1109–1116. Available from: <https://doi.org/10.1111/j.1523-1739.2009.01246.x>

- Fox, R. (2021) *State of Britain's larger moths 2021*. Wareham: Butterfly Conservation.
- Fox, R., Randle, Z., Hill, L., Anders, S., Wiffen, L. & Parsons, M.S. (2011) Moths count: recording moths for conservation in the UK. *Journal of Insect Conservation*, 15(1), 55–68. Available from: <https://doi.org/10.1007/s10841-010-9309-z>
- Frank, K. (1988) Impact of outdoor lighting on moths. *Journal of the Lepidopterists' Society*, 42(2), 51. Available from: <https://doi.org/10.1017/S0252921100003687>
- Fry, R. & Waring, P. (2020) *A Guide to Moth Traps and Their Use*, 3rd ed. Great Britain: Amateur Entomologist's Society.
- Hartig, F. (2024) *DHARMA: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regression Models*. <https://CRAN.R-project.org/package=DHARMA>
- Heath, J. (1966) A comparison of the catches obtained in insect traps fitted with 15 watt and 6 watt "Blacklight" fluorescent tubes. *Bulletin of Entomological Research*, 78, 222–223.
- Hollis, D., McCarthy, M., Kendon, M., Legg, T. & Simpson, I. (2019) HadUK-grid—A new UK dataset of gridded climate observations. *Geoscience Data Journal*, 6(2), 151–159. Available from: <https://doi.org/10.1002/gdj3.78>
- Holyoak, M., Jarosik, V. & Novák, I. (1997) Weather-induced changes in moth activity bias measurement of long-term population dynamics from light trap samples. *Entomologia Experimentalis et Applicata*, 83(3), 329–335. Available from: <https://doi.org/10.1046/j.1570-7458.1997.00188.x>
- Infusino, M., Brehm, G., di Marco, C. & Scalerico, S. (2017) Assessing the efficiency of UV LEDs as light sources for sampling the diversity of macro-moths (Lepidoptera). *European Journal of Entomology*, 114, 25–33. Available from: <https://doi.org/10.14411/eje.2017.004>
- Intachat, J. & Woiwod, I.P. (1999) Trap design for monitoring moth biodiversity in tropical rainforests. *Bulletin of Entomological Research*, 89(2), 153–163. Available from: <https://doi.org/10.1017/S0007485399000243>
- Jonason, D., Franzén, M. & Ranius, T. (2014) Surveying moths using light traps: effects of weather and time of year. *PLoS One*, 9(3), e92453. Available from: <https://doi.org/10.1371/journal.pone.0092453>
- Lenth, R.V. (2024) emmeans: Estimated Marginal Means, aka Least-Squares Means. <https://CRAN.R-project.org/package=emmeans>
- Leong, M., Dunn, R.R. & Trautwein, M.D. (2018) Biodiversity and socioeconomic in the city: A review of the luxury effect. *Biology Letters*, 14(5), 20180082. Available from: <https://doi.org/10.1098/rsbl.2018.0082>
- Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P. & Makowski, D. (2021) Performance: an R package for assessment, comparison and testing of statistical models. *Journal of Open Source Software*, 6(60), 3139. Available from: <https://doi.org/10.21105/joss.03139>
- Marston, C.G., Morton, R.D., O'Neil, A.W. & Rowland, C.S. (2024) Land cover map 2022 (25m rasterised land parcels, GB). *NERC EDS Environmental Information Data Centre*. Available from: <https://doi.org/10.5285/c9449bf5-b8f6-4a1c-b3eb-0d70575cba39>
- McGillycuddy, M., Popovic, G., Bolker, B.M. & Warton, D.I. (2025) Parsimoniously fitting large multivariate random effects in glmmTMB. *Journal of Statistical Software*, 112(1), 1–19. Available from: <https://doi.org/10.18637/jss.v112.i01>
- Merckx, T. & Slade, E.M. (2014) Macro-moth families differ in their attraction to light: implications for light-trap monitoring programmes. *Insect Conservation and Diversity*, 7(5), 453–461. Available from: <https://doi.org/10.1111/icad.12068>
- Mitter, C., Davis, D.R. & Cummings, M.P. (2017) Phylogeny and evolution of Lepidoptera. *Annual Review of Entomology*, 62(1), 265–283. Available from: <https://doi.org/10.1146/annurev-ento-031616-035125>
- Müller, J., Hothorn, T., Yuan, Y., Seibold, S., Mitesser, O., Rothacher, J. et al. (2024) Weather explains the decline and rise of insect biomass over 34 years. *Nature*, 628(8007), 349–354. Available from: <https://doi.org/10.1038/s41586-023-06402-z>
- New, T.R. (1997) Are Lepidoptera an effective 'umbrella group' for biodiversity conservation? *Journal of Insect Conservation*, 1(1), 5–12. Available from: <https://doi.org/10.1023/A:1018433406701>
- Niermann, J. & Brehm, G. (2022) The number of moths caught by light traps is affected more by microhabitat than the type of UV lamp used in a grassland habitat. *European Journal of Entomology*, 119, 36–42. Available from: <https://doi.org/10.14411/eje.2022.004>
- Nowinszky, L. (2007) The hourly distribution of moth species caught by a light-trap. *Applied Ecology and Environmental Research*, 5(1), 103–107. Available from: [https://doi.org/10.15666/aeer/0501\\_103107](https://doi.org/10.15666/aeer/0501_103107)
- O'Connell-Booth, R., Kunin, W.E. & Hassall, C. (2024) PUBLIC – Effect of bulb type on moth trap catch and composition in UK gardens. FigShare. Version 2. <https://doi.org/10.6084/m9.figshare.27727659.v2>
- Oksanen, J. (2022) Package 'vegan'. <https://cran.r-project.org/web/packages/vegan/vegan.pdf>
- Philips Lighting. (2024) HPL-N 125W E27 SG 1CT/24 Standard High Pressure Mercury lamp datasheet. [https://www.lighting.philips.com/api/assets/v1/file/Signify/content/928052007399\\_EU.en\\_AA.PROF.FP/Localized\\_commercial\\_leaflet\\_928052007399\\_en\\_AA.pdf](https://www.lighting.philips.com/api/assets/v1/file/Signify/content/928052007399_EU.en_AA.PROF.FP/Localized_commercial_leaflet_928052007399_en_AA.pdf) [Accessed 1st July 2025].
- Philips Lighting. (2025) Actinic BL TL 6W/10 1FM/10X25CC datasheet. [https://www.lighting.philips.com/api/assets/v1/file/Signify/content/928000501030\\_EU.en\\_AA.PROF.FP/Localized\\_commercial\\_leaflet\\_928000501030\\_en\\_AA.pdf](https://www.lighting.philips.com/api/assets/v1/file/Signify/content/928000501030_EU.en_AA.PROF.FP/Localized_commercial_leaflet_928000501030_en_AA.pdf) [Accessed 1st July 2025].
- Pollard, E. & Yates, T.J. (1993) *Monitoring butterflies for ecology and conservation: the British Butterfly Monitoring Scheme*. London: Chapman and Hall Ltd.
- Puskas, J., Nowinszky, L. & Makra, L. (2006) Joint influence of meteorological events on light trapping of turnip moth (*scotia segetum* Schiff). *Zbornik Matice Srpske Za Prirodne Nauke*, 110, 259–266.
- R Core Team. (2025) 'R: a language and environment for statistical computing', *R Foundation for Statistical Computing*.
- Seibold, S., Gossner, M.M., Simons, N.K., Blüthgen, N., Müller, J., Ambarlı, D. et al. (2019) Arthropod decline in grasslands and forests is associated with landscape-level drivers. *Nature*, 574(7780), 671–674. Available from: <https://doi.org/10.1038/s41586-019-1684-3>
- Singh, R.P., Böttger, D. & Brehm, G. (2022) Moth light traps perform better with vanes: a comparison of different designs. *Journal of Applied Entomology*, 146(10), 1343–1352. Available from: <https://doi.org/10.1111/jen.13068>
- Śmielak, M.K. (2023) Biologically meaningful moonlight measures and their application in ecological research. *Behavioral Ecology and Sociobiology*, 77(2), 21. Available from: <https://doi.org/10.1007/s00265-022-03287-2>
- Somers-Yeates, R., Hodgson, D., McGregor, P.K., Spalding, A. & ffrench-Constant, R.H. (2013) Shedding light on moths: shorter wavelengths attract noctuids more than geometrids. *Biology Letters*, 9(4), 20130376. Available from: <https://doi.org/10.1098/rsbl.2013.0376>
- Sterling, P. & Parsons, M. (2012) *Field Guide to the Micro-moths of Great Britain and Ireland: 2nd edition, 1st ed.* London: Bloomsbury.
- Sterling, P., Parsons, M. & Lewington, R. (2023) *Field guide to the micro-moths of Great Britain and Ireland*, 2nd edition. London: Bloomsbury Wildlife.
- Stöckl, A., Grittner, R., Taylor, G., Rau, C., Bodey, A.J., Kelber, A. et al. (2022) Allometric scaling of a superposition eye optimizes sensitivity and acuity in large and small hawkmoths. *Proceedings of the Royal Society B: Biological Sciences*, 289(1979), 20220758. Available from: <https://doi.org/10.1098/rspb.2022.0758>
- Thomas, J. (2005) Monitoring change in the abundance and distribution of insects using butterflies and other indicator groups. *Philosophical*

- Transactions of the Royal Society, B: Biological Sciences*, 360(1454), 339–357. Available from: <https://doi.org/10.1098/rstb.2004.1585>
- van de Schoot, E., Merckx, T., Ebert, D., Wesselingh, R.A., Altermatt, F. & van Dyck, H. (2024) Evolutionary change in flight-to-light response in urban moths comes with changes in wing morphology. *Biology Letters*, 20(3), 20230486. Available from: <https://doi.org/10.1098/rsbl.2023.0486>
- van Grunsven, R.H.A., Donners, M., Boekee, K., Tichelaar, I., van Geffen, K.G., Groenendijk, D. et al. (2014) Spectral composition of light sources and insect phototaxis, with an evaluation of existing spectral response models. *Journal of Insect Conservation*, 18(2), 225–231. Available from: <https://doi.org/10.1007/s10841-014-9633-9>
- van Langevelde, F., Ettema, J.A., Donners, M., WallisDeVries, M.F. & Groenendijk, D. (2011) Effect of spectral composition of artificial light on the attraction of moths. *Biological Conservation*, 144(9), 2274–2281. Available from: <https://doi.org/10.1016/j.biocon.2011.06.004>
- van Nieuwerkerken, E.J. (2011) Order Lepidoptera. Linnaeus, 1758. *Zootaxa*, 3148(1), 212–221. Available from: <https://doi.org/10.11646/zootaxa.3148.1.41>
- Ver Hoef, J.M. & Boveng, P.L. (2007) Quasi-Poisson vs. negative binomial regression: how should we model Overdispersed count data? *Ecology*, 88(11), 2766–2772. Available from: <https://doi.org/10.1890/07-0043.1>
- Wagner, D.L., Fox, R., Salcido, D.M. & Dyer, L.A. (2021) A window to the world of global insect declines: moth biodiversity trends are complex and heterogeneous. *Proceedings of the National Academy of Sciences*, 118(2), e2002549117. Available from: <https://doi.org/10.1073/pnas.2002549117>
- Waring, P. (1980) Comparison of the Heath and Robinson M.V. Moth traps. *The Entomologists Record*, 92, 283–289.
- Waring, P. & Townsend, M. (2018) *Field guide to the moths of Great Britain and Ireland*, 3rd edition. London: Bloomsbury.
- Warrant, E. & Somanathan, H. (2022) Colour vision in nocturnal insects. *Philosophical Transactions of the Royal Society, B: Biological Sciences*, 377(1862), 20210285. Available from: <https://doi.org/10.1098/rstb.2021.0285>
- Warrant, E.J. (2017) The remarkable visual capacities of nocturnal insects: vision at the limits with small eyes and tiny brains. *Philosophical Transactions of the Royal Society, B: Biological Sciences*, 372(1717), 20160063. Available from: <https://doi.org/10.1098/rstb.2016.0063>
- White, P.J.T., Glover, K., Stewart, J., Rice, A. (2016) The Technical and Performance Characteristics of a Low-Cost, Simply Constructed, Black Light Moth Trap. *Journal of Insect Science*, 16, 25. Available from: <https://doi.org/10.1093/jisesa/iew011>
- Williams, C.B. (1951) Comparing the efficiency of insect traps. *Bulletin of Entomological Research*, 42(3), 513–517. Available from: <https://doi.org/10.1017/S000748530002890X>
- Williams, C.B., French, R.A. & Hosni, M.M. (1955) A second experiment on testing the relative efficiency of insect traps. *Bulletin of Entomological Research*, 46(1), 193–204. Available from: <https://doi.org/10.1017/S0007485300030844>
- Winfree, R., Reilly, J.R. & Genung, M.A. (2025) Biodiversity–ecosystem function research must consider abundance and not just diversity. *Nature Reviews Biodiversity*, 1(6), 347–348. Available from: <https://doi.org/10.1038/s44358-025-00040-1>
- Winfree, R., W. Fox, J., Williams, N.M., Reilly, J.R. & Cariveau, D.P. (2015) Abundance of common species, not species richness, drives delivery of a real-world ecosystem service. *Ecology Letters*, 18(7), 626–635. Available from: <https://doi.org/10.1111/ele.12424>
- Zelený, D. (2022) *Explained variation in constrained ordination*. [https://www.davidzeleny.net/anadat-r/doku.php/en:expl\\_var?do=](https://www.davidzeleny.net/anadat-r/doku.php/en:expl_var?do=) [Accessed 1st July 2023].

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**Table S1.** Extended version of Table 1. Model 1 results showing estimates and significance for each included fixed effect. Natural spline terms are indicated with “ns”.

**Table S2.** Extended version of Table 2. Source, significance, adjusted  $R^2$ , AIC, F values of available variables, produced by OrdR2step with 49,999 permutations. Variables were evaluated OrdR2step with 49,999 permutations.  $R^2$  was adjusted using Holm's correction (adj- $R^2$ ). Degrees of Freedom (Df), AIC (Akaike Information Criterion), F-statistics (F), and F-statistic p-values (Pr(>F)) are provided. GMS variables originate from the Garden Moth Scheme survey dataset. CEH variables were extracted from the 2021 CEH Landcover map.

**Table S3.** Typical minimum and maximum forewing length of moth species examined. Gathered from Cook et al. (2021), with additional data from Waring & Townsend (2018) and Sterling and Parsons (2012). Typical wing size was calculated for analyses by averaging minimum and maximum forewing length.

**Figure S1.** Descriptive statistics showing variation in trap and bulb type among GMS records from Great Britain prior to filtering. Number of samples (single trap nights) across all records grouped by trap type (a) and bulb type (b). Variation in bulb type among Robinson-type traps (c), Heath-type traps (d) and Skinner-type traps (e). Variation in bulb wattage across all Mercury Vapour (MV) bulbs (f) and Actinic bulbs (g).

**Figure S2.** Variation in participant-reported garden habitat features among GMS garden sites in Great Britain prior to filtering. Participants return a garden habitat form as part of the monitoring scheme, detailing presence or distance to the following features, from left to right: (1) presence of a bird table, (2) presence of *Buddleia davidii*, (3) presence of a compost heap, (4) distance to coastline (5) distance to farmland (6) Distance to green space of any kind, (7) distance to street lighting, (8) distance to woodland, (9) distance to open bodies of water of any kind, (10) distance to woodland, (11) garden size among four categories, (12) garden size among four categories, (13) ‘habitat’ type among three broad land-use categories, (14) presence of a native species hedgerow, (15) presence of a lawn over 25 m<sup>2</sup>, (16) presence of a log pile, (17) presence of long grass, (18) presence of a nettle patch, (18) presence of an oak tree over 10 m in height, (19) presence of a pond, (20) presence of a pussy willow (*Salix caprea*), (21) Soil type among three categories, (22) presence of any tree over 10 m in height, (23) presence of a wild flower meadow of any type, (24) presence of a wild honeysuckle (*Lonicera periclymenum*), (25) presence of wild ivy (*Hedera* spp.).

**Figure S3.** Histograms showing variation landcover in a 100-m radius surrounding GMS garden sites in Great Britain prior to filtering. The following aggregated 2022 UKCEH Landcover Classes (Marston et al., 2024) were calculated for each site: built-up, comprised of summed urban and suburban landcover (A); farmland cover, comprised of summed improved grassland, arable and horticultural cover (B); Woodland, comprised of summed coniferous and broadleaf woodland cover (C); Coastal cover, comprised of summed open seawater, littoral sediment, littoral rock, supralittoral sediment, supralittoral rock,

saltwater and saltmarsh cover (D) and Semi-natural cover, comprised of summed neutral grassland, calcareous grassland, acid grassland, heather, heather grassland, bog, fen, marsh and swamp cover (E).

**Figure S4.** Histograms showing variation in rainfall volume (A) and maximum daily temperature (B) conditions across GMS samples. Weather data were collected for each sample from the CEDA Archive HadUK Gridded Climate Observations v1.2.0 dataset (Hollis et al., 2019).

**Figure S5.** Map showing the spatial position of GMS garden sites included in analyses. We exclude sites in Northern Ireland and

site-years which did not complete a full 33–36 week set of trap nights. Colours illustrate variation in the number of years included in analyses at each site.

**How to cite this article:** O'Connell-Booth, R., Hassall, C. & Kunin, W.E. (2025) Effect of bulb type on moth trap catch and composition in UK gardens. *Insect Conservation and Diversity*, 1–14. Available from: <https://doi.org/10.1111/icad.70050>