

An age-associated decline in the role of the sarcoplasmic reticulum and associated calcium-handling proteins sets the pace for sinoatrial node function


Sandra A. Jones¹, Fiona S. Godbeer¹ and Matthew K. Lancaster²

¹Centre for Biomedicine, Hull York Medical School, Faculty of Health Sciences, University of Hull, Kingston-upon-Hull, UK

²School of Biomedical Sciences, Faculty of Biological Sciences, University of Leeds, Leeds, UK

Handling Editors: Eleonora Grandi & Manuel Navedo

The peer review history is available in the Supporting Information section of this article (<https://doi.org/10.1113/JP290221#support-information-section>).



Sandra A. Jones became interested in ageing research during her PhD at the University of Liverpool and this further developed to target the heart during her subsequent postdoctoral work at the University of Leeds. In 2007, she moved to a Lectureship position at the University of Hull to develop her own research group and was later promoted to Senior Lecturer before joining the Hull York Medical School in 2022. Her research revolves around changes in cardiac cellular function with increasing age not only from the perspective of normal physiology, but also uncovering the mechanisms responsible for the ever-increasing age-associated risk of cardiac arrhythmias, particularly atrial fibrillation.



Artificial intelligence generated content disclosure: No AI-assisted technologies were used to generate any content for this manuscript.

Abstract With advancing age, the intrinsic function of the sinoatrial node (SAN) declines, as a result of structural changes and changes in electrical regulation within the constitutive cells of the nodal tissue. This study examined changes to proteins involved in regulating calcium flux balance in the atria and SAN of male rats used as a model of ageing throughout their lifespan at 6, 12 and 24 months of age. Using immunohistochemistry and western blotting, we determined a significant age-dependent decline in the levels of key calcium regulatory proteins within the SAN: $\text{Ca}_v1.2$, PMCA4, RYR2, SERCA2a and phospholamban ($n = 5$; ANOVA, $P < 0.05$). By contrast, levels of NCX protein were significantly elevated by 57.3% ($P = 0.009$) in the oldest group, indicating a potential pronounced change in calcium balance; a difference functionally observed by a steeper dose–response curve to the inhibitory effects of nifedipine. Intrinsic pacemaker beating rate was significantly reduced by 68 beats min^{-1} in the oldest group compared to the youngest, ($n = 6$; ANOVA, $P = 0.022$). Negating sarcoplasmic reticulum calcium cycling and the ‘calcium clock’ using cyclopiazonic acid reduced the intrinsic pacemaker activity of the SAN in young animals to that observed in the oldest group. Under these conditions, spontaneous activity and the response of the SAN to isoprenaline became matched across all age groups. Restoring sarcoplasmic reticulum function to the SAN in the elderly may offer a route to combatting age-related suppression of function, although care should be taken in the use of calcium channel antagonists to avoid precipitating sick-sinus syndrome.

(Received 30 September 2025; accepted after revision 29 January 2026; first published online 18 February 2026)

Corresponding author S. A. Jones: Centre for Biomedicine, Hull York Medical School, Faculty of Health Sciences, University of Hull, Kingston-upon-Hull, HU6 7RX, UK. Email: s.a.jones@hull.ac.uk

Key points

- This study adds understanding and characterisation of the age-dependent progressive reduction in sinoatrial node (SAN) function.
- We compared atria and the SAN across the lifespan of a Han Wistar rat model of ageing with animals studied at 6, 12 and 24 months of age.
- SAN tissue from old rats showed significantly reduced expression of key cellular calcium regulatory proteins compared to nodal tissue from young rats.
- The SAN from old rats exhibited a slowed intrinsic beating rate and an altered response to drugs that modulate cellular calcium handling.
- Eliminating sarcoplasmic reticulum calcium handling makes the SAN of a young animal mimic features of an aged animal.

Introduction

The sinoatrial node (SAN) is a spontaneously active, heterogeneous region of tissue located within the inter-caval region of the right atrium of the heart serving as the heart’s primary pacemaker. The intrinsic automaticity of the SAN and responsiveness to β -adrenergic stimulation declines in an age-correlated manner, contributing to a reduction in functional capacity and ability to respond to stress, as well as potentially increasing the risk of arrhythmias such as atrial fibrillation in later life (Fang et al., 2007; Fleg et al., 2005; Turner et al., 1999). Within 20 years, it is estimated over one-third of the UK population will be ≥ 65 years of age with $>10\%$ diagnosed with atrial arrhythmias, (ONS, 2025) resulting

in spiralling National Health Service costs (Burdett & Lip, 2022). The onset of such arrhythmias is particularly pronounced in males where the impact seems to become evident at an earlier age and so this group has been the initial focus of our study (Keller & Howlett, 2016). The aim of the present study was to add to our knowledge of the age-dependent progressive deterioration in SAN function aiming to identify mechanisms predisposing to rising dysfunction and risk of arrhythmias, with potential routes to preventative and direct treatment.

An unstable membrane voltage drives pacemaking within SAN myocytes. Hyperpolarisation-activated ion channels pass depolarising I_f current contributing to spontaneous depolarisation of SAN myocytes, supplemented in the late phase of diastole by spontaneous

releases of calcium from the sarcoplasmic reticulum (SR) (Bers & Lakatta, 2023). These calcium releases drive a small amount of sodium calcium exchange current (I_{NCX}) enhancing the rate of depolarisation to the membrane threshold for triggering a full action potential. This latter component of depolarisation is referred to as the ‘calcium clock’ and complements the overall ‘membrane clock’ in ensuring a robust capability for the heart rate, as driven by the SAN, to respond dynamically to stress. As such, it is recognised that the balance of calcium ion flux both across the cell membrane and within the SR of the cells of the SAN is a key component controlling SAN pacemaker function (Bers & Lakatta, 2023; Yue et al., 2020).

Adrenergic stimulation predominantly increases the heart rate directly or indirectly by causing elevations in cAMP. Numerous components key to SAN function are sensitive to cAMP and its actions in either directly changing ion-channel kinetics (e.g. by enhancing activation of I_f) or subsequently by its impact on calcium regulation (e.g. by increasing calcium loading of the SR by alterations to phospholamban and the calcium current as well as the overall action potential) (Bers & Lakatta, 2023; Tsutsui et al., 2018). We have previously shown that pacemaker cells from the guinea-pig SAN show an age-associated decline in $Ca_v1.2$ protein expression, (Jones et al., 2007), which is potentially key not only to limiting depolarisation of the cells, but also to reducing calcium-loading of the SR and thus modulating the calcium clock. We have now sought to track age-associated changes in other key calcium handling proteins that impact the SAN ‘calcium clock’ as we aim to further understand age-related changes in the basal function of the SAN and the response to adrenergic stimulation. Our study hypothesis is that a reduced basal beating rate and the dynamic response to adrenergic stimulation in the SAN of aged individuals is the result of a reduced capacity for the dynamic response of the ‘calcium clock’ limiting its ability to facilitate cardiac acceleration.

Methods

Ethical approval

In accordance with our Institution’s animal welfare committee guidelines animals had free access to water and food, appropriate housing, and were monitored to ensure they were healthy and displayed normal daily activities. Animals were killed by anaesthetic overdose (Euthatal; Duggan Veterinary Supplies Limited, Tipperary, Ireland; sodium pentobarbital i.p.) and death confirmed. All animal experimental procedures were performed in accordance with Home Office UK guidelines (United Kingdom Animals (Scientific Procedures) Act, 1986) with ethical review and approval from the local ethics committee (reference number, UoH 25-26.051).

SAN isolated from rats

Male Han Wistar rats (Charles River, Margate, UK) fed standard chow were randomised to differing age categories and raised to their appropriate age in caged social groups. Our rat model of ageing examined rats across their lifespan at the ages of 6 months (young), 12 months, and 24 months (old). Animals were killed at the appropriate age and the heart rapidly excised and immersed in oxygenated Tyrode solution with 95% O_2 /5% CO_2 , pH 7.4, 37°C. For experiments involving mapping the activity of the SAN, the right atrium was dissected from the heart and opened to expose the intercaval region and nodal tissue (Jones et al., 2004). Tissue architecture defined the SA node tissue as detailed in Fig. 1.

Extracellular electrode recordings

We used our published approach for recording sites of initiation and spread of activity across the SAN (Jones et al., 2004). Extracellular bipolar electrodes were positioned on the SAN leading pacemaker site (earliest point of activation) and pectinate muscle of the neighbouring right atrial tissue (Fig. 1) to record nodal activation (Neurolog bridge amplifier; Digitimer, Welwyn Garden City, UK; Digidata Axon 1440A with Clampex 10; Molecular Devices, Wokingham, UK) and determine the intrinsic heart rate (IHR). Movement of one of the electrodes in 1 mm steps across the x and y ranges of the tissue permitted a full map of activation timing to be produced across the tissue. Oxygenated




Figure 1. Endocardial surface of the right atria

Flanked by the right atrial appendage muscle and septum tissue, the dashed lines indicate the location of the crista terminalis (CT) and sinoatrial (SA) node tissue regions, between the upper the superior vena cava (SVC) and lower inferior vena cava (IVC). Scale bar = 5 mm.

Table 1. Primary antibodies

Protein	Catalog. no.	Company	Location	Primary host	Secondary antibody
Ca _v 1.2	ACC-003	Alomone Labs	Jerusalem, Israel	Rabbit	Swine anti-rabbit
Ca _v 1.3	ACC-005	Alomone Labs	Jerusalem, Israel	Rabbit	Swine anti-rabbit
Ca _v 3.1	ACC-021	Alomone Labs	Jerusalem, Israel	Rabbit	Swine anti-rabbit
SERCA2	MA3-910	Thermo Scientific	Altrincham, UK	Mouse	Rabbit anti-mouse
PMCA4	MA1-914	Thermo Scientific	Altrincham, UK	Mouse	Rabbit anti-mouse
RYR2	MA3-916	Thermo Scientific	Altrincham, UK	Mouse	Rabbit anti-mouse
NCX1	MA3-926	Thermo Scientific	Altrincham, UK	Mouse	Rabbit anti-mouse
PLB	A010	Badrilla	Leeds, UK	Mouse	Rabbit anti-mouse
Desmin	M 0760	Dako	Cheadle, UK	Mouse	Rabbit anti-mouse

Tyrode's solution maintained the tissue at a flow rate of 40 mL min⁻¹ with temperature monitored and controlled at 37°C. The kinetics of pacemaking are sensitive to temperature and so it is important for close regulation ($\pm 0.5^\circ\text{C}$) to ensure stable pacemaking in the normal physiological range. Pharmacological agents, isoprenaline hydrochloride, nifedipine and cyclopiazonic acid (CPA) (Sigma-Aldrich, St Louis, MO, USA) were made to 1 mM L⁻¹ stock solutions before dilution to the required concentration in Tyrode's solution. For each drug dose, the preparation was perfused for a minimum of 15 min to allow equilibration and to establish the steady-state IHR prior to recording. All chemicals were purchased from Merck Life Science UK Limited (Gillingham, UK).

Analysis of protein expression

Using our previously published method (Jones et al., 2004), tissue was frozen and a lysate was produced for western blotting. Proteins were separated using SDS-PAGE, transferred to polyvinylidene fluoride and incubated overnight at 4°C with the primary antibodies at 1:1000 dilution followed for 1 h by incubation with the appropriate secondary antibody conjugated to horseradish peroxidase (Dako, Cheadle, UK) (Table 1). A Molecular Imager Chemi Doc XRS+ (Bio-Rad, Hercules, CA, USA) was used to image band densities and associated values determined using ImageJ, version 1.54g (NIH, USA). Density values were normalised to an appropriate housekeeper protein to control for equal protein loading per sample.

As previously described for immunohistochemistry (IHC) (Jones et al., 2004), 10 μm frozen tissue slices were incubated overnight at 4°C with the primary antibodies (dilution 1:250, except for PLB, which was used at 1:100). Tissue was incubated for 2 h with a secondary antibody conjugated to Alexa Fluor 488 antibody (Invitrogen, Paisley, UK) and wheat germ agglutinin conjugated to rhodamine (2BSscientific, Kirtlington, UK). Labelled proteins were imaged using a plan apochromat 40 $\times$ /1.3 oil

immersion M27 lens on a LSM 710 confocal with images captured using Zen, version 3.12 (Zeiss Microscopy, Ltd, Cheadle, UK). Images were processed within the Zen software to allow separation of the fluorescence channels corresponding to the two labels before performing densitometric assessment of fluorescence intensity for regions. To control for background signal, an unlabelled parallel section of tissue from the sample was also measured using the same approach, and the result subtracted. For the protein of interest IHC labelling was collected on the same settings across each age group. Levels of fluorescence labelling are shown in some cases normalised relative to the mean value as determined in tissue from animals at 6 months of age.

Quantitative PCR (qPCR)

We used the RNeasy Fibrous Tissue Mini Kit (Qiagen, Hilden, Germany) to extract mRNA from samples of the rat right atria. Extracted mRNA concentration was assessed by absorbance (NanoDrop 1000; Thermo Fisher Scientific) prior to performing qPCR. Reverse transcriptase (Invitrogen) reverse transcribed RNA to complementary DNA (cDNA). Samples of cDNA were stored at 2–4°C for a maximum of 3 days. We detected the cDNA within each sample using a sequence-specific DNA probe. Primers were Desmin (rat NM_022531.1) and GAPDH (rat NM_017008.3) (Invitrogen). Transcript primers were optimised to ensure a cycle threshold (C_T) value of 29 or less, confirming a strong positive reaction indicative of abundant target nucleic acid. A StepOne Plus qPCR machine (Applied Biosystems, Waltham, MA, USA) was used with StepOne software to quantify expression in each sample.

Statistical analysis

Data are presented as the mean $\pm$ SD. Statistical significance was evaluated by Student's *t* test, one-way ANOVA or two-way repeated-measures (RM) ANOVA

Table 2. Morphological characteristics of the ageing rat model

	6 months	12 months	24 months
Body weight (g)	517 ± 35	556 ± 40	661 ± 125*
Heart weight (g)	1.55 ± 0.20	1.64 ± 0.07	1.63 ± 0.18
Heart: Body ($\times 10^{-3}$)	2.94 ± 0.33	2.96 ± 0.10	2.5 ± 0.27 ^{‡,¥}
Number in group (n)	11	6	11

* $P = 0.000772$, 24 months compared to 6 months of age.
[‡] $P = 0.000481$, 24 months compared to 6 months of age.
[¥] $P = 0.000113$, 24 months compared to 12 months of age.

with Holm–Sidak *post hoc* comparisons as appropriate. Alternatively, a Kruskal–Wallis comparison was used where required and indicated. $P < 0.05$ was considered statistically significant. Curve fits and statistical analysis were performed in Prism, version 10 (GraphPad Software Inc., San Diego, CA, USA) or Excel (Microsoft, Redmond, WA, USA) as appropriate. Unless otherwise stated, ‘n’ is the number of animals. Where appropriate, data are shown relative to that obtained from samples from the youngest age group at 6 months of age. For protein analysis, it is assumed that the 6 months age group possessed 100% of the specific protein of interest to enable calculation of percentage change relative to that age point.

Results

Age-associated changes in morphological characteristics

We compared tissue obtained from healthy male rats across three age groups: 6, 12 and 24 months of age (Table 2). The data showed that rat body weight gradually increased in an age-correlated manner ($r^2 = 0.3966$; $P = 0.0003$ by linear regression). A comparison of rats aged 6 and 24 months revealed a significant increase in body weight from 517 ± 35 g to 661 ± 125 g ($n = 11$; $P = 0.000772$) as the body weight significantly increased between the ages of 6 and 12 months ($P = 0.0416$) and 12 and 24 months ($P = 0.0120$). By contrast, heart weight did not change significantly from 1.55 ± 0.20 g across the age groups of 6, 12 and 24 months (ANOVA, $P = 0.375$) (Table 2).

Rats of 6 months and 12 months of age exhibited no significant difference in their heart-to-body weight ratio (ratio of 6 month rat $2.94 \pm 0.33 \times 10^{-3}$; $n = 6$; t test, $P = 0.244$), whereas rats aged 24 months revealed a significant reduction in their heart-to-body weight ratio ($2.5 \pm 0.27 \times 10^{-3}$) compared to the other age groups of 6 months and 12 months ($n = 11$; ANOVA, 24 vs. 6 months, $P = 0.000481$; 24 vs. 12 months, $P = 0.000113$) (Table 2).

Table 3. Desmin and GAPDH RNA values in right atrial tissue from rats of increasing ages

Animal age (months)	RNA C_T values (mean ± SD)	
	Desmin	GAPDH
6	23.6 ± 0.45	21.8 ± 0.55
12	23.9 ± 0.35	21.8 ± 0.49
24	23.6 ± 0.47	20.8 ± 0.35

Table 4. Desmin protein density

Animal age (months)	Tissue type	Desmin protein (density/pixel)
6	Left atria	3.15 ± 0.25
12	Left atria	3.12 ± 0.19
24	Left atria	3.88 ± 0.58
6	Right atria	3.53 ± 0.71
12	Right atria	2.83 ± 0.40
24	Right atria	3.00 ± 0.59

Confirming a suitable housekeeper reference for our ageing studies

We determined mRNA expression of the muscle-specific protein desmin and metabolic enzyme GAPDH for variance with increasing age (Table 3) using our qPCR protocol. Maximum C_T differences in the right atrial tissue for desmin RNA was 0.3 C_T with no significant changes over the examined age range ($n = 5$; ANOVA, $P = 0.314$), whereas GAPDH significantly reduced by 1.0 C_T with ageing ($n = 5$; ANOVA, 6 vs. 24 months, $P = 0.024$; 12 vs. 24 months, $P = 0.021$). We also examined desmin protein levels in paired right atrial (RA) and left atrial (LA) tissue obtained from rats at 6, 12 and 24 months of age (Table 4). There was no significant change in desmin levels ($n = 5$; ANOVA, $P = 0.691$); thus, desmin was used as the reference housekeeper protein for western blotting in this study.

Age-associated changes in calcium channels in the rat right atria

IHC, as performed on RA tissue from rats aged 6 and 24 months, showed age-associated changes in the proteins $Ca_v1.2$, $Ca_v1.3$ and $Ca_v3.1$ (Fig. 2). Normalised to the young, the old rat $Ca_v1.2$ protein levels significantly declined to $50.2 \pm 4.72\%$, whereas $Ca_v1.3$ and $Ca_v3.1$ levels increased by $81.6 \pm 10.14\%$ and $43 \pm 23.42\%$, respectively (mean ± SD; ANOVA, $P = 0.0001$).

Age-associated changes in Cav1.2 protein levels across the rat atria

RA and LA tissue from rats aged 6, 12 and 24 months were analysed by western blotting (Fig. 3). $\text{Ca}_v1.2$ protein levels in the RA showed a significant drop with increasing age to $53.6 \pm 14.1\%$ of that in the young ($n = 5$; ANOVA, $P = 0.011$), but, in the LA no significant change in expression was seen ($P = 0.051$).

Age-dependent decline in $\text{Ca}_v1.2$ ion channel expression in the SAN

Western blot analysis of $\text{Ca}_v1.2$ channel protein expression within the SAN of rats at 12 months of age showed

no significant differences to that found in the node of those aged 6 months ($100 \pm 24.4\%$; ANOVA, $P = 0.207$) (Fig. 4A). At 24 months, $\text{Ca}_v1.2$ expression levels had significantly declined to $59.6 \pm 18.6\%$ of that observed in the 6 months age group (ANOVA, 12 vs. 24 months, $P = 0.004$; 6 vs. 24 months, $P = 0.0315$) (Fig. 4A). We observed a striated pattern by IHC $\text{Ca}_v1.2$ protein labelling within SAN tissue in all age groups; however, the amount of labelled protein observed significantly declined from 6 months ($100 \pm 2.2\%$) to $55.6 \pm 5.8\%$ at 24 months of age ($P = 2.53 \times 10^{-7}$) (Fig. 4B). The data therefore consistently show a substantial age-dependent decline of $\text{Ca}_v1.2$ protein expression within the SAN region in hearts once animals reach 24 months of age compared to that identified within the young age group (6 months).




Figure 2. Changes in IHC calcium channel label within RA at 24 months of age

A, optical slices of RA tissue from rats at 6 and 24 months of age with fluorescence labelling of $\text{Ca}_v1.2$, $\text{Ca}_v1.3$ and $\text{Ca}_v3.1$. Confocal images, scale bar = 20 μm . B, data shown as the mean $\pm$ SD difference per calcium channel, overlaid with raw data points. Number of rats per group: $\text{Ca}_v1.2$ ($n = 10$), $\text{Ca}_v1.3$ ($n = 5$) and $\text{Ca}_v3.1$ ($n = 9$), with statistically significant changes in each case relative to that seen in tissue from young animals (ANOVA, $P = 0.0001$).

Modulation of $\text{Ca}_v1.2$ function and activity of the SAN

The intrinsic heart rate of the isolated SAN was assessed and impact of blockade of $\text{Ca}_v1.2$ channels determined by application of the selective channel blocker nifedipine across the dose range 0.1–30 μM . Dose–response relationships for the actions of nifedipine in each age group were established using a Hill curve to fit the intrinsic heart rate data and the nifedipine dose, plotted using a standard $\log_{10}$ scale for each concentration trialled (Fig. 4C). The SAN from the youngest age group of 6 months showed a dose-dependent response to nifedipine with Hill slope coefficient of 0.94; the maximum dose of 30 μM nifedipine caused all spontaneous activity to cease. The SAN from the oldest age group of 24 months had a reduced response to 0.1 μM nifedipine, and 0.3 μM caused a dramatic decline in beating rate with complete cessation at 1 μM nifedipine.

The dose–response relationship in the nodes from animals at 24 months of age had an average Hill slope coefficient of 4.2. The IC_{50} for nifedipine in the older age group was not significantly different to that seen in the young (IC_{50} ; 1.14 μM at 6 months vs. 1.23 μM at 24 months). Therefore, although the inhibitory binding of nifedipine for $\text{Ca}_v1.2$ is comparable for each age group, the impact on beating rate within the old age group has an approximately four-fold steeper relationship than that seen at a young age.

Age-dependent changes of calcium handling protein within the SAN

Faced with a reduced capacity for calcium influx, it is reasonable to expect extrusion pathways may be modulated to maintain calcium balance in nodal tissue from elderly animals. Intracellular balance of




Figure 3. Age-associated changes of $\text{Ca}_v1.2$ protein across the atria

A, illustrative blot of $\text{Ca}_v1.2$ protein levels per age group for both the RA and LA. B, data shown as the mean $\pm$ SD per age group, overlaid with raw data points ($n = 5$ animals at each age; ANOVA, $*P = 0.011$ for 24 vs. 6 months).

Ca^{2+} is maintained by extrusion via sodium-calcium exchanger (NCX1) and plasma membrane calcium ATPase 4 (PMCA4) with sarcoendoplasmic reticulum Ca^{2+} ATPase2a (SERCA2a) maintaining internal calcium buffering by pumping calcium back into the SR store,

although this is modulated by the inhibitory action of phospholamban (PLB).

Analysed by western blotting NCX1 protein levels within the SAN did not alter significantly between the ages 6 to 12 months, but, in rats aged 24 months, NCX1




Figure 4. Age-associated changes of $\text{Ca}_v1.2$ protein and activity within the SAN

A, illustrative blot of $\text{Ca}_v1.2$ protein levels in the RA at 6, 12 and 24 months. Data are shown as the mean $\pm$ SD with raw data points ($n = 5$ per group; 12 vs. 24 months $P = 0.004$, 6 vs. 24 months $P = 0.0315$ by ANOVA). B, optical slices from SAN of rats aged 6 and 24 months with $\text{Ca}_v1.2$ protein labelled in green and membranes labelled with wheat germ agglutinin (red). Scale bar = 50 μm . Quantitated labelling density per group with raw data is shown below (mean $\pm$ SD; $n = 5$ animals per age group; 6 vs. 24 months $P = 2.53 \times 10^{-7}$ ANOVA); C, Hill equation fitted to intrinsic heart rate from rats aged 6 months and 24 months with incrementing concentrations of nifedipine (shown as $\log_{10}$) ($n = 6$ per group; $\text{IC}_{50} = 1.14 \mu\text{M}$ at 6 months vs. $1.23 \mu\text{M}$ at 24 months).

levels were significantly increased to $157 \pm 21.5\%$ of that found in tissue from the 6 months group with levels also being elevated significantly relative to that seen at 12 months (ANOVA, 12 vs. 24 months, $P = 0.024$; 6 vs. 24 months, $P = 0.00856$) (Fig. 5B). Analysis of immunolabelling of NCX1 protein within the SAN region identified a similar significant rise with age to $142 \pm 15.6\%$ at 24 months compared with $100 \pm 12.8\%$ at 6 months of age ($P = 0.0008$) (Fig. 6). Using western blotting, we identified PCMA4 calcium pump protein levels within the SAN increased significantly from 6 to 12 months to $122 \pm 17.4\%$ but exhibited a significant decline to $78 \pm 3.0\%$ of the amount at 6 months in rats aged 24 months (6 vs. 12 months, $P = 0.029$; 12 vs. 24 months, $P = 0.002$; 6 vs. 24 months, $P = 0.010$) (Fig. 5C). IHC labelling of PCMA4 within the SAN region also showed a similar significant decline with age to $76.9 \pm 16.3\%$ at 24 months compared with $100 \pm 11.1\%$ at 6 months of age ($P = 0.0157$) (Fig. 6).

Both ryanodine receptor 2 (RYR2) and SERCA2a also showed significant changes in expression with age. Western blotting showed the RYR2 protein levels increased significantly to $135 \pm 18.8\%$ within the SAN from rats aged 12 months compared with tissue from those aged 6 months, but then, in rats aged 24 months, levels significantly declined to $68 \pm 23.2\%$ of the level in tissue from rats at 6 months of age (6 vs. 12 months, $P = 0.014$; 12 vs. 24 months, $P = 0.0005$; 6 vs. 24 months, $P = 0.029$) (Fig. 5D). IHC labelling of RYR2 protein within the SAN region also showed a significant decrease to $72.6 \pm 13.4\%$ at 24 months compared to $100 \pm 13.7\%$ at 6 months of age ($P = 0.006$). SERCA2a levels significantly increased to $144 \pm 39.4\%$ within the SAN from rats aged 12 months compared to those aged 6 months but then declined significantly to $64 \pm 14.7\%$ in tissue from rats at 24 months of age (6 vs. 12 months, $P = 0.035$; 12 vs. 24 months, $P = 0.003$; 6 vs. 24 months, $P = 0.002$) (Fig. 5E). IHC labelling of SERCA2a within the SAN region also showed a significant decrease with age to $71.7 \pm 15.3\%$ at 24 months compared to $100 \pm 14.3\%$ at 6 months of age ($P = 0.008$) (Fig. 6).

Age-dependent changes in expression of phospholamban in the SAN

PLB is expressed as monomer but can dynamically establish a pentameric assembly that disassembles upon phosphorylation in response to adrenergic stimulation, where the monomeric form serves as a more effective regulatory inhibitory protein modulating SERCA pump activity (MacLennan & Kranias, 2003). We quantified monomeric PLB protein levels within the SAN (Fig. 7B) and these did not significantly alter between the ages of 6 and 12 months ($P = 0.38$). Monomeric PLB expression

however significantly fell in SAN tissue from rats aged 24 months to $58.6 \pm 11.3\%$, with this fall being significant compared with levels in SAN from rats at 6 and 12 months of age ($n = 5$; 6 vs. 24 months, $P = 0.0016$; 12 vs. 24 months, $P = 0.0017$).

By comparison, pentameric PLB protein levels increased from $100 \pm 20.8\%$ at 6 months to $119.1 \pm 22.3\%$ at 12 months of age ($P = 0.09$), then, in rats aged 24 months, significantly declined to $72.4 \pm 21.0\%$ (12 vs. 24 months, $P = 0.004$; 6 vs. 24 months, $P = 0.035$) (Fig. 7C). Total PLB levels did not alter from 100% between the ages 6 to 12 months ($P = 0.47$) but significantly decreased to $68.9 \pm 19.3\%$ in rats aged 24 months (12 vs. 24 months, $P = 0.0120$; 6 vs. 24 months, $P = 0.0123$) (Fig. 7D). This age-dependent decrease between young and old rats was also observed in our total PLB protein immunolabelling of the SAN because label density significantly fell from 6 months ($100 \pm 16.5\%$) to $78.7 \pm 15.7\%$ at 24 months (6 vs. 24 months, $P = 0.035$) (Fig. 6). The ratio of SERCA pump protein to PLB monomer (SERCA:PLB) is considered indicative of activity and was determined as 1.0 ± 0.31 at 6 months and significantly rose at 12 months to 1.5 ± 0.148 , indicative of greater tonic SERCA inhibition (6 vs. 12 months, $P = 0.0080$) (Fig. 7E). From 12 months onwards, the SERCA:PLB ratio declined significantly to 1.2 ± 0.17 at 24 months (12 vs. 24 months, $P = 0.0125$). The SERCA:PLB ratio observed in tissue from rats aged 6 months was not significantly different to that observed in 24 month aged animals ($P = 0.10$).

Age-dependent decline in SERCA2a activity of the SAN and pacemaker dynamics

The IHR of the isolated SAN preparation was measured during steady-state spontaneous pacemaking under control conditions and during incrementing β -adrenergic stimulation for each of the three age groups. The data showed an age-associated reduction in basal intrinsic pacemaking (Fig. 8A; see also summary in Table 5) with intrinsic heart rate showing a linearly correlated reduction with age ($r^2 = 0.607$; $P = 0.0017$) falling from 274 ± 33 beats min^{-1} at 6 months to 206 ± 19 beats min^{-1} at 24 months of age. The EC_{50} to isoprenaline, however, remained unchanged with age (13.4 nM at 6 months and 13.7 nM at 24 months) (Fig. 8A). At $1 \mu\text{M}$ isoprenaline, pacemaking differences were abolished across all age groups with intrinsic rates rising to 386 ± 36 and 368 ± 62 beats min^{-1} in the youngest and oldest age groups, respectively, indicating that a maximal response to the highest concentration of isoprenaline remains viable even in the SAN of the oldest rats that we studied.

To assess the functional involvement of the SR during β -adrenergic stimulation across the age groups, $3 \mu\text{M}$




Figure 5. Effect of ageing on levels of calcium handling proteins expressed in the SAN

A, illustrative western blot of nodal tissue examined for each protein of interest per age group with the desmin housekeeper. B–D, mean $\pm$ SD per age group ($n = 5$ animals per group) overlaid with raw data points for nodal expression levels of calcium handling proteins; **NCX1** (B) (ANOVA, 12 vs. 24 months, $*P = 0.02$; 6 vs. 24 months, $\#P = 0.00856$), **PMCA4** (C) (ANOVA, 6 vs. 12 months, $*P = 0.029$; 12 vs. 24 months, $\Delta P = 0.002$; 6 vs. 24 months, $\#P = 0.010$), **RYR2** (D) (ANOVA, 6 vs. 12 months, $*P = 0.014$; 12 vs. 24 months, $\Delta P = 0.0005$; 6 vs. 24 months, $\#P = 0.029$) and **SERCA2A** (E) (6 vs. 12 months, $*P = 0.035$; 12 vs. 24 months, $\Delta P = 0.003$; 6 vs. 24 months, $\#P = 0.002$).




Figure 6. Age-dependent changes of proteins labelled within the sinoatrial region

A, optical slice of each section taken through the SAN regions from rats aged 6 and 24 months were labelled to show expression of specific proteins of interest (green) and all membranes (using wheat germ agglutinin, red).

Confocal images scale bar = 50 μm . *B*, quantified fluorescence from IHC expressed relative to signal density from labelled sections of nodal tissue from rats at 6 months of age (mean $\pm$ SD; $n = 5$ animals; Student's *t* test, NCX1, * $P = 0.0008$; PMCA4, $\Delta P = 0.0157$; RYR2, $\#P = 0.006$; SERCA2A, $\diamond P = 0.008$; total PLB, $\blacktriangle P = 0.035$).

CPA was applied to inhibit SERCA2a and deplete the SR. CPA significantly reduced spontaneous activity at 6 and 12 months ($P = 0.023$ and 0.044 , respectively; RM ANOVA) (Fig. 8A and Table 5). Interestingly, in the oldest group of 24 months of age, CPA did not significantly reduce the basal intrinsic rate ($P = 0.904$). Intrinsic heart rate was no longer significantly different across the age groups in the presence of CPA ($P = 0.307$; RM ANOVA; $n = 6$ in each group). CPA significantly reduced the response to isoprenaline in all age groups. In the presence of CPA, the EC_{50} for isoprenaline did not significantly differ across age groups under control conditions or in the presence of CPA.

Activation maps for each SAN studied were constructed and the leading pacemaker site located during the steady-state response to each condition. Isoprenaline caused a significant average shift in pacemaker location of 3.6 ± 1.6 mm ($P < 0.0001$, $n = 6$ in each group) (Fig. 8B). Application of CPA caused a small but significant shift in the leading pacemaker site (1.2 ± 0.9 mm; $P = 0.0003$) and the significant isoprenaline-dependent shift in the leading pacemaker site was maintained on subsequent application of isoprenaline in the presence of CPA in all age groups (3.78 ± 16 mm; $P < 0.0001$). Comparing between age groups, there was no significant difference in pacemaker shift in response to isoprenaline, CPA or the combination between age groups.

Examining the velocity of conduction between the leading pacemaker site and crista terminalis under control conditions a significant reduction in conduction velocity was seen across the SAN in the oldest group compared with tissue from the other two ages. Average conduction velocity was 4.4 ± 0.25 m.s $^{-1}$ at 6 months and 4.6 ± 0.60 m.s $^{-1}$ at 12 months, falling significantly to 3.8 ± 0.23 m.s $^{-1}$ at 24 months ($P = 0.041$ and 0.015 for 24 vs. 6 months and vs. 12 months, respectively, $n = 6$ in each case).

A dose-dependent significant increase in conduction velocity in response to isoprenaline was observed (Fig. 8C) in all age groups with an average increase in velocity of 3.02 ± 1.07 m.s $^{-1}$ ($P = <0.0001$) in the presence of $1 \mu\text{M}$ isoprenaline. The magnitude of the increase in velocity in response to isoprenaline differed significantly between all age groups, with the largest increase seen in tissue from animals at 6 months of age where velocity increased by 5.1 ± 2.1 m.s $^{-1}$ in response to $1 \mu\text{M}$ isoprenaline, but 3.0 ± 1.1 m.s $^{-1}$ for tissue from animals at 12 months and 1.9 ± 0.5 m.s $^{-1}$ for tissue from animals at 24 months ($P < 0.0001$ for 6 vs. 24 months and $P = 0.001$ for 12 vs. 24 months).

CPA did not significantly impact conduction velocity under control conditions ($P = 0.989$) but did abolish the significant impact of isoprenaline on conduction velocity in SAN tissue from all age groups, with the magnitude of increase in the presence of CPA falling to 1.25 ± 0.91 m.s $^{-1}$; $P = 0.160$). In the presence of CPA, there were no significant differences between age groups in response to isoprenaline.

Discussion

The decline in maximal heart rate and increasing risk of pacemaker dysfunction is a ubiquitous feature of ageing, as has been seen in many animal models and human studies (Peters et al., 2020). Previous studies by ourselves and others have correlated this with morphological changes of the SAN, as well as changes in ion channel expression (Jones et al., 2004, 2007; Moghtadaei et al., 2016; Monfredi & Boyett, 2015). Alongside this, an altered response to adrenergic stimulation has been documented, which may be closely linked to an impaired ability of cAMP to modulate the 'calcium clock' component of SAN pacemaking and so cause acceleration of the SAN (Liu et al., 2014; Segal et al., 2023). In the present study, we have investigated age-associated changes in key components of cellular calcium regulation on pacemaking response, as well as the effect of entirely removing the SR contribution to pacemaking. The results reveal a SAN in the elderly that operates at an intrinsically lower spontaneous frequency, but, when the calcium clock is disrupted, age-associated differences in pacing rate and adrenergic response are removed. The detailed mapping of protein changes and functional adaptations identify a largely balanced scaling of changes with age preserving function and responsiveness in the node but, ultimately, also show that dysregulation of the 'calcium clock' may underlie key age-associated changes in functional capacity, tolerance and response.

Right atria and ageing

Our male Wistar rat model showed that post-maturity with increasing age their body weight increased significantly, as expected, but the heart weight did not change significantly with age. Previous work in rats and other species, including humans, has reported either no change in heart weight simply with age (Hindso et al., 2017) or, more commonly, age-associated hypertrophy that may be secondary to hypertension and body mass increases with age (Linzbach & Akumoa-Boateng, 1973).




Figure 7. Age-dependent changes in phospholamban protein in the SAN

A–D, illustrative western blot and protein levels per age group of total PLB, alongside separated data for pentameric and monomeric forms normalised to desmin (mean $\pm$ SD; $n = 5$; ANOVA). Protein levels declined in SA nodes from rats age 24 months for (B) monomer PLB protein levels (6 vs. 24 months, $*P = 0.0016$; 12 vs. 24 months, $\#P = 0.0017$), (C) pentamer PLB protein levels (12 vs. 24 months, $*P = 0.004$; 6 vs. 24 months, $\#P = 0.035$) and (D) total PLB protein levels (6 vs. 24 months, $*P = 0.0123$; 12 vs. 24 months, $\#P = 0.0120$). E, SERCA:PLB ratio protein levels increased in SA nodes from rats at 12 months of age, then declined back to the 6 month rat levels (6 vs. 12 months, $*P = 0.0080$; 12 vs. 24 months, $\#P = 0.0125$; 6 vs. 24 months, $P = 0.10$).




Figure 8. Isoprenaline-dependent response of nodal function across the age groups

A, percentage change in intrinsic heart rate (beats min^{-1}) (IHR) of the isolated SAN in response to isoprenaline. The beating rate as determined in tissue from rats aged 6 months under control conditions was taken as 100%. Upper traces show the % change (mean $\pm$ SD) for each age group in response to incremental doses of isoprenaline. Lower traces show the effects of isoprenaline in the presence of 3 μM CPA. B, magnitude of shift in the leading

pacemaker site on application of 1 μM isoprenaline, 3 μM CPA or the combination of both. C, conduction velocity across the node from the leading pacemaker site to the crista terminalis during application of different doses of isoprenaline alone or in the presence of 3 μM CPA.

Table 5. Steady-state intrinsic heart rate (IHR) of isolated SAN with and without maximal β -adrenergic stimulation and/or cyclopiazonic acid (CPA) to block SERCA2A

	6 months	12 months	24 months
IHR (beats min^{-1})	274 $\pm$ 33	256 $\pm$ 25	206 $\pm$ 190*
Maximal rate (1 μM isoprenaline)	386 $\pm$ 36 [#]	402 $\pm$ 22 [#]	368 $\pm$ 62 [#]
IHR in CPA	216 $\pm$ 26	204 $\pm$ 9	189 $\pm$ 27
Maximal rate in CPA (1 μM isoprenaline)	258 $\pm$ 30	245 $\pm$ 22	240 $\pm$ 40

* $P = 0.022$ comparing the rate in 24 months vs. 6 months of age.

[#] $P \leq 0.0001$ for rate in 1 μM isoprenaline vs. intrinsic heart rate in the same age group.

Other studies have even indicated potential reductions in cardiac mass relative to lean body mass in old age (Olivetti et al., 1991), but, essentially, it is the metabolically active tissue that is expected to be the key driver of cardiac output.

Across our studies, we have used the muscle-specific protein desmin to normalise our western blot data, (Jones et al., 2004, 2007) because both mRNA and protein levels of desmin do not change with age, as shown in the rat atria (Table 4), whereas the metabolically active protein GAPDH showed significant changes in levels of mRNA with increasing age within the RA (Table 3). Our initial studies of calcium ion channels within the rat RA showed an age-associated significant increase of $\text{Ca}_v1.3$ and $\text{Ca}_v3.1$ protein levels, and a significant decline in $\text{Ca}_v1.2$, whereas $\text{Ca}_v1.2$ did not alter with age in the LA (Figs 2 and 3). Our key focus, however, has been the SAN within the RA and the age-related changes in calcium regulation that may be preponderant there.

Age-dependent changes in calcium regulators

We show a significant loss of $\text{Ca}_v1.2$ channel protein from within the SAN of the rat during advancing age. Such a change will potentially result in reduced excitability of the cells of the node, as well as reduced calcium-loading of the SR, potentially limiting the ability of the SR to modulate pacemaking via the calcium clock. These channels are not only key for the upstroke of the action potential in SAN myocytes once threshold is reached, but also serve to service the calcium loading of the cell, thus directly modulating the functioning of the 'calcium clock', as well as the funny current I_f indirectly (Hagiwara & Irisawa, 1989). Of considerable interest is the observed shift in association between pharmacological modulation of the current through $\text{Ca}_v1.2$ using nifedipine and the ensuing impact on heart rate. Although the EC_{50} does not shift, implying the interaction with the channel in

terms of binding is conserved, the four-fold shift on the Hill coefficient indicates a notable change in the relationship between the current and pacemaking. The elevated sensitivity may in part be simply be a result of the overall drop in channel density, as well as reflecting a shift in the links between membrane calcium flux via $\text{Ca}_v1.2$ and pacemaking. The relationship between pacemaking and this flux is complex to assess because this not only acts as a depolarising influence, but also critically a key loading flux for the SR, which we see is also altering in influence in old age, together with other elements that ensure calcium balance such as NCX. The steep sensitivity implies a reduced capacity to deal with flux variation and still maintain calcium balance within a suitable range to ensure continued stable pacemaking. In the face of an already reduced inward calcium flux and in tandem with reduced SR function, the node experiences a steep reduction in potential depolarising current from both the calcium current and SR calcium releases, with the net impact observed here.

Age and changes in the balance of 'calcium clock' components

Analysing the components of the SAN calcium clock, we see broad depression in the expression of all elements associated directly with the SR and its function. Calcium release and uptake proteins are reduced by a similar magnitude in old age, indicating possible co-regulation to ensure maintained equilibria essential for continued stable function because cumulative imbalance of calcium flux will rapidly lead to arrhythmias or cell death. SERCA2a is key for maintaining loading of the SR with calcium and is a key controller of diastolic calcium in partnership with calcium entry via the L-type calcium channel. Modulating this relationship is the influx of calcium, and the activity of SERCA2a modulated by PLB in response to production of cAMP triggered by adrenergic stimulation (Capel &

Terrar, 2015). Our data shows SERCA, RYR and PLB all fell proportionately in old age, maintaining this control equilibrium and perhaps being more indicative of a general loss in SR density rather than shifting control. As a potential means to ensure maintained calcium balance, even though L-type calcium channel expression is also falling, NCX rises. As such, in old age, the scope for calcium release from the SR to dynamically respond to pacemaking is probably suppressed, although the elevated NCX may be sufficient to facilitate the basal contribution of depolarising current secondary to SR release (Groenke et al., 2013), which is potentially occurring at a reduced density. Previous work in an over-expressor model of NCX showed elevations in NCX to be capable of increasing the dynamic response to adrenergic stimulation (Kaese et al., 2017). Here, we do not see such an enhancement but the elevation in NCX is most probably key in maintaining the response we do see in the face of a falling potential SR component driving pacemaking.

Impact of stopping the 'calcium clock' leaving the membrane one to run the beat

The key role of the calcium clock in regulating the SAN is seen when we effectively remove the function of the SR using CPA to inhibit SERCA function. This only impacted basal pacemaking of the SAN significantly in tissue from the younger groups. By contrast, CPA had insignificant impact on SAN activity in tissue from elderly animals, indicating a key shift in the influence of the calcium clock on activity of the node. One surprising observation is that, despite the apparent lack of a role for the SR in maintaining the pacemaker rate at baseline in the old animals, they do show an apparently normal dose–response to isoprenaline with a maintained EC_{50} , and this response remains comparable between age groups in the presence of CPA. The implication is that the signal transduction processes responsible for the adrenergic response remain perfectly functional in old age. We have not looked in detail at each part of the signal transduction process, but other studies have similarly found preserved potential for a response (Liu et al., 2014) even though kinetics and sensitivity may be subtly altered at various stages of the process from catecholamine interaction with receptors through to cAMP generation (Xiao et al., 1998) and equilibrium with phosphodiesterases (Yaniv et al., 2016). Despite this preserved potential for a response in the isolated SAN, maximal heart rate in the intact organism is depressed with age, indicating key aspects such as neural input and metabolic regulation, which may be differentially modulated or limited in the intact organism, may be key for developing our understanding of why peak heart rate is suppressed in the elderly.

A key observation is that with SR function suppressed the response to isoprenaline at all ages becomes equivalent, although much depressed in range. Thus, it appears, in the male rat at least, modulation of I_f by cAMP is insufficient alone to drive a normal full pacemaker response to adrenergic input. The ~70% reduction in the dynamic response to isoprenaline seen in the SAN with SERCA inhibition present suggests that, in the rat at least, activation of HCN (i.e. hyperpolarisation-activated cyclic nucleotide-gated) channels and the membrane clock can only achieve ~30% of the normal increase in activation frequency. This balance in the dynamic response appears to be preserved across the age range even though basal activity is suppressed in SAN tissue from the oldest animals that we studied, again implying balanced regulation across the age range rather than identifying a distinct imbalance arising that can be targeted to reduce the impact of ageing.

Impact on pacemaker response and conduction

We have conducted our experiments using the intact functioning SAN. A key regulator of pacemaker activity is not just seen at the individual cell level because the activation of the node and its ability to drive the pacing of the heart is dependent on the electrotonic interactions between cells of the leading pacemaker site and surrounding tissue. Changes in pacemaker frequency are commonly associated with shifts in dominant pacemaker location and propagation of activation across the tissue. At its most extreme, there are suggestions that, in some species, including humans, distinct foci for pacemaking depending on frequency may exist (Brennan et al., 2020) representing areas with differing expression of ion channels and/or electrotonic interaction with the surrounding tissue modulated by frequency and focussed autonomic influence. In the isolated nodal preparations that we have used, we did not have focussed autonomic input, although we did still see significant shifts in the location of the leading pacemaker site when isoprenaline was applied, even when SR function was inhibited with CPA. This shift was preserved across all age groups. Despite a significant shift in basal pacemaker frequency in tissue from the younger ages studied, no shift in pacemaker location was observed when SR function was inhibited with CPA. This observation raises questions regarding what may drive pacemaker shift. If shifts in calcium balance and even the depolarisation frequency of the magnitudes we have seen with age and CPA are insufficient to cause pacemaker shift, then this may simply represent a difference in density of expression of adrenergic receptors or other elements of the transduction machinery for actioning the adrenergic response.

An unexpected observation was the shift in conduction velocity across the node with isoprenaline. An increase in conduction velocity is expected as the upstroke velocity and amplitude of the action potential is increased with adrenergic stimulation, although this was only observed in nodal tissue from the younger animals. The failure of isoprenaline to be able to increase conduction velocity in the oldest group may be indicative of the loss of calcium channels impacting the upstroke of the action potential, or equally may reflect loss of connexins key to permit rapid electrical propagation (Jones et al., 2001). Irrespective of the reason, these data indicate a limitation for supporting stable higher-frequency activation of the node and surrounding tissue because of limits in the rate at which the activation can spread to the atria. This has the potential to be a key issue for the development of sick-sinus syndrome and atrial fibrillation, although future work making more-detailed subregional analysis all across the node is required to clearly identify how this failure of conduction is arising.

Clinical implications and implications for human ageing

The identified shifts in calcium regulatory processes and the baseline pacemaker rate, as well as the suppressed ability to enhance conduction velocity, are key issues to consider in the perspective of the aged human heart. Calcium channel blockers such as verapamil are still used widely, mainly for their impact on vascular function, but the changing relationship between calcium balance and pacemaker function does indicate a developing danger. Indeed, the use of verapamil in patients with sick-sinus syndrome has previously been noted to have four to five times the impact on SAN function as it does when applied in control individuals, comprising a magnitude of shift comparable to our identified shift in the slope of the response of the isolated rat SAN to calcium channel block (Carrasco et al., 1978). The obvious clinical implication of this is to use calcium channel modulators with considerable caution in the elderly, paying careful attention to the potential for inducing and exacerbating sick-sinus syndrome.

In the light of broader human ageing, the implications remain that, until we can solve the issue of the continued drop in SAN function and peak heart rate with age, we cannot expect to be able to significantly prolong maximal lifespan without encountering pacemaker problems. Solutions do not appear to simply lie in restoring the response to adrenergic stimulation, but instead probably require maintenance or restoration of key aspects determining conduction through the SAN and capacity within the calcium regulatory processes, membrane and SR associated, aiming to ensure a stable

response and continued activity at suitable rates to meet requirements for adequate cardiac output.

Limitations and future work

One interesting feature of our findings was we did not see a significantly limited response to isoprenaline in aged tissue as generally reported by others in the intact heart and other preparations (Lakatta et al., 1975). In other cases, what has often been seen is a shift in the dose-response curve to adrenergic stimuli, whereas direct elevations of cAMP or saturating doses of agonists show that maximal responses are preserved, as seen here and, for example, in the study by Liu et al. (2014). As such, *in vivo*, a depressed maximal rate is common, but, *ex vivo*, it would appear that any differences in signal transduction arising from alterations to autonomic input or accumulation of agonists can be negated. The implication is that, as the end effector of heart rate elevation, the peak rate of nodal depolarisation remains capable of a full range of function in advanced age. Accordingly, to identify the source of depressed response, we need to look to manipulate the signal transduction pathway if we wish to restore youthful function. Having said this, we do see signs of potential for greater instability because of the changing balance of components regulating intracellular calcium buffering. Future experiments will not only aim to assess this more directly, but also identify steps on the signalling pathway that alter the relationship between cAMP production and activation of the integrated pacemaking mechanism of the calcium and membrane clocks in advanced age. A key feature to permit stable restoration of higher pacemaker rates to the aged SAN though would also appear to be modulation of conduction.

One limitation in translating these findings to the human condition is understanding that the shifts in the impact of differing mechanisms of pacemaker modulation and atrial excitability will vary between species and also ages. For example, in humans, the resting heart rate is normally kept reduced by parasympathetic input and, on removing this and observing the intrinsic rhythm, an acceleration is seen (Christou & Seals, 2008). In other animals, such as the rat, as studied here, the resting heart rate *in vivo* is maintained by tonic sympathetic input and, on removing this, the intrinsic rate is, in contrast to humans, lower than that seen *in vivo* (Di Gennaro et al., 1987). This latter response may also apply to the aged human heart (Christou & Seals, 2008), potentially making the rat heart more in-line with the aged human heart. These indicate a differing role for tonic sympathetic drive across ages and species, which will go hand-in-hand with associated signal transduction, response and the role of pacemaker mechanisms in facilitating the dynamic

response of the heartbeat. Further experiments on human tissue are required to refine our understanding of the relative impact of the membrane and calcium clocks across the SAN and how each is modulated by adrenergic stimulation and age.

Conclusions

In conclusion, our data show a changing relationship between pacemaker function and cellular calcium handling with advancing age. Although the identified changes also show preservation of a good level of calcium handling function and response to isoprenaline, there are indicators of an increased risk of instability and altered responses to physiological and pharmacological interventions. Pharmacological agents modulating the changing dynamics driving the SAN activity are routinely administered in the treatment of hypertension; however, because of an increasing risk of precipitating sinus node dysfunction, greater care is needed in consideration of their use in elderly patients.

References

- Bers, D., & Lakatta, E. G. (2023). What makes the sinoatrial node tick? A question not for the faint of heart. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, **378**(1879), 20220180.
- Brennan, J. A., Chen, Q., Gams, A., Dyavanapalli, J., Mendelowitz, D., Peng, W., & Efimov, I. R. (2020). Evidence of superior and inferior sinoatrial nodes in the mammalian heart. *Journal of the American College of Cardiology-Clinical Electrophysiology*, **6**(14), 1827–1840.
- Burdett, P., & Lip, G. Y. H. (2022). Atrial fibrillation in the UK: Predicting costs of an emerging epidemic recognizing and forecasting the cost drivers of atrial fibrillation-related costs. *European Heart Journal - Quality of Care and Clinical Outcomes*, **8**(2), 187–194.
- Capel, R. A., & Terrar, D. A. (2015). The importance of Ca(2+)-dependent mechanisms for the initiation of the heartbeat. *Frontiers in Physiology*, **6**, 80.
- Carrasco, H. A., Fuenmayor, A., Barboza, J. S., & Gonzalez, G. (1978). Effect of verapamil on normal sinoatrial node function and on sick sinus syndrome. *American Heart Journal*, **96**(6), 760–771.
- Christou, D. D., & Seals, D. R. (2008). Decreased maximal heart rate with aging is related to reduced {beta}-adrenergic responsiveness but is largely explained by a reduction in intrinsic heart rate. *Journal of Applied Physiology (Bethesda, Md : 1985)*, **105**(1), 24–29.
- Di Gennaro, M., Bernabei, R., Sgadari, A., Carosella, L., & Carbonin, P. U. (1987). Age-related differences in isolated rat sinus node function. *Basic Research in Cardiology*, **82**(6), 530–536.
- Fang, M. C., Chen, J., & Rich, M. W. (2007). Atrial fibrillation in the elderly. *The American Journal of Medicine*, **120**(6), 481–487.
- Fleg, J. L., Morrell, C. H., Bos, A. G., Brant, L. J., Talbot, L. A., Wright, J. G., & Lakatta, E. G. (2005). Accelerated longitudinal decline of aerobic capacity in healthy older adults. *Circulation*, **112**(5), 674–682.
- Groenke, S., Larson, E. D., Alber, S., Zhang, R., Lamp, S. T., Ren, X., Nakano, H., Jordan, M. C., Karagueuzian, H. S., Roos, K. P., Nakano, A., Proenza, C., Philipson, K. D., & Goldhaber, J. I. (2013). Complete atrial-specific knockout of sodium-calcium exchange eliminates sinoatrial node pacemaker activity. *PLoS ONE*, **8**(11), e81633.
- Hagiwara, N., & Irisawa, H. (1989). Modulation by intracellular Ca²⁺ of the hyperpolarization-activated inward current in rabbit single sino-atrial node cells. *The Journal of Physiology*, **409**(1), 121–141.
- Hindso, L., Fuchs, A., Kuhl, J. T., Nilsson, E. J., Sigvardsen, P. E., Kober, L., Nordestgaard, B. G., & Kofoed, K. F. (2017). Normal values of regional left ventricular myocardial thickness, mass and distribution-assessed by 320-detector computed tomography angiography in the Copenhagen General Population Study. *The International Journal of Cardiovascular Imaging*, **33**(3), 421–429.
- Jones, S. A., Boyett, M. R., & Lancaster, M. K. (2007). Declining into failure: The age-dependent loss of the L-type calcium channel within the sinoatrial node. *Circulation*, **115**(10), 1183–1190.
- Jones, S. A., Lancaster, M. K., & Boyett, M. R. (2001). Age-associated disappearance of connexin43 within the sinoatrial node. *Circulation*, **104**, 643.
- Jones, S. A., Lancaster, M. K., & Boyett, M. R. (2004). Ageing-related changes of connexins and conduction within the sinoatrial node. *The Journal of Physiology*, **560**(2), 429–437.
- Kaese, S., Bogeholz, N., Pauls, P., Decherling, D., Olligs, J., Kolker, K., Badawi, S., Frommeyer, G., Pott, C., & Eckardt, L. (2017). Increased sodium/calcium exchanger activity enhances beta-adrenergic-mediated increase in heart rate: Whole-heart study in a homozygous sodium/calcium exchanger overexpressor mouse model. *Heart Rhythm*, **14**(8), 1247–1253.
- Keller, K. M., & Howlett, S. E. (2016). Sex differences in the biology and pathology of the aging heart. *The Canadian journal of cardiology*, **32**(9), 1065–1073.
- Lakatta, E. G., Gerstenblith, G., Angell, C. S., Shock, N. W., & Weisfeldt, M. L. (1975). Diminished inotropic response of aged myocardium to catecholamines. *Circulation Research*, **36**(2), 262–269.
- Linzbach, A. J., & Akuamo-Boateng, E. (1973). [Changes in the aging human heart. I. Heart weight in the aged]. *Klinische Wochenschrift*, **51**(4), 156–163.
- Liu, J., Sirenko, S., Juhaszova, M., Sollott, S. J., Shukla, S., Yaniv, Y., & Lakatta, E. G. (2014). Age-associated abnormalities of intrinsic automaticity of sinoatrial nodal cells are linked to deficient cAMP-PKA-Ca²⁺ signaling. *American Journal of Physiology-Heart and Circulatory Physiology*, **306**(10), H1385–H1397.
- MacLennan, D. H., & Kranias, E. G. (2003). Phospholamban: A crucial regulator of cardiac contractility. *Nature Reviews Molecular Cell Biology*, **4**(7), 566–577.

- Moghtadaei, M., Jansen, H. J., Mackasey, M., Rafferty, S. A., Bogachev, O., Sapp, J. L., Howlett, S. E., & Rose, R. A. (2016). The impacts of age and frailty on heart rate and sinoatrial node function. *The Journal of Physiology*, **594**(23), 7105–7126.
- Monfredi, O., & Boyett, M. R. (2015). Sick sinus syndrome and atrial fibrillation in older persons - A view from the sinoatrial nodal myocyte. *Journal of Molecular and Cellular Cardiology*, **83**, 88–100.
- Olivetti, G., Melissari, M., Capasso, J. M., & Anversa, P. (1991). Cardiomyopathy of the aging human heart. Myocyte loss and reactive cellular hypertrophy. *Circulation Research*, **68**(6), 1560–1568.
- ONS. (2025). *National population projections: 2022-based*. Office for National Statistics UK.
- Peters, C. H., Sharpe, E. J., & Proenza, C. (2020). Cardiac pacemaker activity and aging. *Annual Review of Physiology*, **82**(1), 21–43.
- Segal, S., Shemla, O., Shapira, R., Peretz, N. K., Lukyanenko, Y., Brosh, I., Behar, J., Lakatta, E. G., Tsutsui, K., & Yaniv, Y. (2023). cAMP signaling affects age-associated deterioration of pacemaker beating interval dynamics. *Geroscience*, **45**(4), 2589–2600.
- Tsutsui, K., Monfredi, O. J., Sirenko-Tagirova, S. G., Maltseva, L. A., Bychkov, R., Kim, M. S., Ziman, B. D., Tarasov, K. V., Tarasova, Y. S., Zhang, J., Wang, M., Maltsev, A. V., Brennan, J. A., Efimov, I. R., Stern, M. D., Maltsev, V. A., & Lakatta, E. G. (2018). A coupled-clock system drives the automaticity of human sinoatrial nodal pacemaker cells. *Science signaling*, **11**(534), eaap7608.
- Turner, M. J., Mier, C. M., Spina, R. J., Schechtman, K. B., & Ehsani, A. A. (1999). Effects of age and gender on the cardiovascular responses to isoproterenol. *The Journals of Gerontology Series A, Biological Sciences and Medical Sciences*, **54**(9), B393–B400.
- Xiao, R. P., Tomhave, E. D., Wang, D. J., Ji, X., Boluyt, M. O., Cheng, H., Lakatta, E. G., & Koch, W. J. (1998). Age-associated reductions in cardiac beta1- and beta2-adrenergic responses without changes in inhibitory G proteins or receptor kinases. *The Journal of Clinical Investigation*, **101**(6), 1273–1282.
- Yaniv, Y., Ahmet, I., Tsutsui, K., Behar, J., Moen, J. M., Okamoto, Y., Guiriba, T. R., Liu, J., Bychkov, R., & Lakatta, E. G. (2016). Deterioration of autonomic neuronal receptor signaling and mechanisms intrinsic to heart pacemaker cells contribute to age-associated alterations in heart rate variability in vivo. *Aging Cell*, **15**(4), 716–724.
- Yue, X., Hazan, A., Lotteau, S., Zhang, R., Torrente, A. G., Philipson, K. D., Ottolia, M., & Goldhaber, J. I. (2020). Na/Ca exchange in the atrium: Role in sinoatrial node pacemaking and excitation-contraction coupling. *Cell Calcium*, **87**, 102167.

Additional information

Data availability statement

Raw data for activity responses shown in Figs 4 and 8 are available as online Supporting Information.

Competing interests

The authors declare that they have no competing interests.

Author contributions

F.S.G., S.A.J. and M.K.L. were responsible for the data collection process and analysis. S.A.J. and M.K.L. were responsible for manuscript drafting, editing and further data analysis.

Funding

University of Hull (HU): FSG: Studentship University of Hull funded a studentship for FSG, previously known as Miss Fiona Hatch.

Acknowledgements

We thank Dr Jenny Waby for her expertise and assistance to FSG with the desmin mRNA analysis. FSG (née Hatch) thanks the University of Hull for her PhD studentship.

Keywords

ageing, arrhythmia, calcium, sarcoplasmic reticulum, sinoatrial node

Supporting information

Additional supporting information can be found online in the Supporting Information section at the end of the HTML view of the article. Supporting information files available:

Peer Review History

Raw data for Figures 4 and 8