



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/242620/>

Version: Published Version

Article:

Mashiah, L., Medini, O., Araiza Calahorra, A. et al. (2025) Tailored Co-Extruded Cereals for Seniors: a Design Thinking Approach to Functional Food Development. Chemical Engineering Transactions, 118. pp. 127-132. ISSN: 2283-9216

<https://doi.org/10.3303/CET25118022>

© 2025, AIDIC Servizi S.r.l. Reproduced in accordance with the publisher's self-archiving policy.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Tailored Co-Extruded Cereals for Seniors: a Design Thinking Approach to Functional Food Development

Leehen Mashiah^a, Omer Medini^b, Andrea Araiza Calahorra^c, Carmit Shani Levi^a, Anwesha Sarkar^c, Uri Lesmes^{a,*}

^aLaboratory of Chemistry of Foods and Bioactives, Department of Biotechnology and Food Engineering, Technion – Israel Institute of Technology, Haifa, Israel

^bThe International Beer Breweries LTD, Kibbutz, Givat Haim, M.P. Hefer, 3893000, Israel

^cFood Colloids and Bioprocessing Group, School of Food Science and Nutrition, University of Leeds, Leeds, LS2 9JT, UK
lesmesu@bfe.technion.ac.il

The increasing gap between lifespan and health span calls for innovative approaches to promote healthy aging. This study, part of the EAT4AGE project under the Joint Programming Initiative 'A Healthy Diet for a Healthy Life', focuses on developing functional foods for older adults. A co-extruded cereal prototype fortified with Maca root powder and Olive leaf extract was designed to address nutritional needs of older adults, particularly high-quality, easily digestible proteins. Analytical results show the cereals have a rich macronutrient profile with over 12% (w/w) protein, 20% (w/w) fat, and low sugar content (<5% w/w), surpassing commercial alternatives. Texture analyses indicate improved hardness and reduce oral friction. An untrained consumer panel (n=21, Age 73±5) reported high palatability and overall acceptability. The product's digestion was evaluated using an age-tailored *in vitro* digestion model, consistently demonstrating high protein digestibility exceeding 80% across all formulations. The calculated *in vitro* digestible indispensable amino acid score affirms the product's high nutritional quality. This research highlights the potential for designing palatable foods that could contribute to a balanced and sustainable diet for healthy aging.

1. Introduction

The National Institutes of Health (NIH), U.S. observed a significant increase in human lifespan that has not been matched by a corresponding increase in human healthspan (Brett et al., 2019; He et al., n.d.). This trend is accompanied by growing efforts to maximize the potential of plant-based foods for healthier and more sustainable diets (Green et al., 2022; Willett et al., 2019). Consequently, the development of customized nutritional solutions can substantially impact on the independence, quality of life, and healthy aging of older adults by addressing many age-related challenges.

These solutions need to consider relevant age-related physiological declines, such as changes in salivation, reduced jaw strength, swallowing difficulties, slower stomach emptying, decreased secretion of digestive fluids, and altered gut microbiota, along with reduced appetite and early satiety (Biagi et al., 2016; Claesson et al., 2012; Menard et al., 2023; Rémond et al., 2015; Sarkar, 2019). Furthermore, designing foods for older adults requires addressing age-related cognitive decline and changes in eating patterns, where potential sensory deterioration may contribute to malnutrition and insufficient protein intake, negatively affecting health (Dent et al., 2023; Hu, 2024a; Morgan et al., 2023; Rémond et al., 2015). Currently, there is increasing consumer and commercial interest in food and dietary solutions that can effectively meet the unique preferences, acceptance, and needs of seniors (Hu, 2024a; Rémond et al., 2015).

This led to an ERA-Net ERA-HDHL call for the "Development of targeted nutrition for prevention of undernutrition for older adults" (PREVNUT). The EAT4AGE consortium responded to this challenge by targeting the development of palatable nutrient-dense foods. A literature review within this project identified actual nutritional gaps in older adults, with high-quality protein, dietary fibers, and specific micronutrients like calcium and iron set as viable targets (Bruins et al., 2019; Hu, 2024b; Morgan et al., 2023; Phillips, 2021; Putra et al., 2021).

Protein was found to be particularly important as a robust dietary intervention tool to address age-related muscle loss (sarcopenia), promote healthy muscle aging, and even help regulate appetite (Gosby et al., 2011, 2014; Hu, 2024b; Nunes et al., 2022).

This research aimed to design and study the potential digestive fate of a co-extruded cereal product for seniors (age >65). The product was formulated as a functional food with two added bioactive components: Maca root powder and Olive Leaf Extract (OLE) with 20% oleuropein. Three different formulations were developed and evaluated: a control version without any functional ingredients in either the shell or the filling (Ref+ref), a version with 0.03% w/w Maca root powder added to the shell (Maca+ref), and a version with 0.03% w/w Maca and 0.015% w/w OLE incorporated into both the shell and the filling (Maca+OLE). The study focused on the palatability of the functional product, its oral processing, and the potential digestibility of its proteins in healthy seniors. The underlying hypothesis was that seniors have different food preferences and digestive capabilities which affect their perception and breakdown of the tailored cereal product.

2. Materials and Methods

A structured design thinking approach guided all stages of this study, as described by (Mashiah et al., 2025). The process began with empathizing, identifying the nutritional and sensory needs of older adults through literature review. During the define and ideate phases, nutritional gaps, such as protein quality, fiber, and micronutrient deficiencies, were mapped, and functional ingredients (Maca root powder and Olive Leaf Extract) were selected for their relevance to healthy aging. Maca root powder was chosen for its reported antioxidant, hormone-balancing, and immunomodulatory properties beneficial to seniors (Chen et al., 2021; Gan et al., 2017; Lee et al., 2016; Meissner et al., 2006), while Olive Leaf Extract was selected for its demonstrated antidiabetic, antioxidant, and mitochondrial-supporting effects, all of which are supported by clinical and preclinical studies (Martín-Peláez et al., 2017; Stamatopoulos et al., 2014; Sun et al., 2017). Prototyping involved formulating three plant-based co-extruded cereal products using gluten-free flours and a sesame-based filling: a reference product with no functional ingredients, a product with 0.03% w/w Maca in the shell, and a product with 0.03% w/w Maca and 0.015% w/w OLE in both shell and filling. All ingredients were food-grade and used as received. Products were produced via twin-screw co-extrusion (36 mm, 220 rpm, 105°C die temperature), air-dried, and stored at room temperature.

Nutritional composition (protein, fat, minerals) was determined using standard methods (Kjeldahl, CHNS, AOAC). Texture was evaluated by single compression tests with a texture analyzer, and microstructure was assessed by confocal laser scanning microscopy. Tribological properties (bolus lubrication) were measured with a Mini-Traction Machine at 37°C. *In vitro* oral processing and protein digestibility were assessed using models simulating older adults and following the INFOGEST protocol. Sensory analysis included acceptability and comfortability testing with senior volunteers (n=21, Age 73±5 years).

3. Results and Discussion

3.1 Characterization of Cereal Prototypes

The cereal prototypes contained protein content exceeding 12% (w/w), fat content of around 20% (w/w), low sugar content (less than 5%w/w) and dietary fiber content of 17% (w/w), offering a healthier profile than typical commercial cereals consumed by seniors. To address the limited amino acid profile of cereals, chickpea flour was intentionally included in the formulation, providing complementary essential amino acids which is particularly important for older adults at risk of protein malnutrition. However, the success of these healthier food options depends on balancing their nutritional benefits with consumer acceptance based on taste and palatability. The cereal prototypes' texture, measured via hardness and Young's modulus, showed that adding Maca and OLE significantly reduced hardness ($p < 0.05$), potentially improving chewability for older adults (Figure 1. A+B). Young's modulus remained largely unchanged, indicating the overall structural integrity was maintained. Tribological testing showed that the Maca+OLE cereal formulation transitioned through boundary, mixed, and hydrodynamic lubrication regimes, likely due to oil release during mastication. Confocal microscopy revealed smaller, evenly distributed oil droplets and protein fragments, suggesting improved breakdown and potentially enhanced oral processing (Figure 1.C).




Figure 1. Texture Profile Analysis (TPA) for plant-based prototypes- Enhancing oral comfort through functional materials Addition. (A) Strength to strain TPA graph (B) Summary table of textural properties values. Values (mean $\pm$ SD) in the same line followed by a different letter are significantly different ($p < 0.05$). (C) Mean friction coefficients as a function of entrainment speeds and CLSM images of Ref+ref, Maca+ref, Maca+OLE. Protein aggregates are stained fluorescently green using Fast green whilst fat droplets are stained fluorescently red using Nile Red. The friction curve of water and model saliva are added as reference. Data represent average of triplicate on duplicate samples ($n = 2 \times 3$).

3.2 In Vivo Sensory Evaluation


Questionnaires with an elderly panel ($n=21$, Age 73 ± 5 years) assessed food comfort, including ease of chewing, bolus formation, and oral pain. While the untrained and limited-size panel does not allow for definitive sensory conclusions, it provides valuable initial insights from the target population. Importantly, no significant differences were observed between the control (Ref+ref) and the functional products (Maca+ref, Maca+OLE) across all comfortability metrics (Figure 2), indicating that added ingredients did not reduce sensory comfort. Moreover, participants reported high scores for ease of chewing and swallowing, minimal oral pain, and positive textural perceptions described as "crunchy" and "melting" (Figure 2). These findings indicate a favorable overall sensory experience for the target consumer group.

Part 1. General question-


	Ref+ref	Maca+ref	Maca+OLE	P-value
The food is (1-Very uncomfortable 5-Very comfortable)	4.857	4.857	4.800	0.85045*

*Kruskal-Wallis Test. The result is not significant at $p < .05$


Part 2. Bolus formation



Part 3. Mouth pain



Part 4. Texture



Part 5. Flavor




Figure 2. Results from the questionnaire on the perception of food comfortability.

Part 1. General Food Comfort: Rates the overall comfort level of eating the food (scale: "Very Uncomfortable" to "Very Comfortable"). **Part 2. Bolus Formation and Time Needed:** Evaluates the ease of bolus formation (scale: "Impossible" to "Very Easy") and the time required for it (scale: "Impossible" to "Very Brief"). **Part 3. Pain Perception:** Measures the level of pain experienced while eating (scale: "Extremely" to "Not at All"). **Part 4. Texture Perception:** Assesses the perception of food texture (scale: "Extremely" to "Not at All"). **Part 5. Flavor Perception:** Evaluates the intensity of flavor perception (scale: "Extremely" to "Not at All").

3.3 In Vitro Protein Digestion:

Protein digestion was evaluated using a semi-dynamic *in vitro* model simulating young and older adults (Menard et al., 2023). SDS-PAGE showed similar protein breakdown across all formulations (Ref+ref, Maca+ref, Maca+OLE) (Figure 3.2), indicating that the addition of Maca and OLE did not negatively impact protein digestibility, despite concerns that these bioactives might bind proteins and reduce their bioaccessibility. Extruded products (Figure 3.1.B+D) showed higher protein bioaccessibility than pre-extruded (Figure 3.1.A+C) ones, and digestion was slower under elderly conditions. DIAAS analysis identified lysine and tryptophan as limiting amino acids, but overall digestibility exceeded 80%, highlighting the potential of the combined flour formulation to provide a palatable, nutritionally rich, plant-based cereal suitable for older adults. (Orlien et al., 2021; Sá et al., 2020; Sousa et al., 2023).




Figure 3. (1) Comparative SDS-PAGE analyses of digesta samples: Enhanced proteolysis in post-extruded products (Maca+OLE) and reduced proteolysis in senior *in vitro* digestion vs. adult. (A+C) Protein breakdown patterns of the pre-extruded (Mix+maca) (B+D) and post-extruded product (Maca+OLE) under senior (A+B) and adult (C+D) *in vitro* digestion conditions. (2) Consistent protein breakdown pattern: SDS-PAGE analyses of senior digestion for post-extruded products. (A) Ref+ref, (B) Maca+ref, (C) Maca+OLE. (3) High *In vitro* Digestibility highlighted in results. (A) Comparative protein digestibility percentages in different cereal products pre- and post-extrusion under adult and senior IVD models, (B) Table presenting digestibility percentages of individual AAs. Values are expressed as the mean $\pm$ SD. Different letters denote significant differences ($p < 0.05$).

4. Conclusion

This research highlights the potential to fabricate age-tailored functional foods to promote healthy aging. Plant-based co-extruded cereals enriched with Maca root powder and Olive Leaf Extract (OLE), demonstrated high *in vitro* digestibility (>80%) alongside enhanced antioxidant and anti-inflammatory potential. These findings suggest that integrating functional ingredients into familiar food matrices can provide nutritional benefits beyond basic dietary needs. Importantly, the study emphasizes the dual challenge of optimizing both nutritional value and consumer acceptance, underscoring the need for academia-industry collaborations. Ultimately, this study seeks to stimulate innovation in nutritionally optimized products to meet both nutritional requirements and consumer preferences to improve health and quality of life for older adults.

Acknowledgments

We acknowledge the support of the Israeli Ministry of Science and Technology Grant #3-16518 and the EAT4AGE project under umbrella of the European Joint Programming Initiative “A Healthy Diet for a Healthy Life” (JPI HDHL) of the ERA-HDHL (GA N° 696295)”; A.Sarkar and A. Araiza Calahorra also acknowledge the funding from the Biotechnology and Biological Sciences Research Council (BBSRC), UK (BB/V018914/1).

References

- Biagi, E., Franceschi, C., Rampelli, S., Capri, M., Brigidi, P., & Candela Correspondence, M. (2016). Gut Microbiota and Extreme Longevity. *Current Biology*, 26, 1480–1485. <https://doi.org/10.1016/j.cub.2016.04.016>
- Brett, C. E., Dykiert, D., Starr, J. M., & Deary, I. J. (2019). Predicting change in quality of life from age 79 to 90 in the Lothian Birth Cohort 1921. *Quality of Life Research*, 28(3), 737–749. <https://doi.org/10.1007/S11136-018-2056-4/TABLES/4>
- Bruins, M. J., Van Dael, P., & Eggersdorfer, M. (2019). The role of nutrients in reducing the risk for noncommunicable diseases during aging. In *Nutrients* (Vol. 11, Issue 1). MDPI AG. <https://doi.org/10.3390/nu11010085>
- Chen, R., Wei, J., & Gao, Y. (2021). A review of the study of active components and their pharmacology value in *Lepidium meyenii* (Maca). *Phytotherapy Research*, 35(12), 6706–6719. <https://doi.org/10.1002/PTR.7257>
- Claesson, M. J., Jeffery, I. B., Conde, S., Power, S. E., O’connor, E. M., Cusack, S., Harris, H. M. B., Coakley, M., Lakshminarayanan, B., O’sullivan, O., Fitzgerald, G. F., Deane, J., O’connor, M., Harnedy, N., O’connor, K., O’mahony, D., Van Sinderen, D., Wallace, M., Brennan, L., ... O’toole, P. W. (2012). Gut microbiota composition correlates with diet and health in the elderly. *Nature* 2012 488:7410, 488(7410), 178–184. <https://doi.org/10.1038/nature11319>
- Dent, E., Wright, O. R. L., Woo, J., & Hoogendijk, E. O. (2023). Malnutrition in older adults. *The Lancet*, 401(10380), 951–966. [https://doi.org/10.1016/S0140-6736\(22\)02612-5](https://doi.org/10.1016/S0140-6736(22)02612-5)
- Gan, J., Feng, Y., He, Z., Li, X., & Zhang, H. (2017). Correlations between Antioxidant Activity and Alkaloids and Phenols of Maca (*Lepidium meyenii*). *Journal of Food Quality*, 2017. <https://doi.org/10.1155/2017/3185945>
- Gosby, A. K., Conigrave, A. D., Lau, N. S., Iglesias, M. A., Hall, R. M., Jebb, S. A., Brand-Miller, J., Caterson, I. D., Raubenheimer, D., & Simpson, S. J. (2011). Testing protein leverage in lean humans: A randomised controlled experimental study. *PLoS ONE*, 6(10). <https://doi.org/10.1371/journal.pone.0025929>
- Gosby, A. K., Conigrave, A. D., Raubenheimer, D., & Simpson, S. J. (2014). Protein leverage and energy intake. *Obesity Reviews*, 15(3), 183–191. <https://doi.org/10.1111/OBR.12131>
- Green, A., Blattmann, C., Chen, C., & Mathys, A. (2022). The role of alternative proteins and future foods in sustainable and contextually-adapted flexitarian diets. *Trends in Food Science & Technology*, 124, 250–258. <https://doi.org/10.1016/J.TIFS.2022.03.026>
- He, W., Goodkind, D., & Kowal, P. (n.d.). *An Aging World*: 2015.
- Hu, F. B. (2024a). Diet strategies for promoting healthy aging and longevity: An epidemiological perspective. *Journal of Internal Medicine*, 295(4), 508–531. <https://doi.org/10.1111/joim.13728>
- Hu, F. B. (2024b). Diet strategies for promoting healthy aging and longevity: An epidemiological perspective. *Journal of Internal Medicine*, 295(4), 508–531. <https://doi.org/10.1111/JOIM.13728>
- Lee, M. S., Lee, H. W., You, S., & Ha, K. T. (2016). The use of maca (*Lepidium meyenii*) to improve semen quality: A systematic review. In *Maturitas* (Vol. 92, pp. 64–69). Elsevier Ireland Ltd. <https://doi.org/10.1016/j.maturitas.2016.07.013>
- Martín-Peláez, S., Castañer, O., Konstantinidou, V., Subirana, I., Muñoz-Aguayo, D., Blanchart, G., Gaixas, S., de la Torre, R., Farré, M., Sáez, G. T., Nyssönen, K., Zunft, H. J., Covas, M. I., & Fitó, M. (2017). Effect of olive oil phenolic compounds on the expression of blood pressure-related genes in healthy individuals. *European Journal of Nutrition*, 56(2), 663–670. <https://doi.org/10.1007/s00394-015-1110-z>
- Mashiah, L., Lavoisier, A., Gwala, S., Calahorra, A. A., Shani Levi, C., Rødbotten, R., Varela, P., Sarkar, A., Brodkorb, A., Dupont, D., & Lesmes, U. (2025). Strategic Considerations in Designing Food Solutions for Seniors. *Foods* 2025, Vol. 14, Page 396, 14(3), 396. <https://doi.org/10.3390/FOODS14030396>
- Meissner, H. O., Mscisz, A., Reich-Bilinska, H., Mrozikiewicz, P., Bobkiewicz-Kozłowska, T., Kedzia, B., Lowicka, A., & Barchia, I. (2006). Hormone-Balancing Effect of Pre-Gelatinized Organic Maca (*Lepidium peruvianum* Chacon): (III) Clinical responses of early-postmenopausal women to Maca in double blind, randomized, Placebo-controlled, crossover configuration, outpatient study. *International Journal of Biomedical Science: IJBS*, 2(4), 375.

- Menard, O., Lesmes, U., Shani-Levi, C. S., Araiza Calahorra, A., Lavoisier, A., Morzel, M., Rieder, A., Feron, G., Nebbia, S., Mashiah, L., Andres, A., Bornhorst, G., Carrière, F., Egger, L., Gwala, S., Heredia, A., Kirkhus, B., Macierzanka, A., Portman, R., ... Dupont, D. (2023). Static in vitro digestion model adapted to the general older adult population: an INFOGEST international consensus. *Food & Function*. <https://doi.org/10.1039/D3FO00535F>
- Morgan, P. T., Witard, O. C., Højfeldt, G., Church, D. D., Breen, L., & Morgan, P. (2023). Dietary protein recommendations to support healthy muscle ageing in the 21st century and beyond: considerations and future directions. *Proceedings of the Nutrition Society*, 1–14. <https://doi.org/10.1017/S0029665123003750>
- Nunes, E. A., Colenso-Semple, L., McKellar, S. R., Yau, T., Ali, M. U., Fitzpatrick-Lewis, D., Sherifali, D., Gaudichon, C., Tomé, D., Atherton, P. J., Robles, M. C., Naranjo-Modad, S., Braun, M., Landi, F., & Phillips, S. M. (2022). Systematic review and meta-analysis of protein intake to support muscle mass and function in healthy adults. In *Journal of Cachexia, Sarcopenia and Muscle* (Vol. 13, Issue 2, pp. 795–810). John Wiley and Sons Inc. <https://doi.org/10.1002/jcsm.12922>
- Orlien, V., Aalaei, K., Poojary, M. M., Nielsen, D. S., Ahrné, L., & Carrascal, J. R. (2021). Effect of processing on in vitro digestibility (IVPD) of food proteins. *Critical Reviews in Food Science and Nutrition*, 0(0), 1–50. <https://doi.org/10.1080/10408398.2021.1980763>
- Phillips, J. A. (2021). Dietary Guidelines for Americans, 2020–2025. <https://doi.org/10.1177/21650799211026980>, 69(8), 395. <https://doi.org/10.1177/21650799211026980>
- Putra, C., Konow, N., Gage, M., York, C. G., & Mangano, K. M. (2021). Protein Source and Muscle Health in Older Adults: A Literature Review. *Nutrients* 2021, Vol. 13, Page 743, 13(3), 743. <https://doi.org/10.3390/NU13030743>
- Rémond, D., Shahar, D. R., Gille, D., Pinto, P., Kachal, J., Peyron, M. A., Dos Santos, C. N., Walther, B., Bordoni, A., Dupont, D., Tomás-Cobos, L., & Vergères, G. (2015). Understanding the gastrointestinal tract of the elderly to develop dietary solutions that prevent malnutrition. *Oncotarget*, 6(16), 13858. <https://doi.org/10.18632/ONCOTARGET.4030>
- Sá, A. G. A., Moreno, Y. M. F., & Carciofi, B. A. M. (2020). Food processing for the improvement of plant proteins digestibility. *Critical Reviews in Food Science and Nutrition*. <https://doi.org/10.1080/10408398.2019.1688249>
- Sarkar, A. (2019). Oral processing in elderly: understanding eating capability to drive future food texture modifications. *Proceedings of the Nutrition Society*, 78(3), 329–339. <https://doi.org/10.1017/S0029665118002768>
- Sousa, R., Recio, I., Heimo, D., Dubois, S., Moughan, P. J., Hodgkinson, S. M., Portmann, R., & Egger, L. (2023). In vitro digestibility of dietary proteins and in vitro DIAAS analytical workflow based on the INFOGEST static protocol and its validation with in vivo data. *Food Chemistry*, 404, 134720. <https://doi.org/10.1016/J.FOODCHEM.2022.134720>
- Stamatopoulos, K., Katsoyannos, E., & Chatzilazarou, A. (2014). Antioxidant Activity and Thermal Stability of Oleuropein and Related Phenolic Compounds of Olive Leaf Extract after Separation and Concentration by Salting-Out-Assisted Cloud Point Extraction. *Antioxidants* 2014, Vol. 3, Pages 229–244, 3(2), 229–244. <https://doi.org/10.3390/ANTIOX3020229>
- Sun, W., Wang, X., Hou, C., Yang, L., Li, H., Guo, J., Huo, C., Wang, M., Miao, Y., Liu, J., & Kang, Y. (2017). Oleuropein improves mitochondrial function to attenuate oxidative stress by activating the Nrf2 pathway in the hypothalamic paraventricular nucleus of spontaneously hypertensive rats. *Neuropharmacology*, 113, 556–566. <https://doi.org/10.1016/J.NEUROPHARM.2016.11.010>
- Willett, W., Rockström, J., Loken, B., Springmann, M., Lang, T., Vermeulen, S., Garnett, T., Tilman, D., DeClerck, F., Wood, A., Jonell, M., Clark, M., Gordon, L. J., Fanzo, J., Hawkes, C., Zurayk, R., Rivera, J. A., De Vries, W., Majele Sibanda, L., ... Murray, C. J. L. (2019). Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. In *The Lancet* (Vol. 393, Issue 10170, pp. 447–492). Lancet Publishing Group. [https://doi.org/10.1016/S0140-6736\(18\)31788-4](https://doi.org/10.1016/S0140-6736(18)31788-4)