



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/244191/>

Version: Published Version

Article:

Hadji, P., Athanasiadis, A., Brandi, M. et al. (2026) Pregnancy- and lactation-associated osteoporosis: a position statement of the IAPM, IOF, ECTS, ESCEO, IMS, and EMAS. International Journal of Gynecology & Obstetrics. ISSN: 0020-7292

<https://doi.org/10.1002/ijgo.71203>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:

<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

NARRATIVE REVIEW

Pregnancy- and lactation-associated osteoporosis: A position statement of the IAPM, IOF, ECTS, ESCEO, IMS, and EMAS

Peyman Hadji^{1,2}  | Apostolos Athanasiadis³ | Maria-Luisa Brandl^{4,5} |
 Marlene Chakhtoura⁶ | Manju Chandran^{7,8} | Frank Chervenak⁹ | Amos Grünebaum¹⁰ |
 Irene Lambrinoudaki¹¹ | Willem F. Lems¹² | Apostolos M. Mamopoulos¹³ |
 Eugene McCloskey¹⁴ | Rossella E. Nappi¹⁵ | Jessica Pepe¹⁶ | Jean-Yves Reginster¹⁷ |
 René Rizzoli¹⁸ | Petra Stute¹⁹ | Şansın Tuzun²⁰ | Amanda J. Vincent²¹ |
 Maria Belen Zanchetta²² | M. Carola Zillikens²³ | Ioannis Kyvernitis²⁴

Correspondence

Peyman Hadji, Head of Department,
 Frankfurt Centre for Bone Health and
 Endocrinology, Goethestraße 23, 60313,
 Frankfurt, Germany and Philipps-
 University of Marburg, Marburg, Germany.
 Email: p.hadji@outlook.de

Abstract

Pregnancy- and lactation-associated osteoporosis (PLO) is a syndrome characterized by fragility fractures (most commonly vertebral, often multiple) occurring in late pregnancy or the early postpartum period. This position statement summarizes the current knowledge of PLO, and the recommended procedure for assessment, diagnosis and treatment based on a review of published evidence. As PLO is a rare condition, controlled, comparative clinical studies are limited. Individual studies were reviewed, focusing on design, size, follow-up, evaluation of safety, and factors that impact PLO outcomes. Diagnosis of PLO should include a detailed clinical examination, a review of medical and treatment history, and laboratory tests to rule out secondary osteoporosis. Imaging techniques can be used to inform the appropriate treatment pathway, which will depend on whether the patient is antepartum or postpartum. Management of PLO and fractures includes calcium/vitamin D intake/supplementation, considering stopping breastfeeding, and analgesia, if necessary. Treatment with bone-specific agents should be evaluated on an individual basis to reduce subsequent fracture risk, noting that their use in PLO patients is off label. If pharmacological treatments are used, they must be given alongside effective contraceptive measures. Bisphosphonates can pass the placental barrier and are detectable in bone for years, so adverse effects on future pregnancies, although not yet reported, cannot be excluded. Preferred treatment options are teriparatide/abaloparatide or romosozumab. Second-choice treatments are denosumab followed by bisphosphonates or bisphosphonates alone. This position statement includes a pragmatic

Abbreviations: ECTS, European Calcified Tissue Society; EMAS, European Menopause and Andropause Society; ESCEO, European Society for Clinical and Economics Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases; IAPM, International Academy of Perinatal Medicine; IMS, International Menopause Society; IOF, International Osteoporosis Foundation.

For affiliations refer to page 9.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2026 The Author(s). *International Journal of Gynecology & Obstetrics* published by John Wiley & Sons Ltd on behalf of International Federation of Gynecology and Obstetrics.

approach for identifying women with PLO and suggests developing an individualized treatment plan that might include pharmaceutical intervention.

KEYWORDS

bone mineral density, fractures, lactation, pregnancy, pregnancy- and lactation-associated osteoporosis

1 | INTRODUCTION

Osteoporosis is a systemic skeletal disease characterized by low bone mineral density (BMD) microarchitectural deterioration of bone tissue with a consequent increase in bone fragility and susceptibility to fractures.¹ The World Health Organization (WHO) defines the BMD threshold for osteoporosis as ≥ 2.5 standard deviations below the young adult mean peak bone mass when measured at the femoral neck and spine using dual-energy X-ray absorptiometry (DXA).²

Around 18.3% of the world's population is estimated to have osteoporosis with women having a higher prevalence (23.1%) than men (11.7%).^{3,4} A systematic review and meta-analysis of population-based studies that used the WHO criteria for osteoporosis found an overall global prevalence of 19.7%.⁵ The prevalence of osteoporosis is known to increase with age and is particularly high in postmenopausal women, with some reports estimating that around 50% of women over 50 years of age will suffer from osteoporotic fractures in their lifetime, twice the incidence seen in men of the same age.⁶ The primary causes of morbidity, mortality, and healthcare costs due to osteoporosis are osteoporotic fractures, so the prompt identification and appropriate management of individuals at increased risk for fractures is critical.

In addition to low BMD, several other independent risk factors for osteoporosis have been identified.^{1,5,7} These include age, gender, premature menopause, a family history of osteoporosis or hip fracture in a parent, low body weight, lifestyle factors (e.g., low intake of calcium and vitamin D, inactivity, smoking, and excessive alcohol intake), and certain medical conditions (e.g., diabetes mellitus, rheumatoid arthritis, inflammatory bowel disease, breast cancer, and chronic liver and kidney disease). In clinical practice, fracture risk assessment tools, including FRAX (developed by the Metabolic Bone Disease Unit at the University of Sheffield in 2008), which combine BMD measurements and assessment of a range of possible clinical risk factors, are commonly used to determine 10-year fracture risk in individuals over the age of 40 years.^{8,9}

From a clinical management perspective, it is important to differentiate between primary and secondary osteoporosis, a distinction that is dependent on their underlying causes. Primary osteoporosis is the most common form and includes idiopathic osteoporosis occurring in both women and men, usually as a result of natural aging or due to hormonal changes, for example low estrogen levels in postmenopausal women.¹ In primary osteoporosis, no other disease or external factor can be identified as the cause of bone loss. Secondary osteoporosis is less common, but well documented, and arises as a result of other

medical conditions or due to treatment with medications that cause bone loss or hormonal changes. Examples include endocrine disorders (e.g., diabetes mellitus type I, hyperthyroidism, hyperparathyroidism, and premature ovarian insufficiency), chronic diseases such as rheumatoid arthritis or Parkinson's disease, long-term use of certain drugs (e.g., glucocorticoids, depot medroxyprogesterone acetate [DMPA], chemotherapy, aromatase inhibitors used to treat breast cancer in women, and heparin).^{1,10,11} Other factors include malabsorption of calcium and vitamin D, anorexia nervosa, prolonged immobilization, and possibly advanced maternal age at first pregnancy.¹²

A rare form of osteoporosis, and the focus of this joint position statement, is pregnancy- and lactation-associated osteoporosis (PLO).¹³⁻¹⁶ PLO can be defined as a clinical syndrome characterized by fragility fractures (most commonly vertebral, often multiple) occurring in late pregnancy or the early postpartum period, usually, but not always, accompanied by low BMD for age. As PLO can have a significant negative impact on bone health and substantially reduce long-term quality of life,¹⁷ there is a need to raise awareness and recognition of PLO among all healthcare professionals who care for women through pregnancy and in the postpartum period to ensure prompt referral for specialist evaluation. Clear strategies are needed to assist clinicians with assessment and diagnosis of PLO and to direct appropriate treatment. This joint position statement, developed by several interdisciplinary obstetrics and bone societies, provides a review of the available literature regarding what we currently know about PLO and its management and proposes algorithms to help clinicians determine the most appropriate pathways for diagnosis and treatment of women in their care.

2 | PHYSIOLOGY OF NORMAL BONE TURNOVER

To provide important context to the discussion of PLO and its management, key drivers of the physiology of normal bone turnover in premenopausal women and the influence of estrogen and other hormones on bone health during pregnancy and lactation are briefly reviewed.

2.1 | Bone turnover in premenopausal women

During adolescence in girls, estrogen levels rise significantly and this change affects bone development as well as reproductive health.

The increase in estrogen levels triggers the onset of puberty and the start of menstruation. Throughout adolescence, estrogen levels continue to increase steadily, resulting in the acceleration of bone growth and the closure of epiphyseal growth plates.¹⁸ This period is recognized as critical for achieving peak bone mass in adulthood. Estrogen levels eventually reach a steady state with the establishment of regular menstrual cycles and bone turnover becomes balanced, with consolidation of bone mineral density and architecture until adulthood. Typically, peak bone mass is achieved by the end of the second decade of life, but it varies by skeletal site, sex, and population, and while much of bone accrual occurs by late adolescence, peak or plateau BMD might not be reached until the third decade of life or later.^{19–23}

2.2 | Influence of pregnancy on bone health

During pregnancy, estrogen levels increase significantly, produced primarily by the ovaries and placenta, and peak in the third trimester, to support the growth and development of the fetus and prepare the mother's body for childbirth and lactation.²⁴ To meet the requirement of the developing fetal skeleton during pregnancy and its growth postpartum, maternal metabolism undergoes some significant adaptations, particularly with respect to calcium homeostasis^{25–27} (Figure 1).

During pregnancy, approximately 30g of calcium is transferred from the maternal skeleton to the fetus, and most of this (around 80%) occurs during the third trimester.²⁸ To compensate for such requirement, and to support fetal bone accretion during

pregnancy, a twofold increase in maternal fractional calcium absorption is observed, resulting from a twofold to fivefold increase in 1,25-dihydroxyvitamin D (calcitriol) levels derived from maternal kidneys.^{25,29} Both estrogen and prolactin, which are elevated in the third trimester, promote calcium absorption indirectly by increasing the production of 1,25-dihydroxyvitamin D. This is balanced by an increase in urinary calcium excretion through the kidneys; however, renal conservation of calcium is also observed.^{30,31} Other maternal compensatory mechanisms of calcium homeostasis and bone turnover during pregnancy include elevated estradiol (from the ovaries and placenta), which influences calcium absorption and bone metabolism,^{28,30} PTH-rP, and osteoprotegerin (OPG) levels. OPG inhibits bone resorption by binding to receptor activator of nuclear factor kappa B (RANK) ligand (RANKL) and blocking the signaling pathway that regulates osteoclastogenesis and osteoclast activation and inhibiting excessive bone resorption.^{32,33}

The International Osteoporosis Foundation (IOF) and the International Federation of Clinical Chemistry and Laboratory Medicine recommend serum/plasma procollagen type I N-propeptide (P1NP) and serum β isomerized C-terminal telopeptide of type I collagen (β -CTX-I) as reference bone turnover markers for osteoporosis.³⁴ During pregnancy, particularly in the first two trimesters, the levels of these bone turnover markers are low to normal but increase rapidly in the third trimester.³⁵ Studies using DXA and quantitative ultrasound (QUS) have demonstrated relatively stable BMD levels in pregnant women during the first trimester with some decreases in the second trimester, but the most significant decreases are observed in the third trimester.^{36–41} However, the degree of bone loss varies by location. For example, one study found a significant




FIGURE 1 Pathophysiology of the calcium pathway in normal, pregnant and lactating women.

decrease in BMD at the ultra-distal radius (approximately 2%), spine (approximately 1.8%), and total hip (approximately 3.2%) and a larger decrease (4.2%) at the trochanteric region.⁴² Breasail et al., using peripheral quantitative computed tomography (pQCT) and HR-pQCT scans of tibia and radius, found significantly greater decreases in total BMD, trabecular volumetric (v) BMD and cortical vBMD in pregnant versus non-pregnant women, as well as pregnancy-related decreases in bone microarchitecture.⁴³ Bone microarchitecture deterioration was observed in women with PLO, with the trabecular compartment being the most affected. It showed significantly lower trabecular density, number, and thickness, with subsequent greater trabecular separation and network heterogeneity. Concerning the cortical compartment, its density and thickness were also deteriorated but to a lesser extent. For most women, the changes in BMD are temporary, and it usually returns to its pre-pregnancy level after 1 year of weaning.^{44–46}

2.3 | Influence of lactation on bone health

During lactation, estrogen levels decline rapidly due to rising levels of prolactin, which suppresses hypothalamic gonadotropin-releasing hormone.⁴⁷ This reduces the release of the pituitary gonadotropins, luteinizing hormone and follicle-stimulating hormone, thereby inhibiting ovarian estrogen production. The rising prolactin levels, as well as suckling, stimulate milk production and promote parathyroid hormone-related protein (PTHrP) secretion from mammary tissue. PTHrP acts synergistically with low estrogen to mobilize calcium from the maternal skeleton, contributing to bone loss during breastfeeding.⁴⁷ It is estimated that women lose 300 to 400 mg of calcium daily through breast milk, and this calcium demand is met by a 5–10% loss of skeletal mineral content during 6 months of exclusive lactation.^{27,48} The hypoestrogenic state and the increased demand for calcium are key drivers of this increased bone resorption during lactation⁴⁹ (Figure 1). Calcitriol levels also remain high during the postpartum period, which can help offset skeletal calcium mobilization. Further, insufficient levels of vitamin D, due to factors such as lack of exposure to sunlight or poor dietary intake, can contribute to low calcium absorption and exacerbate bone loss during lactation.^{50,51}

Bone turnover studies in both experimental animals and nursing women have largely also found that levels of bone resorption markers, such as collagen type 1 c-telopeptide (CTX) and n-telopeptide (NTX), are elevated during lactation.⁴⁷ Recent studies in animal models suggest that brain-derived cellular communication network factor 3 (CCN3) secreted from KISS1 neurons of the arcuate nucleus (ARC/KISS1) can act as a potent osteoanabolic factor that can build bone in females during lactation.⁵² While these findings are interesting, translational clinical studies are needed to establish its role in lactating women.

In general, studies suggest that bone loss during lactation is transitory and usually there is a recovery of BMD at all sites studied, depending on the return of menstruation after weaning.⁵³ A meta-analysis of studies evaluating BMD during lactation found that

the duration of lactation was also an important factor in outcomes: at 12 months postpartum, the group of women who breastfed for >6 months had significant losses at the lumbar spine, while those who breastfed for a shorter duration (≤ 4 months) had recovered their BMD at that timepoint.⁵⁴ Overall, however, accumulated evidence suggests that lactation does not have any long-term negative effects on the maternal skeleton^{49,55,56} and does not translate into an increased risk of fragility fractures.^{46,55,57,58}

Despite known compensatory mechanisms that help promote a restoration of bone after weaning, there is a subgroup of women who suffer from a severe form of PLO.^{13,59,60} These women present with more severe bone microarchitecture impairment than control women undergoing a similar profound process of trabecular bone resorption during lactation, which is in line with the hypothesis that bone quality might in fact be deteriorating before gestation in most patients, and consequently they are unable to meet the increased calcium requirements.^{61,62} It is worth noting that, although not always available, HR-pQCT is a useful noninvasive tool for assessing bone microarchitecture, offering new insights into the pathophysiology of PLO.⁶³

3 | DEVELOPMENT OF THIS POSITION STATEMENT ON PREGNANCY- AND LACTATION-ASSOCIATED OSTEOPOROSIS

Searches were undertaken of PubMed and MEDLINE (National Library of Medicine, Bethesda, MD), as well as other databases, to identify clinical trials, systematic reviews, meta-analyses, narrative reviews, observational studies, case-control studies regarding PLO and its management up to 2025 with the following search terms: [pregnancy OR lactation] AND fracture, osteoporosis, BMD, QUS, high-resolution peripheral quantitative computed tomography (HR-pQCT), and bone turnover markers. In addition, the Cochrane Register of Controlled Trials and database of ongoing and unpublished trials were searched (www.clinicaltrials.gov). The retrieved literature, including primary studies, published reviews and case reports, was examined and interpreted in the context of existing consensus documents and clinical experience but were not assessed or graded for level of evidence. These findings were used to propose the most appropriate diagnostic methodology, determine the choice of therapy, and define follow-up/monitoring procedures.

Over the past decade, several reviews and meta-analyses^{13–15,26,27,61,64–68} as well as retrospective cohort analyses^{59,69,70} have provided helpful summaries of the available clinical evidence about PLO and best practice insights into its management. In developing this position statement, we have reviewed these previous publications alongside evaluation of recent data, individual clinical studies, and case reports to develop suggested algorithms for assessment, diagnosis, and treatment of PLO that can be used in routine clinical practice. PLO is a condition that requires an individualized approach to care and multidisciplinary team input, with the wellbeing of both the mother and unborn child taken into consideration.

4 | INCIDENCE, RISK FACTORS AND PATHOPHYSIOLOGY OF PREGNANCY- AND LACTATION-ASSOCIATED OSTEOPOROSIS

4.1 | Incidence

Pregnancy- and lactation-associated osteoporosis is a rare condition and, coupled with the fact that there is a lack of robust epidemiological studies in this setting, definitive figures for incidence and prevalence are difficult to establish.⁵⁹ A retrospective single-center study that evaluated medical records over a 20-year period estimated a PLO prevalence of 4.1 in every 100 000 women of reproductive age.⁵⁹ In addition, nationwide data from Japan suggest a rate of 4.5/10 000, but this included fractures occurring within 2 years of obstetric hospitalization.⁷¹ A recent systematic literature review and meta-analysis aimed to estimate the prevalence of PLO in postpartum women within 1 year of delivery.⁷² Eight observational studies were identified and seven of these used DXA to assess BMD. However, they used different parameters to assess osteoporosis (*T*-score, *Z*-score, or a combination of both) and different timeframes for evaluation, highlighting the challenge of comparing data across studies. With these limitations in mind, the estimated prevalence of PLO defined by DXA was 5% at the lumbar spine and 12% at the femoral neck.⁷²

4.2 | Risk factors

Potential risk factors for PLO often overlap with those described earlier for idiopathic osteoporosis.¹⁵ A large observational case-control study that directly compared 102 women with PLO and 102 healthy controls found that those with PLO had significantly more dental problems in childhood, were less physically active both before and after puberty, and were less mobile during pregnancy.⁶⁰ Other potential risk factors that have been reported include a family history of osteoporosis and osteoporotic fracture, the presence of conditions such as type 1 diabetes mellitus or rheumatoid arthritis, and a history of anorexia nervosa.^{15,17,27,69,73} There is also an epidemiological relationship with ovulation disorders, a history of infertility treatment, and advanced maternal age.^{12,72}

Medications that might contribute to osteoporosis risk in premenopausal women include thromboprophylactic agents such as low-molecular-weight heparin (which is widely used in pregnancy), corticosteroids, tamoxifen, aromatase inhibitors, anticonvulsants, DMPA, and luteinizing hormone-releasing hormone agonists.^{59,64,72}

Other possible underlying causes for PLO include previously undiagnosed metabolic bone disease or monogenic forms of osteoporosis, such as osteogenesis imperfecta, and the presence of genetic mutations such as *COL1A1*, *COL1A2*, *RPDV*, *LRP5*, *Wnt1* and *LPR5/WINT1*.^{15,74-80} If available, genetic testing should ideally be undertaken in women with suspected PLO, to determine if they have any previously unrecognized skeletal rare disorders.⁷⁸ In addition, as bones and teeth share some characteristics of development and

mineralization, it has been suggested that the presence of dental disorders might be a potential marker for congenital rare bone conditions, with phenotypes often present in PLO.⁸¹

4.3 | Clinical presentation

Women subsequently diagnosed with PLO generally present with severe pain, most commonly back pain, often combined with height loss.⁸² These symptoms occur predominantly in the last trimester of pregnancy and during lactation, mostly during the first pregnancy, rarely in the second or third. Many patients report multiple consultations before finally being correctly diagnosed with PLO, which might indicate a high rate of underdiagnosis and underscores the need to raise awareness of the condition and provide guidance for assessment and management. Vertebral fractures are the most common type of fracture observed in PLO and are generally the underlying cause of the low back pain and height loss.⁸² Patients often present with multiple vertebral fractures with, one review reporting a mean of 2-5.8 vertebral fractures per patient.^{66,83} Commonly, fractures occur in the thoracic and/or in the lumbar spine, indicating the need for evaluation of both if PLO is suspected. Non-vertebral fractures might occur but are less common; they can include those of the hip, femoral neck, sacrum, radius, humerus, ribs, and feet.^{15,66} Some patients might experience hip pain accompanied by gait impairment, which might result in increasing immobility and/or fractures. In these patients, transient osteoporosis of the hip (TOH) should be considered as an alternative diagnosis. This is a recognized cause of hip fracture in pregnancy but is distinct from PLO and is characterized by localized edema visible on magnetic resonance imaging (MRI) and not skeletal resorption.^{16,84-86} In TOH, pain is localized to the hip, often unilaterally, with no other skeletal sites involved, and it generally resolves spontaneously postpartum.

4.4 | Pathophysiology

Studies at the tissue level suggest that the characteristics of bone structure and metabolism seem to differ between women with PLO and those without PLO or with idiopathic premenopausal osteoporosis. In a case-control study, women with PLO were found to have a more severe clinical presentation (significantly more total fractures, more vertebral fractures, and higher prevalence of multiple fractures), with more profound BMD deficits and lower tissue-level rates of mineral apposition and bone formation⁸⁷ than the comparative cohorts. In addition, they had lower serum levels of the bone turnover markers P1NP and CTX than the cohort with idiopathic osteoporosis. In a retrospective single-center study, BMD assessments using DXA, bone microarchitecture analysis using HR-pQCT, and measurements of bone turnover marker in seven women with PLO found severe deterioration of bone trabecular and cortical microarchitecture compared with a control group of healthy lactating women.⁶²

Data on the long-term outcomes of women with PLO are limited, but a single-center prospective cohort study that investigated subsequent fracture risk of 107 patients with PLO found that after 6 years of follow-up, almost 25% of these patients had sustained a subsequent fracture, and this fracture risk correlated with the number of fractures at time of PLO diagnosis.⁸⁸

5 | RECOMMENDATIONS FOR ASSESSMENT AND DIAGNOSIS OF PREGNANCY- AND LACTATION-ASSOCIATED OSTEOPOROSIS

The proposed pathway for assessment and diagnosis of women with suspected PLO will differ depending on whether they are antepartum or postpartum (Figure 2). Women with severe and persistent back pain during pregnancy or lactation should be referred for further specialist diagnostic procedures. For all patients, initial assessment should include a detailed review of their family and medical history to evaluate osteoporosis-related risk factors, especially of secondary causes of osteoporosis. This should include discussion of any previous and current pharmacological treatments, plus a physical examination, including documentation of height for clinical signs of fracture.

Women with clinical signs of fracture, which might include a height loss of ≥ 3 cm, who are currently pregnant should be referred for MRI, but this should be performed without the use of contrast

agents because they are contraindicated during pregnancy. If no fractures are present, conservative management is recommended, but if a recent peripheral or vertebral fragility fracture (Genant grade II) is observed, it is recommended that the patient's medical history and medical treatment are reviewed again to check for any oversight and a full laboratory work-up undertaken. Routine blood tests are needed to identify any possible secondary cause of osteoporosis. These should include C-reactive protein, complete blood count, calcium, phosphate, alkaline phosphatase, glucose, liver function tests, creatinine, thyroid-stimulating hormone, estrogen, 25-hydroxyvitamin D, parathyroid hormone, and protein electrophoresis. Additionally, the assessment of biochemical markers of bone turnover (CTX and P1NP) might be useful. This laboratory work-up should preferably be performed after delivery (post weaning) as some laboratory values will be affected by pregnancy. In terms of imaging techniques, although DXA is the recognized gold standard for the diagnosis of osteoporosis, the use of ionizing radiation is not recommended in pregnancy due to possible deleterious effects on the fetus. As an alternative, the use of QUS could be considered in antepartum women. This inexpensive, radiation-free method provides information on bone properties that are somewhat different to BMD and that may include bone quality.³⁸

In the case of postpartum women with persistent severe back pain, conventional X-ray optionally followed by MRI of the thoracic and lumbar spine can be used to determine the presence of fracture. If fragility fractures are present, further assessment follows the same pathway as for antepartum women, other than DXA may be used for imaging.




FIGURE 2 Assessment and diagnosis of women with suspected pregnancy- and lactation-associated osteoporosis (PLO). *Consider referral to an orthopedic specialist or traumatologist. **Should include C-reactive protein, complete blood count, calcium, phosphate, alkaline phosphatase, glucose, liver function tests, creatinine, thyroid-stimulating hormone, estrogen, 25-hydroxyvitamin D, parathyroid hormone, and protein electrophoresis. Additionally, the assessment of biochemical markers of bone turnover (CTX, P1NP) might be useful. ***Including vertebral fracture assessment, if available. DXA, dual energy X-ray absorptiometry; MRI, magnetic resonance imaging; PLO, pregnancy- and lactation-associated osteoporosis.

6 | RECOMMENDATIONS FOR TREATMENT AND FOLLOW-UP OF PREGNANCY- AND LACTATION-ASSOCIATED OSTEOPOROSIS

There is currently no standard of care for PLO. In fact, a recent systematic review and meta-analysis of the comparative effectiveness of interventions currently used for PLO concluded that due to heterogeneity of studies and lack of robust comparative data, no definitive conclusions can be drawn regarding the optimal therapeutic choice in this setting.⁶⁷ This position statement has been developed based on the available clinical evidence and expert opinion. The treatment and follow-up recommendations below are not intended as formal evidence-based guidelines and, due to the relative rarity of this condition and lack of controlled studies, the available data supporting these has not been assessed and graded for level of evidence.

None of the available pharmacological therapies for osteoporosis is licensed for use in PLO, and so use of any of them in this setting is considered off label. Data on their effectiveness in improving BMD and bone health is mostly drawn from studies in women with postmenopausal osteoporosis and a few observational trials in PLO.

Selection of the most appropriate management approach in women with a confirmed diagnosis of PLO is dependent on pregnancy status (pregnant or postpartum) and the number of vertebral

or non-vertebral fragility fractures present (Figure 3). The initial approach in all cases should be to check there is adequate dietary intake of calcium and vitamin D, with supplementation (calcium: 1000–2000 mg/day; vitamin D 1000–2000 IU/day) to ensure a sufficient vitamin D level of >30 ng/mL. However, while most data from randomized controlled trials indicate that vitamin D supplementation during pregnancy has positive effects on the BMD of offspring during childhood, it does not appear to have any specific beneficial effects on the maternal skeleton.⁸⁹ Patients with severe and persistent back pain during pregnancy and/or lactation with no vertebral or non-vertebral fragility fracture present should be referred to a specialist for further diagnosis/treatment.

In women with a single vertebral or non-vertebral fragility fracture and any T-score, if available, pain medication and later physiotherapy can be employed, if necessary; in some cases, conservative/operative treatment might also need to be considered. Patients might be encouraged to consider limiting/stopping breastfeeding, which should result in the resumption of menses and a regular menstrual cycle. Patients should be informed about effective methods of contraception (e.g., continuous use of the combined oral contraceptive pill or estradiol-containing oral contraceptives). If after weaning a regular menstrual cycle does not return or there are symptoms of estrogen deficiency, a gynecological consultation should be considered. Further follow-up with BMD and other laboratory assessments




FIGURE 3 Management of women diagnosed with pregnancy- and lactation-associated osteoporosis (PLO) based on the number of fragility fractures. *If after weaning a regular menstrual cycle does not resume, a gynecological consultation should be considered. BMD, bone mineral density; IU, international units; MRI, magnetic resonance imaging.

may be scheduled for 6 months' to 2 years' time, as BMD should at least partially recover. The role of hormone replacement therapy (HRT)/menopausal hormone therapy (MHT) for the treatment of PLO is unknown. HRT/MHT could be considered in the setting of lactation-associated amenorrhea and estrogen deficiency if the woman did not want to cease breastfeeding. Limited data indicated that transdermal estrogen replacement therapy does not adversely affect lactation in women with established lactation.⁹⁰ Medical treatment with osteoporosis therapies can be considered, but these may generally be reserved for women with ≥ 2 vertebral or non-vertebral fragility fractures, regardless of T-score if available. Avoidance of medical treatment is the ideal scenario because all are unlicensed in PLO and their effects on the fetus in future potential pregnancies are not well established. As a result, it is recommended that all medical treatments should be combined with contraceptives, with the exception of the use of DMPA. Although treatment with bisphosphonates has been shown to improve BMD in women with PLO,^{67,91,92} they should be avoided where possible as they persist in bone for many years, are able to cross the placenta barrier, and might affect future pregnancies.

It is recommended that in postpartum women with ≥ 2 vertebral or non-vertebral fragility fractures during pregnancy and/or lactation, the first choice of treatment is teriparatide/abaloparatide or perhaps romosozumab. Teriparatide and abaloparatide are both osteoanabolic agents that mimic parathyroid hormone and are indicated for the treatment of severe osteoporosis in postmenopausal women.^{93,94} These are good therapeutic options as they increase bone turnover and BMD, particularly at the spine, where the majority of fractures occur in PLO. The efficacy of teriparatide (a recombinant human parathyroid hormone [PTH 1–34] analog) has been demonstrated in postmenopausal women with multiple vertebral fractures. In the Fracture Prevention Trial (FPT), it was effective in reducing the risk of vertebral and non-vertebral fractures and increasing BMD in postmenopausal women with osteoporosis.^{95,96} The FPT follow-up study showed that 18 months after discontinuation of teriparatide treatment, the reduction in vertebral fracture risk was still evident.⁹⁷

There have been some small studies and case reports of the successful use of teriparatide in PLO, as summarized in a review by Ali and colleagues.⁹⁸ In a patient with severe PLO and 11 vertebral fractures, teriparatide treatment increased her BMD and she had no further fractures.⁹⁹ In a multicenter retrospective cohort study, 19 women with PLO were treated with teriparatide (plus calcium and vitamin D supplementation) for up to 18 months. Teriparatide treatment resulted in significantly greater increases in lumbar spine BMD compared with supplementation alone at 12 and 24 months.¹⁰⁰ A retrospective study evaluated 47 patients with PLO who had suffered multiple vertebral fractures (mean: 4.4 fractures) postpartum and had been referred to the German Reference Center for PLO in Frankfurt between 2004 and 2018.¹⁰¹ Treatment with teriparatide for 24 months significantly increased BMD, predominantly at the spine, and no significant decrease in BMD was observed in the 12 months following cessation of

treatment without any follow-up treatment. When used in PLO and after completion of teriparatide treatment with no subsequent fractures, no follow-up treatment is regularly necessary if the patient has resumed a normal menstrual cycle, as endogenous estrogen is a potent antiresorptive.

Romosozumab is also a possible treatment option in PLO in patients who have either vertebral or non-vertebral fragility fractures, but it has rarely been applied in PLO and should be used with caution.^{102–105} Aside from these observations from individual case reports, there is only one single-arm study to date that has evaluated the effect of romosozumab treatment for up to 12 months in a cohort of 29 women with PLO. Romosozumab was associated with large BMD increases at 6 and 12 months, particularly at the lumbar spine, in this cohort. While these findings are promising, they require confirmation in further trials and evaluation of longer-term outcomes.¹⁰⁶

Denosumab, an antiresorptive agent that blocks RANKL, is a possible second-choice treatment option in this setting. Denosumab reduced the risk of vertebral, nonvertebral, and hip fractures and increased BMD across skeletal sites compared to placebo in clinical trials of postmenopausal women with osteoporosis or low BMD.¹⁰⁷ However, denosumab is associated with a rebound phenomenon whereby cessation of therapy is associated with a rapid increase in markers of bone turnover, loss of BMD, and an increased risk of fractures, in particular multiple vertebral fractures.¹⁰⁸ As a result, sequential treatment with bisphosphonates is usually recommended, but as previously highlighted, they should be avoided, if possible, in PLO. In a case report, denosumab was used successfully to treat a woman with severe PLO and vertebral fractures.¹⁰⁹ After 18 months of treatment with denosumab, a clinically significant increase of BMD was observed at the lumbar spine, femoral neck, and total hip of 32.2%, 13.0%, and 11.5%, respectively, with no further subsequent fractures.

7 | RESEARCH GAPS IN PREGNANCY- AND LACTATION-ASSOCIATED OSTEOPOROSIS

While our understanding of the pathophysiology and treatment of PLO has expanded over the past decade, there are still several gaps in our knowledge regarding this relatively rare but debilitating condition.¹¹⁰ The epidemiology of PLO has not been widely investigated and studies on the long-term outcomes of women who experience PLO are limited, with only a few studies to date addressing the important clinical question of whether an increased risk of fracture in subsequent pregnancies exists. Similarly, it is still unclear if women who have had PLO in the past are at an increased risk of osteoporosis-related fractures after menopause compared to women who have not had PLO. In the absence of available clinical evidence, ideally an intensified follow-up should be considered in women with a history of PLO (DXA at the time of menopause as well as 2–3 years later). Depending on the individual BMD,

microarchitectural deterioration and existing risk factors, shared decision-making regarding a treatment plan is advised.

Further outstanding clinical questions relate to the optimal contraceptive method during or after weaning, which is still a matter of debate, and the procedure that should be followed after teriparatide treatment in women with PLO and more than two fragility fractures. Is endogenous estradiol in regularly menstruating women sufficient or should a follow-up treatment be advised?

8 | CONCLUSION

PLO is a rare disease in pregnant and lactating women which, in the majority of cases, is identified late, highlighting the need for vigilance among clinicians to ensure that it is correctly identified, properly diagnosed and effectively managed. PLO often presents with multiple – predominantly spinal – fractures that lead to considerable pain and discomfort as well as having severe long-term consequences for the women affected. This position statement includes a pragmatic approach for identifying women with PLO in clinical practice and suggests developing an individualized treatment plan which may include pharmaceutical intervention.

AUTHOR CONTRIBUTIONS

Peyman Hadji: Conceptualization, Writing – original draft, Writing – review and editing. All other authors: Writing – review and editing. All authors meet the qualifications for authorship as detailed in the ICMJE specifications.

AFFILIATIONS

¹Philipps-Universität Marburg, Marburg, Germany

²Frankfurt Centre for Bone Health and Endocrinology, Frankfurt, Germany

³Department of Obstetrics, Gynecology and Maternal Fetal Medicine, Aristotle University of Thessaloniki, Thessaloniki, Greece

⁴Fondazione FIRMO, Florence, Italy

⁵Division of Endocrinology and Metabolism, McMaster University, Hamilton, Canada

⁶American University of Beirut, Beirut, Lebanon

⁷Osteoporosis and Bone Metabolism Unit, Department of Endocrinology, Singapore General Hospital, Singapore, Singapore

⁸DUKE NUS Medical School, Singapore

⁹Northwell Health Zucker School of Medicine, New York, New York, USA

¹⁰Donald and Barbara Zucker School of Medicine at Hofstra/Northwell, Hempstead, New York, USA

¹¹Second Department of Obstetrics and Gynecology, Medical School, National and Kapodistrian University of Athens, Athens, Greece

¹²Department of Rheumatology, Amsterdam UMC, Amsterdam, the Netherlands

¹³School of Medicine, Aristotle University of Thessaloniki, Thessaloniki, Greece

¹⁴Division of Clinical Medicine, School of Medicine and Population Health, and Mellanby Centre for Musculoskeletal Research, University of Sheffield, Sheffield, UK

¹⁵Department of Clinical, Surgical, Diagnostic and Pediatric Sciences, University of Pavia, and Research Center for Reproductive Medicine, Gynecological Endocrinology and Menopause, IRCCS S. Matteo Foundation, Pavia, Italy

¹⁶Department of Medical and Cardiovascular Sciences, Sapienza University of Rome, Rome, Italy

¹⁷Protein Research Chair, Biochemistry Department, College of Science, King Saud University, Riyadh, Kingdom of Saudi Arabia

¹⁸Geneva University Hospitals and Faculty of Medicine, Geneva, Switzerland

¹⁹Department of Obstetrics and Gynecology, University Clinic Inselspital Bern, Bern, Switzerland

²⁰Department of Physical Medicine and Rehabilitation, Faculty of Medicine, Istanbul University-Cerrahpaşa, Cerrahpaşa, Turkey

²¹Department of Endocrinology, Monash Health and Monash Centre for Health Research and Implementation, Monash University, Clayton, Victoria, Australia

²²IDIM, Institute of Metabolic Research, Universidad del Salvador, Buenos Aires, Argentina

²³Department of Internal Medicine, Erasmus MC, Erasmus University Medical Center, Rotterdam, the Netherlands

²⁴Section for Prenatal Diagnostics and Fetal Therapy, Asklepios Clinics Hamburg, Hamburg, Germany

ACKNOWLEDGMENTS

Editorial assistance in the development of the manuscript was provided by Dr. Karen Wolstencroft. Open Access funding enabled and organized by Projekt DEAL.

FUNDING INFORMATION

No specific grant was received from funding agencies in the public, commercial, or non-profit sectors for the development of this position statement. The contribution of JYR to this manuscript has been funded by the Distinguished Scientist Fellowship Program (DSFP) of the King Saud University, Riyadh, Kingdom of Saudi Arabia.

CONFLICT OF INTEREST STATEMENT

Peyman Hadji has received speaker and consultancy fees/advisory board honoraria from Agita, Amgen, Celtrion, Fresenius, Hexal, Gedeon Richter, Novartis, Samsung, Sandoz, Stada, Theramax, and UCB. **Apostolos Athanasiadis** has no COI regarding this work. **Maria-Luisa Brandi** has received honoraria from Amgen, Ascendis, Bruno Farmaceutici, Calcilytix and Kyowa Kirin; grants and/or speaker fees from Alexion, Amgen, Amolyt, Bruno Farmaceutici, CoGeDi, Echolight, Gedeon Richter, Kyowa Kirin, Monte Rosa Therapeutics and UCB; consultancy fees from Aboca, Alexion, Amolyt, Bruno Farmaceutici, Calcilytix, Echolight, Enterabio, Kyowa Kirin, Personal Genomics, and Septerna. **Marlene Chakhtoura** has no COI regarding this work. **Manju Chandran** has received honoraria and travel grants from Amgen and Promedius. **Frank**

Chervenak has no COI regarding this work. **Amos Grünebaum** has no COI regarding this work. **Ioannis Kyvernitakis** has no COI regarding this work. **Irene Lambrinoudaki** has no COI regarding this work. **Willem F Lems** has received speaker fees/advisory board honoraria from Amgen, Theramax, UCB, Pfizer, Novartis and Galapagos. **Apostolos M. Mamopoulos** is a member of the advisory board of Pfizer Pharmaceuticals for the RSV vaccination program, and a member of the advisory board for ITF Pharmaceuticals for the treatment of hyperemesis gravidarum. **Eugene McCloskey** has no COI regarding this work. **Rossella E. Nappi** has ongoing consultancy relationships with Abbott, Astellas, Bayer HealthCare AG, Besins Healthcare, Biocodex, Exeltis, Fidia, Gedeon Richter, Merck & Co, Novo Nordisk, Organon & Co, Shionogi Limited, Theramex, Viatrix, and Vichy Laboratories. She has no COI regarding this work other than she is the current President of the International Menopause Society (IMS). **Jessica Pepe** has no COI regarding this work. **Jean-Yves Reginster** has no COI regarding this work other than he is President of the European Society for Clinical and Economics Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (ESCEO). **René Rizzoli** has no COI regarding this work. **Petra Stute** has no COI regarding this work. **Şansin Tuzun** has no COI regarding this work. **Amanda J Vincent** has received consulting fees or honoraria from Theramex, Astellas, Besins, IQ Fertility, Alcimed and Dietitians Australia and is a board member of the International Menopause Society. **Maria Belen Zanchetta** has no COI regarding this work. **M. Carola Zillikens** has received honoraria for lectures or consulting from Amgen, Kyowa Kirin, and Theramex and research support from Kyowa Kirin for her institution.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Peyman Hadji <https://orcid.org/0000-0003-4787-3151>

REFERENCES

1. Morin SN, Leslie WD, Schousboe JT. Osteoporosis: a review. *JAMA*. 2025;334(10):894-907.
2. Kanis JA, McCloskey EV, Johansson H, Oden A, Melton LJ 3rd, Khaltayev N. A reference standard for the description of osteoporosis. *Bone*. 2008;42(3):467-475.
3. Salari N, Darvishi N, Bartina Y, et al. Global prevalence of osteoporosis among the world older adults: a comprehensive systematic review and meta-analysis. *J Orthop Surg Res*. 2021;16(1):669.
4. Safiri S, Amiri F, Karamzad N, et al. Global Health effects of osteopenia and osteoporosis, 1990-2021. *Health Sci Rep*. 2025;8(10):e71378.
5. Xiao PL, Cui AY, Hsu CJ, et al. Global, regional prevalence, and risk factors of osteoporosis according to the World Health Organization diagnostic criteria: a systematic review and meta-analysis. *Osteoporos Int*. 2022;33(10):2137-2153.
6. Liang H, Chen S, Shi M, et al. Global epidemiology and burden of osteoporosis among postmenopausal women: insights from the global burden of disease study 2021. *NPJ Aging*. 2025;11(1):78.

7. Anagnostis P, Siolos P, Gkekas NK, et al. Association between age at menopause and fracture risk: a systematic review and meta-analysis. *Endocrine*. 2019;63(2):213-224.
8. Zerikly R, Demetriou EW. Use of fracture risk assessment tool in clinical practice and fracture risk assessment tool future directions. *Womens Health*. 2024;20:17455057241231387.
9. Schini M, Johansson H, Harvey NC, Lorentzon M, Kanis JA, McCloskey EV. An overview of the use of the fracture risk assessment tool (FRAX) in osteoporosis. *J Endocrinol Investig*. 2024;47(3):501-511.
10. Ebeling PR, Nguyen HH, Aleksova J, Vincent AJ, Wong P, Milat F. Secondary osteoporosis. *Endocr Rev*. 2022;43(2):240-313.
11. Hadji P, Aapro M, Al-Dagri N, et al. Management of aromatase inhibitor-associated bone loss (AIBL) in women with hormone-sensitive breast cancer: an updated joint position statement of the IOF, CABS, ECTS, IEG, ESCEO, IMS, and SIOG. *J Bone Oncol*. 2025;53:100694.
12. Ahn E, Lee J, Park YS, Noh HM, Kim BH. Association between delivery at an advanced maternal age and osteoporosis in elderly Korean women. *J Bone Miner Metab*. 2015;33(6):666-673.
13. Orhadjie E, Makaram N, Berg K, Hauser B, Ralston SH. Clinical presentation, risk factors and management of pregnancy-associated osteoporosis: a systematic review and meta-analysis. *Osteoporos Int*. 2025;36(6):981-993.
14. Qian Y, Wang L, Yu L, Huang W. Pregnancy- and lactation-associated osteoporosis with vertebral fractures: a systematic review. *BMC Musculoskelet Disord*. 2021;22(1):926.
15. Hardcastle SA. Pregnancy and lactation associated osteoporosis. *Calcif Tissue Int*. 2022;110(5):531-545.
16. Kovacs CS, Ralston SH. Presentation and management of osteoporosis presenting in association with pregnancy or lactation. *Osteoporos Int*. 2015;26(9):2223-2241.
17. Gehlen M, Lazarescu AD, Hinz C, et al. Long-term outcome of patients with pregnancy and lactation-associated osteoporosis (PLO) with a particular focus on quality of life. *Clin Rheumatol*. 2019;38(12):3575-3583.
18. Riggs BL, Khosla S, Melton LJ 3rd. The assembly of the adult skeleton during growth and maturation: implications for senile osteoporosis. *J Clin Invest*. 1999;104(6):671-672.
19. Chevalley T, Rizzoli R. Acquisition of peak bone mass. *Best Pract Res Clin Endocrinol Metab*. 2022;36(2):101616.
20. Xue S, Kemal O, Lu M, Lix LM, Leslie WD, Yang S. Age at attainment of peak bone mineral density and its associated factors: the National Health and nutrition examination survey 2005-2014. *Bone*. 2020;131:115163.
21. Berger C, Goltzman D, Langsetmo L, et al. Peak bone mass from longitudinal data: implications for the prevalence, pathophysiology, and diagnosis of osteoporosis. *J Bone Miner Res*. 2010;25(9):1948-1957.
22. Makker A, Mishra G, Singh BP, Tripathi A, Singh MM. Normative bone mineral density data at multiple skeletal sites in Indian subjects. *Arch Osteoporos*. 2008;3(1):25-37.
23. Shivane VK, Sarathi V, Lila AR, et al. Peak bone mineral density and its determinants in an Asian Indian population. *J Clin Densitom*. 2012;15(2):152-158.
24. Berkane N, Liere P, Oudinet JP, et al. From pregnancy to preeclampsia: a key role for estrogens. *Endocr Rev*. 2017;38(2):123-144.
25. Gak N, Abbara A, Dhillon WS, Keen R, Comninou AN. Current and future perspectives on pregnancy and lactation-associated osteoporosis. *Front Endocrinol (Lausanne)*. 2024;15:1494965.
26. Chandran M, Tse Tan SY. Disorders of bone and mineral metabolism in pregnancy and lactation: a case based clinical review. *Osteoporos Sarcopenia*. 2025;11(1):1-8.
27. Kovacs CS. Pregnancy and lactation associated bone fragility. *J Endocr Soc*. 2025;9(9):bvaf126.

28. Kovacs CS. Maternal mineral and bone metabolism during pregnancy, lactation, and post-weaning recovery. *Physiol Rev*. 2016;96(2):449-547.
29. Kovacs CS. The role of vitamin D in pregnancy and lactation: insights from animal models and clinical studies. *Annu Rev Nutr*. 2012;32:97-123.
30. Kovacs CS. Calcium and phosphate metabolism and related disorders during pregnancy and lactation. In: Feingold KR, Ahmed SF, Anawalt B, et al., eds. *Endotext* [Internet]. MDText.com, Inc.; 2000. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK279173/>.
31. Cooper MS. Disorders of calcium metabolism and parathyroid disease. *Best Pract Res Clin Endocrinol Metab*. 2011;25(6):975-983.
32. Styczynska H, Lis K, Sobanska I, Pater A, Pollak J, Mankowska A. Bone turnover markers and Osteoprotegerin in uncomplicated pregnancy. *EJIFCC*. 2009;19(4):193-202.
33. Hong JS, Santolaya-Forgas J, Romero R, et al. Maternal plasma osteoprotegerin concentration in normal pregnancy. *Am J Obstet Gynecol*. 2005;193(3 Pt 2):1011-1015.
34. Bhattoa HP, Vasikaran S, Trifonidi I, et al. Update on the role of bone turnover markers in the diagnosis and management of osteoporosis: a consensus paper from the European Society for Clinical and Economic Aspects of osteoporosis, osteoarthritis and musculoskeletal diseases (ESCEO), international osteoporosis foundation (IOF), and International Federation of Clinical Chemistry and Laboratory Medicine (IFCC). *Osteoporos Int*. 2025;36(4):579-608.
35. Hellmeyer L, Ziller V, Anderer G, Ossendorf A, Schmidt S, Hadji P. Biochemical markers of bone turnover during pregnancy: a longitudinal study. *Exp Clin Endocrinol Diabetes*. 2006;114(9):506-510.
36. Ulrich U, Miller PB, Eyre DR, Chesnut CH 3rd, Schlebusch H, Soules MR. Bone remodeling and bone mineral density during pregnancy. *Arch Gynecol Obstet*. 2003;268(4):309-316.
37. Hellmeyer L, Ossendorf A, Ziller V, Tekesin I, Schmidt S, Hadji P. Quantitative ultrasonometry of the phalanges during pregnancy: a longitudinal study. *Climacteric*. 2006;9(6):446-451.
38. Hellmeyer L, Hahn B, Fischer C, et al. Quantitative ultrasonometry during pregnancy and lactation: a longitudinal study. *Osteoporos Int*. 2015;26(3):1147-1154.
39. Hellmeyer L, Hadji P, Ziller V, Wagner U, Schmidt S. Pregnancy-associated osteoporosis. *Geburtshilfe Frauenheilkd*. 2004;65:38-45.
40. To WW, Wong MW. Persistence of back pain symptoms after pregnancy and bone mineral density changes as measured by quantitative ultrasound—a two year longitudinal follow up study. *BMC Musculoskelet Disord*. 2011;12:55.
41. Kalkwarf HJ, Specker BL. Bone mineral changes during pregnancy and lactation. *Endocrine*. 2002;17(1):49-53.
42. Moller UK, Vieth Streym S, Mosekilde L, Rejnmark L. Changes in bone mineral density and body composition during pregnancy and postpartum. A controlled cohort study. *Osteoporos Int*. 2012;23(4):1213-1223.
43. Breasail OM, Prentice A, Ward K. Pregnancy-related bone mineral and microarchitecture changes in women aged 30 to 45 years. *J Bone Miner Res*. 2020;35(7):1253-1262.
44. Polatti F, Capuzzo E, Viazzo F, Colleoni R, Klersy C. Bone mineral changes during and after lactation. *Obstet Gynecol*. 1999;94(1):52-56.
45. de Bakker CMJ, Tseng WJ, Li Y, et al. Reproduction differentially affects trabecular bone depending on its mechanical versus metabolic role. *J Biomech Eng*. 2017;139(11):1110061.
46. Egund L, Malmgren L, Woolf AD, FE MG, Akesson KE. Recovery of BMD after pregnancy and breastfeeding—a 10-yr prospective observational study of 25-yr-old women. *J Bone Miner Res*. 2025;40(12):1360-1369.
47. Athonvarangkul D, Wysolmerski JJ. Crosstalk within a brain-breast-bone axis regulates mineral and skeletal metabolism during lactation. *Front Physiol*. 2023;14:1121579.
48. Kovacs CS. Calcium and bone metabolism during pregnancy and lactation. *J Mammary Gland Biol Neoplasia*. 2005;10(2):105-118.
49. Kovacs CS. The skeleton is a storehouse of mineral that is plundered during lactation and (fully?) replenished afterwards. *J Bone Miner Res*. 2017;32(4):676-680.
50. Dura-Trave T, Gallinas-Victoriano F. Pregnancy, breastfeeding, and vitamin D. *Int J Mol Sci*. 2023;24(15):11881.
51. Brannon PM, Picciano MF. Vitamin D in pregnancy and lactation in humans. *Annu Rev Nutr*. 2011;31:89-115.
52. Babey ME, Krause WC, Chen K, et al. A maternal brain hormone that builds bone. *Nature*. 2024;632(8024):357-365.
53. Grizzo FMF, Alarcao ACJ, Dell' Agnolo CM, et al. How does women's bone health recover after lactation? A systematic review and meta-analysis. *Osteoporos Int*. 2020;31(3):413-427.
54. Oboh I, Coleman C, Cremona A. The influence of lactation and its duration on bone mineral density in pregnancy and postpartum: a systematic review with meta-analysis. *Clin Nutr ESPEN*. 2021;46:121-132.
55. Cooke-Hubley S, Gao Z, Mugford G, et al. Parity and lactation are not associated with incident fragility fractures or radiographic vertebral fractures over 16 years of follow-up: Canadian multicentre osteoporosis study (CaMos). *Arch Osteoporos*. 2019;14(1):49.
56. Hadji P, Ziller V, Kalder M, et al. Influence of pregnancy and breastfeeding on quantitative ultrasonometry of bone in postmenopausal women. *Climacteric*. 2002;5(3):277-285.
57. Liu XS, Wang L, de Bakker CMJ, Lai X. Mechanical regulation of the maternal skeleton during reproduction and lactation. *Curr Osteoporos Rep*. 2019;17(6):375-386.
58. de Bakker CMJ, Burt LA, Gabel L, Hanley DA, Boyd SK. Associations between breastfeeding history and early postmenopausal bone loss. *Calcif Tissue Int*. 2020;106(3):264-273.
59. Orhadje E, Berg K, Hauser B, Ralston SH. Clinical features, incidence and treatment outcome in pregnancy-associated osteoporosis: a single-Centre experience over two decades. *Calcif Tissue Int*. 2023;113(6):591-596.
60. Hadji P, Boekhoff J, Hahn M, Hellmeyer L, Hars O, Kyvernitis I. Pregnancy-associated osteoporosis: a case-control study. *Osteoporos Int*. 2017;28(4):1393-1399.
61. Scioscia MF, Zanchetta MB. Recent insights into pregnancy and lactation-associated osteoporosis (PLO). *Int J Women's Health*. 2023;15:1227-1238.
62. Scioscia MF, Vidal M, Sarli M, et al. Severe bone microarchitecture impairment in women with pregnancy and lactation-associated osteoporosis. *J Endocr Soc*. 2021;5(5):bvab031.
63. Agarwal S, El-Najjar D, Kondapalli A, et al. HR-pQCT reveals marked trabecular and cortical structural deficits in women with pregnancy and lactation-associated osteoporosis (PLO). *J Bone Miner Res*. 2024;40(1):38-49.
64. Langdahl BL. Osteoporosis in premenopausal women. *Curr Opin Rheumatol*. 2017;29(4):410-415.
65. Yang B, Kong L, Hao D. Challenges in the Management of Pregnancy and Lactation Associated Osteoporosis: literature review and retrospective study. *Curr Stem Cell Res Ther*. 2021;16(6):688-694.
66. Carsote M, Turtorea MR, Valea A, Buescu C, Nistor C, Turtorea IF. Bridging the gap: pregnancy-and lactation-associated osteoporosis. *Diagnostics (Basel)*. 2023;13(9):1615.
67. Anagnostis P, Lampropoulou-Adamidou K, Bosdou JK, et al. Comparative effectiveness of therapeutic interventions in pregnancy and lactation-associated osteoporosis: a systematic review and meta-analysis. *J Clin Endocrinol Metab*. 2024;109(3):879-901.
68. Issa Z, Saad R, Lama Y, et al. Pregnancy and lactation associated osteoporosis: a systematic review. *Calcif Tissue Int*. 2026;117(1):42.
69. Laroche M, Talibart M, Cormier C, Roux C, Guggenbuhl P, Degboe Y. Pregnancy-related fractures: a retrospective study of a French

- cohort of 52 patients and review of the literature. *Osteoporos Int*. 2017;28(11):3135-3142.
70. Miles B, Panchbhavi M, Mackey JD. Analysis of fracture incidence in 135 patients with pregnancy and lactation osteoporosis (PLO). *Cureus*. 2021;13(10):e19011.
 71. Toba M, Terauchi M, Moriwaki M, Obayashi S, Miyasaka N, Fushimi K. Fractures within 2 years of an obstetric hospitalization: analysis of nationwide administrative data in Japan. *J Bone Miner Metab*. 2022;40(5):748-754.
 72. Kasahara K, Tanaka-Mizuno S, Tsuji S, et al. Pregnancy and lactation-associated osteoporosis as a major type of premenopausal osteoporosis: a retrospective cohort study based on real-world data. *BMC Pregnancy Childbirth*. 2024;24(1):301.
 73. Tuna F, Akleyek C, Ozdemir H, Demirbag Kabayel D. Risk factors, fractures, and management of pregnancy-associated osteoporosis: a retrospective study of 14 Turkish patients. *Gynecol Endocrinol*. 2020;36(3):238-242.
 74. Hardcastle SA, Yahya F, Bhalla AK. Pregnancy-associated osteoporosis: a UK case series and literature review. *Osteoporos Int*. 2019;30(5):939-948.
 75. Butscheidt S, Delsmann A, Rolvien T, et al. Mutational analysis uncovers monogenic bone disorders in women with pregnancy-associated osteoporosis: three novel mutations in LRP5, COL1A1, and COL1A2. *Osteoporos Int*. 2018;29(7):1643-1651.
 76. Butscheidt S, Tsourdi E, Rolvien T, et al. Relevant genetic variants are common in women with pregnancy and lactation-associated osteoporosis (PLO) and predispose to more severe clinical manifestations. *Bone*. 2021;147:115911.
 77. Fu Z, Ni X, Han S, et al. Gene variants and bisphosphonates treatment in pregnancy and lactation-associated osteoporosis (PLO): a retrospective study of 22 Chinese patients. *Calcif Tissue Int*. 2025;116(1):70.
 78. Lynch L, Shea PR, Vena N, et al. Genetic contributors to osteoporosis in pregnancy and lactation associated osteoporosis (PLO). *Bone*. 2026;204:117744.
 79. Koumakis E, Cormier-Daire V, Dellal A, et al. Osteogenesis imperfecta: characterization of fractures during pregnancy and postpartum. *Orphanet J Rare Dis*. 2022;17(1):22.
 80. Cook FJ, Mumm S, Whyte MP, Wenkert D. Pregnancy-associated osteoporosis with a heterozygous deactivating LDL receptor-related protein 5 (LRP5) mutation and a homozygous methylenetetrahydrofolate reductase (MTHFR) polymorphism. *J Bone Miner Res*. 2014;29(4):922-928.
 81. Kovacs CS, Chaussain C, Osdoby P, Brandi ML, Clarke B, Thakker RV. The role of biomineralization in disorders of skeletal development and tooth formation. *Nat Rev Endocrinol*. 2021;17(6):336-349.
 82. Orhadje E, Berg K, Whitehead K, Hauser B, Ralston SH. Height loss, back pain and severe pain are highly predictive of pregnancy associated osteoporosis. *Osteoporos Int*. 2025;36(9):1719-1724.
 83. Kondapalli AV, Kamanda-Kosseh M, Williams JM, et al. Clinical characteristics of pregnancy and lactation associated osteoporosis: an online survey study. *Osteoporos Int*. 2023;34(8):1477-1489.
 84. Galanis A, Dimopoulou S, Karampinas P, et al. The correlation between transient osteoporosis of the hip and pregnancy: a review. *Medicine (Baltimore)*. 2023;102(41):e35475.
 85. Hadji P, Boekhoff J, Hahn M, Hellmeyer L, Hars O, Kyvernitis I. Pregnancy-associated transient osteoporosis of the hip: results of a case-control study. *Arch Osteoporos*. 2017;12(1):11.
 86. Quaresima P, Angeletti M, Luziatelli D, et al. Pregnancy associated transient osteoporosis of the hip (PR-TOH): a non-obstetric indication to caesarean section. A case report with literature review. *Eur J Obstet Gynecol Reprod Biol*. 2021;262:28-35.
 87. Cohen A, Kamanda-Kosseh M, Dempster DW, et al. Women with pregnancy and lactation-associated osteoporosis (PLO) have low bone remodeling rates at the tissue level. *J Bone Miner Res*. 2019;34(9):1552-1561.
 88. Kyvernitis I, Reuter TC, Hellmeyer L, Hars O, Hadji P. Subsequent fracture risk of women with pregnancy and lactation-associated osteoporosis after a median of 6 years of follow-up. *Osteoporos Int*. 2018;29(1):135-142.
 89. Moon RJ, D'Angelo S, Curtis EM, et al. Pregnancy vitamin D supplementation and bone mineral density of the mother: a post hoc analysis of the MAVIDOS randomised placebo-controlled trial. *Osteoporos Int*. 2025;36(9):1583-1591.
 90. Estradiol. National Institute of Child Health and Human Development; 2026.
 91. Sokal A, Elefant E, Leturcq T, Beghin D, Mariette X, Seror R. Pregnancy and newborn outcomes after exposure to bisphosphonates: a case-control study. *Osteoporos Int*. 2019;30(1):221-229.
 92. Grizzo FM, da Silva Martins J, Pinheiro MM, Jorgetti V, Carvalho MD, Pelloso SM. Pregnancy and lactation-associated osteoporosis: bone Histomorphometric analysis and response to treatment with zoledronic acid. *Calcif Tissue Int*. 2015;97(4):421-425.
 93. Ebina K, Etani Y, Noguchi T, Nakata K, Okada S. Clinical effects of teriparatide, abaloparatide, and romosozumab in postmenopausal osteoporosis. *J Bone Miner Metab*. 2025;43(1):3-9.
 94. Tabatabai L, Cosman F, Curtis JR, et al. Comparative effectiveness of Abaloparatide and teriparatide in women 50 years of age and older: update of a real-world retrospective analysis. *Endocr Pract*. 2025;31(2):159-168.
 95. Neer RM, Arnaud CD, Zanchetta JR, et al. Effect of parathyroid hormone (1-34) on fractures and bone mineral density in postmenopausal women with osteoporosis. *N Engl J Med*. 2001;344(19):1434-1441.
 96. Brixen KT, Christensen PM, Ejersted C, Langdahl BL. Teriparatide (biosynthetic human parathyroid hormone 1-34): a new paradigm in the treatment of osteoporosis. *Basic Clin Pharmacol Toxicol*. 2004;94(6):260-270.
 97. Lindsay R, Scheele WH, Neer R, et al. Sustained vertebral fracture risk reduction after withdrawal of teriparatide in postmenopausal women with osteoporosis. *Arch Intern Med*. 2004;164(18):2024-2030.
 98. Ali DS, Khan AA, Brandi ML. Effective strategies for pregnancy and lactation-associated osteoporosis: teriparatide use in focus. *Endocrine*. 2024;86(2):459-469.
 99. Winarno AS, Kyvernitis I, Hadji P. Successful treatment of 1-34 parathyroid hormone (PTH) after failure of bisphosphonate therapy in a complex case of pregnancy associated osteoporosis and multiple fractures. *Z Geburtshilfe Neonatol*. 2014;218(4):171-173.
 100. Lampropoulou-Adamidou K, Trovas G, Triantafyllopoulos IK, et al. Teriparatide treatment in patients with pregnancy- and lactation-associated osteoporosis. *Calcif Tissue Int*. 2021;109(5):554-562.
 101. Hadji P, Mouzakiti N, Kyvernitis I. Effect of teriparatide on subsequent fracture and bone mineral density in 47 women with pregnancy- and lactation-associated osteoporosis and vertebral fractures. *Geburtshilfe Frauenheilkd*. 2022;82(6):619-626.
 102. Goshima A, Tsuboi H, Kaito T, Iwasaki M. Effective management of pregnancy- and lactation-associated osteoporosis using romosozumab: a clinical case report. *JBMR Plus*. 2025;9(10):z1af141.
 103. Nomura H, Nomura S. Pregnancy and lactation-related osteoporosis associating multiple vertebral fragility fractures treated with romosozumab: a case report. *AME Case Rep*. 2025;9:68.
 104. Kaneuchi Y, Iwabuchi M, Hakozaaki M, Yamada H, Konno SI. Pregnancy and lactation-associated osteoporosis successfully treated with Romosozumab: a case report. *Medicina (Kaunas)*. 2022;59(1):19.
 105. Lee S, Hong N, Cho SJ, Shin S, Rhee Y. Effect of romosozumab on bone mineral density and trabecular bone score

- in premenopausal women with low bone mass. *Osteoporos Int.* 2025;36(2):323-331.
106. Lynch LK, Kamanda-Kosseh M, Agarwal S, et al. *Romsozumab Is Associated with Large BMD Increases in Premenopausal Idiopathic Osteoporosis*. American Society for Bone and Mineral Research Annual Meeting; 2025.
107. Deeks ED. Denosumab: a review in postmenopausal osteoporosis. *Drugs Aging.* 2018;35(2):163-173.
108. Anastasilakis AD, Makras P, Yavropoulou MP, Tabacco G, Naciu AM, Palermo A. Denosumab discontinuation and the rebound phenomenon: a narrative review. *J Clin Med.* 2021;10(1):152.
109. Stumpf U, Kraus M, Hadji P. Influence of denosumab on bone mineral density in a severe case of pregnancy-associated osteoporosis. *Osteoporos Int.* 2021;32(11):2383-2387.
110. Mobasheri A, Reginster JY, Chandran M, McSwan J. Bone health and pregnancy: osteoporosis and beyond. *Case Rep Womens Health.* 2025;47:e00725.

How to cite this article: Hadji P, Athanasiadis A, Brandi M-L, et al. Pregnancy- and lactation-associated osteoporosis: A position statement of the IAPM, IOF, ECTS, ESCEO, IMS, and EMAS. *Int J Gynecol Obstet.* 2026;00:1-13. doi:[10.1002/ijgo.71203](https://doi.org/10.1002/ijgo.71203)