White Rose Research Online

eprints@whiterose.ac.uk
https://eprints.whiterose.ac.uk
Universities of Leeds, Sheffield and York

Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:
https://eprints.whiterose.ac.uk/id/eprint/247170/

Version: Accepted Version

Article:
Bavula, V.V. (2026) Ore sets, denominator sets and the left regular left quotient ring of a ring. Journal of Algebra and Its Applications. ISSN: 0219-4988

https://doi.org/10.1142/s0219498826420065

© 2026 The Authors. Except as otherwise noted, this author-accepted version of a journal article published in Journal of Algebra and Its Applications is made available via the University of Sheffield Research Publications and Copyright Policy under the terms of the Creative Commons Attribution 4.0 International License (CC-BY 4.0), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/

Reuse
This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here: https://creativecommons.org/licenses/

Takedown
If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

UNIVERSITY OF LEEDS
University of Sheffield
UNIVERSITY of York

Ore sets, denominator sets and the left regular left quotient ring of a ring

V. V. Bavula

Dedicated to Professor T. Rizvi

Abstract

The aim of the papers is to describe the left regular left quotient ring $'Q(R)$ and the right regular right quotient ring $Q'(R)$ for the following algebras R: $\mathbb{S}_n = \mathbb{S}_1^{\otimes n}$ is the algebra of one-sided inverses, where $\mathbb{S}_1 = K\langle x, y \,|\, yx = 1\rangle$, $\mathcal{I}_n = K\langle \partial_1, \ldots, \partial_n, \int_1, \ldots, \int_n\rangle$ is the algebra of scalar integro-differential operators and the Jacobian algebra $\mathbb{A}_1 = K\langle x, \partial, (\partial x)^{-1}\rangle$. The sets of left and right regular elements of the algebras $\mathbb{S}_1$, $\mathcal{I}_1$, $\mathbb{A}_1$ and $\mathbb{I}_1 = K\langle x, \partial, \int\rangle$ are described. A progress is made on the following conjecture, [12]:

$$'Q(\mathbb{I}_n) \simeq Q(A_n) \text{ where } \mathbb{I}_n = K\Big\langle x_1, \ldots, x_n, \partial_1, \ldots, \partial_n, \int_1, \ldots, \int_n \Big\rangle$$

is the algebra of polynomial integro-differential operators and $Q(A_n)$ is the classical quotient ring (of fractions) of the n'th Weyl algebra A_n, i.e. a criterion is given when the isomorphism holds. We produce several general constructions of left Ore and left denominator sets that appear naturally in applications and are of independent interest and use them to produce explicit left denominator sets that give the localization ring isomorphic to $'Q(\mathbb{S}_n)$ or $'Q(\mathbb{I}_n)$ or $'Q(\mathbb{A}_n)$ where $\mathbb{A}_n := \mathbb{A}_1^{\otimes n}$. Several characterizations of one-sided regular elements of a ring are given in module-theoretic and one-sided-ideal-theoretic way.

Key Words: Ore set, denominator set, the Jacobian algebra, the Weyl algebra, the algebra of polynomial integro-differential operators, left regular element, the left regular left quotient ring of a ring, Goldie's Theorem.

Mathematics subject classification 2020: 16S85, 16P50, 16U20, 13B30, 16D30.

Contents

1 Introduction 1
2 Ore sets, denominator sets and left or right regular elements of a ring 4
3 The rings $'Q(\mathbb{S}_n)$, $'Q(\mathcal{I}_n)$, $Q'(\mathbb{S}_n)$ and $Q'(\mathcal{I}_n)$ 11
4 The rings $'Q(\mathbb{I}_n)$ and $Q'(\mathbb{I}_n)$ 18
5 The rings $'Q(\mathbb{A}_n)$ and $Q'(\mathbb{A}_n)$ 24

1 Introduction

In this paper, module means a *left* module. The following notation will remain fixed throughout the paper (if it is not stated otherwise):

- R is a unital ring and $R^\times$ be its group of units,
- $\mathcal{C} = \mathcal{C}_R$ is the set of *regular* elements of the ring R (i.e. $\mathcal{C}$ is the set of non-zero-divisors of the ring R);
- $'\mathcal{C}_R$ is the set of *left regular* elements of the ring R, i.e. $'\mathcal{C}_R := \{c \in R \,|\, \ker(\cdot c) = 0\}$ where $\cdot c : R \to R$, $r \mapsto rc$;

- $\mathcal{C}'_R := \{c \in R \,|\, \ker(c\cdot) = 0\}$ is the set of *right regular* elements of R;
- $Q = Q_{l,cl}(R) := \mathcal{C}_R^{-1}R$ is the *left quotient ring* (the *classical left ring of fractions*) of the ring R (if it exists, i.e. if $\mathcal{C}_R$ is a left Ore set) and $Q^\times$ is the group of units of Q;
- $\mathrm{Ore}_l(R) := \{S \,|\, S \text{ is a left Ore set in } R\}$;
- $\mathrm{Den}_l(R) := \{S \,|\, S \text{ is a left denominator set in } R\}$;
- $\mathrm{Ass}_l(R) := \{\mathrm{ass}(S) \,|\, S \in \mathrm{Den}_l(R)\}$ where $\mathrm{ass}(S) := \{r \in R \,|\, sr = 0 \text{ for some } s = s(r) \in S\}$;
- $\mathrm{Den}_l(R, \mathfrak{a})$ is the set of left denominator sets S of R with $\mathrm{ass}(S) = \mathfrak{a}$ where $\mathfrak{a}$ is an ideal of R;
- $S_\mathfrak{a} = S_\mathfrak{a}(R) = S_{l,\mathfrak{a}}(R)$ is the *largest element* of the poset $(\mathrm{Den}_l(R, \mathfrak{a}), \subseteq)$ and $Q_\mathfrak{a}(R) := Q_{l,\mathfrak{a}}(R) := S_\mathfrak{a}^{-1}R$ is the *largest left quotient ring associated with* $\mathfrak{a}$. The fact that $S_\mathfrak{a}$ exists is proven in [11, Theorem 2.1];
- In particular, $S_0 = S_0(R) = S_{l,0}(R)$ is the largest element of the poset $(\mathrm{Den}_l(R, 0), \subseteq)$, i.e. the *largest regular left Ore set* of R, and $Q_l(R) := S_0^{-1}R$ is the *largest left quotient ring* of R [11];
- $'S(R) = 'S_l(R)$ is the *largest left denominator set* in $'\mathcal{C}_R$ and $'Q(R) := 'Q_l(R) := 'S_l(R)^{-1}R$ is the *left regular left quotient ring* of R;
- $'\mathfrak{a} := \mathrm{ass}_R('S_l(R))$ and $'\pi : R \to \overline{R}' := R/'\mathfrak{a}$, $r \mapsto \overline{r} := r + '\mathfrak{a}$;
- $S'(R) = S'_r(R)$ is the *largest right denominator set* in $\mathcal{C}'_R$ and $Q'(R) := Q'_r(R) := RS'_r(R)^{-1}$ is the *right regular right quotient ring* of R.

Semisimplicity criteria for the ring $'Q_{l,cl}(R)$. For each element $r \in R$, let $r\cdot : R \to R$, $x \mapsto rx$ and $\cdot r : R \to R$, $x \mapsto xr$. The sets $'\mathcal{C}_R := \{r \in R \,|\, \ker(\cdot r) = 0\}$ and $\mathcal{C}'_R := \{r \in R \,|\, \ker(r\cdot) = 0\}$ are called the *sets of left and right regular elements* of R, respectively. Their intersection $\mathcal{C}_R = '\mathcal{C}_R \cap \mathcal{C}'_R$ is the *set of regular elements* of R. The rings $Q_{l,cl}(R) := \mathcal{C}_R^{-1}R$ and $Q_{r,cl}(R) := R\mathcal{C}_R^{-1}$ are called the *classical left and right quotient rings* of R, respectively. Goldie's Theorem states that the ring $Q_{l,cl}(R)$ is a semisimple Artinian ring iff the ring R is semiprime, $\mathrm{udim}(\mathbb{R}) < \infty$ and the ring R satisfies the a.c.c. on left annihilators (udim stands for the uniform dimension). In [9], four more new criteria are given based on different ideas, [9, Theorems 3.1, 4.1, 5.1, 6.2].

In [12], the rings $'Q_{l,cl}(R) := '\mathcal{C}_R^{-1}R$ (the *classical left regular left quotient ring* of R) and $Q'_{r,cl}(R) := R{\mathcal{C}'_R}^{-1}$ (the *classical right regular right quotient ring* of R) are introduced and studied, and several semisimplicity criteria for them are given ([12, Theorems 1.1, 3.1, 3.3, 3.4, 3.5]). The ring $'Q_{l,cl}(R)$ is a semisimple Artinian ring iff the ring $'Q_l(R)$ is so ([12, Theorem 4.3]).

A subset S of a ring R is called a *multiplicative set* if $1 \in S$, $SS \subseteq S$ and $0 \notin S$. Suppose that S and T are multiplicative sets in R such that $S \subseteq T$. The multiplicative subset S of T is called *dense* (or *left dense*) in T if for each element $t \in T$ there exists an element $r \in R$ such that $rt \in S$. For a left ideal I of R, let

$$'\mathcal{C}_I := \{i \in I \,|\, \cdot i : I \to I, \;\; x \mapsto xi \;\text{ is an injection}\}.$$

For a nonempty subset S of a ring R, let $\mathrm{ass}_R(S) := \{r \in R \,|\, sr = 0 \text{ for some } s \in S\}$. Let us mention a semisimplicity criteria for the ring $'Q_{l,cl}(R)$ that is used in the paper (see the proofs of Theorem 4.4 and Theorem 5.4).

Theorem 1.1. *([12, Theorems 1.1]) Let R be a ring, $'\mathcal{C} = '\mathcal{C}_R$ and $\mathfrak{a} := \mathrm{ass}_R('\mathcal{C})$. The following statements are equivalent.*

1. *$'Q := 'Q_{l,cl}(R)$ is a semisimple Artinian ring.*

2. (a) $\mathfrak{a}$ *is a semiprime ideal of* R,
   (b) *the set* $'\overline{\mathcal{C}} := \pi('\mathcal{C})$ *is a dense subset of* $'\mathcal{C}_{\overline{R}}$ *where* $\pi : R \to \overline{R} := R/\mathfrak{a}$, $r \mapsto \overline{r} := r + \mathfrak{a}$,
   (c) $\mathrm{udim}(_{\overline{R}}\overline{R}) < \infty$, *and*
   (d) $'\mathcal{C}_V \neq \emptyset$ *for all uniform left ideals* V *of* $\overline{R}$.

3. $\mathfrak{a}$ *is a semiprime ideal of* R, $'\overline{\mathcal{C}}$ *is a dense subset of* $\mathcal{C}_{\overline{R}}$ *and* $Q_{l,cl}(\overline{R})$ *is a semisimple Artinian ring.*

If one of the equivalent conditions holds then $'\overline{\mathcal{C}} \in \mathrm{Den}_l(\overline{R}, 0)$, $'\overline{\mathcal{C}}$ *is a dense subset of* $\mathcal{C}_{\overline{R}}$ *and* $'Q \simeq {'\overline{\mathcal{C}}}^{-1}\overline{R} \simeq Q_{l,cl}(\overline{R})$. *Furthermore, the ring* $'Q$ *is a simple ring iff the ideal* $\mathfrak{a}$ *is a prime ideal.*

The left regular left quotient ring $'Q_l(R)$ of a ring R and its semisimplicity criteria. Let R be a ring. In general, the classical left quotient ring $Q_{l,cl}(R)$ does not exists, i.e. the set of regular elements $\mathcal{C}_R$ of R is not a left Ore set. The set $\mathcal{C}_R$ contains the *largest left Ore set* denoted by $S_l(R)$ and the ring $Q_l(R) := S_l(R)^{-1}R$ is called the *(largest) left quotient ring* of R, [11]. Clearly, if $\mathcal{C}_R$ is a left Ore set then $\mathcal{C}_R = S_l(R)$ and $Q_{l,cl}(R) = Q_l(R)$. Similarly, the set $'\mathcal{C}_R$ of left regular elements of the ring R is not a left denominator set, in general, and so in this case the classical left regular left quotient ring $'Q_{l,cl}(R)$ does not exist. The set $'\mathcal{C}_R$ contains the *largest* left denominator set $'S_l(R)$ ([12, Lemma 4.1.(1)]) and the ring $'Q_l(R) := {'S_l(R)}^{-1}R$ is called the *left regular left quotient ring* of R, [12]. If $'\mathcal{C}_R$ is a left denominator set then $'\mathcal{C}_R = {'S_l(R)}$ and $'Q_{l,cl}(R) = {'Q_l(R)}$.

The main difficulty in constructing the rings $'Q(R)$ and $Q'(R)$ is to find descriptions of the sets $'S(R)$ and $S'(R)$. The main idea in constructing the rings $'Q(R)$ and $Q'(R)$ is to find larger or smaller or other denominator sets that give the *same* localization as the sets $'S(R)$ and $S'(R)$ do. In order to do so, we produce several constructions of Ore or denominator sets (that satisfy various conditions, appear naturally in applications and are of independent interest) and use them in the paper.

The paper is organized as follows. In Section 2, we present several results on and constructions of left Ore and denominator sets of a ring (Proposition 2.7) and give a sufficient condition for the sets $'\mathcal{C}_R^{le}$, ${\mathcal{C}'_R}^{re}$ and $\mathcal{C}_R^e$ being denominator sets (Proposition 2.10). Lemma 2.5 and Corollary 2.6 give equivalent conditions to the left Ore condition. Lemma 2.16 makes connection between the sets of right or left regular elements of a ring and sets of module monomorphisms. Lemma 2.17 is an application of Lemma 2.16 for one-sided ideals. For a module M and its submodule N, Lemma 2.18 makes connections between the sets $'\mathcal{C}_M$, $\mathcal{C}'_M$ and $\mathcal{C}_M$ and $'\mathcal{C}_N$, $\mathcal{C}'_N$ and $\mathcal{C}_N$, respectively. Corollary 2.19 and Corollary 2.20 are applications of the above result to one-sided essential ideals. These two corollaries are used in proofs.

In Section 3, the rings $'Q(\mathbb{S}_n)$ and $Q'(\mathbb{S}_n)$ are described (Theorem 3.11 and Corollary 3.15.(1)) where $\mathbb{S}_n := \mathbb{S}_1^{\otimes n}$ is the algebra of one-sided inverses and $\mathbb{S}_1 := K\langle x, y \,|\, yx = 1\rangle$. It is proven that $'\mathcal{C}_{\mathbb{S}_n} = {'S(\mathbb{S}_n)}$ and $'Q_{l,cl}(\mathbb{S}_n) = {'Q(\mathbb{S}_n)}$ (Corollary 3.13), $\mathcal{C}'_{\mathbb{S}_n} = S'(\mathbb{S}_n)$ and $Q'_{l,cl}(\mathbb{S}_n) = Q'(\mathbb{S}_n)$ (Corollary 3.15.(2)). The algebra $\mathbb{S}_n$ is a non-commutative, non-Noetherian, central, prime, catenary algebra; its ideals commute and satisfy the ascending chain condition; its classical Krull dimension is $2n$ but the weak and the global dimensions are n, [4]. The same results hold for the algebra of scalar integro-differential operators (K is a field of characteristic zero),

$$\mathcal{I}_n := K\Big\langle \partial_1, \dots, \partial_n, \int_1, \dots, \int_n \Big\rangle,$$

see Corollary 3.18.

Let K be a field of characteristic zero. The algebra $A_n = K\langle x_1, \dots, x_n, \partial_1, \dots, \partial_n\rangle$ is called the n'th Weyl algebra. It is canonically isomorphic to the algebra of polynomial differential operators ($\partial_i = \frac{\partial}{\partial x_i}$). The algebra A_n is a Noetheian domain. Hence, by Goldie's Theorem it (left and right) classical quotient ring $Q(A_n)$ is a division ring.

The algebra

$$\mathbb{I}_n := K\Big\langle x_1, \dots, x_n, \partial_1, \dots, \partial_n, \int_1, \dots, \int_n \Big\rangle$$

is called the algebra of polynomial integro-differential operators. The algebra $\mathbb{I}_n$ is a prime, central, catenary, non-Noetherian algebra of classical Krull dimension n and of Gelfand-Kirillov dimension $2n$, [5]. In [12], explicit descriptions of the sets ${}'\mathcal{C}_{\mathbb{I}_1}$ and $\mathcal{C}'_{\mathbb{I}_1}$ are given ([12, Theorem 6.7]). These descriptions are far from being trivial or obvious. It is also proven that

$$'Q_{l,cl}(\mathbb{I}_1) \simeq Q(A_1),$$

see [12, Theorem 6.5.(1)]. In [12], it is conjectured that

$$'Q_{l,cl}(\mathbb{I}_n) \simeq Q(A_n).$$

More information on the algebras of integro-differential operators (not necessarily polynomial) can be found in the following recent papers [1, 2, 17, 18, 25, 25, 26, 27].

In Section 4, we make progress on the conjecture. Namely, Theorem 4.6 is a criterion for the ring ${}'Q(\mathbb{I}_n)$ being isomorphic to quotient ring $Q(A_n)$. Despite the fact that there are no descriptions yet for the set ${}'\mathcal{C}_{\mathbb{I}_n}$ and $\mathcal{C}'_{\mathbb{I}_n}$ where $n \geq 2$, Theorem 4.4 provides explicit left denominators sets $S \in \mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$ such that $S^{-1}\mathbb{I}_n \simeq {}'Q(\mathbb{I}_n)$.

Definition 1.2. *([3]) The* **Jacobian algebra** $\mathbb{A}_n$ *is the subalgebra of* $\mathrm{End}_K(P_n)$ *generated by the Weyl algebra* A_n *and the elements* $H_1^{-1}, \dots, H_n^{-1} \in \mathrm{End}_K(P_n)$ *where*

$$H_1 := \partial_1 x_1, \dots, H_n := \partial_n x_n.$$

The algebra $\mathbb{I}_n$ properly contains the algebras A_n, $\mathbb{I}_n$ and $\mathcal{I}_n$.

In Section 5, a criterion is given for ${}'Q(\mathbb{A}_n) \simeq Q(A_n)$ (Theorem 5.4). As a corollary it is shown that

$$'Q(\mathbb{A}_1) \simeq Q(A_1) \text{ and } Q'(\mathbb{A}_1) \simeq Q(A_1),$$

see Theorem 5.5 and Corollary 5.6. The sets ${}'\mathcal{C}_{\mathbb{A}_1}$ and $\mathcal{C}'_{\mathbb{A}_1}$ are described (Theorem 5.7). There are no descriptions yet of the sets ${}'\mathcal{C}_{\mathbb{A}_n}$ and $\mathcal{C}'_{\mathbb{A}_n}$ for $n \geq 2$ but Theorem 5.4 provides explicit left denominators sets $S \in \mathrm{Den}_l(\mathbb{A}_n, \mathfrak{a}_n)$ such that $S^{-1}\mathbb{A}_n \simeq {}'Q(\mathbb{A}_n)$.

2 Ore sets, denominator sets and left or right regular elements of a ring

The aim of this section is to present several results on and constructions of left Ore and denominator sets of a ring. Several characterizations of one-sided regular elements of a ring are given in module-theoretic and one-sided-ideal-theoretic way. These results are used in the paper and are of independent interest.

Ore and denominator sets, localization of a ring at a denominator set. Let R be a ring. A subset S of R is called a *multiplicative set* if $SS \subseteq S$, $1 \in S$ and $0 \notin S$. A multiplicative subset S of R is called a *left Ore set* if it satisfies the *left Ore condition*: for each $r \in R$ and $s \in S$,

$$Sr \bigcap Rs \neq \emptyset.$$

Let $\mathrm{Ore}_l(R)$ be the set of all left Ore sets of R. For $S \in \mathrm{Ore}_l(R)$, $\mathrm{ass}_l(S) := \{r \in R \,|\, sr = 0 \text{ for some } s \in S\}$ is an ideal of the ring R.

A left Ore set S is called a *left denominator set* of the ring R if $rs = 0$ for some elements $r \in R$ and $s \in S$ implies $tr = 0$ for some element $t \in S$, i.e., $r \in \mathrm{ass}_l(S)$. Let $\mathrm{Den}_l(R)$ (resp., $\mathrm{Den}_l(R, \mathfrak{a})$) be the set of all left denominator sets of R (resp., such that $\mathrm{ass}_l(S) = \mathfrak{a}$). For $S \in \mathrm{Den}_l(R)$, let

$$S^{-1}R = \{s^{-1}r \,|\, s \in S, r \in R\}$$

be the *left localization* of the ring R at S (the *left quotient ring* of R at S). By definition, in Ore's method of localization one can localize *precisely* at the left denominator sets. In a similar way, right

Ore and right denominator sets are defined. Let $\mathrm{Ore}_r(R)$ and $\mathrm{Den}_r(R)$ be the set of all right Ore and right denominator sets of R, respectively. For $S \in \mathrm{Ore}_r(R)$, the set $\mathrm{ass}_r(S) := \{r \in R \,|\, rs = 0 \text{ for some } s \in S\}$ is an ideal of R. For $S \in \mathrm{Den}_r(R)$,

$$RS^{-1} = \{rs^{-1} \,|\, s \in S, r \in R\}$$

is the *right localization* of the ring R at S.

Given ring homomorphisms $\nu_A : R \to A$ and $\nu_B : R \to B$. A ring homomorphism $f : A \to B$ is called an *R-homomorphism* if $\nu_B = f\nu_A$. A left and right Ore set is called an *Ore set*. Let $\mathrm{Ore}(R)$ and $\mathrm{Den}(R)$ be the set of all Ore and denominator sets of R, respectively. For $S \in \mathrm{Den}(R)$,

$$S^{-1}R \simeq RS^{-1}$$

(an R-isomorphism) is the *localization* of the ring R at S, and $\mathrm{ass}(S) := \mathrm{ass}_l(S) = \mathrm{ass}_r(S)$.

The ring $R\langle S^{-1}\rangle$ and the ideal $\mathrm{ass}_R(S)$. Let R be a ring and S be a subset of R. Let $R\langle X_S\rangle$ be a ring freely generated by the ring R and a set $X_S = \{x_s \,|\, s \in S\}$ of free noncommutative indeterminates (indexed by the elements of the set S). Let I_S be the ideal of $R\langle X_S\rangle$ generated by the set $\{sx_s - 1, x_s s - 1 \,|\, s \in S\}$ and

$$R\langle S^{-1}\rangle := R\langle X_S\rangle / I_S. \tag{1}$$

The ring $R\langle S^{-1}\rangle$ is called the **localization of R at S**. Let $\mathrm{ass}(S) = \mathrm{ass}_R(S)$ be the kernel of the ring homomorphism

$$\sigma_S : R \to R\langle S^{-1}\rangle, \quad r \mapsto r + I_S. \tag{2}$$

The map $\pi_S : R \to \overline{R} := R/\mathrm{ass}_R(S)$, $r \mapsto \overline{r} := r + \mathrm{ass}_R(S)$ is an epimorphism. The ideal $\mathrm{ass}_R(S)$ of R has a complex structure, its description is given in [13, Proposition 2.12] when $R\langle S^{-1}\rangle = \{\overline{s}^{-1}\overline{r} \,|\, s \in S, r \in R\}$ is a ring of left fractions. We identify the factor ring $\overline{R}$ with its isomorphic copy in the ring $R\langle S^{-1}\rangle$ via the monomorphism

$$\overline{\sigma_S} : \overline{R} \to R\langle S^{-1}\rangle, \quad r + \mathrm{ass}_R(S) \mapsto r + I_S. \tag{3}$$

Clearly, $\overline{S} := (S + \mathrm{ass}_R(S))/\mathrm{ass}_R(S) \subseteq \mathcal{C}_{R\langle S^{-1}\rangle}$. [14, corollary 2.2] shows that the rings $R\langle S^{-1}\rangle$ and $\overline{R}\langle \overline{S}^{-1}\rangle$ are R-isomorphic. For $S = \emptyset$, $R\langle \emptyset^{-1}\rangle := R$ and $\mathrm{ass}_R(\emptyset) := 0$.

Definition 2.1. *A subset S of a ring R is called a* **localizable set** *of R if $R\langle S^{-1}\rangle \neq \{0\}$. Let $\mathrm{L}(R)$ be the set of localizable sets of R and*

$$\mathrm{ass}\,\mathrm{L}(R) := \{\mathrm{ass}_R(S) \,|\, S \in \mathrm{L}(R)\}. \tag{4}$$

For an ideal $\mathfrak{a}$ of R, let $\mathrm{L}(R, \mathfrak{a}) := \{S \in \mathrm{L}(R) \,|\, \mathrm{ass}_R(S) = \mathfrak{a}\}$. Then

$$\mathrm{L}(R) = \coprod_{\mathfrak{a} \in \mathrm{ass}\,\mathrm{L}(R)} \mathrm{L}(R, \mathfrak{a}) \tag{5}$$

is a disjoint union of non-empty sets. The set $(\mathrm{L}(R), \subseteq)$ is a partially ordered set (poset) w.r.t. inclusion $\subseteq$, and $(\mathrm{L}(R, \mathfrak{a}), \subseteq)$ is a sub-poset of $(\mathrm{L}(R), \subseteq)$ for every $\mathfrak{a} \in \mathrm{ass}\,\mathrm{L}(R)$.

Proposition 2.2 is the *universal property of localization*.

Proposition 2.2. *Let R be a ring, $S \in \mathrm{L}(R)$, and $\sigma_S : R \to R\langle S^{-1}\rangle$, $r \mapsto r + \mathrm{ass}_R(S)$. Let $f : R \to A$ be a ring homomorphism such that $f(S) \subseteq A^\times$. Then there is a unique R-homomorphism $f' : R\langle S^{-1}\rangle \to A$ such $f = f'\sigma_S$, i.e. the diagram below is commutative*

$$\begin{array}{ccc} R & \overset{\sigma_S}{\to} & R\langle S^{-1}\rangle \\ & \overset{f}{\searrow} & \downarrow{\scriptstyle \exists!\, f'} \\ & & A \end{array}$$

Every Ore set is a localizable set. Let S be an Ore set of the ring R. Theorem 2.3 states that every Ore set is localizable, gives an explicit description of the ideal $\mathrm{ass}_R(S)$ and the ring $R\langle S^{-1}\rangle$. Theorem 2.3 also states that the ring $R\langle S^{-1}\rangle$ is R-isomorphic to the localization $\overline{S}^{-1}\overline{R}$ of the ring $\overline{R}$ at the denominator set $\overline{S}$ of $\overline{R}$.

Theorem 2.3. *Let R be a ring and $S \in \mathrm{Ore}(R)$.*

1. *[7, Theorem 4.15] Every Ore set is a localizable set.*
2. *[13, Theorem 1.6.(1)] $\mathfrak{a} := \{r \in R \,|\, srt = 0$ for some elements $s, t \in S\}$ is an ideal of R such that $\mathfrak{a} \neq R$.*
3. *[13, Theorem 1.6.(2)] Let $\pi : R \to \overline{R} := R/\mathfrak{a}$, $r \mapsto \overline{r} = r + \mathfrak{a}$. Then $\overline{S} := \pi(S) \in \mathrm{Den}(\overline{R}, 0)$, $\mathfrak{a} = \mathfrak{a}(S) = \mathrm{ass}_R(S)$, $S \in \mathrm{L}(R, \mathfrak{a})$, and $S^{-1}R \simeq \overline{S}^{-1}\overline{R}$, an R-isomorphism. In particular, every Ore set is localizable.*

Equivalent conditions of the left Ore condition and applications. Let M be an R-module and $\mathbb{E}_M$ be the set of essential submodules of M.

Lemma 2.4. *Let M and M' be R-modules.*

1. *For all $f \in \mathrm{Hom}_R(M, M')$, $f^{-1}(\mathbb{E}_{M'}) := \{f^{-1}(L') \,|\, L' \in \mathbb{E}_{M'}\} \subseteq \mathbb{E}_M$.*
2. *If $M \subseteq M'$ then $\mathbb{E}_M = \{M \cap L' \,|\, L' \in \mathbb{E}_{M'}\}$.*

Proof. 1. Suppose that $f^{-1}(L') \notin \mathbb{E}_M$ for some $L' \in \mathbb{E}_{M'}$. Then we can choose a nonzero submodule, say N, of M such that $N \cap f^{-1}(L') \neq \{0\}$. Hence, $f(N) \neq \{0\}$ and $f(N) \cap L' = \{0\}$, a contradiction (since $L' \in \mathbb{E}_{M'}$).

2. By statement 1, $\{M \cap L' \,|\, L' \in \mathbb{E}_{M'}\} \subseteq \mathbb{E}_M$. Given $L \in \mathbb{E}_M$, we have to show that $L = M \cap L'$ for some $L' \in \mathbb{E}_{M'}$. Let C be the complement of the submodule $L \subseteq M'$. Then the direct sum $L' := L \oplus C$ is an essential submodule of M' such that $M \cap L' = L \oplus (M \cap C) = L$, as required (since L is an essential submodule of M). □

For a ring R, its left ideal L and an element $r \in R$, the set $(L : r) := \{r' \in R \,|\, r'r \in L\}$ is a left ideal of R.

Lemma 2.6 gives equivalent conditions to the left Ore condition. They are used in constructions of (new large) classes of left Ore sets by strengthening some of them (Proposition 2.7).

Lemma 2.5. *Suppose that S be a multiplicative subset of a ring R and $\mathfrak{a} := \mathrm{ass}_l(S) := \{r \in R \,|\, sr = 0$ for some element $s \in S\}$. Then the following statements are equivalent:*

1. *$S \in \mathrm{Ore}_l(R)$*
2. *The set $\mathfrak{a}$ is an ideal of R, $\overline{S} := S + \mathfrak{a} \in \mathrm{Ore}_l(\overline{R})$ where $\overline{R} := R/\mathfrak{a}$.*
3. *The set $\mathfrak{a}$ is an ideal of R, $\overline{R}\overline{s}$ is an essential left ideal of $\overline{R}$ for all $\overline{s} \in \overline{S}$, and $\overline{S} \cap (\overline{R}\overline{s} : \overline{r}) \neq \emptyset$ for all $\overline{s} \in \overline{S}$ and $\overline{r} \in \overline{R}$. Furthermore, for all $\overline{r} \in \overline{R}$ and $\overline{s} \in \overline{S}$, the left ideal $(\overline{R}\overline{s} : r)$ is an essential left ideal of $\overline{R}$ (Lemma 2.4.(1)).*
4. *For all left ideals L of R such that $L \not\subseteq \mathfrak{a}$ and all $s \in S$, $L \cap Rs \neq \{0\}$, and $S \cap (Rs : r) \neq \emptyset$ for all $s \in S$ and $r \in R \backslash \mathfrak{a}$. Furthermore, for all $r \in R \backslash \mathfrak{a}$ and $s \in S$, the left ideal $(Rs : r)$ is an essential left ideal of R (Lemma 2.4.(1)).*

Proof. $(1 \Rightarrow 2)$ The implication is well-known (and easy to prove).

$(2 \Rightarrow 3)$ Suppose that $0 \neq \overline{r} \in \overline{R}$. Then $0 \notin \overline{S}\overline{r}$ (since $\overline{S} \subseteq \mathcal{C}_{\overline{R}}$). Hence the left Ore condition for $\overline{S} \in \mathrm{Ore}_l(\overline{R})$ implies that $\overline{S}\overline{r} \cap \overline{R}\overline{s} \neq \{0\}$ for all $\overline{s} \in \overline{S}$. Hence $\overline{R}\overline{r} \cap \overline{R}\overline{s} \neq \{0\}$ for all $\overline{s} \in \overline{S}$. Therefore, $\overline{R}\overline{s}$ is an essential left ideal of $\overline{R}$ for all $\overline{s} \in \overline{S}$ and $\overline{S} \cap (\overline{R}\overline{s} : \overline{r}) \neq \emptyset$ for all $0 \neq \overline{r} \in \overline{R}$ and $\overline{s} \in \overline{S}$.

If $\overline{r} = 0$ then $\overline{S} \cap (\overline{R}\overline{s} : 0) = \overline{S} \cap \overline{R} = \overline{S} \neq \emptyset$ for all $\overline{s} \in \overline{S}$.

$(1 \Rightarrow 4)$ Suppose that $r \in R\backslash\mathfrak{a}$. Then $0 \notin Sr$. Hence the left Ore condition for $S \in \mathrm{Ore}_l(R)$ implies that $Sr \cap Rs \neq \{0\}$ for all $s \in \overline{S}$. Hence, $Rr \cap Rs \neq \{0\}$ for all $s \in \overline{S}$. Therefore, for all left ideals L of R such that $L \not\subseteq \mathfrak{a}$ and all $s \in S$, $L \cap Rs \neq \{0\}$, and $S \cap (Rs : r) \neq \emptyset$ for all $r \in R\backslash\mathfrak{a}$.

$(4 \Rightarrow 1)$ If $r \in \mathfrak{a}$ then $s'r = 0$ for some element $s \in S$, and so $s'r = 0 = 0s$ for all elements $s \in S$.

If $r \notin \mathfrak{a}$ then $S \cap (Rs : r) \neq \emptyset$ for all $s \in S$, and so $s'r = r's$ for for some elements $s' \in S$ and $r' \in R$ (that depend on the pair (s, r)). It follows that $S \in \mathrm{Ore}_l(R)$. □

Corollary 2.6. *Suppose that S is a multiplicative subset of a ring R such that $S \subseteq \mathcal{C}_R$. Then $S \in \mathrm{Ore}_l(R)$ iff Rs is an essential left ideal of R for all $s \in S$, and $S \cap (Rs : r) \neq \emptyset$ for all $s \in S$ and $r \in R$. Furthermore, for all $r \in R$ and $s \in S$, the left ideal $(Rs : r)$ is an essential left ideal of R.*

Proof. The corollary follows from the equivalence $(1 \Leftrightarrow 3)$ of Lemma 2.5. □

We strengthen the second condition of Corollary 2.6 to obtain Proposition 2.7.

Proposition 2.7. *Suppose that S is a multiplicative subset of a ring R such that*

(A) For every $s \in S$, the left ideal Rs of R is essential, and

(B) For all essential left ideals L of R, $S \cap L \neq \emptyset$.

Then $S \in \mathrm{Ore}_l(R)$.

Proof. We have to show that the left Ore condition holds for the set S: $Sr \cap Rs \neq \emptyset$ for all $s \in S$ and $r \in R$. By the statement (A), Rs is an essential left ideal of R (Lemma 2.4.(1)). Then $(Rs : r)$ is also an essential left ideal of R. By the statement (B), $S \cap (Rs : r) \neq \emptyset$, i.e. $s'r = r's$ for some elements $s' \in S$ and $r' \in R$, as required. □

Recall that a ring R is called a **left Goldie ring** if R has finite left uniform dimension and satisfies the a.c.c. on left annihilators. The following example shows that Proposition 2.7 covers a lot of ground.

Example 2.8. *It is known that the conditions (A) and (B) of Proposition 2.7 hold for all semiprime left Goldie rings R and $S = \mathcal{C}_R$ (the statement (A) follows at once from the fact that the ring R has finite left uniform dimension and the statement (B) is [22, Proposition 2.3.5.(ii)]). By Proposition 2.7, $\mathcal{C}_R \in \mathrm{Den}_l(R, 0)$. This fact is the crucial step in the proof of Goldie's Theorem.*

Corollary 2.9. *Suppose that S is a multiplicative subset of a ring R such that the conditions (A) and (B) of Proposition 2.7 and their right analogues hold. Then:*

1. *$S \in \mathrm{Ore}(R)$.*
2. *$\mathfrak{a} := \{r \in R \,|\, srt = 0$ for some elements $s, t \in S\}$ is an ideal of R such that $\mathfrak{a} \neq R$. Let $\pi : R \to \overline{R} := R/\mathfrak{a}$, $r \mapsto \overline{r} = r + \mathfrak{a}$. Then $\overline{S} := \pi(S) \in \mathrm{Den}(\overline{R}, 0)$, $\mathfrak{a} = \mathfrak{a}(S) = \mathrm{ass}_R(S)$, $S \in \mathrm{L}(R, \mathfrak{a})$, and $R\langle S^{-1}\rangle \simeq \overline{S}^{-1}\overline{R}$, an R-isomorphism.*

Proof. 1. Statement 1 follows from Proposition 2.7.

2. Statement 2 follows from Theorem 2.3. □

The denominator sets $'\mathcal{C}_R^{le}$, ${\mathcal{C}'_R}^{re}$ and $\mathcal{C}_R^e$. For a ring R, let R^{le} and R^{re} be the sets that contain elements $r \in R$ such that the left ideal Rr and the ideal rR are essential, respectively. Let $R^e := R^{le} \cap R^{re}$. Each element $r \in R$ determines two maps $r\cdot : R \to R$, $r' \mapsto rr'$ and $\cdot r : R \to R$, $r' \mapsto r'r$. An element $r \in R$ is called a *left* (resp., *right*) *regular* if $\ker(\cdot r) = 0$ (resp., $\ker(r\cdot) = 0$). The sets of all left and right regular elements of the ring R are denoted by $'\mathcal{C}_R$ and $\mathcal{C}'_R$, respectively. Let $'\mathcal{C}_R^{le} := {}'\mathcal{C}_R \cap R^{le}$, ${\mathcal{C}'_R}^{re} := \mathcal{C}'_R \cap R^{re}$ and $\mathcal{C}_R^e := {}'\mathcal{C}_R^{le} \cap {\mathcal{C}'_R}^{re} = \mathcal{C}_R \cap R^e$.

Proposition 2.10. *1. The sets $'\mathcal{C}_R^{le}$, ${\mathcal{C}'_R}^{re}$ and $\mathcal{C}_R^e$ are multiplicative sets of R.*

2. *Suppose that the set* ${}'\mathcal{C}_R^{le}$ *meets all the essential left ideals of the ring* R. *Then* ${}'\mathcal{C}_R^{le} \in \mathrm{Den}_l(R)$.

3. *Suppose that the set* ${\mathcal{C}'_R}^{re}$ *meets all the essential right ideals of the ring* R. *Then* ${\mathcal{C}'_R}^{re} \in \mathrm{Den}_r(R)$.

4. *Suppose that the set* $\mathcal{C}_R^e$ *meets all the essential left ideals and essential right ideals of the ring* R. *Then* $\mathcal{C}_R^e \in \mathrm{Den}(R, 0)$.

Proof. 1. It suffices to show that the set ${}'\mathcal{C}_R^{le}$ is a multiplicative set since then by symmetry the set ${\mathcal{C}'_R}^{re}$ is also multiplicative. These two results imply that $\mathcal{C}_R^e$ is a multiplicative set.

Clearly, $1 \in {}'\mathcal{C}_R^{le}$. Let $s, t \in {}'\mathcal{C}_R^{le}$. Then $st \in {}'\mathcal{C}_R$. It remain to show that $st \in R^{le}$. The left ideal Rs of R is an essential ideal. Then the R-module Rst is an essential submodule of Rt (since $t \in {}'\mathcal{C}$). The inclusions of essential left ideals of R, $Rst \subseteq Rt \subseteq R$, imply that the left ideal Rst of R is essential, as required.

2. By statement 1, the set ${}'\mathcal{C}_R^{le}$ is a multiplicative set of R. Since ${}'\mathcal{C}_R^{le} \subseteq {}'\mathcal{C}_R$, it remains to show that ${}'\mathcal{C}_R^{le} \in \mathrm{Ore}_l(R)$. This follows from Proposition 2.7 (since the conditions (A) and (B) of Proposition 2.7 are satisfied).

3. Statement 3 follows from statement 2 when apply to the opposite ring of the ring R.

4. Statement 4 follows from statements 2 and 3. □

For two multiplicative sets S and T of R, let ST be a multiplicative submonoid of $(R, \cdot)$ generated by S and T. Clearly, the product is commutative, $ST = TS$, associative and $\{1\}S = S$. The product ST is a multiplicative set iff $0 \notin ST$.

Lemma 2.11. *([8, Lemma 2.4])*

1. *Let* $S, T \in \mathrm{Ore}_l(R)$. *If* $0 \notin ST$ *then* $ST \in \mathrm{Ore}_l(R)$.

2. *Let* $S, T \in \mathrm{Den}_l(R)$. *If* $0 \notin ST$ *then* $ST \in \mathrm{Den}_l(R)$.

3. *Statements 1 and 2 hold also for Ore sets and denominator sets, respectively.*

Lemma 2.12. *Let* S *be a multiplicative set of a ring* R. *Then:*

1. *The set* S *contains the largest left/right/left and right Ore set which is denoted by* $S^{lO}/S^{rO}/S^O$.

2. *The set* S *contains the largest left/right/left and right denominator set which is denoted by* $S^{ld}/S^{rd}/S^d$.

3. *In the set* ${}'\mathcal{C}_R$ *every left Ore set is a left denominator set, and vice versa.*

4. *In the set* $\mathcal{C}'_R$ *every right Ore set is a right denominator set, and vice versa.*

5. *In the set* $\mathcal{C}_R$ *every Ore set is a denominator set, and vice versa.*

Proof. 1. Statement 1 follows from Lemma 2.11.(1) and the largest left/right/left and right Ore set of S is a union of all left/right/left and right Ore sets in S.

2. Statement 2 follows from Lemma 2.11.(2) and the largest left/right/left and right denominator set of S is a union of all left/right/left and right denominator sets in S.

3-5. Statements 3-5 are obvious. □

In view of Proposition 2.10.(1) and Lemma 2.12, we have the following definitions.

Definition 2.13. *Let* ${}'\mathcal{C}_R^{leO}$ *be the largest left Ore/denominator set in the multiplicative set* ${}'\mathcal{C}_R^{le}$. *Let* ${}'\mathcal{C}_R^{lee}$ *be the largest multiplicative subset of the multiplicative set* ${}'\mathcal{C}_R^{le}$ *that meets all the essential left ideals of* R. *Let* ${\mathcal{C}'_R}^{reO}$ *be the largest right Ore/denominator set in the multiplicative set* ${\mathcal{C}'_R}^{re}$. *Let* ${\mathcal{C}'_R}^{ree}$ *be the largest multiplicative subset of the multiplicative set* ${\mathcal{C}'_R}^{re}$ *that meets all the essential right ideals of* R.

Lemma 2.14 describes the sets ${}'\mathcal{C}_R^{leO}$, ${\mathcal{C}'_R}^{reO}$, ${}'\mathcal{C}_R^{lee}$ and ${\mathcal{C}'_R}^{ree}$.

Lemma 2.14. *1. The set ${}'\mathcal{C}_R^{leO}$ (resp., ${\mathcal{C}'_R}^{reO}$) is the union of all left (resp., right) Ore/denominator sets in ${}'\mathcal{C}_R^{leO}$ (resp., ${\mathcal{C}'_R}^{re}$). In particular, ${}'\mathcal{C}_R^{leO} \neq \emptyset$ and ${\mathcal{C}'_R}^{reO} \neq \emptyset$.*

2. ${}'\mathcal{C}_R^{lee} \in \{\emptyset, {}'\mathcal{C}_R^{le}\}$ and ${\mathcal{C}'_R}^{ree} \in \{\emptyset, {\mathcal{C}'_R}^{re}\}$. If ${}'\mathcal{C}_R^{lee} = {}'\mathcal{C}_R^{le}$ (resp., ${\mathcal{C}'_R}^{ree} = {\mathcal{C}'_R}^{re}$) then ${}'\mathcal{C}_R^{lee} = {}'\mathcal{C}_R^{le} \in \mathrm{Den}_l(R)$ (resp., ${\mathcal{C}'_R}^{ree} = {\mathcal{C}'_R}^{re} \in \mathrm{Den}_r(R)$).

Proof. 1. By Proposition 2.10.(1), the sets ${}'\mathcal{C}_R^{le}$ and ${\mathcal{C}'_R}^{re}$ are multiplicative sets of R. Now, statement 1 follows from Lemma 2.12.(1,2).

2. Suppose that ${}'\mathcal{C}_R^{lee} \neq \emptyset$. Notice that ${}'\mathcal{C}_R^{lee} \subseteq {}'\mathcal{C}_R^{le}$. Then clearly ${}'\mathcal{C}_R^{lee} = {}'\mathcal{C}_R^{le}$. Similarly, suppose that ${\mathcal{C}'_R}^{ree} \neq \emptyset$. Notice that ${\mathcal{C}'_R}^{ree} \subseteq {\mathcal{C}'_R}^{re}$. Then ${\mathcal{C}'_R}^{ree} = {\mathcal{C}'_R}^{re}$. The equalities in statements 2 follows from Proposition 2.10.(2,3). □

By Lemma 2.14, the sets ${}'\mathcal{C}_R^{leO}$ and ${}'\mathcal{C}_R^{lee}$ are left denominator sets of the ring R such that ${}'\mathcal{C}_R^{lee} \subseteq {}'\mathcal{C}_R^{leO}$ provided ${}'\mathcal{C}_R^{lee} \neq \emptyset$. By Lemma 2.14, the sets ${\mathcal{C}'_R}^{reO}$ and ${\mathcal{C}'_R}^{ree}$ are right denominator sets of the ring R such that ${\mathcal{C}'_R}^{ree} \subseteq {\mathcal{C}'_R}^{reO}$ provided ${}'\mathcal{C}_R^{ree} \neq \emptyset$.

Definition 2.15. *Let ${}'Q(R)^{leO} := \big({}'\mathcal{C}_R^{leO}\big)^{-1}R$, ${}'Q(R)^{lee} := \big({}'\mathcal{C}_R^{lee}\big)^{-1}R$, ${Q'}^{reO} := R\big({\mathcal{C}'_R}^{reO}\big)^{-1}$ and ${Q'}^{ree} := R\big({\mathcal{C}'_R}^{ree}\big)^{-1}$.*

Recall that ${}'S_l(R)$ is the largest left denominator set in ${}'\mathcal{C}_R$ and ${}'Q_l(R) := {}'S_l(R)^{-1}R$ is the left regular left quotient ring of R, ${}'\mathfrak{a} := \mathrm{ass}_R({}'S_l(R))$ and ${}'\pi : R \to \overline{R}' := R/{}'\mathfrak{a}$, $r \mapsto \overline{r} := r + {}'\mathfrak{a}$.

If ${}'\mathcal{C}_R^{lee} \neq \emptyset$ and ${\mathcal{C}'_R}^{ree} \neq \emptyset$ then, by Proposition 2.10.(2,3),

$$R^\times \subseteq {}'\mathcal{C}_R^{lee} = {}'\mathcal{C}_R^{le} \subseteq {}'\mathcal{C}_R^{leO} \subseteq {}'S_l(R) \subseteq {}'\mathcal{C}_R \text{ and } R^\times \subseteq {\mathcal{C}'_R}^{ree} = {\mathcal{C}'_R}^{re} \subseteq {\mathcal{C}'_R}^{reO} \subseteq S'_r(R) \subseteq \mathcal{C}'_R. \quad (6)$$

Hence there are R-homomorphisms:

$${}'Q(R)^{leO} \to {}'Q(R)^{lee}, \ \ s^1r \mapsto s^{-1}r \text{ and } {Q'}^{reO} \to {Q'}^{ree}, \ \ rt^{-1} \mapsto rt^{-1} \quad (7)$$

with kernels $\mathrm{ass}_l({}'\mathcal{C}_R^{leO})/\mathrm{ass}_l({}'\mathcal{C}_R^{lee})$ and $\mathrm{ass}_r({\mathcal{C}'_R}^{reO})/\mathrm{ass}_r({\mathcal{C}'_R}^{ree})$, respectively.

The left/right/two-sided regular sets of a ring R and monomorphism of R-modules. For a right R-module M_R, let ${}'\mathcal{C}_M := \{r \in R \,|\, \ker(\cdot r_M) = 0\}$ where $\cdot r_M : M \to M, \ \ m \to mr$. Similarly, for a left R-module ${}_RM$, let $\mathcal{C}'_M := \{r \in R \,|\, \ker(r_M \cdot) = 0\}$ where $r_M \cdot : M \to M, \ \ m \to rm$. For an R-bimodule M, let $\mathcal{C}_M := {}'\mathcal{C}_M \cap \mathcal{C}'_M$.

Lemma 2.16 makes connections between the sets ${}'\mathcal{C}_R$, $\mathcal{C}'_R$ and $\mathcal{C}_R$ and ${}'\mathcal{C}_M$, $\mathcal{C}'_M$ and $\mathcal{C}_M$, respectively (under faithfulness condition).

Lemma 2.16. *1. If M_R is a faithful right R-module then ${}'\mathcal{C}_M \subseteq {}'\mathcal{C}_R$.*

2. If ${}_RM$ is a faithful left R-module then $\mathcal{C}'_M \subseteq \mathcal{C}'_R$.

3. If ${}_RM_R$ is an R-module which is faithful as a left and right R-module then $\mathcal{C}_M \subseteq \mathcal{C}_R$.

4. If I is an ideal of the ring R which is faithful as a left and right R-module then $\mathcal{C}_I \subseteq \mathcal{C}_R$.

Proof. 1. Suppose that $c \in {}'\mathcal{C}_M$ but $c \notin {}'\mathcal{C}_R$. Then $dc = 0$ or some nonzero element $d \in R$. By the assumption, M_R is a faithful right R-module. Hence, $m' := md \neq 0$ for some element $m \in M$. Then $0 \neq m'c = mdc = 0$, a contradiction, and statement 1 follows.

2. By symmetry, statement 2 follows from statement 1.

3. Statement 3 follows from statements 1 and 2.

4. Statement 4 is a particular case of statement 3. □

Applying Lemma 2.16 for faithful (one-sided) ideals I of a ring R we obtain even tighter connections between the sets ${}'\mathcal{C}_R$, $\mathcal{C}'_R$ and $\mathcal{C}_R$ and ${}'\mathcal{C}_I$, $\mathcal{C}'_I$ and $\mathcal{C}_I$, respectively.

Lemma 2.17. *1. If I_R is a faithful right ideal of R then ${}'\mathcal{C}_R = {}'\mathcal{C}_I$.*

2. *If ${}_RI$ is a faithful left ideal of R then $\mathcal{C}'_R = \mathcal{C}'_I$.*

3. *If I is an ideal of R which is faithful as a left and right R-module then $\mathcal{C}_R = \mathcal{C}_I$.*

Proof. 1. By Lemma 2.16.(1), ${}'\mathcal{C}_R \supseteq {}'\mathcal{C}_I$. The opposite inclusion follows from the inclusion $I_R \subseteq R$.
2. By symmetry, statement 2 follows from statement 1.
3. Statement 3 follows from statements 1 and 2. □

For a module M and its submodule N, Lemma 2.18 makes connections between the sets ${}'\mathcal{C}_M$, $\mathcal{C}'_M$ and $\mathcal{C}_M$ and ${}'\mathcal{C}_N$, $\mathcal{C}'_N$ and $\mathcal{C}_N$, respectively (under essentiality condition).

Lemma 2.18. *1. If M_R is a right R-module and N_R is an essential submodule of M then ${}'\mathcal{C}_M = {}'\mathcal{C}_N$.*

2. *If ${}_RM$ is a left R-module and ${}_RN$ is an essential submodule of M then $\mathcal{C}'_M = \mathcal{C}'_N$.*

3. *If ${}_RM_R$ is an R-bimodule and ${}_RN_R$ is an R-sub-bimodule such that ${}_RN$ and N_R are essential submodules of ${}_RM$ and M_R, respectively. Then $\mathcal{C}_M = \mathcal{C}_N$.*

Proof. 1.
Suppose that $c \in {}'\mathcal{C}_M$ but $c \notin {}'\mathcal{C}_R$. Then $dc = 0$ or some nonzero element $d \in R$. By the assumption, M_R is a faithful right R-module. Hence, $m' := md \neq 0$ for some element $m \in M$. Then $0 \neq m'c = mdc = 0$, a contradiction, and statement 1 follows.
2. By symmetry, statement 2 follows from statement 1.
3. Statement 3 follows from statements 1 and 2. □

Corollary 2.19 is a particular case of Lemma 2.18.

Corollary 2.19. *1. If I_R is an essential right ideal of R then ${}'\mathcal{C}_R = {}'\mathcal{C}_I$.*

2. *If ${}_RI$ is an essential left ideal of R then $\mathcal{C}'_R = \mathcal{C}'_I$.*

3. *If I is an ideal of R which is essential as a left and right R-module then $\mathcal{C}_R = \mathcal{C}_I$.*

Proof. 1. By Lemma 2.16.(1), ${}'\mathcal{C}_R \supseteq {}'\mathcal{C}_I$. The opposite inclusion follows from the inclusion $I_R \subseteq R$.
2. By symmetry, statement 2 follows from statement 1.
3. Statement 3 follows from statements 1 and 2. □

Let $\mathrm{ICS}({}_RM)$ (resp., $\mathrm{ICS}(M_R)$) be the set of isomorphism classes of simple submodules of a semisimple left (resp. right) R-module M. Every semisimple left (resp. right) R-module M is a unigue direct sum of its isotypic components, $M = \oplus_{[V]\in \mathrm{ICS}({}_RM)} M_{[V]}$ (resp., $M = \oplus_{[U]\in \mathrm{ICS}(M_R)} M_{[U]}$), where $M_{[V]}$ (resp., $M_{U]}$) is the sum of all simple submodules of M that are isomorphic to the module V (resp., U).

Corollary 2.20. *1. If I_R is an essential right ideal of R which is a semisimple right R-module then ${}'\mathcal{C}_R = \bigcap_{[U]\in \mathrm{ICS}(I_R)} {}'\mathcal{C}_U$.*

2. *If ${}_RI$ is an essential left ideal of R which is a semisimple left R-module then $\mathcal{C}'_R = \bigcap_{[V]\in \mathrm{ICS}({}_RI)} \mathcal{C}'_V$.*

3. *If I is an ideal of R which is essential and semisimple as a left and right R-module and then $\mathcal{C}_R = \bigcap_{[U]\in \mathrm{ICS}(I_R),[V]\in \mathrm{ICS}({}_RI)} {}'\mathcal{C}_U \cap \mathcal{C}'_V$.*

Proof. 1 and 2. Statements 1 and 2 follows from Statements 1 and 2 of Corollary 2.19, respectively.
3. Statement 3 follows from statements 1 and 2. □

3 The rings $'Q(\mathbb{S}_n)$, $'Q(\mathcal{I}_n)$, $Q'(\mathbb{S}_n)$ and $Q'(\mathcal{I}_n)$

In this section, Theorem 3.11 and Corollary 3.15.(1) describe the algebras $'Q(\mathbb{S}_n)$ and $Q'(\mathbb{S}_n)$, respectively. Similarly, Corollary 3.18 describes the algebras $'Q(\mathcal{I}_n)$ and $Q'(\mathcal{I}_n)$. Theorem 3.16 and Theorem 3.17 describe the sets $'\mathcal{C}_{\mathbb{S}_1}$ and $\mathcal{C}'_{\mathbb{S}_1}$.

The algebra $\mathbb{S}_n$ of one-sided inverses of a polynomial algebra. We collect some results on the algebras $\mathbb{S}_n$ from [4] that are used in the proofs later.

Definition 3.1. *([4]) The* **algebra of one-sided inverses** *of $P_n = K[x_1, \dots, x_n]$, $\mathbb{S}_n$, is an algebra generated over a field K by $2n$ elements $x_1, \dots, x_n, y_n, \dots, y_n$ subject to the defining relations:*

$$y_1x_1 = \cdots = y_nx_n = 1, \;\; [x_i, y_j] = [x_i, x_j] = [y_i, y_j] = 0 \;\text{ for all } i \neq j,$$

where $[a, b] := ab - ba$, the commutator of elements a and b.

By the very definition, the algebra $\mathbb{S}_n$ is obtained from the polynomial algebra P_n by adding commuting, left (or right) inverses of its canonical generators. Clearly, $\mathbb{S}_n = \mathbb{S}_1(1) \otimes \cdots \otimes \mathbb{S}_1(n) \simeq \mathbb{S}_1^{\otimes n}$ where $\mathbb{S}_1(i) := K\langle x_i, y_i \,|\, y_ix_i = 1\rangle \simeq \mathbb{S}_1$ and

$$\mathbb{S}_n = \bigoplus_{\alpha,\beta\in\mathbb{N}^n} Kx^\alpha y^\beta$$

where $x^\alpha := x_1^{\alpha_1} \cdots x_n^{\alpha_n}$, $\alpha = (\alpha_1, \dots, \alpha_n)$, $y^\beta := y_1^{\beta_1} \cdots y_n^{\beta_n}$ and $\beta = (\beta_1, \dots, \beta_n)$. In particular, the algebra $\mathbb{S}_n$ contains two polynomial subalgebras $P_n = K[x_1, \dots, x_n]$ and $\mathbb{Y}_n := K[y_1, \dots, y_n]$ and is equal, as a vector space, to their tensor product $P_n \otimes \mathbb{Y}_n$. Note that also the Weyl algebra A_n is a tensor product (as a vector space) $P_n \otimes K[\partial_1, \dots, \partial_n]$ of two polynomial subalgebras.

When $n = 1$, we usually drop the subscript '1' if this does not lead to confusion. So, $\mathbb{S}_1 = K\langle x, y \,|\, yx = 1\rangle = \bigoplus_{i,j\geq 0} Kx^iy^j$. For each natural number $d \geq 1$, let $M_d(K) := \bigoplus_{i,j=0}^{d-1} KE_{ij}$ be the algebra of d-dimensional matrices where $\{E_{ij}\}$ are the matrix units, and

$$M_\infty(K) := \varinjlim M_d(K) = \bigoplus_{i,j\in\mathbb{N}} KE_{ij}$$

be the algebra (without 1) of infinite dimensional matrices. The algebra $\mathbb{S}_1$ contains the ideal $F := \bigoplus_{i,j\in\mathbb{N}} KE_{ij}$, where

$$E_{ij} := x^iy^j - x^{i+1}y^{j+1}, \;\; i, j \geq 0. \tag{8}$$

For all natural numbers i, j, k, and l, $E_{ij}E_{kl} = \delta_{jk}E_{il}$ where δ_{jk} is the Kronecker delta function. The ideal F is an algebra (without 1) isomorphic to the algebra $M_\infty(K)$ via $E_{ij} \mapsto E_{ij}$. For all $i, j \geq 0$,

$$xE_{ij} = E_{i+1,j}, \;\; yE_{ij} = E_{i-1,j} \quad (E_{-1,j} := 0), \tag{9}$$

$$E_{ij}x = E_{i,j-1}, \;\; E_{ij}y = E_{i,j+1} \quad (E_{i,-1} := 0). \tag{10}$$

The algebra

$$\mathbb{S}_1 = K \oplus xK[x] \oplus yK[y] \oplus F \tag{11}$$

is the direct sum of vector spaces. Then

$$\mathbb{S}_1/F \simeq L_1 := K[x, x^{-1}], \;\; x \mapsto x, \;\; y \mapsto x^{-1}, \tag{12}$$

since $yx = 1$, $xy = 1 - E_{00}$ and $E_{00} \in F$.

The algebra $\mathbb{S}_n = \bigotimes_{i=1}^n \mathbb{S}_1(i)$ contains the ideal

$$F_n := F^{\otimes n} = \bigoplus_{\alpha,\beta\in\mathbb{N}^n} KE_{\alpha\beta}, \;\; \text{where } \; E_{\alpha\beta} := \prod_{i=1}^n E_{\alpha_i\beta_i}(i), \;\; E_{\alpha_i\beta_i}(i) := x_i^{\alpha_i}y_i^{\beta_i} - x_i^{\alpha_i+1}y_i^{\beta_i+1}.$$

Note that $E_{\alpha\beta}E_{\gamma\rho} = \delta_{\beta\gamma}E_{\alpha\rho}$ for all elements $\alpha, \beta, \gamma, \rho \in \mathbb{N}^n$ where $\delta_{\beta\gamma}$ is the Kronecker delta function; $F_n = \bigotimes_{i=1}^n F(i)$ and $F(i) := \bigoplus_{s,t\in\mathbb{N}} KE_{st}(i)$.

The involution η on $\mathbb{S}_n$. The algebra $\mathbb{S}_n$ admits the *involution*

$$\eta : \mathbb{S}_n \to \mathbb{S}_n, \;\; x_i \mapsto y_i, \;\; y_i \mapsto x_i, \;\; i = 1, \dots, n. \tag{13}$$

It is a K-algebra anti-isomorphism ($\eta(ab) = \eta(b)\eta(a)$ for all $a, b \in \mathbb{S}_n$) such that $\eta^2 = \mathrm{id}_{\mathbb{S}_n}$, the identity map on $\mathbb{S}_n$. So, the algebra $\mathbb{S}_n$ is *self-dual* (i.e. it is isomorphic to its opposite algebra, $\eta : \mathbb{S}_n \simeq \mathbb{S}_n^{op}$). The involution η acts on the 'matrix' ring F_n as the transposition,

$$\eta(E_{\alpha\beta}) = E_{\beta\alpha}. \tag{14}$$

The canonical generators x_i, y_j $(1 \leq i, j \leq n)$ determine the ascending filtration $\{\mathbb{S}_{n,\leq i}\}_{i\in\mathbb{N}}$ on the algebra $\mathbb{S}_n$ in the obvious way (i.e. by the total degree of the generators): $\mathbb{S}_{n,\leq i} := \bigoplus_{|\alpha|+|\beta|\leq i} Kx^\alpha y^\beta$ where $|\alpha| = \alpha_1 + \cdots + \alpha_n$ ($\mathbb{S}_{n,\leq i}\mathbb{S}_{n,\leq j} \subseteq \mathbb{S}_{n,\leq i+j}$ for all $i, j \geq 0$). Then $\dim(\mathbb{S}_{n,\leq i}) = \binom{i+2n}{2n}$ for $i \geq 0$, and so the Gelfand-Kirillov dimension $\mathrm{GK}(\mathbb{S}_n)$ of the algebra $\mathbb{S}_n$ is equal to $2n$. It is not difficult to show that the algebra $\mathbb{S}_n$ is neither left nor right Noetherian. Moreover, it contains infinite direct sums of left and right ideals (see [4]).

The set of height 1 primes of $\mathbb{S}_n$. Consider the ideals of the algebra $\mathbb{S}_n$:

$$\mathfrak{p}_1 := F \otimes \mathbb{S}_{n-1}, \;\; \mathfrak{p}_2 := \mathbb{S}_1 \otimes F \otimes \mathbb{S}_{n-2}, \dots, \mathfrak{p}_n := \mathbb{S}_{n-1} \otimes F.$$

Then $\mathbb{S}_n/\mathfrak{p}_i \simeq \mathbb{S}_{n-1} \otimes (\mathbb{S}_1/F) \simeq \mathbb{S}_{n-1} \otimes K[x_i, x_i^{-1}]$ and $\bigcap_{i=1}^n \mathfrak{p}_i = \prod_{i=1}^n \mathfrak{p}_i = F^{\otimes n} = F_n$. Clearly, $\mathfrak{p}_i \not\subseteq \mathfrak{p}_j$ for all $i \neq j$.

- *The set $\mathcal{H}_1$ of height one prime ideals of the algebra $\mathbb{S}_n$ is $\{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$.*

Let $\mathfrak{a}_n := \mathfrak{p}_1 + \cdots + \mathfrak{p}_n$. Then the factor algebra

$$\mathbb{S}_n/\mathfrak{a}_n \simeq (\mathbb{S}_1/F)^{\otimes n} \simeq L_n := \bigotimes_{i=1}^n K[x_i, x_i^{-1}] = K[x_1, x_1^{-1}, \dots, x_n, x_n^{-1}] \tag{15}$$

is a Laurent polynomial algebra in n variables, and so $\mathfrak{a}_n$ is a prime ideal of height and co-height n of the algebra $\mathbb{S}_n$.

$$S_y := \{y^\alpha \,|\, \alpha \in \mathbb{N}^n\} \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n) \;\text{ an }\; S_y^{-1}\mathbb{S}_n \simeq \mathbb{S}_n/\mathfrak{a}_n = L_n.$$

The proof of the following statements can be found in [4].

- *The algebra $\mathbb{S}_n$ is central, prime and catenary. Every nonzero ideal of $\mathbb{S}_n$ is an essential left and right submodule of $\mathbb{S}_n$.*
- *The ideals of $\mathbb{S}_n$ commute ($IJ = JI$); and the set of ideals of $\mathbb{S}_n$ satisfy the a.c.c..*
- *The classical Krull dimension* $\mathrm{cl.Kdim}(\mathbb{S}_n)$ *of $\mathbb{S}_n$ is $2n$.*
- *Let I be an ideal of $\mathbb{S}_n$. Then the factor algebra $\mathbb{S}_n/I$ is left (or right) Noetherian iff $\mathfrak{a}_n \subseteq I$.*

Proposition 3.2. [4, Corollary 2.2] *The polynomial algebra P_n is the only (up to isomorphism) simple faithful left $\mathbb{S}_n$-module.*

In more detail, ${}_{\mathbb{S}_n}P_n \simeq \mathbb{S}_n/(\sum_{i=0}^n \mathbb{S}_n y_i) = \bigoplus_{\alpha\in\mathbb{N}^n} Kx^\alpha\overline{1}$, $\overline{1} := 1 + \sum_{i=1}^n \mathbb{S}_n y_i$; and the action of the canonical generators of the algebra $\mathbb{S}_n$ on the polynomial algebra P_n is given by the rule:

$$x_i * x^\alpha = x^{\alpha+e_i}, \;\; y_i * x^\alpha = \begin{cases} x^{\alpha-e_i} & \text{if } \alpha_i > 0, \\ 0 & \text{if } \alpha_i = 0, \end{cases} \;\text{ and }\; E_{\beta\gamma} * x^\alpha = \delta_{\gamma\alpha}x^\beta, \tag{16}$$

where the set $e_1 := (1, 0, \dots, 0), \dots, e_n := (0, \dots, 0, 1)$ is the canonical basis for the free $\mathbb{Z}$-module $\mathbb{Z}^n$. We identify the algebra $\mathbb{S}_n$ with its image in the algebra $\mathrm{End}_K(P_n)$ of all the K-linear maps from the vector space P_n to itself, i.e. $\mathbb{S}_n \subset \mathrm{End}_K(P_n)$.

Corollary 3.3. *The polynomial algebra $P'_n := \eta(P_n) = K[y_1, \dots, y_n]$ is the only (up to isomorphism) simple faithful right $\mathbb{S}_n$-module.*

Proof. In view of the involution η, the corollary follows from Proposition 3.2. □

In more detail, $(P'_n)_{\mathbb{S}_n} \simeq \mathbb{S}_n/(\sum_{i=0}^n x_i\mathbb{S}_n) = \bigoplus_{\alpha\in\mathbb{N}^n} \widetilde{1}Ky^\alpha$, $\widetilde{1} := 1 + \sum_{i=1}^n x_i\mathbb{S}_n$; and the action of the canonical generators of the algebra $\mathbb{S}_n$ on the polynomial algebra P'_n is given by the rule:

$$y^\alpha * y_i = y^{\alpha+e_i}, \;\; y^\alpha * x_i = \begin{cases} y^{\alpha-e_i} & \text{if } \alpha_i > 0, \\ 0 & \text{if } \alpha_i = 0, \end{cases} \text{ and } y^\alpha * E_{\beta\gamma} = \delta_{\alpha\beta}y^\gamma. \tag{17}$$

Proposition 3.4. *([4, Proposition 3])*

1. ${}_{\mathbb{S}_n}F_n \simeq P_n^{(\mathbb{N}^n)}$.
2. $(F_n)_{\mathbb{S}_n} \simeq (P'_n)^{(\mathbb{N}^n)}$.

Constructions of Ore and denominator sets. Below we collect and prove some useful results that are used in the proofs. They also are of independent interest.

Lemma 3.5. *([12, Lemma 2.5]) Suppose that $S, T \in \mathrm{Den}_l(R)$ and $S \subseteq T$. Then the map $\varphi : S^{-1}R \to S^{-1}T$, $s^{-1}r \mapsto s^{-1}r$, is a ring homomorphism (where $s \in S$ and $r \in R$).*

1. *φ is a monomorphism iff $\mathrm{ass}_R(S) = \mathrm{ass}_R(T)$.*
2. *φ is a epimorphism iff for each $t \in T$ there exists an element $r \in R$ such that $rt \in S + \mathrm{ass}_R(T)$.*
3. *φ is a isomorphism iff $\mathrm{ass}_R(S) = \mathrm{ass}_R(T)$ and for each element $t \in T$ there exists an element $r \in R$ such that $rt \in S$.*
4. *If, in addition, $T \subseteq {}'\mathcal{C}_R$, then φ is a isomorphism iff $\mathrm{ass}_R(S) = \mathrm{ass}_R(T)$ and for each element $t \in T$ there exists an element $r \in {}'\mathcal{C}_R$ such that $rt \in S$.*

Lemma 3.6. *([12, Lemma 6.1]) Suppose that $T \in \mathrm{Den}_l(R)$ and S be a multiplicative set of R such that $S \subseteq T$, $\mathrm{ass}_R(S) = \mathrm{ass}_R(T)$ and for each element $t \in T$ there exists an element $r \in R$ such that $rt \in S + \mathrm{ass}_R(T)$. Then $S \in \mathrm{Den}_l(R)$ and $S^{-1}R \simeq T^{-1}R$.*

Definition 3.7. *The pair (S, T) that satisfies the conditions of Lemma 3.6 is called a* **left localization pair** *of the ring R.*

Lemma 3.8 provides sufficient conditions for the pre-image of a left Ore/denominator set being a left Ore/denominator set.

Lemma 3.8. *Let R be a ring, $\mathfrak{a}$ be an ideal of R, $\pi_\mathfrak{a} : R \to \overline{R} := R/\mathfrak{a}$, $r \mapsto \overline{r} = r + \mathfrak{a}$, $\overline{\mathfrak{b}}$ be an ideal of $\overline{R}$, $\mathfrak{b} = \pi_\mathfrak{a}^{-1}(\overline{\mathfrak{b}})$, $\overline{S}$ be a multiplicative subset of $\overline{R}$ and $S = \pi_\mathfrak{a}^{-1}(\overline{S})$.*

1. *If $\overline{S} \in \mathrm{Ore}_l(\overline{R}, \overline{\mathfrak{b}})$ and $\mathrm{ass}_l(S) = \mathfrak{b}$ then $S \in \mathrm{Ore}_l(R, \mathfrak{b})$.*
2. *If $\overline{S} \in \mathrm{Den}_l(\overline{R}, \overline{\mathfrak{b}})$ and $\mathrm{ass}_l(S) = \mathfrak{b}$ then $S \in \mathrm{Den}_l(R, \mathfrak{b})$.*
3. *If $\overline{S} \in \mathrm{Den}(\overline{R}, \overline{\mathfrak{b}})$ and $\mathrm{ass}_l(S) = \mathrm{ass}_r(S) = \mathfrak{b}$ then $S \in \mathrm{Den}(R, \mathfrak{b})$.*
4. *If $\overline{S} \in \mathrm{Ore}_l(\overline{R}, \overline{\mathfrak{b}})$ and $T \subseteq S$ for some $T \in \mathrm{Ore}_l(R, \mathfrak{b})$ then $S \in \mathrm{Ore}_l(R, \mathfrak{b})$.*
5. *If $\overline{S} \in \mathrm{Den}_l(\overline{R}, \overline{\mathfrak{b}})$ and $T \subseteq S$ for some $T \in \mathrm{Ore}_l(R, \mathfrak{b})$ then $S \in \mathrm{Den}_l(R, \mathfrak{b})$.*

Proof. Clearly, the set S is a multiplicative subset in R.

1. We have to show that the left Ore condition holds for the multiplicative subset S of the ring R. For elements $s \in S$ and $r \in R$, $\overline{s'}\overline{r} = \overline{r'}\overline{s}$ for some elements $s' \in S$ and $r' \in R$ (since $\overline{S} \in \mathrm{Ore}_l(\overline{R}, \overline{\mathfrak{b}})$). Therefore, $d := s'r - r's \in \mathfrak{b}$. By the assumption $\mathrm{ass}_l(S) = \mathfrak{b}$. So, there is an element $s'' \in S$, such that $0 = s''d = s''s'r - s''r's$, i.e. the left Ore condition holds for S.

2. By statement 1, $S \in \mathrm{Ore}_l(R, \mathfrak{b})$. It remains to show that $\mathrm{ass}_r(S) \subseteq \mathrm{ass}_l(S) = \mathfrak{b}$. Suppose that $r \in \mathrm{ass}_r(S)$, i.e. $rs = 0$ for some element $s \in S$. Then $\overline{rs} = 0$. It follows that $\overline{r} \in \overline{\mathfrak{b}}$ (since $\overline{S} \in \mathrm{Den}_l(\overline{R}, \overline{\mathfrak{b}})$), and so $r \in \pi_{\mathfrak{a}}^{-1}(\overline{\mathfrak{b}}) = \mathfrak{b} = \mathrm{ass}_l(S)$. Hence, $\mathrm{ass}_r(S) \subseteq \mathrm{ass}_l(S) = \mathfrak{b}$.

3. Statement 3 follows from statement 2 and its right analogue.

4. Since $\overline{S} \in \mathrm{Ore}_l(\overline{R}, \overline{\mathfrak{b}})$, $\mathrm{ass}_l(S) \subseteq \mathfrak{b}$. Since $T \subseteq S$ and $T \in \mathrm{Ore}_l(R, \mathfrak{b})$, $\mathfrak{b} = \mathrm{ass}_l(T) \subseteq \mathrm{ass}_l(S)$. Hence, $\mathrm{ass}_l(S) = \mathfrak{b}$. Now, $S \in \mathrm{Ore}_l(R, \mathfrak{b})$, by statement 1.

5. By statement 4, $S \in \mathrm{Ore}_l(R, \mathfrak{b})$. It remains to show that $\mathrm{ass}_r(S) \subseteq \mathrm{ass}_l(S) = \mathfrak{b}$. Suppose that $r \in \mathrm{ass}_r(S)$, i.e. $rs = 0$ for some element $s \in S$. Then $\overline{rs} = 0$ and so $\overline{r} \in \overline{\mathfrak{b}}$ (since $\overline{S} \in \mathrm{Den}_l(\overline{R}, \overline{\mathfrak{b}})$), and so $r \in \pi_{\mathfrak{a}}^{-1}(\overline{\mathfrak{b}}) = \mathfrak{b} = \mathrm{ass}_l(S)$. Therefore, $\mathrm{ass}_r(S) \subseteq \mathrm{ass}_l(S) = \mathfrak{b}$. □

Now, we obtain a useful corollary.

Corollary 3.9. *Let R be a ring, $\mathfrak{a}$ be an ideal of R and $\pi_{\mathfrak{a}} : R \to \overline{R} := R/\mathfrak{a}$, $r \mapsto \overline{r} = r + \mathfrak{a}$.*

1. *If $S \in \mathrm{Ore}_l(R, \mathfrak{a})$ then $S + \mathfrak{a} \in \mathrm{Ore}_l(R, \mathfrak{a})$.*
2. *If $S \in \mathrm{Den}_l(R, \mathfrak{a})$ then $S + \mathfrak{a} \in \mathrm{Den}_l(R, \mathfrak{a})$ and $(S + \mathfrak{a})^{-1}R \simeq S^{-1}R$.*
3. *If $S \in \mathrm{Den}(R, \mathfrak{a})$ then $S + \mathfrak{a} \in \mathrm{Den}(R, \mathfrak{a})$ and $(S + \mathfrak{a})^{-1}R \simeq S^{-1}R$.*

Proof. 1 and 2. We keep the notation of Lemma 3.8. Suppose that $S \in \mathrm{Ore}_l(R, \mathfrak{a})/\mathrm{Den}_l(R, \mathfrak{a})$. Then $S' := S + \mathfrak{a} = \pi_{\mathfrak{a}}^{-1}(S)$ is a multiplicative subset of R that contains S. Therefore, $\mathfrak{a} = \mathrm{ass}_l(S) \subseteq \mathrm{ass}_l(S')$. Let $r \in \mathrm{ass}_l(S')$. Then $(s + a)r = 0$ for some elements $s \in S$ and $a \in \mathfrak{a}$, and so $sr = -ar \in \mathfrak{a}$. There is an element $t \in S$ such that $tsr = 0$, i.e. $r \in \mathfrak{a}$. Therefore, $\mathrm{ass}_l(S) = \mathfrak{a} = \mathrm{ass}_l(S')$. Clearly, $\overline{S} := \pi_{\mathfrak{a}}(S) \in \mathrm{Ore}_l(\overline{R}, 0)/\mathrm{Den}_l(\overline{R}, 0)$. Now, by Lemma 3.8.(1,2), $S' \in \mathrm{Ore}_l(R, \mathfrak{a})/\mathrm{Den}_l(R, \mathfrak{a})$. If $S' \in \mathrm{Den}_l(R, \mathfrak{a})$ then ${S'}^{-1}R \simeq \pi_{\mathfrak{a}}(S')^{-1}\overline{R} = \pi_{\mathfrak{a}}(S)^{-1}\overline{R} \simeq S^{-1}R$.

3. Statement 3 follows from statement 2 and its right analogue. □

Lemma 3.10 provides sufficient conditions for a left Ore/denominator of a larger ring being a left Ore/denominator set of a smaller ring which contains it.

Lemma 3.10. *Let R be a subring of a ring R' and S be a multiplicative subset of R such that the left R-module R'/R is S-torsion (for each $r' \in R'$ there is an element $s \in S$ such that $sr' \in R$).*

1. *If $S \in \mathrm{Ore}_l(R', \mathfrak{a}')$ then $S \in \mathrm{Ore}_l(R, R \cap \mathfrak{a}')$.*
2. *If $S \in \mathrm{Den}_l(R', \mathfrak{a}')$ then $S \in \mathrm{Den}_l(R, R \cap \mathfrak{a}')$ and $S^{-1}R \simeq S^{-1}R'$.*

Proof. 1. (i) $S \in \mathrm{Ore}_l(R)$: For each element $s \in S$ and $r \in R$, $s'r = r's$ for some elements $s' \in S$ and $r' \in R'$ (since $S \in \mathrm{Ore}_l(R', \mathfrak{a}')$). By the assumption, the left R-module R'/R is S-torsion, and so $tr' \in R$ for some element $t \in S$. Now, $ts'r = tr's$ where $ts' \in S$ and $tr' \in R$, and the $S \in \mathrm{Ore}_l(R)$.

(ii) $\mathrm{ass}_{l,R}(S) = R \cap \mathfrak{a}'$: Since $R \subseteq R'$ and $S \in \mathrm{Ore}_l(R', \mathfrak{a}')$, $\mathrm{ass}_{l,R}(S) = R \cap \mathrm{ass}_{l,R'}(S) = R \cap \mathfrak{a}'$.

2. (i) $S \in \mathrm{Den}_l(R, R \cap \mathfrak{a}')$: By statement 1, $S \in \mathrm{Ore}_l(R, R \cap \mathfrak{a}')$. It remain to show that $\mathrm{ass}_r(S) \subseteq R \cap \mathfrak{a}'$. Given an element $r \in R$ such that $rs = 0$ for some $s \in S$. Then $r \in \mathfrak{a}'$ (since $S \in \mathrm{Den}_l(R', \mathfrak{a}')$), and so $r \in R \cap \mathfrak{a}'$, as required.

(ii) $S^{-1}R \simeq S^{-1}R'$: By the statement (i), the map $\phi : S^{-1}R \to S^{-1}R'$, $s^{-1}r \mapsto s^{-1}r$ is a ring homomorphism. Suppose that $a := s^{-1}r \in \ker(\phi)$. Then $r \in \mathfrak{a}'$, and so $r \in R \cap \mathfrak{a}'$, i.e. $a = 0$. So, the map ϕ is a monomorphism. It remains to show that the map ϕ is an epimorphism. Given element $s^{-1}r' \in S^{-1}R'$ where $s \in S$ and $r' \in R'$. Since the R-module R'/R is S-torsion, there is an element $t \in S$ such that $r := tr' \in R$. Then $s^{-1}r' = (ts)^{-1}tr' = (ts)^{-1}r \in S^{-1}R$. Therefore, the map ϕ is epimorphism and the statement (ii) follows. □

The ring $'Q(\mathbb{S}_n)$. Recall that $\mathbb{Y}_n = K[y_1, \ldots, y_n]$ is a polynomial algebra. Let $\mathbb{Y}_n^0 := \mathbb{Y}_n \backslash \{0\}$, $'\mathbb{Y}_n^0 := \mathbb{Y}_n \cap {}'\mathcal{C}_{\mathbb{S}_n}$,

$$'\mathbb{Y}_n := \mathbb{Y}_n \cap {}'S(\mathbb{S}_n) \text{ and } \widetilde{'\mathbb{Y}_n} := \{c \in \mathbb{Y}_n \,|\, y^\alpha c \in {}'S(\mathbb{S}_n) \text{ for some } \alpha \in \mathbb{N}^n\},$$

$\mathcal{T} = \pi_{\mathfrak{a}_n}^{-1}(L_n \backslash \{0\}) = \mathbb{S}_n \backslash \mathfrak{a}_n$ and $\mathcal{S} := \pi_{\mathfrak{a}_n}^{-1}(\mathbb{Y}_n \backslash \{0\}) = \mathbb{Y}_n \backslash \{0\} + \mathfrak{a}_n$ where $\pi_{\mathfrak{a}_n} : \mathbb{S}_n \to \mathbb{S}_n/\mathfrak{a}_n = L_n$, $r \mapsto \overline{r} := r + \mathfrak{a}_n$. Clearly, $'\mathbb{Y}_n \subseteq {}'\mathbb{Y}_n^0 \subseteq \mathbb{Y}_n^0$ and $\mathcal{S} \subseteq \mathcal{T}$.

Theorem 3.11 describes the algebra $'Q(\mathbb{S}_n)$.

Theorem 3.11. *1.* $'Q(\mathbb{S}_n) \simeq K(y_1, \ldots, y_n)$.

2. $'\mathbb{Y}_n \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ *and* $'\mathbb{Y}_n^{-1}\mathbb{S}_n \simeq {}'Q(\mathbb{S}_n)$. *Furthermore, the subset* $'\mathbb{Y}_n$ *of* $\mathbb{Y}_n$ *is a left denominator set of* $\mathbb{S}_n$ *which is the largest left denominator set that is contained in the multiplicative set* $\mathbb{Y}_n \cap {}'\mathcal{C}_{\mathbb{S}_n}$.

3. $\mathcal{S}, \mathcal{T} \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ *and* $'\mathbb{Y}_n^{-1}\mathbb{S}_n \simeq \mathcal{T}^{-1}\mathbb{S}_n \simeq \mathcal{S}^{-1}\mathbb{S}_n \simeq K(y_1, \ldots, y_n)$.

Proof. 1. Statement 1 follows from statements 2 and 3.

2. (i) *The pair* $('\mathbb{Y}_n, {}'S(\mathbb{S}_n))$ *is a left localization pair of the ring* $\mathbb{S}_n$: By the definition, the set $'\mathbb{Y}_n$ is a multiplicative subset of $'S(\mathbb{S}_n) \subseteq \mathbb{S}_n$. It follows from the inclusions $S_y \subseteq {}'\mathbb{Y}_n \subseteq {}'S(\mathbb{S}_n)$ that

$$\mathfrak{a}_n = \mathrm{ass}_l(S_y) \subseteq \mathrm{ass}_l('\mathbb{Y}_n) \subseteq \mathrm{ass}_l('S(\mathbb{S}_n)) = \mathfrak{a}_n,$$

and so $\mathrm{ass}_l('\mathbb{Y}_n) = \mathfrak{a}_n = \mathrm{ass}_l('S(\mathbb{S}_n))$. For each element $s \in {}'S(\mathbb{S}_n)$, there is an element $\alpha \in \mathbb{N}^n$ such that $y^\alpha s \in {}'\mathbb{Y}_n$. Notice that $y^\alpha \in S_y \subseteq {}'\mathbb{Y}_n$, and the statement (i) follows.

(ii) $'\mathbb{Y}_n \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ *and* $'\mathbb{Y}_n^{-1}\mathbb{S}_n \simeq {}'Q(\mathbb{S}_n)$: The statement (i) follows from the statement (i) and Lemma 3.6 where $T = {}'S(\mathbb{S}_n) \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ and $S = {}'\mathbb{Y}_n$.

(iii) *The subset* $'\mathbb{Y}_n$ *of* $\mathbb{S}_n$ *is a left denominator set of* $\mathbb{S}_n$ *which is the largest left denominator set that is contained in the multiplicative set* $\mathbb{Y}_n \cap {}'\mathcal{C}_{\mathbb{S}_n}$: Let T be a left denominator set of $\mathbb{S}_n$ which is the largest left denominator set that is contained in the multiplicative set $\mathbb{Y}_n \cap {}'\mathcal{C}_{\mathbb{S}_n}$. By the statement (ii), $'\mathbb{Y}_n \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$. Clearly,

$$'\mathbb{Y}_n = \mathbb{Y}_n \cap {}'S(\mathbb{S}_n) \subseteq \mathbb{Y}_n \cap {}'\mathcal{C}_{\mathbb{S}_n},$$

and so $'\mathbb{Y}_n \subseteq T$. Since $T \subseteq {}'\mathcal{C}_{\mathbb{S}_n}$ and $'S(\mathbb{S}_n)$ is a the largest left denominator set in ${}'\mathcal{C}_{\mathbb{S}_n}$, we have the inclusion $T \subseteq \mathbb{Y}_n \cap {}'S(\mathbb{S}_n) = {}'\mathbb{Y}_n$. Therefore, $T = {}'\mathbb{Y}_n$.

3. (i) $\mathcal{T} \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ *and* $\mathcal{T}^{-1}\mathbb{S}_n \simeq K(y_1, \ldots, y_n)$: Since $S_y \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ and $S_y \subseteq \mathcal{T}$, $\mathfrak{a}_n = \mathrm{ass}_l(S_y) \subseteq \mathrm{ass}_l(\mathcal{T})$. The opposite inclusion follows from the fact that the factor ring $\mathbb{S}_n/\mathfrak{a}_n \simeq K[y_1^{\pm 1}, \ldots, y_n^{\pm 1}]$ is a domain. So, $\mathrm{ass}_l(\mathcal{T}) = \mathfrak{a}_n$. Notice that

$$\pi_{\mathfrak{a}_n}(\mathcal{T}) = K[y_1^{\pm 1}, \ldots, y_n^{\pm 1}] \backslash \{0\} \in \mathrm{Den}_l(K[y_1^{\pm 1}, \ldots, y_n^{\pm 1}], 0).$$

By Lemma 3.9.(2), $\mathcal{T} \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ and

$$\mathcal{T}^{-1}\mathbb{S}_n \simeq \pi_{\mathfrak{a}_n}(\mathcal{T})^{-1}(\mathbb{S}_n/\mathfrak{a}_n) \simeq \pi_{\mathfrak{a}_n}(\mathcal{T})^{-1}(K[y_1^{\pm 1}, \ldots, y_n^{\pm 1}]) \simeq K(y_1, \ldots, y_n).$$

(ii) *The pair* $('\mathbb{Y}_n, \mathcal{T})$ *is a left localization pair of the ring* $\mathbb{S}_n$: By statement 2, $'\mathbb{Y}_n \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$. For each element $t \in \mathcal{T}$, there is an element $\alpha \in \mathbb{N}^n$ such that $y^\alpha t \in {}'\mathbb{Y}_n$. Notice that $y^\alpha \in S_y \subseteq {}'\mathbb{Y}_n$, and the statement (ii) follows.

(iii) $'\mathbb{Y}_n^{-1}\mathbb{S}_n \simeq \mathcal{T}^{-1}\mathbb{S}_n$: The statement (iii) follows from the statement (ii) and Lemma 3.6 where $T = \mathcal{T} \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ and $S = {}'\mathbb{Y}_n$.

(iv) *The pair* $(\mathcal{S}, \mathcal{T})$ *is a left localization pair of the ring* $\mathbb{S}_n$: By the statement (i), $\mathcal{T} \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$. By the definition, the set $\mathcal{S}$ is a multiplicative subset of $\mathbb{S}_n$. The inclusions $S_y \subseteq \mathcal{S} \subseteq \mathcal{T}$ imply that

$$\mathfrak{a}_n = \mathrm{ass}_l(S_y) \subseteq \mathrm{ass}_l(\mathcal{S}) \subseteq \mathrm{ass}_l(\mathcal{T}) = \mathfrak{a}_n,$$

and so $\mathrm{ass}_l(\mathcal{S}) = \mathfrak{a}_n = \mathrm{ass}_l(\mathcal{T})$. For each element $t \in \mathcal{T}$, there is an element $\alpha \in \mathbb{N}^n$ such that $y^\alpha t \in \mathcal{S}$. Notice that $y^\alpha \in S_y \subseteq \mathcal{S}$, and the statement (iv) follows.

(v) $\mathcal{S}^{-1}\mathbb{S}_n \simeq \mathcal{T}^{-1}\mathbb{S}_n$: The statement (v) follows from the statement (iv) and Lemma 3.6 where $T = \mathcal{T} \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ and $S = \mathcal{S}$.

Now, statement 3 follows. □

Theorem 3.12. *([12, Theorem 4.3]) Let R be a ring. Then*

1. *$'Q_l(R)$ is a left Artinian ring iff $'Q_{l,cl}(R)$ is a left Artinian ring. If one of the equivalent conditions holds then $'S_l(R) = {}'\mathcal{C}_R$ and $'Q_l(R) = 'Q_{l,cl}(R)$.*
2. *$'Q_l(R)$ is a semisimple Artinian ring iff $'Q_{l,cl}(R)$ is a semisimple Artinian ring. If one of the equivalent conditions holds then $'S_l(R) = {}'\mathcal{C}_R$ and $'Q_l(R) = 'Q_{l,cl}(R)$.*

Corollary 3.13. *$'\mathcal{C}_{\mathbb{S}_n} = 'S(\mathbb{S}_n)$ and $'Q_{l,cl}(\mathbb{S}_n) = 'Q(\mathbb{S}_n) \simeq K(y_1, \dots, y_n)$.*

Proof. By Theorem 3.11, the ring $'Q(\mathbb{S}_n) \simeq K(y_1, \dots, y_n)$ is a field. In particular, it is a semisimple Artinian ring. Now, the corollary follows from Theorem 3.12.(2). □

Theorem 3.11 and Corollary 3.14 produce explicit left denominators sets $S \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ such that $S^{-1}\mathbb{I}_n \simeq 'Q(\mathbb{S}_n)$. By Theorem 3.11 and Corollary 3.14, there are inclusions in the set $\mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ apart from S_y:

$$S_y \subseteq {}'\mathbb{Y}_n \subseteq {}'S(\mathbb{S}_n) \subseteq {}'S(\mathbb{S}_n) + \mathfrak{a}_n \subseteq \mathcal{T},\ {}'\mathbb{Y}_n \subseteq \widetilde{{}'\mathbb{Y}_n} \subseteq {}'\mathbb{Y}_n + \mathfrak{a}_n \subseteq {}'S(\mathbb{Y}_n) + \mathfrak{a}_n,\ {}'\mathbb{Y}_n + \mathfrak{a}_n \subseteq \mathcal{S} \subseteq \mathcal{T}. \tag{18}$$

Corollary 3.14. *Let $S = \widetilde{{}'\mathbb{Y}_n}, {}'\mathbb{Y}_n + \mathfrak{a}_n, {}'S(\mathbb{Y}_n) + \mathfrak{a}_n$. Then $S \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ an $S^{-1}\mathbb{S}_n \simeq {}'Q(\mathbb{S}_n) \simeq K(y_1, \dots, y_n)$.*

Proof. By Theorem 3.11.(2) and Corollary 3.9.(2), ${}'\mathbb{Y}_n + \mathfrak{a}_n, {}'S(\mathbb{Y}_n) + \mathfrak{a}_n \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$. Now, for $S = {}'\mathbb{Y}_n + \mathfrak{a}_n, {}'S(\mathbb{Y}_n) + \mathfrak{a}_n$, the corollary follows from Corollary 3.9.(2). It remains to consider the case when $S = \widetilde{{}'\mathbb{Y}_n}$.

(i) *The pair $(\widetilde{{}'\mathbb{Y}_n}, {}'\mathbb{Y}_n + \mathfrak{a}_n)$ is a left localization pair of the ring $\mathbb{S}_n$*: The inclusions ${}'\mathbb{Y}_n \subseteq \widetilde{{}'\mathbb{Y}_n} \subseteq {}'\mathbb{Y}_n + \mathfrak{a}_n$ yield

$$\mathfrak{a}_n = \mathrm{ass}_l({}'\mathbb{Y}_n) \subseteq \mathrm{ass}_l(\widetilde{{}'\mathbb{Y}_n}) \subseteq \mathrm{ass}_l({}'\mathbb{Y}_n + \mathfrak{a}_n) = \mathfrak{a}_n.$$

Therefore, $\mathrm{ass}_l(\widetilde{{}'\mathbb{Y}_n}) = \mathfrak{a}_n = \mathrm{ass}_l({}'\mathbb{Y}_n + \mathfrak{a}_n)$. For each element $s \in {}'\mathbb{Y}_n + \mathfrak{a}_n$, there is an element $\alpha \in \mathbb{N}^n$ such that $y^\alpha s \in {}'\mathbb{Y}_n \subseteq \widetilde{{}'\mathbb{Y}_n}$. Notice that $y^\alpha \in S_y \subseteq {}'\mathbb{Y}_n \subseteq \widetilde{{}'\mathbb{Y}_n}$, and the statement (i) follows.

(ii) $\widetilde{{}'\mathbb{Y}_n} \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$ and $\widetilde{{}'\mathbb{Y}_n}^{-1}\mathbb{S}_n \simeq ({}'\mathbb{Y}_n + \mathfrak{a}_n)^{-1}\mathbb{S}_n \simeq K(y_1, \dots, y_n)$: The statement (ii) follows from the statement (i) and Lemma 3.6 where $S = \widetilde{{}'\mathbb{Y}_n}$ and $T = {}'\mathbb{Y}_n + \mathfrak{a}_n \in \mathrm{Den}_l(\mathbb{S}_n, \mathfrak{a}_n)$. □

Corollary 3.15. *1. $Q'(\mathbb{S}_n) \simeq \eta('Q(\mathbb{S}_n)) = K(x_1, \dots, x_n)$.*

2. $\mathcal{C}'_{\mathbb{S}_n} = S'(\mathbb{S}_n)$ and $Q'_{l,cl}(\mathbb{S}_n) = Q'(\mathbb{S}_n) \simeq K(x_1, \dots, x_n)$.

Proof. 1. Since η is an involution of the algebra $\mathbb{S}_n$, $Q'(\mathbb{S}_n) \simeq \eta('Q(\mathbb{S}_n))$. Since $\eta(y_i) = x_i$ for all $i = 1, \dots, n$, $\eta('Q(\mathbb{S}_n)) = \eta(K(y_1, \dots, y_n) = K(x_1, \dots, x_n)$, by Theorem 3.11.(1).

2. By statement 1, the ring $Q'(\mathbb{S}_n) \simeq K(x_1, \dots, x_n)$ is a field. In particular, it is a semisimple Artinian ring, and so statement 2 follows from Theorem 3.12.(2). □

Descriptions of the sets $'\mathcal{C}_{\mathbb{S}_1}$ and $\mathcal{C}'_{\mathbb{S}_1}$.

(i) $'\mathcal{C}_{\mathbb{S}_1} \subseteq \mathbb{S}_1 \backslash F$: It is obvious that every element of the ideal $F = \oplus_{i,j\in\mathbb{N}} KE_{ij} \simeq M_\infty(K)$ is a left and right zero divisor of the algebra F (without 1) and of $\mathbb{S}_1$.

(ii) $\mathbb{Y}_1^0 \subseteq {}'\mathcal{C}_{\mathbb{S}_1}$: The ideal F is an essential right ideal of the algebra $\mathbb{S}_1$ such that $F_{\mathbb{S}_1} \simeq (P'_1)^{(\mathbb{N})}$ is a semisimple right $\mathbb{S}_1$-module (Proposition 3.4). By Corollary 2.20.(1), $'\mathcal{C}_{\mathbb{S}_1} = {}'\mathcal{C}_{P'_1}$ where $P'_1 = K[y]$ is the only simple faithful right $\mathbb{S}_1$-module. Now, the statement (ii) follows from (17).

(iii) *For each nonzero element $d \in \mathbb{S}_1 \backslash F$, $\partial^i d \in \mathbb{Y}_1^0$ for some $i \in \mathbb{N}$*: The statement (iii) follows (11).

(iv) *For each nonzero element $d \in \mathbb{S}_1 \backslash F$, $\partial^i d \in {}'\mathcal{C}_{\mathbb{S}_1}$ for some $i \in \mathbb{N}$*: The statement (iii) follows from the statements (ii) and (iii).

Then the well-defined map

$$d : \mathbb{S}_1\backslash F \rightarrow \mathbb{N}, \;\; a \mapsto d(a) := \min\{i \in \mathbb{N} \,|\, \partial^i a \in {}'\mathcal{C}_{\mathbb{S}_1}\} \tag{19}$$

is called the *left regularity degree function* and the natural number $d(a)$ is called the *left regularity degree* of a. For each element $a \in \mathbb{S}_1\backslash F$, $d(a)$ can be found in finitely many steps. Now, Theorem 3.16.(1) follows. Then Theorem 3.16.(2) follows from Theorem 3.16.(1).

Theorem 3.16. *1.* ${}'\mathcal{C}_{\mathbb{S}_1} = \{\partial^{d(a)} a \,|\, a \in \mathbb{S}_1\backslash F\}$.

2. $\mathcal{C}'_{\mathbb{S}_1} = \eta({}'\mathcal{C}_{\mathbb{S}_1})$.

By (11), each element $a \in \mathbb{S}_1$ is a unique sum $a = \sum_{i=0}^{l} \lambda_{-i} y^i + \sum_{j=1}^{m} \lambda_j x^j + a_F$ where $\lambda_k \in K$, $k = -l, \ldots, m$ and $a_F \in F$. Let $a_y := \sum_{i=0}^{l} \lambda_{-i} y^i$. The integer

$$s(a_F) := \begin{cases} \min\{n \in \mathbb{N} \,|\, a_F \in \bigoplus_{i,j=0}^{n} K e_{ij}\} & \text{if } a_F \neq 0, \\ -1 & \text{if } a_F = 0. \end{cases}$$

is called the *size* of the element $a_F \in F$. The integer $s(a) := s(a_F)$ is called the *size* of the element a. For each $i \in \mathbb{N}$, let $P'_{1,\leq i} := \{a \in P'_1 \,|\, \deg_y(a) \leq i\}$ where $\deg_y$ is the degree of the polynomial $a \in P'_1$ in the variable y.

Theorem 3.17. *1.* ${}'\mathcal{C}_{\mathbb{S}_1} = \{a \in \mathbb{S}_1\backslash(xK[x] + F) \,|\, \cdot a : P'_{1,\leq s(a)} \rightarrow P'_{1,\leq s(a)+\deg_y(a_y)}, \; p \mapsto pa$ *is an injection*$\}$.

2. $\mathcal{C}'_{\mathbb{S}_1} = \eta({}'\mathcal{C}_{\mathbb{S}_1})$ *where* η *is the involution of the algebra* $\mathbb{S}_1$*, see (13).*

Proof. 1. (i) ${}'\mathcal{C}_{\mathbb{S}_1} \cap (xK[x] + F) = \emptyset$: Suppose that $a \in xK[x]$. Then $1 \in \ker(\cdot a)$. Recall that ${}'\mathcal{C}_{\mathbb{S}_1} \subseteq \mathbb{S}_1\backslash F$ (see the statement (i) in the proof of Theorem 3.16). So, it remains to consider the case when $a \in (xK[x] + F)\backslash(xK[x] \cup F)$. Then the map $\cdot a : P'_{1,\leq s(a)+1} \rightarrow P'_{1,\leq s(a)}$, $p \mapsto pa$ is a well-defined map. Since

$$\dim_K(P'_{1,\leq s(a)+1}) = s(a) + 2 > s(a) + 1 = \dim_K(P'_{1,\leq s(a)}),$$

$\ker(\cdot a) \neq 0$. Therefore, ${}'\mathcal{C}_{\mathbb{S}_1} \cap (xK[x] + F) = \emptyset$.

(ii) *For each element* $a \in \mathbb{S}_1\backslash(xK[x] + F)$, $\ker_{P'_1}(\cdot a) \subseteq P'_{1,\leq s(a)}$: Since $a \in \mathbb{S}_1\backslash(xK[x] + F)$, $a_y := \sum_{i=0}^{l} \lambda_{-i} y^i \neq 0$ where $\lambda_{-i} \in K$. Suppose that $\lambda_{-l} \neq 0$. Suppose that $p \in \ker_{P'_1}(\cdot a)\backslash P'_{1,\leq s(a)}$, i.e. $\deg_y(p) > s(a)$. Then

$$\deg(pa) = l + \deg_y(p),$$

a contradiction (since $pa = 0$).

Now, statement 1 follows from statements (i) and (ii).

2. Statement 2 follows from statement 1. □

The algebras $\mathcal{I}_n$ of scalar integro-differential operators. In the next section, we will see that the algebra $\mathbb{I}_n$ of polynomial integro-differential operators contains the algebra of scalar integro-differential operators, [5]:

$$\mathcal{I}_n := K\Big\langle \partial_1, \ldots, \partial_n, \int_1, \ldots, \int_n \Big\rangle.$$

The algebra $\mathcal{I}_n$ is canonically isomorphic to the algebra $\mathbb{S}_n$, [5, Eq. (9)], see (20) for an explicit isomorphism.

Corollary 3.18. *1.* ${}'\mathcal{C}_{\mathcal{I}_n} = {}'S(\mathcal{I}_n)$ *and* ${}'Q_{l,cl}(\mathcal{I}_n) = {}'Q(\mathcal{I}_n) \simeq {}'Q(\mathbb{S}_n) \simeq K(y_1, \ldots, y_n)$.

2. $\mathcal{C}_{\mathcal{I}_n} = S'(\mathcal{I}_n)$ *and* $Q'_{l,cl}(\mathcal{I}_n) = Q'(\mathcal{I}_n) \simeq Q'(\mathbb{S}_n) \simeq K(x_1, \ldots, x_n)$.

Proof. 1. The algebras $\mathcal{I}_n$ and $\mathbb{S}_n$ are isomorphic (see (20)) and statement 1 follows from Theorem 3.11 and Corollary 3.13.

2. Statement 2 follows from Corollary 3.15. □

Corollary 3.19. ${}'\mathcal{C}_{\mathcal{I}_1} = \xi({}'\mathcal{C}_{\mathbb{S}_1})$ *and* $\mathcal{C}'_{\mathcal{I}_1} = \xi(\mathcal{C}'_{\mathbb{S}_1})$.

4 The rings $'Q(\mathbb{I}_n)$ and $Q'(\mathbb{I}_n)$

The aim of the section is to prove Theorem 4.4, to obtain a description of the sets $'\mathcal{C}_{\mathbb{I}_1}$ and $\mathcal{C}'_{\mathbb{I}_1}$ (Theorem 4.7), and to prove Theorem 4.6 which is a criterion for $'Q(\mathbb{I}_n) \simeq Q(I_n)$.

The rings $\mathbb{I}_n$ of integro-differential operators and the Jacobian algebras $\mathbb{A}_n$. In this section the following notation is fixed: K is a field of characteristic zero and K^* is its group of units; $P_n := K[x_1, \dots, x_n]$ is a polynomial algebra over K; $\partial_1 := \frac{\partial}{\partial x_1}, \dots, \partial_n := \frac{\partial}{\partial x_n}$ are the partial derivatives (K-linear derivations) of P_n; $\mathrm{End}_K(P_n)$ is the algebra of all K-linear maps from P_n to P_n; the subalgebra $A_n := K\langle x_1, \dots, x_n, \partial_1, \dots, \partial_n\rangle$ of $\mathrm{End}_K(P_n)$ is called the n'th **Weyl algebra**.

Definition 4.1. *([3]) The* **Jacobian algebra** $\mathbb{A}_n$ *is the subalgebra of* $\mathrm{End}_K(P_n)$ *generated by the Weyl algebra* A_n *and the elements* $H_1^{-1}, \dots, H_n^{-1} \in \mathrm{End}_K(P_n)$ *where*

$$H_1 := \partial_1 x_1, \dots, H_n := \partial_n x_n.$$

Clearly, $\mathbb{A}_n = \bigotimes_{i=1}^n \mathbb{A}_1(i) \simeq \mathbb{A}_1^{\otimes n}$ where $\mathbb{A}_1(i) := K\langle x_i, \partial_i, H_i^{-1}\rangle \simeq \mathbb{A}_1$. The algebra $\mathbb{A}_n$ contains all the integrations $\int_i : P_n \to P_n$, $p \mapsto \int p\, dx_i$, i.e.

$$\int_i = x_i H_i^{-1} : x^\alpha \mapsto (\alpha_i + 1)^{-1} x_i x^\alpha.$$

The algebra $\mathbb{A}_n$ contains the **algebra of polynomial integro-differential operators**, [5]:

$$\mathbb{I}_n := K\Big\langle x_1, \dots, x_n, \partial_1, \dots, \partial_n, \int_1, \dots, \int_n \Big\rangle.$$

Notice that $\mathbb{I}_n = \bigotimes_{i=1}^n \mathbb{I}_1(i) \simeq \mathbb{I}_1^{\otimes n}$ where $\mathbb{I}_1(i) := K\langle x_i, \partial_i, \int_i\rangle \simeq \mathbb{I}_1$. The algebra $\mathbb{I}_n$ contains the **algebra of scalar integro-differential operators**, [5]:

$$\mathcal{I}_n := K\Big\langle \partial_1, \dots, \partial_n, \int_1, \dots, \int_n \Big\rangle.$$

The algebra $\mathcal{I}_n$ is canonically isomorphic to the algebra $\mathbb{S}_n$, [5, Eq. (9)]:

$$\xi : \mathbb{S}_n \to \mathcal{I}_n, \quad x_i \mapsto \int_i, \quad y_i \mapsto \partial_i, \quad i = 1, \dots, n. \tag{20}$$

For the reader's convenience we collect some known results on the algebras $\mathbb{I}_n$ and $\mathbb{A}_n$ from the papers [3, 5] that are used later in the paper. The algebra $\mathbb{I}_n$ is a prime, central, catenary, non-Noetherian algebra of classical Krull dimension n and of Gelfand-Kirillov dimension $2n$, [5]. Since $x_i = \int_i H_i$, where $H_i := \partial_i x_i$, the algebra $\mathbb{I}_n$ is generated by the elements $\{\partial_i, H_i, \int_i \mid i = 1, \dots, n\}$, and $\mathbb{I}_n = \bigotimes_{i=1}^n \mathbb{I}_1(i)$ where

$$\mathbb{I}_1(i) := K\Big\langle \partial_i, H_i, \int_i \Big\rangle = K\Big\langle \partial_i, x_i, \int_i \Big\rangle \simeq \mathbb{I}_1.$$

When $n = 1$ we usually drop the subscript '1' in ∂_1, $\int_1$, H_1, and x_1. The algebra $\mathbb{I}_1 = \oplus_{i\in\mathbb{Z}} \mathbb{I}_{1,i}$ is a $\mathbb{Z}^n$-graded algebra where

$$\mathbb{I}_{1,i} = \begin{cases} \int^i D_1 & \text{if } i \geq 1, \\ D_1 & \text{if } i = 0, \\ D_1 \partial^{-i} & \text{if } i \leq -1, \end{cases} \tag{21}$$

where $D_1 := K[H] \oplus \bigoplus_{i\in\mathbb{N}} K e_{ii}$ is a commutative, not Noetherian, not finitely generated algebra and $K[H]$ is a polynomial algebra in the variable H and $H = \partial x = x\partial + 1$.

The following elements of the algebra $\mathbb{I}_1 = K\langle \partial, H, \int\rangle$,

$$e_{ij} := \int^i \partial^j - \int^{i+1} \partial^{j+1}, \quad i, j \in \mathbb{N}, \tag{22}$$

satisfy the relations: $e_{ij}e_{kl} = \delta_{jk}e_{il}$ where δ_{ij} is the Kronecker delta. The matrices of the linear maps $e_{ij} \in \mathrm{End}_K(K[x])$ with respect to the basis $\{x^{[s]} := \frac{x^s}{s!}\}_{s\in\mathbb{N}}$ of the polynomial algebra $K[x]$ are the elementary matrices, i.e.

$$e_{ij} * x^{[s]} = \begin{cases} x^{[i]} & \text{if } j = s, \\ 0 & \text{if } j \neq s. \end{cases}$$

The direct sum $F := \bigoplus_{i,j\in\mathbb{N}} Ke_{ij}$ is the only proper (hence maximal) ideal of the algebra $\mathbb{I}_1$. As an algebra without 1 it is isomorphic to the algebra without 1 of infinite dimensional matrices $M_\infty(K) := \varinjlim M_d(K) = \bigoplus_{i,j\in\mathbb{N}} KE_{ij}$ via $e_{ij} \mapsto E_{ij}$ where E_{ij} are the matrix units. For all $i, j \in \mathbb{N}$,

$$\int e_{ij} = e_{i+1,j}, \;\; e_{ij}\int = e_{i,j-1}, \;\; \partial e_{ij} = e_{i-1,j}, \;\; e_{ij}\partial = e_{i,j+1}, \tag{23}$$

where $e_{-1,j} := 0$ and $e_{i,-1} := 0$.

$$\mathbb{I}_1 = \bigoplus_{i\geq 1} K[H]\partial^i \oplus K[H] \oplus \bigoplus_{i\in\geq 1} K[H]\int^i \oplus F \tag{24}$$

and $K[H]\partial^i = \partial^i K[H]$ and $K[H]\int^i = \int^i K[H]$ for all $i \geq 1$. The algebra $\mathbb{I}_1$ is generated by the elements ∂, $\int$ and H subject to the following defining relations (Proposition 2.2, [5]):

$$\partial\int = 1, \;\; \Big[H, \int\Big] = \int, \;\; [H, \partial] = -\partial, \;\; H\Big(1 - \int\partial\Big) = \Big(1 - \int\partial\Big)H = 1 - \int\partial.$$

The algebra $\mathbb{I}_n = \otimes_{i=1}^n \mathbb{I}_1(i) = \oplus_{\alpha\in\mathbb{Z}^n}\mathbb{I}_{n,\alpha}$ is a $\mathbb{Z}^n$-graded algebra where $\mathbb{I}_{n,\alpha} := \otimes_{k=1}^n \mathbb{I}_{1,\alpha_k}(k)$ for $\alpha = (\alpha_1, \ldots, \alpha_n)$. The algebra $\mathbb{I}_n$ contains the ideal

$$F_n := F^{\otimes n} = \bigotimes_{i=1}^n F(i) = \bigoplus_{\alpha,\beta\in\mathbb{N}^n} Ke_{\alpha\beta}$$

where $e_{\alpha\beta} := \prod_{i=1}^n e_{\alpha_i\beta_i}(i)$, $e_{\alpha_i\beta_i}(i) := \int_i^{\alpha_i}\partial_i^{\beta_i} - \int_i^{\alpha_i+1}\partial_i^{\beta_i+1}$ and $F(i) = \bigoplus_{s,t\in\mathbb{N}} Ke_{st}(i)$.

Lemma 4.2. *1.* ([5, Corollary 3.3.(2)]) *The set of height one prime ideals of the algebra $\mathbb{I}_n$ is $\{\mathfrak{p}_1 := F \otimes \mathbb{I}_{n-1}, \mathfrak{p}_1 := \mathbb{I}_1 \otimes F \otimes \mathbb{I}_{n-2}, \ldots, \mathfrak{p}_n := \mathbb{I}_{n-1} \otimes F\}$.*

2. ([5, Corollary 3.3.(3)]) *Each ideal of the algebra $\mathbb{I}_n$ is an idempotent ideal ($\mathfrak{a}^2 = \mathfrak{a}$).*

3. ([5, Lemma 5.2.(2)]) *Each nonzero ideal of the algebra $\mathbb{I}_n$ is an essential left and right submodule of $\mathbb{I}_n$.*

4. ([5, Corollary 3.3.(8)]) *The ideal $\mathfrak{a}_n := \mathfrak{p}_1 + \cdots + \mathfrak{p}_n$ is the largest (i.e. the only maximal) ideal of $\mathbb{I}_n$ and $F_n = F^{\otimes n} = \cap_{i=1}^n \mathfrak{p}_i$ is the smallest nonzero ideal of $\mathbb{I}_n$.*

5. ([5, Proposition 3.8]) *The polynomial algebra P_n is the only (up to isomorphism) faithful simple left $\mathbb{I}_n$-module and ${}_{\mathbb{I}_n}P_n \simeq \mathbb{I}_n/\mathbb{I}_n(\partial_1, \ldots, \partial_n)$* ([6, Proposition 3.4.(3)])*.*

6. ([5, Lemma 5.2.(1)]) *For all nonzero ideals $\mathfrak{a}$ of the algebra $\mathbb{I}_n$,* $\mathrm{l.ann}_{\mathbb{I}_n}(\mathfrak{a}) = \mathrm{r.ann}_{\mathbb{I}_n}(\mathfrak{a}) = 0$.

7. ([5, Lemma 5.2.(2)]) *Each nonzero ideal of the algebra $\mathbb{I}_n$ is an essential left and right submodule of $\mathbb{I}_n$.*

The involution $*$ on the algebra $\mathbb{I}_n$. The algebra $\mathbb{I}_n$ admits the involution:

$$* : \mathbb{I}_n \to \mathbb{I}_n, \;\; \partial_i \mapsto \int_i, \;\; \int_i \mapsto \partial_i, \;\; H_i \mapsto H_i, \;\; i = 1, \ldots, n, \tag{25}$$

i.e. it is a K-algebra *anti-isomorphism* ($(ab)^* = b^*a^*$) such that $* \circ * = \mathrm{id}_{\mathbb{I}_n}$. Therefore, the algebra $\mathbb{I}_n$ is *self-dual*, i.e. is isomorphic to its *opposite* algebra $\mathbb{I}_n^{op}$. As a result, the left and the right properties of the algebra $\mathbb{I}_n$ are the same. For all elements $\alpha, \beta \in \mathbb{N}^n$,

$$e_{\alpha\beta}^* = e_{\beta\alpha}. \tag{26}$$

The involution $*$ can be extended to an involution of the algebra $\mathbb{A}_n$ by setting

$$x_i^* = H_i\partial_i, \;\; \partial_i^* = \int_i, \;\; (H_i^{\pm 1})^* = H_i^{\pm 1}, \;\; i = 1, \ldots, n.$$

Note that $y_i^* = (H_i^{-1}\partial_i)^* = \int_i H_i^{-1} = x_i H_i^{-2}$, $A_n^* \not\subseteq A_n$, but $\mathcal{I}_n^* = \mathcal{I}_n$ where

$$\mathcal{I}_n := K\Big\langle \partial_1, \ldots, \partial_n \int_1, \ldots, \int_n \Big\rangle$$

is the algebra of integro-differential operators with constant coefficients.

For a subset S of a ring R, the sets $\mathrm{l.ann}_R(S) := \{r \in R \,|\, rS = 0\}$ and $\mathrm{r.ann}_R(S) := \{r \in R \,|\, Sr = 0\}$ are called the *left* and the *right annihilators* of the set S in R. Using the fact that the algebra $\mathbb{I}_n$ is a GWA and its $\mathbb{Z}^n$-grading, we see that

$$\mathrm{l.ann}_{\mathbb{I}_n}\Big(\int_i\Big) = \bigoplus_{k\in\mathbb{N}} K e_{k0}(i) \bigotimes \bigotimes_{i\neq j} \mathbb{I}_1(j), \;\; \mathrm{r.ann}_{\mathbb{I}_n}\Big(\int_i\Big) = 0. \tag{27}$$

$$\mathrm{r.ann}_{\mathbb{I}_n}(\partial_i) = \bigoplus_{k\in\mathbb{N}} K e_{0k}(i) \bigotimes \bigotimes_{i\neq j} \mathbb{I}_1(j), \;\; \mathrm{l.ann}_{\mathbb{I}_n}(\partial_i) = 0. \tag{28}$$

Let $\mathfrak{a}$ be an ideal of the algebra $\mathbb{I}_n$. The factor algebra $\mathbb{I}_n/\mathfrak{a}$ is a Noetherian algebra iff $\mathfrak{a} = \mathfrak{a}_n$ (Proposition 4.1, [5]). The factor algebra $B_n := \mathbb{I}_n/\mathfrak{a}_n$ is isomorphic to the skew Laurent polynomial algebra

$$\bigotimes_{i=1}^n K[H_i][\partial_i, \partial_i^{-1}; \tau_i] = \mathcal{P}_n[\partial_1^{\pm 1}, \ldots, \partial_n^{\pm 1}; \tau_1, \ldots, \tau_n],$$

via $\partial_i \mapsto \partial_i$, $\int_i \mapsto \partial_i^{-1}$, $H_1 \mapsto H_i$ (and $x_i \mapsto \partial_i^{-1} H_i$) where $\mathcal{P}_n := K[H_1, \ldots, H_n]$ and $\tau_i(H_i) = H_i + 1$. We identify these two algebras via this isomorphism. It is obvious that

$$B_n = \bigotimes_{i=1}^n K[H_i][z_i, z_i^{-1}; \sigma_i] = \mathcal{P}_n[z_1^{\pm 1}, \ldots, z_n^{\pm 1}; \sigma_1, \ldots, \sigma_n],$$

where $z_i := \partial_i^{-1}$ and $\sigma_i = \tau_i^{-1} : H_i \mapsto H_i - 1$. The algebra B_n is also the left (but not right) localization of the algebra $\mathbb{I}_n$ at the multiplicatively closed set

$$S_\partial := S_{\partial_1, \ldots, \partial_n} := \{\partial_1^{\alpha_1} \cdots \partial_n^{\alpha_n} \,|\, (\alpha_i) \in \mathbb{N}^n\}, \;\; \text{and} \;\; B_n \simeq S_\partial^{-1}\mathbb{I}_n.$$

The algebra B_n contains the algebra $\Delta_n := \mathcal{P}_n[\partial_1, \ldots, \partial_n; \tau_1, \ldots, \tau_n]$ which is a skew polynomial ring.

Using the involution on the algebra $*$ on $\mathbb{I}_n$, the polynomial algebra P_n can be seen as the *right* $\mathbb{I}_n$-module by the rule

$$pa := a^*p \;\; \text{for all} \;\; p \in P_n \;\; \text{and} \;\; a \in \mathbb{I}_n.$$

By Lemma 4.2.(5), $P_n = K[x_1, \ldots, x_n]$ is the only faithful, simple, right $\mathbb{A}_n$-module. Let $P_n' := (P_n)^* = K[\partial_1, \ldots, \partial_n]$, a polynomial algebra in n variables. Clearly,

$$(P_n)_{\mathbb{A}_n} \simeq (\mathbb{I}_n/\mathbb{I}_n(H_1{-}1, \ldots, H_n{-}1, \partial_1, \ldots, \partial_n))^* = \mathbb{I}_n/\Big(H_1{-}1, \ldots, H_n{-}1, \int_1, \ldots, \int_n\Big)\mathbb{I}_n \simeq P_n'\widetilde{1} \simeq (P_n')_{P_n'}$$

where $\widetilde{1} := 1 + \Big(\int_1, \ldots, \int_n\Big)\mathbb{I}_n$. So, P_n' *is the only faithful, simple, right* $\mathbb{I}_n$*-module.*

Lemma 4.3. *1.* ${}_{\mathbb{I}_n}F_n \simeq P_n^{(\mathbb{N}^n)}$ *is a direct sum of* N^n *copies of the simple faithful left* $\mathbb{I}_n$*-module* P_n*.*

2. $(F_n)_{\mathbb{I}_n} \simeq (P_n')^{(\mathbb{N}^n)}$ *is a direct sum of* N^n *copies of the simple faithful right* $\mathbb{I}_n$*-module* P_n'*.*

Proof. 1. ${}_{\mathbb{I}_n}F_n = \oplus_{\alpha,\beta\in\mathbb{N}^n} KE_{\alpha\beta} = \oplus_{\beta\in\mathbb{N}^n}\big(\oplus_{\alpha,\in\mathbb{N}^n} KE_{\alpha\beta}\big) \simeq \oplus_{\beta\in\mathbb{N}^n} P_n \simeq P_n^{(\mathbb{N}^n)}$.

2. Statement 2 follows from statement 1 by applying the involution $*$ and using the fact that $F_n^* = F_n$. □

The ring $'Q(\mathbb{I}_n)$**.** Recall that $\Delta_n = \mathcal{P}_n[\partial_1,\dots,\partial_n;\tau_1,\dots,\tau_n]$. Let $\Delta_n^0 := \Delta_n\backslash\{0\}$, $'\Delta_n^0 := \Delta_n \cap {}'\mathcal{C}_{\mathbb{I}_n}$,

$$'\Delta_n := \Delta_n \cap {}'S(\mathbb{I}_n) \text{ and } \widetilde{'\Delta_n} := \{c \in \Delta_n \,|\, \partial^\alpha c \in {}'S(\mathbb{I}_n) \text{ for some } \alpha \in \mathbb{N}^n\}.$$

Notice that $'\Delta_n \subseteq {}'\Delta_n^0 \subseteq \Delta_n^0$. Theorem 4.4 produces explicit left denominators sets $S \in \mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$ such that $S^{-1}\mathbb{I}_n \simeq {}'Q(\mathbb{I}_n)$. By Theorem 4.4, there are inclusions in the set $\mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$ apart from S_∂:

$$S_\partial \subseteq {}'\Delta_n \subseteq {}'S(\mathbb{I}_n) \subseteq {}'S(\mathbb{I}_n) + \mathfrak{a}_n \text{ and } {}'\Delta_n \subseteq \widetilde{'\Delta_n} \subseteq' \Delta_n + \mathfrak{a}_n \subseteq {}'S(\mathbb{I}_n) + \mathfrak{a}_n. \tag{29}$$

Theorem 4.4. *1.* $'\Delta_n \in \mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$ *and* $'\Delta_n^{-1}\mathbb{I}_n \simeq {}'Q(\mathbb{I}_n)$*. Furthermore, the subset* $'\Delta_n$ *of* $\mathbb{I}_n$ *is a left denominator set of* $\mathbb{I}_n$ *which is the largest left denominator set that is contained in the multiplicative set* $'\mathcal{C}_{\mathbb{I}_n} \cap \Delta_n$*.*

2. $'\Delta_n + \mathfrak{a}_n, {}'S(\mathbb{I}_n) + \mathfrak{a}_n \in \mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$ *and*

$$\big('\Delta_n + \mathfrak{a}_n\big)^{-1}\mathbb{I}_n \simeq \big({}'S(\mathbb{I}_n) + \mathfrak{a}_n\big)^{-1}\mathbb{I}_n \simeq {}'Q(\mathbb{I}_n).$$

3. $'\Delta_n = \overline{'\Delta_n}, \overline{'\Delta_n + \mathfrak{a}_n}, \overline{'S(\mathbb{I}_n) + \mathfrak{a}_n} \in \mathrm{Den}_l(L_n, 0)$ *and*

$$'\Delta_n^{-1}L_n \simeq \overline{'\Delta_n + \mathfrak{a}_n}^{-1}L_n \simeq \overline{'S(\mathbb{I}_n) + \mathfrak{a}_n}^{-1}L_n \simeq {}'Q(\mathbb{I}_n)$$

where $\overline{S} := \pi_{\mathfrak{a}_n}(S)$ *and* $\pi_{\mathfrak{a}_n} : \mathbb{I}_n \to L_n = \mathbb{I}_n/\mathfrak{a}_n$*,* $r \mapsto \overline{r} := r + \mathfrak{a}_n$*.*

4. $\widetilde{'\Delta_n} \in \mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$ *and* $\widetilde{'\Delta_n}^{-1}\mathbb{I}_n \simeq {}'Q(\mathbb{I}_n)$*.* $\overline{\widetilde{'\Delta_n}} \in \mathrm{Den}_l(L_n, 0)$ *and* $\overline{\widetilde{'\Delta_n}}^{-1}L_n \simeq {}'Q(\mathbb{I}_n)$*.*

5. $\widetilde{'\Delta_n} + \mathfrak{a}_n \in \mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$ *and* $\big(\widetilde{'\Delta_n} + \mathfrak{a}_n\big)^{-1}\mathbb{I}_n \simeq {}'Q(\mathbb{I}_n)$*.*

6. $'\Delta_n \in \mathrm{Den}_l(\Delta_n, 0)$ *and* $'\Delta_n^{-1}\Delta_n \simeq {}'Q(\mathbb{I}_n)$*.*

7. $''\Delta_n := {}'S(\mathbb{I}_n) \cap ({}'\mathcal{C}_{\Delta_n+\mathfrak{a}_n} + \mathfrak{a}_n) + \mathfrak{a}_n \in \mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$*,* $'\Delta_n \subseteq ''\Delta_n$ *and* $''\Delta_n^{-1}\mathbb{I}_n \simeq {}'Q(\mathbb{I}_n)$*.*

Proof. 1. (i) $'\Delta_n \in \mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$ *and* $'\Delta_n^{-1}\mathbb{I}_n \simeq {}'Q(\mathbb{I}_n)$: By the definition, the set $'\Delta_n$ is a multiplicative subset of ${}'S(\mathbb{I}_n) \subseteq \mathbb{I}_n$. It follows from the inclusions $S_\partial \subseteq {}'\Delta_n \subseteq {}'S(\mathbb{I}_n)$ that

$$\mathfrak{a}_n = \mathrm{ass}_l(S_\partial) \subseteq \mathrm{ass}_l('\Delta_n) \subseteq \mathrm{ass}_l({}'S(\mathbb{I}_n)) = \mathfrak{a}_n,$$

and so $\mathrm{ass}_l('\Delta_n) = \mathfrak{a}_n = \mathrm{ass}_l({}'S(\mathbb{I}_n))$. For each element $s \in {}'S(\mathbb{I}_n)$, there is an element $\alpha \in \mathbb{N}^n$ such that $\partial^\alpha s \in {}'\Delta_n$. Notice that $\partial^\alpha \in S_\partial \subseteq {}'\Delta_n$. Now, the statement (i) follows from Lemma 3.6 where $T = {}'S(\mathbb{I}_n) \in \mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$ and $S = {}'\Delta_n$.

(ii) *The subset* $'\Delta_n$ *of* $\mathbb{I}_n$ *is a left denominator set of* $\mathbb{I}_n$ *which is the largest left denominator set that is contained in the multiplicative set* $'\mathcal{C}_{\mathbb{I}_n} \cap \Delta_n$: Let T be a left denominator set of $\mathbb{I}_n$ which is the largest left denominator set that is contained in the multiplicative set $'\mathcal{C}_{\mathbb{I}_n} \cap \Delta_n$. By the statement (i), $'\Delta_n \in \mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$. Clearly, $'\Delta_n \subseteq {}'\mathcal{C}_{\mathbb{I}_n} \cap \Delta_n$ and so $'\Delta_n \subseteq T$. Since $T \subseteq {}'\mathcal{C}_{\mathbb{I}_n}$ and ${}'S(\mathbb{I}_n)$ is a the largest left denominator set in ${}'\mathcal{C}_{\mathbb{I}_n}$, we have the inclusion $T \subseteq \Delta_n \cap {}'S(\mathbb{I}_n) = {}'\Delta_n$. Therefore, $T = {}'\Delta_n$.

2. By statement 1, ${}'\Delta_n, {}'S(\mathbb{I}_n) \in \mathrm{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$ and ${}'\Delta_n^{-1}\mathbb{I}_n \simeq {}'S(\mathbb{I}_n)^{-1}\mathbb{I}_n = {}'Q(\mathbb{I}_n)$. Now, statement 2 follows from Corollary 3.9.(2).

3. Statement 3 follows at once from statement 2 (If $S \in \mathrm{Den}_l(R, \mathfrak{a})$ then $\overline{S} := S + \mathfrak{a} \in \mathrm{Den}_l(\overline{R}, 0)$ and $S^{-1}R \simeq \overline{S}^{-1}\overline{R}$ where $\overline{R} := R/\mathfrak{a}$).

4. By the definition, the set $\widetilde{{}'\Delta_n}$ is a multiplicative set such that ${}'\Delta_n \subseteq \widetilde{{}'\Delta_n} \subseteq {}'S(\mathbb{I}_n)$. Therefore, $\mathfrak{a}_n = \mathrm{ass}_l({}'\Delta_n) \subseteq \mathrm{ass}_l(\widetilde{{}'\Delta_n}) \subseteq \mathrm{ass}_l({}'S(\mathbb{I}_n)) = \mathfrak{a}_n$, and so $\mathrm{ass}_l(\widetilde{{}'\Delta_n}) = \mathfrak{a}_n$. Now, the first part of statement 4 follows from the definition of the set $\widetilde{{}'\Delta_n}$ and Lemma 3.6 where $S = \widetilde{{}'\Delta_n}$ and $T = {}'S(\mathbb{I}_n)$.

The second part of statement 4 follows from the first one.

5. Statement 5 follows from statement 4.

6. By statement 3, ${}'\Delta_n \in \mathrm{Den}_l(L_n, 0)$ and ${}'\Delta_n^{-1}L_n \simeq {}'Q(\mathbb{I}_n)$. Now, statement 6 follows from Lemma 3.10.(2) where $R = \Delta_n$, $R' = L_n$ and $S = {}'\Delta_n$ (The R-module R'/R is S_∂-torsion. Hence it is also ${}'\Delta_n$-torsion as $S_\partial \subseteq {}'\Delta_n$).

7. By the definition, the subset ${}''\Delta_n$ of $\mathbb{I}_n$ is a multiplicative set such that ${}'\Delta_n \subseteq {}''\Delta_n \subseteq {}'S(\mathbb{I}_n)$, see statements 1 and 2. Hence, $\mathfrak{a}_n = \mathrm{ass}_l({}'\Delta_n) \subseteq \mathrm{ass}_l({}''\Delta_n) \subseteq \mathrm{ass}_l({}'S(\mathbb{I}_n)) = \mathfrak{a}_n$, and so $\mathrm{ass}_l({}''\Delta_n) = \mathfrak{a}_n = \mathrm{ass}_l({}'S(\mathbb{I}_n))$. Notice that $S_\partial \subseteq {}'\Delta_n \subseteq {}''\Delta_n$ and for each element $s \in {}''\Delta_n$ there is an element $\alpha \in \mathbb{N}^n$ such that $\partial^\alpha s \in {}'\Delta_n \subseteq {}''\Delta_n$. Now, statement 7 follows from Lemma 3.6 where $S = {}''\Delta_n$ and $T = {}'S(\mathbb{I}_n)$. □

In order to prove Theorem 4.6, we need the following lemma which is a characterization of the set ${}'\mathcal{C}_{\mathbb{I}_n}$.

Lemma 4.5. *Let $a \in \mathbb{I}_n$. Then the following statements are equivalent:*

1. $a \in {}'\mathcal{C}_{\mathbb{I}_n}$.
2. $a \in {}'\mathcal{C}_{F_n}$.
3. $a \in {}'\mathcal{C}_{P'_n}$ *where* $P'_n = K[\partial_1, \ldots, \partial_n]$ *is the only simple, faithful, right* $\mathbb{I}_n$*-module.*
4. $a^* \in \mathcal{C}'_{P_n}$ *where* $P_n = K[x_1, \ldots, x_n]$ *is the only simple, faithful, left* $\mathbb{I}_n$*-module.*

Proof. ($1 \Leftrightarrow 2$) The equivalence follows from Corollary 2.19.(1) and the fact that every nonzero ideal is an essential right ideal of the algebra $\mathbb{I}_n$ (Lemma 4.2.(7)).

($2 \Leftrightarrow 3$) The equivalence follows from Corollary 2.20.(1) and Lemma 4.3.(1).

($3 \Leftrightarrow 4$) The equivalence follows from the fact that $P'_n = P_n^*$. □

By applying the involution $*$ to Lemma 4.5, we obtain a similar characterization of right regular elements of the ring $\mathbb{I}_n$ ($a \in {}'\mathcal{C}_{\mathbb{I}_n}$ iff $a^* \in \mathcal{C}'_{\mathbb{I}_n}$).

Criterion for ${}'Q(\mathbb{I}_n) \simeq Q(A_n)$. The algebras Δ_n and B_n are Noetherian domains. By Goldie's Theorem, their quotient rings are division rings. It follows from the inclusions $A_n \subseteq B_n \subseteq Q(A_n)$ and $\Delta_n \subseteq S_\partial^{-1}\Delta_n \simeq B_n \subseteq Q(A_n)$ that

$$Q(\Delta_n) = Q(B_n) = Q(A_n). \tag{30}$$

Theorem 4.6. *The following statements are equivalent:*

1. ${}'Q(\mathbb{I}_n) \simeq Q(A_n)$.
2. ${}'Q_{l,cl}(\mathbb{I}_n) \simeq Q(A_n)$.
3. *The set* $\overline{{}'S(\mathbb{I}_n)}$ *is dense in* $B_n \backslash \{0\}$.
4. *The set* $\overline{{}'\mathcal{C}_{\mathbb{I}_n}}$ *is dense in* $B_n \backslash \{0\}$.
5. *The set* ${}'\Delta_n$ *is dense in* $B_n \backslash \{0\}$.

6. *The set* $'\Delta_n$ *is left dense in* Δ_n^0.

7. *For each element* $s \in \Delta_n^0$, *there is an element* $s' \in \Delta_n^0$ *such that* $s's \in {}'\mathcal{C}_{F_n}$.

8. *For each element* $s \in \Delta_n^0$, *there is an element* $s' \in \Delta_n^0$ *such that* $s's \in {}'\mathcal{C}_{P'_n}$ *where* $P'_n = K[\partial_1, \dots, \partial_n]$ *is the unique simple faithful right* $\mathbb{I}_n$*-module.*

Proof. $(1 \Leftrightarrow 2 \Leftrightarrow 3 \Leftrightarrow 4)$ The equivalence $(1 \Leftrightarrow 2)$ follows from Theorem 3.12.(2). The equivalences $(1 \Leftrightarrow 3)$ and $(2 \Leftrightarrow 4)$follows from Theorem 1.1 and Theorem 3.12.(2) (since $\operatorname{ass}_l({}'S(\mathbb{I}_n)) = \mathfrak{a}_n$ is a prime ideal of $\mathbb{I}_n$, the algebra $\mathbb{I}_n/\mathfrak{a}_n = B_n$ is a domain and $Q(\mathbb{I}_n/\mathfrak{a}_n) = Q(B_n)$ is a division ring).

$(1 \Leftrightarrow 5)$ By Theorem 4.4.(1), $'\Delta_n \in \operatorname{Den}_l(\mathbb{I}_n, \mathfrak{a}_n)$ and $'\Delta_n^{-1}\mathbb{I}_n \simeq {}'Q(\mathbb{I}_n)$. Since the set $'\Delta_n$ is dense in ${}'S(\mathbb{I}_n)$, the equivalence $(1 \Leftrightarrow 5)$ holds iff the equivalence $(1 \Leftrightarrow 3)$ holds.

$(5 \Leftrightarrow 6)$ Recall that $'\Delta_n, \Delta_n^0 \in \operatorname{Den}_l(\Delta_n, 0)$, $'\Delta_n \subseteq \Delta_n^0$ and $Q(\Delta_n) = Q(A_n)$, see (30). By [12, Lemma 3.5.(3)], $'\Delta_n^{-1}\Delta_n \simeq (\Delta_n^0)^{-1}\Delta_n = Q(A_n)$ iff the set $'\Delta_n$ is left dense in Δ_n^0.

$(6 \Leftrightarrow 7)$ By Lemma 4.5, the inclusion $s's \in {}'\mathcal{C}_{F_n}$ is equivalent to the inclusion $s's \in {}'\mathcal{C}_{\mathbb{I}_n}$. Hence, statement 7 is equivalent to the statement that for each element $s \in \Delta_n^0$, there is an element $s' \in \Delta_n^0$ such that $s's \in {}'\mathcal{C}_{\mathbb{I}_n}$, i.e. $s's \in \Delta_n \cap {}'\mathcal{C}_{\mathbb{I}_n} = {}'\Delta_n$, i.e. it is equivalent to statement 6.

$(7 \Leftrightarrow 8)$ The equivalence follows from Lemma 4.5. □

Description of the set $'\mathcal{C}_{\mathbb{A}_1}$**.** By (24), each element $a \in \mathbb{I}_1$ is a unique sum $a = \sum_{i=0}^{l} d_{-i}\partial^i + \sum_{j=1}^{m} \int^j d_j + a_F$ where $d_k \in K[H]$, $k = -l, \dots, m$ and $a_F \in F$. Let $a_\partial := \sum_{i=0}^{l} d_{-i}\partial^i$. The integer

$$s(a_F) := \begin{cases} \min\{n \in \mathbb{N} \,|\, a_F \in \bigoplus_{i,j=0}^{n} Ke_{ij}\} & \text{if } a_F \neq 0, \\ -1 & \text{if } a_F = 0. \end{cases}$$

is called the *size* of the element $a_F \in F$. The integer $s(a) := s(a_F)$ is called the *size* of the element a. For each $i \in \mathbb{N}$, let $P'_{1,\leq i} := \{a \in P'_1 \,|\, \deg_y(a) \leq i\}$ where $\deg_y \partial$ is the degree of the polynomial $a \in P'_1 = K[\partial]$ in the variable ∂.

For all polynomials $p \in K[H]$, $\partial p = \tau(p)\partial$ where $\tau \in \operatorname{Aut}_K(K[H])$ and $\tau(H) = H + 1$. For each nonzero polynomials $p \in K[H]$, let

$$\mu(p) := \min\{i \in \mathbb{N} \,|\, \text{the polynomial } \tau^i(p) \in K[H] \text{ has no root in } \mathbb{N}_+\}.$$

Let $\Psi := \bigoplus_{i\geq 1} \int^i K[H] \oplus F$. If $a = \sum_{i=0}^{n} d_{-i}\partial^i + \sum_{j=1}^{m} \int^j d_j + a_F \in \mathbb{I}_1 \backslash \Psi$ then $a_\partial \neq 0$ and so $d_{-n} \neq 0$ where $n = \deg_\partial(a_\partial)$. Let

$$\mu(a) := \mu(d_{-n}) \text{ and } \nu(a) := \max\{s(a), \mu(a)\}.$$

Theorem 4.7. *1.* $'\mathcal{C}_{\mathbb{I}_1} = \{a \in \mathbb{I}_1 \backslash \Psi \,|\, \cdot a : P'_{1,\leq \nu(a)} \to P'_{1,\leq \nu(a)+\deg_y(a_y)},\ p \mapsto pa \text{ is an injection}\}$.

2. $\mathcal{C}'_{\mathbb{I}_1} = {}'\mathcal{C}^*_{\mathbb{I}_1}$ *where* $*$ *is the involution of the algebra* $\mathbb{I}_1$, *see (25).*

Proof. 1. (i) $'\mathcal{C}_{\mathbb{I}_1} \cap \Psi = \emptyset$: Suppose that $a \in \Psi$. Then the map $\cdot a : P'_{1,\leq s(a)+1} \to P'_{1,\leq s(a)}$, $p \mapsto pa$ is a well-defined map. Since

$$\dim_K(P'_{1,\leq s(a)+1}) = s(a) + 2 > s(a) + 1 = \dim_K(P'_{1,\leq s(a)}),$$

$\ker(\cdot a) \neq 0$, $a \notin {}'\mathcal{C}_{\mathbb{I}_1}$. Therefore, $'\mathcal{C}_{\mathbb{I}_1} \cap \Psi = \emptyset$.

(ii) *For each element* $a \in \mathbb{I}_1 \backslash \Psi$, $\ker_{P'_1}(\cdot a) \subseteq P'_{1,\leq \nu(a)}$: Since $a \in \mathbb{I}_1 \backslash \Psi$, $a_y := \sum_{i=0}^{l} d_{-i}\partial^i \neq 0$ where $d_{-i} \in K[H]$. Suppose that $d_{-l} \neq 0$. Suppose that $p \in \ker_{P'_1}(\cdot a) \backslash P'_{1,\leq \nu(a)}$, i.e. $\deg_y(p) > \nu(a)$. Then

$$\deg(pa) = l + \deg_y(p),$$

a contradiction (since $pa = 0$).

Now, statement 1 follows from statements (i) and (ii).

2. Statement 2 follows from statement 1. □

Lemma 4.8 provides examples of rings R such that ${}'\mathcal{C}_R^{lee} = \emptyset$, $\mathcal{C}'^{ree}_R = \emptyset$ and $\mathcal{C}_R^{ee} = \emptyset$.

Lemma 4.8. *Let R be either $\mathbb{I}_n$ or $\mathbb{S}_n$. Then:*

1. *${}'\mathcal{C}_R \cap F_n = \emptyset$, $\mathcal{C}'_R \cap F_n = \emptyset$ and $\mathcal{C}_R \cap F = \emptyset$.*
2. *${}'\mathcal{C}_R^{lee} = \emptyset$, $\mathcal{C}'^{ree}_R = \emptyset$ and $\mathcal{C}_R^{ee} = \emptyset$.*

Proof. 1. Clearly, every element of the ideal F is a left and right zero-divisor, and statement 1 follows.

2. The ideal F_n is a left and right essential ideal of R, and so statement 2 follows from statement 1. □

5 The rings ${}'Q(\mathbb{A}_n)$ and $Q'(\mathbb{A}_n)$

The aim of the section is to prove Theorem 5.1 and Theorem 5.5. Theorem 5.4 a criterion for ${}'Q(\mathbb{A}_n) \simeq Q(A_n)$. As a corollary we obtain that ${}'Q(\mathbb{A}_1) \simeq Q(A_1)$ (Theorem 5.5). Theorem 5.8 and Theorem 5.7 describe the set ${}'\mathcal{C}_{\mathbb{A}_1}$. At the beginning of the section, we recall necessary facts about the Jacobian algebras $\mathbb{A}_n$ that are used in the proofs. The details can found in [3].

The Weyl algebra $A_n = A_n(K)$ is a simple, Noetherian domain of Gelfand-Kirillov dimension $\operatorname{GK}(A_n) = 2n$. The Jacobian algebra $\mathbb{A}_n$ is neither left nor right Noetherian, it contains infinite direct sums of nonzero left and right ideals. This means that adding the inverses of the commuting regular elements $H_1, \dots, H_n$ to the Weyl algebra A_n is neither a left nor right Ore localization of the algebra Weyl A_n. This fact is a prime reason why the properties of the Jacobian algebras are almost opposite to the ones of the Weyl algebras.

The algebra $\mathbb{A}_n$ is a central, prime algebra of Gelfand-Kirillov dimension $3n$ ([3, Corollary 2.7]). The canonical involution θ of the Weyl algebra A_n can be uniquely extended to the algebra $\mathbb{A}_n$ (see (32)). So, the algebra $\mathbb{A}_n$ is self-dual ($\mathbb{A}_n \simeq \mathbb{A}_n^{op}$) and its left and right algebraic properties are the same. Note that the *Fourier transform* on the Weyl algebra A_n cannot be lifted to $\mathbb{A}_n$. Many properties of the algebra $\mathbb{A}_n = \mathbb{A}_1^{\otimes n}$ are determined by properties of $\mathbb{A}_1$. When $n = 1$ we usually drop the subscript '1' in x_1, ∂_1, H_1, etc. The algebra $\mathbb{A}_1$ contains the only proper ideal $F = \oplus_{i,j\in\mathbb{N}} KE_{ij}$ where

$$E_{ij} := \begin{cases} x^{i-j}(x^j \frac{1}{\partial^j x^j}\partial^j - x^{j+1}\frac{1}{\partial^{j+1}x^{j+1}}\partial^{j+1}) & \text{if } i \geq j, \\ (\frac{1}{\partial x}\partial)^{j-i}(x^j \frac{1}{\partial^j x^j}\partial^j - x^{j+1}\frac{1}{\partial^{j+1}x^{j+1}}\partial^{j+1}) & \text{if } i < j. \end{cases}$$

As a ring without 1, the ring F is canonically isomorphic to the ring $M_\infty(K) := \varinjlim M_d(K) = \oplus_{i,j\in\mathbb{N}} KE_{ij}$ of infinite-dimensional matrices where E_{ij} are the matrix units ($F \to M_\infty(K)$, $E_{ij} \mapsto E_{ij}$). This is a very important fact as we can apply concepts of finite-dimensional linear algebra (like trace, determinant, etc) to integro-differential operators which is not obvious from the outset. This fact is crucial in finding an inversion formula for elements of $\mathbb{A}_1^*$.

Notice that $\mathbb{A}_n = \otimes_{i=1}^n \mathbb{A}_1(i) \simeq \mathbb{A}_1^{\otimes n}$ where $\mathbb{A}_1(i) := K\langle x_i, \partial_i, H_i^{\pm 1}\rangle$ and $H_i = \partial_i x_i$. The algebra $\mathbb{A}_n = \oplus_{\alpha\in\mathbb{Z}^n} \mathbb{A}_{n,\alpha}$ is a $\mathbb{Z}^n$-graded algebra where $\mathbb{A}_{n,\alpha} := \otimes_{k=1}^n \mathbb{A}_{1,\alpha_k}(k)$ for $\alpha = (\alpha_1, \dots, \alpha_n)$. For $n = 1$, ([3, Theorem 2.3]),

$$\mathbb{A}_{1,i} = \begin{cases} x^i\mathbb{D}_1 & \text{if } i \geq 1, \\ \mathbb{D}_1 & \text{if } i = 0, \\ \mathbb{D}_1\partial^{-i} & \text{if } i \leq -1, \end{cases} \tag{31}$$

where $\mathbb{D}_1 := L \oplus (\oplus_{i,j\geq 1} Kx^i H^{-j}\partial^i)$ is a *commutative, non-Noetherian* algebra and

$$L := K[H^{\pm 1}, (H+1)^{-1}, (H+2)^{-1}, \dots] \text{ and } H = \partial x = x\partial + 1.$$

This gives a 'compact' K-basis for the algebra $\mathbb{A}_1$ (and $\mathbb{A}_n$). This basis 'behaves badly' under multiplication. A more conceptual ('multiplicatively friendly') basis is given in [3, Theorem 2.5], see also [3, Corollary 2.4] below.

- ([3, Corollary 2.4]) $\mathbb{D}_1 = L \oplus (\oplus_{i\geq 1, j\geq 0} K\rho_{ji})$ *where* $\rho_{ji} := x^i \frac{1}{H^j \partial^i x^i} \partial^i$.
- For all $i \geq 1$ and $j \geq 0$, $\partial^i \rho_{ji} = \frac{1}{H^j} \partial^i$ (a direct computation).
- ([3, Corollary 2.7.(10)]) $P_n = K[x_1, \ldots, x_n]$ *is the only faithful, simple* $\mathbb{A}_n$*-module.*
- ([3, Corollary 3.5]) $\mathfrak{p}_1 := F \otimes \mathbb{A}_{n-1}, \mathfrak{p}_2 := \mathbb{A}_1 \otimes F \otimes \mathbb{A}_{n-2}, \ldots, \mathfrak{p}_n := \mathbb{A}_{n-1} \otimes F$, *are precisely the prime ideals of height 1 of* $\mathbb{A}_n$.
- ([3, Corollary 3.15]) $\mathfrak{a}_n := \mathfrak{p}_1 + \cdots + \mathfrak{p}_n$ *is the only prime ideal of* $\mathbb{A}_n$ *which is completely prime;* $\mathfrak{a}_n$ *is the only ideal* $\mathfrak{a}$ *of* $\mathbb{A}_n$ *such that* $\mathfrak{a} \neq \mathbb{A}_n$ *and* $\mathbb{A}_n/\mathfrak{a}$ *is a Noetherian (resp. left Noetherian, resp. right Noetherian) ring.*
- ([3, Theorem 3.1.(2)]) *Each ideal* I *of* $\mathbb{A}_n$ *is an idempotent ideal* ($I^2 = I$).
- ([3, Corollary 2.7.(4)]) *The ideal* $F_n := F^{\otimes n}$ *is the smallest nonzero ideal of the algebra* $\mathbb{A}_n$.
- ([3, Corollary 2.7.(8)]) ${}_{\mathbb{A}_n}F^{\otimes n} \simeq P_n^{(\mathbb{N}^n)}$ *is a faithful, semi-simple, left* $\mathbb{A}_n$*-module;* $F^{\otimes n}_{\mathbb{A}_n} \simeq P_n^{(\mathbb{N}^n)}{}_{\mathbb{A}_n}$ *is a faithful, semi-simple, right* $\mathbb{A}_n$*-module;* ${}_{\mathbb{A}_n}F^{\otimes n}_{\mathbb{A}_n}$ *is a faithful, simple* $\mathbb{A}_n$*-bimodule.*

Recall that $\mathcal{P}_n$ is a polynomial algebra $K[H_1, \ldots, H_n]$ in n indeterminates and $\sigma = (\sigma_1, ..., \sigma_n)$ is an n-tuple of commuting automorphisms of $\mathcal{P}_n$ where $\sigma_i(H_i) = H_i - 1$ and $\sigma_i(H_j) = H_j$, for $i \neq j$. By [6, Theorem 2.2], $\mathfrak{a}_n$ is the only maximal ideal of the Jacobian algebra $\mathbb{A}_n$. The factor algebra $\mathcal{A}_n := \mathbb{A}_n/\mathfrak{a}_n$ is the skew Laurent polynomial algebra

$$\begin{array}{rcl} \mathcal{A}_n & := & \mathcal{L}_n[\partial_1^{\pm 1}, \ldots, \partial_n^{\pm 1}; \tau_1, \ldots, \tau_n] = \mathcal{L}_n[x_1^{\pm 1}, \ldots, x_n^{\pm 1}; \sigma_1, \ldots, \sigma_n], \\ \mathcal{L}_n & := & K[H_1^{\pm 1}, (H_1 \pm 1)^{-1}, (H_1 \pm 2)^{-1}, \ldots, H_n^{\pm 1}, (H_n \pm 1)^{-1}, (H_n \pm 2)^{-1}, \ldots], \end{array}$$

where $\tau_i(H_j) = H_j + \delta_{ij}$, δ_{ij} is the Kronecker delta, $z_i = \partial_i^{-1}$ and $\sigma_i = \tau_i^{-1}$. The algebra B_n is a subalgebra of $\mathcal{A}_n$. Let S_n be a multiplicative submonoid of $\mathcal{P}_n$ generated by the elements $H_i + j$, $i = 1, \ldots, n$, and $j \in \mathbb{Z}$. Then S_n is an Ore set for the Weyl algebra A_n, the algebras B_n and the polynomial algebra $\mathcal{P}_n$ such that $\mathcal{A}_n := S_n^{-1}A_n \simeq S_n^{-1}B_n$ and

$$S_n^{-1}\mathcal{P}_n = K[H_1^{\pm 1}, (H_1 \pm 1)^{-1}, (H_1 \pm 2)^{-1}, \ldots, H_n^{\pm 1}, (H_n \pm 1)^{-1}, (H_n \pm 2)^{-1}, \ldots].$$

We identify the Weyl algebra A_n with a subalgebra of $\mathcal{A}_n$ via the monomorphism

$$A_n \to \mathcal{A}_n, \;\; x_i \mapsto x_i, \;\; \partial_i \mapsto H_i x_i^{-1}, \;\; i = 1, \ldots, n.$$

The Weyl algebra A_n is a Noetherian domain. So, by Goldie's Theorem, the (left and right) quotient ring of A_n, $Q(A_n)$, is a division ring. Then the algebra $\mathcal{A}_n$ is a K-subalgebra of $Q(A_n)$ generated by the elements x_i, x_i^{-1}, H_i and H_i^{-1}, $i = 1, \ldots, n$ since

$$(H_i \pm j)^{-1} = x_i^{\mp j} H_i^{-1} x_i^{\pm j}, \;\; i = 1, \ldots, n \;\text{ and }\; j \in \mathbb{N}.$$

Clearly, $\mathcal{A}_n \simeq \mathcal{A}_1^{\otimes n}$.

The involution θ on $\mathbb{A}_n$. The Weyl algebra A_n admits the *involution*

$$\theta : A_n \to A_n, \;\; x_i \mapsto \partial_i, \;\; \partial_i \mapsto x_i, \;\; i = 1, \ldots, n.$$

The involution θ is uniquely extended to an involution of $\mathbb{A}_n$ by the rule

$$\theta : \mathbb{A}_n \to \mathbb{A}_n, \;\; x_i \mapsto \partial_i, \;\; \partial_i \mapsto x_i, \;\; \theta(H_i^{-1}) = H_i^{-1}, \;\; i = 1, \ldots, n. \tag{32}$$

Uniqueness is obvious: $\theta(H_i) = \theta(\partial_i x_i) = \theta(x_i)\theta(\partial_i) = \partial_i x_i = H_i$ and so $\theta(H_i^{-1}) = H_i^{-1}$. So, the algebra $\mathbb{A}_n$ is self-dual and left and right algebraic properties of the algebra $\mathbb{A}_n$ are the same.

The polynomial algebra P_n is a left A_n-module where $\partial_i * f := \frac{\partial f}{\partial x_i}$ for all $i = 1, \dots, n$ and $f \in P_n$. The left A_n-module P_n is isomorphic to the A_n-module

$$A_n/A_n(\partial_1, \dots, \partial_n) \simeq P_n\overline{1} \simeq {}_{P_n}P_n \text{ where } \overline{1} := 1 + A_n(\partial_1, \dots, \partial_n).$$

The maps $H_i : P_n \to P_n$ are invertible for all $i = 1, \dots, n$ since $H_i x^\alpha \overline{1} = (\alpha_i + 1)x^\alpha$ for all $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ where $x^\alpha := \prod_{i=1}^n x_i^{\alpha_i}$. Therefore, the polynomial algebra P_n is also a left $\mathbb{A}_n$-module which is isomoprphic to

$$\mathbb{A}_n/\mathbb{A}_n(H_1 - 1, \dots, H_n - 1, \partial_1, \dots, \partial_n) \simeq P_n\overline{1}, \text{ where } \overline{1} := 1 + \mathbb{A}_n(H_1 - 1, \dots, H_n - 1, \partial_1, \dots, \partial_n),$$

by [3, Theorem 2.3]. Using the involution θ on $\mathbb{A}_n$, the polynomial algebra P_n can be seen as the *right* $\mathbb{A}_n$-module by the rule

$$pa := \theta(a)p \text{ for all } p \in P_n \text{ and } a \in \mathbb{A}_n.$$

By [3, Corollary 2.7.(10)]), $P_n = K[x_1, \dots, x_n]$ *is the only faithful, simple, right* $\mathbb{A}_n$*-module.* Let $P_n' := \theta(P_n) = K[\partial_1, \dots, \partial_n]$, a polynomial algebra in n variables. Clearly,

$$(P_n)_{\mathbb{A}_n} = \theta(P_n\overline{1}) = \widetilde{1}\theta(P_n) = \widetilde{1}P_n' \simeq \mathbb{A}_n/(H_1 - 1, \dots, H_n - 1, x_1, \dots, x_n)\mathbb{A}_n \simeq (P_n')_{P_n'}$$

where $\widetilde{1} := 1 + (H_1 - 1, \dots, H_n - 1, \partial_1, \dots, \partial_n)\mathbb{A}_n$.

For $n = 1$, the set F is the only proper ideal of $\mathbb{A}_1$, hence $\theta(F) = F$. Moreover,

$$\theta(E_{ij}) = \frac{i!}{j!} E_{ji} \tag{33}$$

where $0! := 1$. The ring $F = \oplus_{i,j\in\mathbb{N}} KE_{ij}$ is equal to the matrix ring $M_\infty(K) := \cup_{d\geq 1} M_d(K)$ where $M_d(K) := \oplus_{0\leq i,j\leq d-1} KE_{ij}$. The ring $F = M_\infty(K)$ admits the canonical involution which is the transposition $(\cdot)^t : E_{ij} \mapsto E_{ji}$. Let $D_!$ be the infinite diagonal matrix $\text{diag}(0!, 1!, 2!, \dots)$. Then, for $u \in F = M_\infty(K)$,

$$\theta(u) = D_!^{-1} u^t D_!. \tag{34}$$

Note that $D_! \notin M_\infty(K)$. For $n \geq 1$, $F_n := F^{\otimes n} = \oplus_{\alpha,\beta\in\mathbb{N}^n} KE_{\alpha\beta} = M_\infty(K)^{\otimes n}$ where $E_{\alpha\beta} := \otimes_{i=1}^n E_{\alpha_i\beta_i}$. By (33),

$$\theta(E_{\alpha\beta}) = \frac{\alpha!}{\beta!} E_{\beta\alpha}, \tag{35}$$

$$\theta(F^{\otimes n}) = F^{\otimes n}. \tag{36}$$

Let $D_{n,!} := D_!^{\otimes n}$. Then, for $u \in F^{\otimes n}$,

$$\theta(u) = D_{n,!}^{-1} u^t D_{n,!} \tag{37}$$

where $(\cdot)^t : M_\infty(K)^{\otimes n} \to M_\infty(K)^{\otimes n}$, $E_{\alpha\beta} \mapsto E_{\beta\alpha}$, is the transposition map.

Consider the bilinear, symmetric, nondegenerate form $(\cdot, \cdot) : P_n \times P_n \to K$ given by the rule $(x^\alpha, x^\beta) := \alpha!\delta_{\alpha,\beta}$ for all $\alpha, \beta \in \mathbb{N}^n$. Then, for all $p, q \in P_n$ and $a \in \mathbb{A}_n$,

$$(p, aq) = (\theta(a)p, q). \tag{38}$$

The Weyl algebra A_n admits, the so-called, *Fourier transform*, which is the K-algebra automorphism

$$\mathcal{F} : A_n \to A_n, \quad x_i \mapsto \partial_i, \quad \partial_i \mapsto -x_i \text{ for } i = 1, \dots, n.$$

Since $\mathcal{F}(H_i) = -(H_i - 1)$, H_i is a unit of $\mathbb{A}_n$ and $H_i - 1$ is not, one *cannot* extend the Fourier transform to $\mathbb{A}_n$.

For all $i = 1, \dots, n$, $\partial_i \in {}'\mathcal{C}_{\mathbb{A}_n}$. Recall that if $n = 1$ then $\partial^i \rho_{ji} = \frac{1}{H^j}\partial^i$ for all $i \geq 1$ and $j \geq 0$. It follows that

$$S_\partial := S_{\partial_1,\dots,\partial_n} := \{\partial_1^{\alpha_1} \cdots \partial_n^{\alpha_n} \mid (\alpha_i) \in \mathbb{N}^n\} \in \text{Den}_l(\mathbb{A}_n, \mathfrak{a}_n), \quad S_\partial \subseteq {}'\mathcal{C}_{\mathbb{A}_n} \text{ and } S_\partial^{-1}\mathbb{A}_n \simeq \mathcal{A}_n.$$

The ring $'Q(\mathbb{A}_n)$. The ring

$$\begin{array}{rcl}\Delta_n & := & \mathcal{L}_n^+[\partial_1,\dots,\partial_n;\tau_1,\dots,\tau_n], \text{ where}\\ \mathcal{L}_n^+ & := & K[H_1^{\pm 1},(H_1+1)^{-1},(H_1+2)^{-1},\dots,H_n^{\pm 1},(H_n+1)^{-1},(H_n+2)^{-1},\dots]\simeq L^{\otimes n},\end{array}$$

is a skew polynomial ring where $\tau_i(H_j)=H_i+\delta_{ij}$, $\Delta_n^0:=\Delta_n\backslash\{0\}$, $'\Delta_n^0:=\Delta_n\cap{}'\mathcal{C}_{\mathbb{A}_n}$,

$$'\Delta_n:=\Delta_n\cap{}'S(\mathbb{A}_n) \text{ and } \widetilde{'\Delta_n}:=\{c\in\Delta_n\,|\,\partial^\alpha c\in{}'S(\mathbb{A}_n) \text{ for some } \alpha\in\mathbb{N}^n\}.$$

Notice that $'\Delta_n\subseteq{}'\Delta_n^0\subseteq\Delta_n^0$. Theorem 5.1 produces explicit left denominators sets $S\in \mathrm{Den}_l(\mathbb{A}_n,\mathfrak{a}_n)$ such that $S^{-1}\mathbb{A}_n\simeq{}'Q(\mathbb{A}_n)$. By Theorem 5.1, there are inclusions in the set $\mathrm{Den}_l(\mathbb{A}_n,\mathfrak{a}_n)$ apart from S_∂:

$$S_\partial\subseteq{}'\Delta_n\subseteq{}'S(\mathbb{A}_n)\subseteq{}'S(\mathbb{A}_n)+\mathfrak{a}_n \text{ and } '\Delta_n\subseteq\widetilde{'\Delta_n}\subseteq'\,\Delta_n+\mathfrak{a}_n\subseteq{}'S(\mathbb{A}_n)+\mathfrak{a}_n. \tag{39}$$

Theorem 5.1. *1. $'\Delta_n\in\mathrm{Den}_l(\mathbb{A}_n,\mathfrak{a}_n)$ and $'\Delta_n^{-1}\mathbb{A}_n\simeq{}'Q(\mathbb{A}_n)$. Furthermore, the subset $'\Delta_n$ of $\mathbb{A}_n$ is a left denominator set of $\mathbb{A}_n$ which is the largest left denominator set that is contained in the multiplicative set $'\mathcal{C}_{\mathbb{A}_n}\cap\Delta_n$.*

2. $'\Delta_n+\mathfrak{a}_n,{}'S(\mathbb{A}_n)+\mathfrak{a}_n\in\mathrm{Den}_l(\mathbb{A}_n,\mathfrak{a}_n)$ and

$$\left('\Delta_n+\mathfrak{a}_n\right)^{-1}\mathbb{A}_n\simeq\left('S(\mathbb{A}_n)+\mathfrak{a}_n\right)^{-1}\mathbb{A}_n\simeq{}'Q(\mathbb{A}_n).$$

3. $'\Delta_n=\overline{'\Delta_n},\overline{'\Delta_n+\mathfrak{a}_n},\overline{'S(\mathbb{A}_n)+\mathfrak{a}_n}\in\mathrm{Den}_l(\mathcal{L}_n,0)$ and

$$'\Delta_n^{-1}\mathcal{L}_n\simeq\overline{'\Delta_n+\mathfrak{a}_n}^{-1}\mathcal{L}_n\simeq\overline{'S(\mathbb{A}_n)+\mathfrak{a}_n}^{-1}\mathcal{L}_n\simeq{}'Q(\mathbb{A}_n)$$

where $\overline{S}:=\pi_{\mathfrak{a}_n}(S)$ and $\pi_{\mathfrak{a}_n}:\mathbb{A}_n\to\mathcal{L}_n=\mathbb{A}_n/\mathfrak{a}_n$, $r\mapsto\overline{r}:=r+\mathfrak{a}_n$.

4. $\widetilde{'\Delta_n}\in\mathrm{Den}_l(\mathbb{A}_n,\mathfrak{a}_n)$ and $\widetilde{'\Delta_n}^{-1}\mathbb{A}_n\simeq{}'Q(\mathbb{A}_n)$. $\widetilde{\overline{'\Delta_n}}\in\mathrm{Den}_l(\mathcal{L}_n,0)$ and $\widetilde{\overline{'\Delta_n}}^{-1}\mathcal{L}_n\simeq{}'Q(\mathbb{A}_n)$.

5. $\widetilde{'\Delta_n}+\mathfrak{a}_n\in\mathrm{Den}_l(\mathbb{A}_n,\mathfrak{a}_n)$ and $\left(\widetilde{'\Delta_n}+\mathfrak{a}_n\right)^{-1}\mathbb{A}_n\simeq{}'Q(\mathbb{A}_n)$.

6. $'\Delta_n\in\mathrm{Den}_l(\Delta_n,0)$ and $'\Delta_n^{-1}\Delta_n\simeq{}'Q(\mathbb{A}_n)$.

7. $''\Delta_n:={}'S(\mathbb{A}_n)\cap({}'\mathcal{C}_{\Delta_n+\mathfrak{a}_n}+\mathfrak{a}_n)+\mathfrak{a}_n\in\mathrm{Den}_l(\mathbb{A}_n,\mathfrak{a}_n)$, $'\Delta_n\subseteq''\Delta_n$ and $''\Delta_n^{-1}\mathbb{A}_n\simeq{}'Q(\mathbb{A}_n)$.

Proof. 1. (i) $'\Delta_n\in\mathrm{Den}_l(\mathbb{A}_n,\mathfrak{a}_n)$ and $'\Delta_n^{-1}\mathbb{A}_n\simeq{}'Q(\mathbb{A}_n)$: By the definition, the set $'\Delta_n$ is a multiplicative subset of $'S(\mathbb{A}_n)\subseteq\mathbb{A}_n$. It follows from the inclusions $S_\partial\subseteq{}'\Delta_n\subseteq{}'S(\mathbb{A}_n)$ that

$$\mathfrak{a}_n=\mathrm{ass}_l(S_\partial)\subseteq\mathrm{ass}_l('\Delta_n)\subseteq\mathrm{ass}_l('S(\mathbb{A}_n))=\mathfrak{a}_n,$$

and so $\mathrm{ass}_l('\Delta_n)=\mathfrak{a}_n=\mathrm{ass}_l('S(\mathbb{A}_n))$. For each element $s\in{}'S(\mathbb{A}_n)$, there is an element $\alpha\in\mathbb{N}^n$ such that $\partial^\alpha s\in{}'\Delta_n$. Notice that $\partial^\alpha\in S_\partial\subseteq{}'\Delta_n$. Now, the statement (i) follows from Lemma 3.6 where $T={}'S(\mathbb{A}_n)\in\mathrm{Den}_l(\mathbb{A}_n,\mathfrak{a}_n)$ and $S={}'\Delta_n$.

(ii) *The subset $'\Delta_n$ of $\mathbb{A}_n$ is a left denominator set of $\mathbb{A}_n$ which is the largest left denominator set that is contained in the multiplicative set $'\mathcal{C}_{\mathbb{A}_n}\cap\Delta_n$*: Let T be a left denominator set of $\mathbb{A}_n$ which is the largest left denominator set that is contained in the multiplicative set $'\mathcal{C}_{\mathbb{A}_n}\cap\Delta_n$. By the statement (i), $'\Delta_n\in\mathrm{Den}_l(\mathbb{A}_n,\mathfrak{a}_n)$. Clearly, $'\Delta_n\subseteq{}'\mathcal{C}_{\mathbb{A}_n}\cap\Delta_n$ and so $'\Delta_n\subseteq T$. Since $T\subseteq{}'\mathcal{C}_{\mathbb{A}_n}$ and ${}'S(\mathbb{A}_n)$ is a the largest left denominator set in $'\mathcal{C}_{\mathbb{A}_n}$, we have the inclusion $T\subseteq\Delta_n\cap{}'S(\mathbb{A}_n)={}'\Delta_n$. Therefore, $T={}'\Delta_n$.

2. By statement 1, $'\Delta_n,{}'S(\mathbb{A}_n)\in\mathrm{Den}_l(\mathbb{A}_n,\mathfrak{a}_n)$ and $'\Delta_n^{-1}\mathbb{A}_n\simeq{}'S(\mathbb{A}_n)^{-1}\mathbb{A}_n={}'Q(\mathbb{A}_n)$. Now, statement 2 follows from Corollary 3.9.(2).

3. Statement 3 follows at once from statement 2 (If $S\in\mathrm{Den}_l(R,\mathfrak{a})$ then $\overline{S}:=S+\mathfrak{a}\in\mathrm{Den}_l(\overline{R},0)$ and $S^{-1}R\simeq\overline{S}^{-1}\overline{R}$ where $\overline{R}:=R/\mathfrak{a}$).

4. By the definition, the set $\widetilde{{}'\Delta_n}$ is a multiplicative set such that ${}'\Delta_n \subseteq \widetilde{{}'\Delta_n} \subseteq {}'S(\mathbb{A}_n)$. Therefore, $\mathfrak{a}_n = \mathrm{ass}_l({}'\Delta_n) \subseteq \mathrm{ass}_l(\widetilde{{}'\Delta_n}) \subseteq \mathrm{ass}_l({}'S(\mathbb{A}_n)) = \mathfrak{a}_n$, and so $\mathrm{ass}_l(\widetilde{{}'\Delta_n}) = \mathfrak{a}_n$. Now, the first part of statement 4 follows from the definition of the set $\widetilde{{}'\Delta_n}$ and Lemma 3.6 where $S = \widetilde{{}'\Delta_n}$ and $T = {}'S(\mathbb{A}_n)$.

The second part of statement 4 follows from the first one.

5. Statement 5 follows from statement 4.

6. By statement 3, ${}'\Delta_n \in \mathrm{Den}_l(\mathcal{L}_n, 0)$ and ${}'\Delta_n^{-1}\mathcal{L}_n \simeq {}'Q(\mathbb{A}_n)$. Now, statement 6 follows from Lemma 3.10.(2) where $R = \Delta_n$, $R' = \mathcal{L}_n$ and $S = {}'\Delta_n$ (The R-module R'/R is S_∂-torsion. Hence it is also ${}'\Delta_n$-torsion as $S_\partial \subseteq {}'\Delta_n$).

7. By the definition, the subset ${}''\Delta_n$ of $\mathbb{A}_n$ is a multiplicative set such that ${}'\Delta_n \subseteq {}''\Delta_n \subseteq {}'S(\mathbb{A}_n)$, see statements 1 and 2. Hence, $\mathfrak{a}_n = \mathrm{ass}_l({}'\Delta_n) \subseteq \mathrm{ass}_l({}''\Delta_n) \subseteq \mathrm{ass}_l({}'S(\mathbb{A}_n)) = \mathfrak{a}_n$, and so $\mathrm{ass}_l({}''\Delta_n) = \mathfrak{a}_n = \mathrm{ass}_l({}'S(\mathbb{A}_n))$. Notice that $S_\partial \subseteq {}'\Delta_n \subseteq {}''\Delta_n$ and for each element $s \in {}''\Delta_n$ there is an element $\alpha \in \mathbb{N}^n$ such that $\partial^\alpha s \in {}'\Delta_n \subseteq {}''\Delta_n$. Now, statement 7 follows from Lemma 3.6 where $S = {}''\Delta_n$ and $T = {}'S(\mathbb{A}_n)$. □

In order to prove Theorem 5.4, we need the following two lemmas that are also interesting on their own.

Lemma 5.2. *1. Every nonzero ideal of the algebra $\mathbb{A}_n$ has zero left and right annihilator.*

2. Every nonzero ideal of the algebra $\mathbb{A}_n$ is an essential left and right ideal of $\mathbb{A}_n$.

Proof. The ideal F_n is the smallest nonzero ideal of the algebra $\mathbb{A}_n$. So, it suffices to prove statements 1 and 2 for the ideal F_n. Suppose that the left or right annihilator of F_n is a nonzero ideal of $\mathbb{A}_n$. Hence, it contains the idempotent ideal F_n, and so their product, which is the zero ideal, contains $F_n^2 = F_n \neq 0$, a contradiction.

Let I and J be left and right ideals of $\mathbb{A}_n$, respectively. By statement 1, $I \supseteq F_n I \neq 0$ and $J \supseteq J F_n \neq 0$, and statement 2 follows. □

Lemma 5.3 gives a characterization of left regular elements of the ring $\mathbb{A}_n$. By applying the involution θ, we obtain a similar characterization of right regular elements of the ring $\mathbb{A}_n$ ($a \in {}'\mathcal{C}_{\mathbb{A}_n}$ iff $\theta(a) \in \mathcal{C}'_{\mathbb{A}_n}$).

Lemma 5.3. *Let $a \in \mathbb{A}_n$. Then the following statements are equivalent:*

1. $a \in {}'\mathcal{C}_{\mathbb{A}_n}$.
2. $a \in {}'\mathcal{C}_{F_n}$.
3. $a \in {}'\mathcal{C}_{P'_n}$ *where* $P'_n = K[\partial_1, \ldots, \partial_n]$ *is the only simple, faithful, right* $\mathbb{A}_n$*-module.*
4. $\theta(a) \in \mathcal{C}'_{P_n}$ *where* $P_n = K[x_1, \ldots, x_n]$ *is the only simple, faithful, left* $\mathbb{A}_n$*-module.*

Proof. ($1 \Leftrightarrow 2$) The equivalence follows from Corollary 2.19.(1) and Lemma 5.2.(2).

($2 \Leftrightarrow 3$) The equivalence follows from Corollary 2.20.(1) and the fact that $(F_n)_{\mathbb{A}_n} \simeq (P'_n)^{(\mathbb{N}^n)}$ ([3, Corollary 2.7.(8)]).

($3 \Leftrightarrow 4$) The equivalence follows from the fact that $P'_n = \theta(P_n)$. □

Criterion for ${}'Q(\mathbb{A}_n) \simeq Q(A_n)$. The algebras Δ_n, B_n and $\mathcal{A}_n$ are Noetherian domains. By Goldie's Theorem, their quotient rings are division rings. It follows from the inclusions $A_n \subseteq B_n \subseteq \mathcal{A}_n \subseteq Q(A_n)$ and $\Delta_n \subseteq S_\partial^{-1}\Delta_n \simeq \mathcal{A}_n \subseteq Q(A_n)$ that

$$Q(\Delta_n) = Q(B_n) = Q(\mathcal{A}_n) = Q(A_n). \tag{40}$$

Theorem 5.4. *The following statements are equivalent:*

1. ${}'Q(\mathbb{A}_n) \simeq Q(A_n)$.
2. ${}'Q_{l,cl}(\mathbb{A}_n) \simeq Q(A_n)$.

3. *The set* $\overline{{}'S(\mathbb{A}_n)}$ *is left dense in* $\mathcal{A}_n\backslash\{0\}$.
4. *The set* $\overline{{}'\mathcal{C}_{\mathbb{A}_n}}$ *is left dense in* $\mathcal{A}_n\backslash\{0\}$.
5. *The set* ${}'\Delta_n$ *is left dense in* $\mathcal{A}_n\backslash\{0\}$.
6. *The set* ${}'\Delta_n$ *is left dense in* Δ_n^0.
7. *For each element* $s \in \Delta_n^0$, *there is an element* $s' \in \Delta_n^0$ *such that* $s's \in {}'\mathcal{C}_{F_n}$.
8. *For each element* $s \in \Delta_n^0$, *there is an element* $s' \in \Delta_n^0$ *such that* $s's \in {}'\mathcal{C}_{P'_n}$ *where* $P'_n = K[\partial_1, \dots, \partial_n]$ *is the unique simple faithful right* $\mathbb{A}_n$*-module.*

Proof. $(1 \Leftrightarrow 2 \Leftrightarrow 3 \Leftrightarrow 4)$ The equivalence $(1 \Leftrightarrow 2)$ follows from Theorem 3.12.(2). The equivalences $(1 \Leftrightarrow 3)$ and $(2 \Leftrightarrow 4)$ follows from Theorem 1.1 and Theorem 3.12.(2) (since $\mathrm{ass}_l({}'S(\mathbb{A}_n)) = \mathfrak{a}_n$ is a prime ideal of $\mathbb{A}_n$, the algebra $\mathbb{A}_n/\mathfrak{a}_n = \mathcal{A}_n$ is a domain and $Q(\mathbb{A}_n/\mathfrak{a}_n) = Q(\mathcal{A}_n)$ is a division ring).

$(1 \Leftrightarrow 5)$ By Theorem 5.1.(1), ${}'\Delta_n \in \mathrm{Den}_l(\mathbb{A}_n, \mathfrak{a}_n)$ and ${}'\Delta_n^{-1}\mathbb{A}_n \simeq {}'Q(\mathbb{A}_n)$. Since the set ${}'\Delta_n$ is dense in ${}'S(\mathbb{A}_n)$, the equivalence $(1 \Leftrightarrow 5)$ holds iff the equivalence $(1 \Leftrightarrow 3)$ holds.

$(5 \Leftrightarrow 6)$ Recall that ${}'\Delta_n, \Delta_n^0 \in \mathrm{Den}_l(\Delta_n, 0)$, ${}'\Delta_n \subseteq \Delta_n^0$ and $Q(\Delta_n) = Q(A_n)$, see (40). By [12, Lemma 3.5.(3)], ${}'\Delta_n^{-1}\Delta_n \simeq (\Delta_n^0)^{-1}\Delta_n = Q(A_n)$ iff the set ${}'\Delta_n$ is left dense in Δ_n^0.

$(6 \Leftrightarrow 7)$ By Lemma 5.3, the inclusion $s's \in {}'\mathcal{C}_{F_n}$ is equivalent to the inclusion $s's \in {}'\mathcal{C}_{\mathbb{A}_n}$. Hence, statement 7 is equivalent to the statement that for each element $s \in \Delta_n^0$, there is an element $s' \in \Delta_n^0$ such that $s's \in {}'\mathcal{C}_{\mathbb{A}_n}$, i.e. $s's \in \Delta_n \cap {}'\mathcal{C}_{\mathbb{A}_n} = {}'\Delta_n$, i.e. it is equivalent to statement 6.

$(7 \Leftrightarrow 8)$ The equivalence follows from Lemma 5.3. □

The ring ${}'Q(\mathbb{A}_1)$**.** As an application of Theorem 5.4 we obtain Theorem 5.5.

Theorem 5.5. ${}'Q(\mathbb{A}_1) \simeq Q(A_1)$.

Proof. (i) *A nonzero rational function* $\phi \in \mathcal{L}_1$ *belongs to the set* ${}'\mathcal{C}_{\mathbb{A}_1}$ *iff it has no root in the set* $\mathbb{N}_+ = \{1, 2, \dots\}$: The right $\mathbb{A}_1$-module F is an essential right ideal of the algebra $\mathbb{A}_1$ ([3, Corollary 2.7.(6)]). Therefore, $\phi \in {}'\mathcal{C}_{\mathbb{A}_1}$ iff the map $\cdot\phi_F : F \to F$, $f \mapsto f\phi$ is an injection. The right $\mathbb{A}_1$-module $F = \oplus_{i\in\mathbb{N}} E_{i0}\mathbb{A}_1 \simeq \left(E_{00}\mathbb{A}_1\right)^{(\mathbb{N})}$ is a direct sum of countably many copies of the right $\mathbb{A}_1$-module

$$E_{00}\mathbb{A}_1 = E_{00}F \simeq E_{00}K[\partial] \simeq K[\partial]_{K[\partial]} \text{ and } E_{00}\partial^i H = E_{00}\partial^i(i+1) \text{ for all } i \geq 0,$$

and the statement (i) follows.

(ii) *For each nonzero rational function* $\phi \in \mathcal{L}_1$, $\tau^i(\phi) \in {}'\mathcal{C}_{\mathbb{A}_1}$ *for all* $i \gg 1$ *where* $\tau(H) = H + 1$: The statement (ii) follows from the statement (i).

(iii) *For each nonzero element* $d \in \Delta_n$, $\partial^i d \in {}'\Delta_n$ *for all* $i \gg 1$: The element d is a unique sum $\phi_n\partial^n + \phi_{n+1}\partial^{n+1} + \cdots + \phi_m\partial^m$ where $\phi_i \in \mathcal{L}_1$ and $\phi_n \neq 0$. By the statement (ii), $\tau^i(\phi_n) \in {}'\mathcal{C}_{\mathbb{A}_1}$ for some $i \geq 0$. Therefore, the map $\cdot\tau^i(\phi_n) : F \to F$, $f \mapsto f\tau^i(\phi_n)$ is an injections. Hence, the map $\cdot\partial^i d : F \to F$, $f \mapsto f\partial^i d$ is also an injections since

$$\partial^i d = \tau(\phi_n)^i\partial^{n+i} + \cdots + \tau(\phi_m)^i\partial^{m+i} \text{ and } \tau(\phi_n)^i\partial^{n+i} \in {}'\mathcal{C}_{\mathbb{A}_1}.$$

Therefore, $\partial^i d \in {}'\mathcal{C}_{\mathbb{A}_1}$.

(iv) *The set* ${}'\Delta_1$ *is left dense in* $\mathcal{A}_1\backslash\{0\}$: Clearly, the set ${}'\Delta_1\backslash\{0\}$ is left dense in $\mathcal{A}_1\backslash\{0\}$. Now, the statement (iv) follows from the statement (iii).

The theorem follows from Theorem 5.4 and the fact that the set ${}'\Delta_1$ is left dense in $\mathcal{A}_1\backslash\{0\}$, the statement (iv). □

Corollary 5.6. $Q'(\mathbb{A}_1) \simeq Q(A_1)$.

Proof. The result follows from Theorem 5.5: $Q'(\mathbb{A}_1) = \theta({}'Q(\mathbb{A}_1)) \simeq \theta(Q(A_1) = Q(\theta(A_1)) = Q(A_1)$. □

Descriptions of the sets ${}'\mathcal{C}_{\mathbb{A}_1}$ and $\mathcal{C}'_{\mathbb{A}_1}$.

(i) ${}'\mathcal{C}_{\mathbb{A}_1} \subseteq \mathbb{A}_1\backslash F$: It is obvious that every element of the ideal $F = \oplus_{i,j\in\mathbb{N}} KE_{ij} \simeq M_\infty(K)$ is a left and right zero divisor of the algebra F (without 1) and of $\mathbb{A}_1$.

(ii) *For each nonzero element $d \in \mathbb{A}_1\backslash F$, $\partial^i d \in \Delta_1^0$ for some $i \in \mathbb{N}$*: The statement (ii) follows (31) and the equality $\mathbb{D}_1 = L \oplus (\oplus_{i,j\geq 1} Kx^i H^{-j}\partial^i)$.

(iii) *For each nonzero element $d \in \mathbb{A}_1\backslash F$, $\partial^i d \in {}'\mathcal{C}_{\mathbb{A}_1}$ for some $i \in \mathbb{N}$*: The statement (iii) follows from the statement (ii) and the statement (iii) of the proof of Theorem 5.5.

Then the well-defined map

$$d : \mathbb{A}_1\backslash F \to \mathbb{N}, \quad a \mapsto d(a) := \min\{i \in \mathbb{N} \,|\, \partial^i a \in {}'\mathcal{C}_{\mathbb{A}_1}\} \tag{41}$$

is called the *left regularity degree function* and the natural number $d(a)$ is called the *left regularity degree* of a. For each element $a \in \mathbb{A}_1\backslash F$, $d(a)$ can be found in finitely many steps, see the proof of Theorem 5.5. Now, Theorem 5.7.(1) follows. Then Theorem 5.7.(2) follows from Theorem 5.7.(1).

Theorem 5.7. *1.* ${}'\mathcal{C}_{\mathbb{A}_1} = \{\partial^{d(a)} a \,|\, a \in \mathbb{A}_1\backslash F\}$.

2. $\mathcal{C}'_{\mathbb{A}_1} = \theta({}'\mathcal{C}_{\mathbb{A}_1})$.

By (31), each element $a \in \mathbb{A}_1$ is a unique sum $a = \sum_{i=0}^{l} d_{-i}\partial^i + \sum_{j=1}^{m} x^j d_j + a_F$ where $d_k \in \mathbb{D}_1$, $k = -l, \ldots, m$ and $a_F \in F$. Let $a_\partial := \sum_{i=0}^{l} d_{-i}\partial^i$. The integer

$$s(a_F) := \begin{cases} \min\{n \in \mathbb{N} \,|\, a_F \in \bigoplus_{i,j=0}^{n} Ke_{ij}\} & \text{if } a_F \neq 0, \\ -1 & \text{if } a_F = 0. \end{cases}$$

is called the *size* of the element $a_F \in F$. The integer $s(a) := s(a_F)$ is called the *size* of the element a. For each $i \in \mathbb{N}$, let $P'_{1,\leq i} := \{a \in P'_1 \,|\, \deg_y(a) \leq i\}$ where $\deg_y \partial$ is the degree of the polynomial $a \in P'_1 = K[\partial]$ in the variable ∂. Let

$$L^\perp := \bigoplus_{i,j\geq 1} Kx^i H^{-j}\partial^i \;\text{ and }\; \Xi := \Big(\bigoplus_{i\geq 0} L^\perp\partial^i \oplus \bigoplus_{i\geq 1} x^i\mathbb{D}_1\Big) \bigcup \Big(\bigoplus_{i\geq 1} x^i\mathbb{D}_1 + F\Big).$$

Then $\mathbb{D}_1 = L \oplus L^\perp$. For a nonzero element $l^\perp = \sum_{i,j\geq 1} \lambda_{ij} x^i H^{-j}\partial^i \in L^\perp$, let

$$\phi(l^\perp) := \sum_{i,j\geq 1} \lambda_{ij} \frac{(H-i)(H-i+1)\cdots(H-1)}{(H-i)^j} \;\; \delta(l^\perp) := \max\{i \geq 1 \,|\, \lambda_{ij} \neq 0 \;\text{ for some }\; j \geq 1\}$$

and $\delta(0) := 0$. Then for all $k \geq \delta(l^\perp)$,

$$\partial^k l^\perp = \tau(\phi(l^\perp))\partial^k \;\text{ and }\; \tau(\phi(l^\perp)) \in L$$

where $\tau(H) = H + 1$. For each nonzero element $d = l + l^\perp \in \mathbb{D}_1 = L \oplus D^\perp$, where $l \in \Lambda$ and $l^\perp \in L^\perp$, let

$$\mu(d) := \min\{i \geq \delta(l^\perp) \,|\, \text{the rational function } \tau^i(l + \phi(l^\perp)) \in L \text{ has no root in } \mathbb{N}_+\}.$$

If $a = \sum_{i=0}^{n} d_{-i}\partial^i + \sum_{j=1}^{m} x^j d_j + a_F \in \mathbb{A}_1\backslash\Xi$ then $a_\partial \neq 0$ and so $d_{-n} \neq 0$ where $n = \deg_\partial(a_\partial)$. Let

$$\mu(a) := \mu(d_{-n}) \;\text{ and }\; \nu(a) := \max\{s(a), \mu(a)\}.$$

Theorem 5.8. *1.* ${}'\mathcal{C}_{\mathbb{A}_1} = \{a \in \mathbb{A}_1\backslash\Xi \,|\, \cdot a : P'_{1,\leq\nu(a)} \to P'_{1,\leq\nu(a)+\deg_y(a_y)},\ p \mapsto pa \text{ is an injection}\}$.

2. $\mathcal{C}'_{\mathbb{A}_1} = \theta({}'\mathcal{C}_{\mathbb{A}_1})$ *where θ is the involution of the algebra $\mathbb{A}_1$, see (32).*

Proof. 1. (i) ${}'\mathcal{C}_{\mathbb{A}_1} \cap \Xi = \emptyset$: Let $a \in \Xi$. We have to show that $a \notin {}'\mathcal{C}_{\mathbb{A}_1}$. Suppose that $a \in \bigoplus_{i\geq 0} L^{\perp}\partial^i \oplus \bigoplus_{i\geq 1} x^i\mathbb{D}_1$. Then $1 \in \ker(\cdot a)$, and so $a \notin {}'\mathcal{C}_{\mathbb{A}_1}$.

Suppose that $a \in \bigoplus_{i\geq 1} x^i\mathbb{D}_1 + F$. Then the map $\cdot a : P'_{1,\leq s(a)+1} \to P'_{1,\leq s(a)}$, $p \mapsto pa$ is a well-defined map. Since

$$\dim_K(P'_{1,\leq s(a)+1}) = s(a) + 2 > s(a) + 1 = \dim_K(P'_{1,\leq s(a)}),$$

$\ker(\cdot a) \neq 0$, $a \notin {}'\mathcal{C}_{\mathbb{A}_1}$. Therefore, ${}'\mathcal{C}_{\mathbb{A}_1} \cap \Xi = \emptyset$.

(ii) *For each element* $a \in \mathbb{A}_1\backslash\Xi$, $\ker_{P'_1}(\cdot a) \subseteq P'_{1,\leq \nu(a)}$: Since $a \in \mathbb{A}_1\backslash\Xi$, $a_y := \sum_{i=0}^{l} d_{-i}\partial^i \neq 0$ where $d_{-i} \in \mathbb{D}_1$. Suppose that $d_{-l} \neq 0$. Suppose that $p \in \ker_{P'_1}(\cdot a)\backslash P'_{1,\leq \nu(a)}$, i.e. $\deg_y(p) > \nu(a)$. Then

$$\deg(pa) = l + \deg_y(p),$$

a contradiction (since $pa = 0$).

Now, statement 1 follows from statements (i) and (ii).

2. Statement 2 follows from statement 1. □

Theorem 5.9. *1.* $(1-\sigma)^d(\phi) = d!\lambda_d$ *where* $(1-\sigma)^d := \prod_{i=1}^{n}(1-\sigma_i)^{d_i}$ *and* $d! = d_1!\cdots d_n!$.

2. $\sum_{\alpha\in\Pi_d} \mathcal{P}_n\sigma^{\alpha}(\phi) = \mathcal{P}_n$, *i.e.* $\bigcap_{\alpha\in\Pi_d} V(\sigma^{\alpha}(\phi)) = \emptyset$.

3. For every automorphism $\tau \in \mathrm{Aut}_K(\mathcal{P}_n)$, *the automorphisms* $\sigma'_1 = \tau\sigma_1\tau^{-1}, \ldots, \sigma'_n = \tau\sigma_n\tau^{-1} \in \mathrm{Aut}_K(\mathcal{P}_n)$ *commute and* $\sum_{\alpha\in\Pi_d} \mathcal{P}_n\sigma'^{\alpha}(\tau(\phi)) = \mathcal{P}_n$, *i.e.* $\bigcap_{\alpha\in\Pi_d} V(\sigma'^{\alpha}(\tau(\phi))) = \emptyset$.

Proof. 1. Since $\mathcal{P}_n = \mathcal{P}_{n-1} \otimes K[H_n]$, $\phi = \phi_{n-1}H_n^{d_n} + \psi_{n-1}H_n^{d_n-1} + \cdots$ where $\phi_{n-1}, \psi_{n-1}, \ldots \in \mathcal{P}_{n-1}$. Then

$$(1-\sigma_n)^{d_n}(\phi) = d_n!\phi_{n-1} \in \mathcal{P}_{n-1}$$

and the leading term of the polynomial $d_n!\phi_{n-1} \in \mathcal{P}_{n-1}$ is $d_n\lambda_d H_1^{d_1}\cdots H_{n-1}^{d_{n-1}}$. Now, the result follows by induction on n (or by repeating the above computation $n-1$ more times).

2. By statement 1, $K^{\times} \ni d!\lambda_d = (1-\sigma)^d(\phi) \in \sum_{\alpha\in\Pi_d} \mathcal{P}_n\sigma^{\alpha}(\phi)$, and statement 2 follows.

3. Clearly, the automorphisms $\sigma'_1, \ldots, \sigma'_n$ commute and

$$\mathcal{P}_n = \tau(\mathcal{P}_n) = \tau\Big(\sum_{\alpha\in\Pi_d} \mathcal{P}_n\sigma^{\alpha}(\phi)\Big) = \sum_{\alpha\in\Pi_d} \mathcal{P}_n\sigma'^{\alpha}(\tau(\phi)).$$

□

Corollary 5.10. *Let* $\mathcal{P}_n = K[H_1, \ldots, H_n]$ *be a polynomial algebra over a field* K *of characteristic zero and* $\sigma_i \in \mathrm{Aut}_K(\mathcal{P}_n)$ *where* $\sigma_i(H_j) = H_j - \mu_i\delta_{ij}$ *for* $i, j = 1, \ldots, n$, $\mu_i \in K^{\times}$ *and* δ_{ij} *is the Kronecker delta. Let* $\phi \in \mathcal{P}_n\backslash\{0\}$ *and* $\lambda_d H^d$ *be the leading term of the polynomial* ϕ *with respect to the lexicographic ordering* $H_1 < \cdots < H_n$ *where* $\lambda_d \in K^{\times}$ *and* $d = (d_1, \ldots, d_n) \in \mathbb{N}^n$. *Then:*

1. $(1-\sigma)^d(\phi) = d!\mu^d\lambda_d$ *where* $(1-\sigma)^d := \prod_{i=1}^{n}(1-\sigma_i)^{d_i}$, $d! = d_1!\cdots d_n!$ *and* $\mu^d := \mu_1^{d_1}\cdots\mu_n^{d_n}$.

2. $\sum_{\alpha\in\Pi_d} \mathcal{P}_n\sigma^{\alpha}(\phi) = \mathcal{P}_n$, *i.e.* $\bigcap_{\alpha\in\Pi_d} V(\sigma^{\alpha}(\phi)) = \emptyset$.

Proof. Repeat the proofs of statement 1 and 2 of Theorem 5.9 and making an obvious adjustments. □

Licence. For the purpose of open access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author Accepted Manuscript version arising from this submission.

Disclosure statement. No potential conflict of interest was reported by the author.

Data availability statement. Data sharing not applicable – no new data generated.

References

[1] F. Abadie, M. Dokuchaev and R. Exel, Strong equivalence of graded algebras, *J. Algebra* **659** (2024), 818–858.

[2] M. Barkatou and T. Cluzeau, Thomas, On the computation of rational solutions of linear integro-differential equations with polynomial coefficients, *J. Symbolic Comput.* **121** (2024), Paper No. 102252, 19 pp.

[3] V. V. Bavula, The Jacobian algebras, *J. Pure Appl. Algebra* **213** (2009), 664-685; Arxiv:math.RA/0704.3850.

[4] V. V. Bavula, The algebra of one-sided inverses of a polynomial algebra, *J. Pure Appl. Algebra*, **214** (2010), 1874–1897; ArXiv:math.AG/0903.0641.

[5] V. V. Bavula, The algebra of integro-differential operators on a polynomial algebra, *Journal of the London Math. Soc.*, **83** (2011) no. 2, 517-543; ArXiv:math.RA/0912.1723.

[6] V. V. Bavula, The group of automorphisms of the algebra of polynomial integro-differential operators, *J. of Algebra*, **348** (2011) 233–263; Arxiv:math.AG/ 0912.2537.

[7] V. V. Bavula, The algebra of integro-differential operators on an affine line and its modules, *J. Pure Appl. Algebra*, **217** (2013) 495-529. Arxiv:math.RA: 1011.2997.

[8] V. V. Bavula, The largest strong left quotient ring of a ring, *J. Algebra*, **439** (2015) 1-32.

[9] V. V. Bavula, New criteria for a ring to have a semisimple left quotient ring, *Journal of Alg. and its Appl.*, **14** (2015) no. 6, 1550090, 28pp; Arxiv:math.RA:1303.0859.

[10] V. V. Bavula, The group of automorphisms of the algebra of one-sided inverses of a polynomial algebra, II, *Proceedings of the Edinburgh Math. Soc. (Series 2)*, **58** (2015) no. 3, 543-580; Arxiv:math.AG:0906.3733.

[11] V. V. Bavula, The largest left quotient ring of a ring, *Comm. in Algebra*,**44** (2016) no. 8, 3219-3261. Arxiv:math.RA:1101.5107.

[12] V. V. Bavula, The classical left regular left quotient ring of a ring and its semisimplicity criteria, *J. Algebra Appl.*, **16** (2017), no. 6, 1750103, 30 pp. Arxiv:math.RA:1504.07257.

[13] V. V. Bavula, Localizable sets and the localization of a ring at a localizable set, *J. Algebra*, **610** (2022), no. 15, 38–75. (ArXiv:2112.13447).

[14] V. V. Bavula, General theory of localizations of rings and modules, ArXiv:2312.17145.

[15] R. Baer, A unified theory of projective spaces and finite Abelian groups, *Trans. Amer. Math. Soc.*, **52** (1942) 283—343.

[16] H. Cartan and S. Eilenberg, Homological algebra, Princeton University Press, 1956.

[17] L. Guo, G. Regensburger and M. Rosenkranz, On integro-differential algebras, *J. Pure Appl. Algebra* **218** (2014), no.3, 456-–473.

[18] A. Haghany and A. Kassaian, Study of the algebra of smooth integro-differential operators with applications., *J. Algebra Appl.* **18** (2019), no. 1, 1950009, 27 pp.

[19] I. Kaplansky, Infinite Abelian groups, University of Michigan Press, 1954.

[20] P. A. Krylov and A. A. Tuganbaev, Modules over discrete valuation domains. I, *Journal of Mathematical Sciences*, **145** (2007) no. 4, 4997–5115.

[21] W. Liebert, Characterization of the endomorphism rings of divisible torsion modules and reduced complete torsion-free modules over complete discrete valuation rings, *Pacific J. Math.*, **37** (1971) no. 1, 141–170.

[22] J. C. McConnell and J. C. Robson, Noncommutative Noetherian rings, Wiley, Chichester, 1987.

[23] W. S. Martindale, Prime rings satisfying a generalized polynomial identity, *J. Algebra*, **12** (1969) 576-584.

[24] E. Matlis, Injective modules over Noetherian rings, *Pacific J. Math.*, **8** (1958) no. 3, 511–528.

[25] H. Qiu, S. Zhengand and Y. Dan, Construction of free quasi-idempotent differential Rota-Baxter algebras by Gröbner-Shirshov bases, *Comm. Algebra* **52** (2024), no.6, 2404–2421.

[26] C. Raab and G. Regensburger, The fundamental theorem of calculus in differential rings. *Adv. Math.* **447** (2024), Paper No. 109676, 50 pp.

[27] Y. Wang, M. Li and D. Wu, K-comultiplication semimodules. *Comm. Algebra* **52** (2024), no. 6, 2670–2691.

V. V. Bavula
School of Mathematical and Physical Sciences
Division of Pure Mathematics
University of Sheffield
Hicks Building
Sheffield S3 7RH
UK
email: v.bavula@sheffield.ac.uk