



This is a repository copy of *Measuring vertebrate telomeres: applications and limitations* .

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/353/>

Article:

Nakagawa, S., Gemmell, N.J. and Burke, T. (2004) Measuring vertebrate telomeres: applications and limitations. *Molecular Ecology*, 13 (9). pp. 2523-2533. ISSN 0962-1083

<https://doi.org/10.1111/j.1365-294X.2004.02291.x>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

INVITED REVIEW

Measuring vertebrate telomeres: applications and limitations

SHINICHI NAKAGAWA,* NEIL J. GEMMELL† and TERRY BURKE*

*Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK, †Department of Biological Sciences, University of Canterbury, Private Bag 4800, Christchurch, New Zealand

Abstract

Telomeres are short tandem repeated sequences of DNA found at the ends of eukaryotic chromosomes that function in stabilizing chromosomal end integrity. *In vivo* studies of somatic tissue of mammals and birds have shown a correlation between telomere length and organismal age within species, and correlations between telomere shortening rate and lifespan among species. This result presents the tantalizing possibility that telomere length could be used to provide much needed information on age, ageing and survival in natural populations where longitudinal studies are lacking. Here we review methods available for measuring telomere length and discuss the potential uses and limitations of telomeres as age and ageing estimators in the fields of vertebrate ecology, evolution and conservation.

Keywords: age, ageing, biological age, chronological age, estimator, telomere length, telomere shortening

Received 28 April 2004; revision received 22 June 2004; accepted 22 June 2004

Introduction

We all know that age affects the way organisms function and is critically important in determining an individual's physiological state and response to environmental conditions (Ricklefs 1977; Stearns & Koella 1986; Stearns 1989; Lindstrom 1999; Ricklefs & Wikelski 2002). Age affects risk of injury and disease, probability of survival and reproductive potential, and is a major determinant of population growth potential (Stearns & Koella 1986; Albon *et al.* 2000; Coulson *et al.* 2001; Kern *et al.* 2001). Despite its importance, age structure is seldom incorporated into population models, yet we know that knowledge of age provides important insights into population and evolutionary processes that would otherwise remain hidden (e.g. how reproductive output, offspring sex and offspring viability vary with age) (Stearns & Koella 1986; Stearns 1989; Charnov 1991; Lindstrom 1999; Kern *et al.* 2001).

In most natural populations the age of individuals is rarely known because these data are impractical or, as yet, impossible to obtain. Extensive longitudinal study of marked

individuals provides one means of obtaining such data, but generally these data come at significant cost and cover only a fraction of the study population. Morphological correlates of age may also be useful, but these approaches (e.g. counting otolith growth rings in fish, measuring wax plugs from the inner ears of whales) often require destructive sampling. Recently it has been suggested that telomere length may provide a powerful new tool for estimating age in natural populations where longitudinal data are limited (Hausmann & Vleck 2002; Vleck *et al.* 2003).

Telomeres are short tandem repeated sequences of DNA found at the ends of eukaryotic chromosomes that function in stabilizing chromosomal end integrity (Box 1, Fig. 1). Hausmann and colleagues (Hausmann & Vleck 2002; Hausmann *et al.* 2003a; Vleck *et al.* 2003) have demonstrated recently that it might be possible to estimate the age of animals by measuring telomere lengths because telomeres in somatic tissues normally shorten with age. They have shown that telomere (or terminal) restriction fragments (TRF) from blood samples in some avian species shortened with age [e.g. zebra finches, *Taeniopygia guttata* (Hausmann & Vleck 2002) and common terns, *Sterna hirundo* (Hausmann *et al.* 2003a)]. They also found that the rate of telomere shortening (telomere length rate of change; TROC) was

Correspondence: Neil J. Gemmell. Fax: +64 3364 2590; E-mail: neil.gemmell@canterbury.ac.nz

Box 1. Telomeres, factors regulating telomere length, and aging

Telomeres comprise many copies of an evolutionarily conserved DNA repeat, found at the natural ends of all linear eukaryotic chromosomes. The repeats consist of a short G-rich sequence; in vertebrates, the telomeric repeat (TTAGGG)_n is conserved (Meyne *et al.* 1989) (Fig. 1). This natural end, along with associated proteins, provides chromosome stability, preventing degradation and chromosome fusion (while also anchoring chromosomes in the nuclear matrix); chromosome ends created by breakage are prone to fuse with other chromosomes (Blackburn 1991; Greider 1996). Telomere-associated proteins such as TRF1 and TRF2 are also involved in the regulation of telomere length (Fig. 1) (Broccoli *et al.* 1997; van Steensel & de Lange 1997; Karlseder *et al.* 2002). Telomeres play an essential part in DNA replication. At each cell division, a small number of telomeric repeats is lost because DNA replication is incomplete at the 3' end of the double strands (i.e. the end-replication problem) (Watson 1972), leaving G-rich strand overhangs [whose length is a determinant in the rate of telomere shortening (Huffman *et al.* 2000)].

Telomerase is a ribonucleoprotein reverse transcriptase that restores telomere repeats (Lingner *et al.* 1997). Tel-

omerase activity is found in continually proliferating germ cells and stem cells, but for other types of somatic cell, various or zero activity levels are observed in different species (Prowse & Greider 1995; Venkatesan & Price 1998). The shortening of telomeres has been suggested to be one of the main mechanisms underlying ageing and age-related diseases, because loss of telomere function can lead to genome instability and cell replicative senescence (Harley *et al.* 1992; Campisi 1996). Mechanistic explanations of ageing involve the accumulation of mutations in genes, the shortening of telomeres and damage in mitochondrial DNA, in all of which oxidative damage plays a crucial role (Goyns 2002) [for free radical theories of ageing, see Finkel & Holbrook (2000)]. Oxidative stress incurred by reactive oxygen species (ROS) increases the rate of telomere shortening per cell division and also the rate of cell turnover (Fig. 1).

Interestingly, in humans, telomere shortening (Benetos *et al.* 2001) occurs at different rates in males and females, presumably as a response to different levels of oestrogen between the sexes, which can stimulate telomerase activity and attenuate ROS (Aviv 2002b). Sex differences in telomere shortening have not been observed in any other species, but the possibility needs to be considered if telomere length is to become a tool useful for molecular ecological studies.




Fig. 1 Vertebrate telomere structure (region codes are used in Table 1) and factors regulating telomere length: direct factors (large arrows), indirect factors (small arrows), factors accelerating telomere shortening (black arrows) and factors attenuating telomere shortening (grey arrows). TRF1 and TRF2 are telomere-associated proteins and ROS (reactive oxygen species) are the main source of oxidative stress. Adapted and modified from Aviv (2002b) and Sedivy *et al.* (2003).

correlated with lifespan (also known as a maximum lifespan) in birds and mammals (Hausmann *et al.* 2003b; Vleck *et al.* 2003) (Box 2). They suggested that variation in telomere length rate of change (TROC) might be a molecular mechanism underlying the evolution of species lifespan (or ageing), which appear to be species-specific and genetically determined (Campisi 1996; Turker & Martin 1998).

Inevitably, for many wild populations information on age is lacking, as the time and effort usually required to determine individual ages and age structure in study populations is often prohibitive. Nevertheless, the insight that such knowledge provides into the ontogeny of life history traits and population dynamics is recognized widely (Coulson *et al.* 2001; Vleck *et al.* 2003). Telomere length assays may provide a direct and cost-effective approach to obtaining immediate knowledge of age, age structure and ageing in wild populations and represent a major leap forward in the molecular arsenal at the disposal of ecologists. This technology may prove to be as revolutionary as earlier DNA-based approaches to parentage analysis (Jones & Ardren 2003) and molecular sexing (Ellegren & Sheldon 1997). We anticipate that just like these earlier approaches, telomere

Box 2. Telomere length, telomere length rate of change and lifespan in birds and mammals

Maximum telomere length and the telomere length rate of change (TROC) differ among species, as does maximum lifespan among species (Table 2 and Fig. 2). Haussmann and colleagues (Haussmann *et al.* 2003b; Vleck *et al.* 2003) found that telomere length at a given life stage did not correlate with lifespan but TROC correlated with lifespan in birds and mammals (Fig. 2c). Telomere length and TROC vary among individuals of the same species and among tissues from an individual (Table 1). Inter-species, interindividual and intertissue differences are accounted for mainly by factors such as different rates of cell replication, levels of telomerase activity and levels of oxidative stress (Fig. 1) (Aviv 2002b). It is particularly interesting that the telomeres of Leach's storm-petrels (*Oceanodroma leucorhoa*), an unusually long-lived species, lengthen with age (Fig. 2b), because lengthening of telomeres (i.e. telomerase activity) is associated usually with cancer [cancer cells are immortalized through telomerase activity or alternative

processes that lengthen telomeres (Henson *et al.* 2002; Blasco 2003)]. It is proposed that the low or zero telomerase activity found in somatic cells may have been selected for to reduce the frequency of cancer (Harley *et al.* 1994). How Leach's storm-petrels avoid the tumour-susceptibility imposed by telomerase activity is of considerable potential interest in medical research (Haussmann *et al.* 2003b).

Despite the apparently adverse conditions promoting ageing in birds, such as high oxygen consumption and high body temperature, avian species tend to live longer than mammals of comparable body sizes. For a given body size, birds produce less reactive oxygen species (ROS), which are one of the main sources of oxidative damage, and are more tolerant to oxidative damage than mammals (Ogburn *et al.* 2001). Also, in several mammalian species the production of ROS and levels of oxidative damage are higher in short-lived species than long-lived ones (Barja & Herrero 2000). Clarification of the relationships between telomerase activity and both oxidative damage and lifespan in a range of avian species is currently under investigation (Vleck *et al.* 2003).

Table 1 Selected methods available for measuring telomeric regions

Method ^a	Hybridization ^b or PCR-based method?	No. of cells required ^c	Telomeric regions measured ^d	Appropriate for blood samples?	Average of single or all sample cells?	Average of single or all chromosomes?	Refs
TRF analysis	Hybridization	Large	2, 3, 4 (1) ^e	Yes	All	All	Harley <i>et al.</i> (1990)
Q-PCR	PCR-based	Small	2, 3, 4	Yes	All	All	Cawthon (2002)
Q-FISH	Hybridization	Intermediate	2, 3, 4	No	Single	Both	Zijlmans <i>et al.</i> (1997)
Flow-FISH	Hybridization	Intermediate	2, 3, 4	Yes	Single	All	Rufer <i>et al.</i> (1998)
STELA	PCR-based	Small	2, 3 (1) ^e	Yes	All	Single	Baird <i>et al.</i> (2003)
T-OLA	Hybridization	Large	4	Yes	All	All	Cimino-Reale <i>et al.</i> (2001)

^aAbbreviations: terminal (or telomere) restriction fragment (TRF), quantitative fluorescence *in situ* hybridization (Q-FISH), flow cytometry method using fluorescence *in situ* hybridization (Flow-FISH), telomeric-oligonucleotide ligation assays (T-OLA), real time kinetic quantitative polymerase chain reaction (Q-PCR), single telomere length analysis (STELA).

^bMethods using probes hybridize TTAGGG-repeat sequence.

^cCell quantities for assays are taken from Sedivy *et al.* (2003). As a guide, TRF requires 5–10 µg of DNA, while STELA requires 50–100 ng of DNA.

^dRegion codes are from Fig. 1: 1 = subtelomeric, 2 = imperfect telomeric repeats, 3 = telomeric repeats, 4 = G-rich single-stranded 3'-overhang.

^ePart of the region 1 (subtelomeric region) is measured.

length assays will open up new avenues and research opportunities in the fields of ecology, evolution and conservation.

In this article, we review the methods available for measuring telomere length and TROC and explore the use and limitations of vertebrate telomere assays in the field of ecology, evolution and conservation.

Measuring telomere length and TROC

Telomere restriction fragment assays

There are a number of techniques available for measuring telomere length (Table 1). The telomere (terminal) restriction

Table 2 Lifespan, telomere length, and telomere length rate of change (TROC) in birds and mammals from selected published literature^a

Species	Maximum lifespan (years) ^b	Maximum observed telomere length (bp) ^c	TROC (bp/year)	Tissue sampled	Refs
Zebra finch (<i>Taeniopygia guttata</i>)	5	9300	-515	Erythrocyte	Hausmann <i>et al.</i> (2003b)
Tree swallow (<i>Tachycineta bicolor</i>)	11	17 300	-391	Erythrocyte	Hausmann <i>et al.</i> (2003b)
Adélie penguin (<i>Pygoscelis adeliae</i>)	20	9500	-235	Erythrocyte	Hausmann <i>et al.</i> (2003b)
Common tern (<i>Sterna hirundo</i>)	26	9800	-57	Erythrocyte	Hausmann <i>et al.</i> (2003b)
Leach's storm-petrel (<i>Ocenodroma leucorhoa</i>)	36	20 000	75 ^d	Erythrocyte	Hausmann <i>et al.</i> (2003b)
Western wild mice (<i>Mus spretus</i>)	3.5	9500	-600	Spleen	Coviello-McLaughlin & Prowse (1997)
Cattle (<i>Bos taurus</i>)	30	22 000	-230	Leucocytes	Miyashita <i>et al.</i> (2002)
Cynomolgus monkey (<i>Macaca fascicularis</i>)	37	16 500	-63	Leucocytes	Lee <i>et al.</i> (2002)
Human (<i>Homo sapiens</i>)	110	10 000	-33	Leucocytes	Hastie <i>et al.</i> (1990)
Human (<i>Homo sapiens</i>)	110	10 500	-15	Fibroblasts	Allsopp <i>et al.</i> (1992)
Human (<i>Homo sapiens</i>)	110	9000	-68	Stem cells	Vaziri <i>et al.</i> (1994)
Human (<i>Homo sapiens</i>)	110	20 000	71 ^d	Sperm	Allsopp <i>et al.</i> (1992)

^aModified from Hausmann *et al.* (2003b).

^bFrom Hausmann *et al.* (2003b) and references therein.

^cThese are approximate values observed in the study and the values come from different age stages (mostly at the age of zero).

^dTelomere length increased with age in these cells.

fragment (TRF) analysis (Harley *et al.* 1990) and related methods such as the telomere amount and length assay (TALA) (Gan *et al.* 2001) are relatively easy methods and are probably the most widely used in telomere research. In these methods, average lengths of TRFs (created by particular restriction enzymes and hybridized with a radioactive oligonucleotide) are measured (see Harley *et al.* 1990). One major drawback of this approach is that the lengths of TRFs usually differ between cells and chromosomes because cells may differ in age and different chromosomes have different restriction sites relative to telomeric ends. Consequently, TRF assays produce autoradiographic smears which have some element of subjectivity in their analyses, although there is software available to remove some arbitrary decision making (Grant *et al.* 2001). Furthermore, the TRF method requires relatively large amounts of DNA (10 µg) and time, and TRF length differences between individuals can differ as much as 5%, depending on which restriction enzyme is used (indicating that subtelomeric restriction site polymorphisms and/or subtelomeric length polymorphisms may exist) (Cawthon 2002).

Quantitative polymerase chain reaction (PCR) assays

Real-time kinetic quantitative PCR (Q-PCR) measurement of telomere length (Cawthon 2002) avoids many of the problems encountered by TRF analysis by using ingeniously designed primers that hybridize to vertebrate telomeric regions without generating primer dimer-derived products. This method measures relative telomere lengths by determining the factor by which a sample DNA differs from an arbitrary reference DNA in its ratio of telomere repeat copy number (T) to single gene copy number (S) (i.e. relative T/S ratio). The T/S ratio of one individual relative to the T/S ratio of another should reflect relative telomere length differences between individuals. This method is very useful for determining interindividual differences in telomere length within a species and may also be used to measure relative telomere length among species if those species share the same single copy reference sequence. In theory, any single copy gene sequence can be used for standardization, so the technique should be applicable to species even where genetic data are limiting.




Fig. 2 Telomere length (measured by telomere restriction fragment length) as a function of age (a) in tree swallows (slope = -391 ± 65 (SE) bpy^{-1} , $r^2 = 0.34$) and (b) in Leach's storm-petrels (slope = 75 ± 10 (SE) bpy^{-1} , $r^2 = 0.66$). (c) The telomere length rate of change as a function of maximum lifespan in birds (slope = 19.5 ± 1.2 (SE) bpy^{-1} , $r^2 = 0.99$): zebra finches (black circle), tree swallows (triangle), Adélie penguins (square), common terns (diamond) and Leach's storm-petrels (open circle). Adapted from Haussmann *et al.* (2003b).

Despite its strengths, Q-PCR has two problems: (1) it does not provide absolute telomere lengths and (2) if target species have interstitial telomeric sequences [birds have telomeric repeats in the centromeric region (Venkatesan & Price 1998)], Q-PCR will measure interstitial telomeric

repeats as well as terminal telomeric repeats. The former problem is resolved by using both TRF analysis (Harley *et al.* 1990) and Q-PCR for a portion of the samples so that the change in length in base pairs (bp) of DNA per unit change in the T/S ratio can be determined, but the latter will be problematic if there is large variation in interstitial telomeric sequences among individuals or species. However, this is not a critical issue if one is interested in measuring only telomere length rate of change, because the extent of interstitial telomeric repeats should not change. Furthermore, it may be possible to estimate the extent of interstitial telomeric sequence by using Q-PCR in combination with TRF analysis and Bal31 exonuclease digestion, which will preferentially digest terminal telomeric sequences.

Fluorescence in situ hybridization (FISH) methods

Quantitative fluorescence *in situ* hybridization (Q-FISH) (Zijlmans *et al.* 1997) and flow cytometry methods using fluorescence *in situ* hybridization (Flow-FISH) (Rufer *et al.* 1998) are also often used to measure telomere length. As their name suggests, fluorescent dyes are used to visualize telomeres which are quantified by measuring light intensity. The FISH methods are able to measure telomere lengths of single cells. Flow-FISH uses fluorescence-activated cell sorting (FACS) to sort different types of cells, while Q-FISH provides measurements of telomere lengths of individual chromosomes. There is some evidence that the shortest telomere, rather than the average, is critical for loss of telomere function (Hemann *et al.* 2001), although average length might correlate with shortest telomere length. However, because Q-FISH visualizes telomeres in metaphase chromosomes tissue types used need to be proliferating cells, which limits its suitability for studies of wild populations, as blood or tissues would need to be collected and processed with a view to generating adequate numbers of metaphase chromosomes. Furthermore, the FISH methods, especially Q-FISH (Sedivy *et al.* 2003), are difficult to establish in nonspecialist laboratories so we think that there will be few cases where the high-resolution approaches of the FISH methods would be applied in ecological, evolutionary and conservational work.

Single telomere length analysis (STELA)

STELA (single telomere length analysis), a recently devised PCR-based method that measures the lengths of individual telomeres (Baird *et al.* 2003), also merits consideration. Baird *et al.* (2003) reported marked telomere length differences between alleles (some are much shorter than others) in human sex chromosomes and speculated that these differences might account for significant differences in the rate of ageing among human populations (Perls *et al.* 2002). Currently, this method is not readily applicable to most

species because the DNA sequences of subtelomeric regions need to be known. However, with sequence information coming on line from a host of genome projects, this approach will soon be applicable to primates, rodents, ungulates, carnivores, galliform birds and frog species related closely to *Xenopus*. We think that a generalized STELA assay for vertebrate telomeres might not be far away.

Telomeric-oligonucleotide ligation assays (T-OLA)

T-OLA measure the length of the G-rich telomeric-3'-overhang by ligating telomeric repeat oligonucleotides hybridized to the overhang (Cimino-Reale *et al.* 2001). Telomere shortening is proportional to the size of G-rich overhangs in human tissues (Huffman *et al.* 2000), and this observation might extend to other vertebrates. Although it has yet to be tested explicitly, given the correlation between TROC and lifespan (Hausmann *et al.* 2003b), it might be expected that long-lived species (i.e. species with slow telomere shortening) will have short G-rich overhangs while short-lived species will have long G-rich overhangs (Huffman *et al.* 2000). However, the abundance of short overhangs may also be proportional to the number of cells that have entered cellular senescence (Stewart *et al.* 2003). Consequently, it is currently unclear whether short overhangs mean slow telomere shortening (perhaps predicting longer lifespans) or many senescent cells (perhaps predicting shorter remaining life).

Sampling for telomere assays

In order for telomere assays to become a common tool in vertebrate ecology we will need to be able to sample somatic cells that are subject to constant division, and to possess a widely applicable method which can be used to measure telomere length efficiently in many individuals. Hausmann & Vleck (2002) suggest blood as an excellent candidate tissue for two reasons: (1) blood samples can be obtained easily from a wide variety of species and are often collected for other reasons and (2) telomeres in blood cells may shorten at a greater rate than those of other tissues (Hastie *et al.* 1990) because of their high turnover rate (Chang & Harley 1995). In nonmammalian vertebrates, blood samples of 50 µl may contain adequate numbers of nucleated red blood cells, from which telomere length can be obtained. For mammals, whose erythrocytes lack nuclei, telomere length can be measured in white blood cells, albeit with a need to draw larger blood volumes to account for the smaller numbers of white vs. red blood cells. However, there is no reason to restrict ourselves only to the use of blood samples; tissue biopsy samples taken routinely for many large vertebrates (e.g. Lambertsen 1987; Gemmell & Majluf 1997) might well prove to be adequate for telomere analyses.

Telomere shortening among and within species

Hausmann and colleagues (Hausmann *et al.* 2003b; Vleck *et al.* 2003) demonstrated that the rate of shortening varies among species, and that lifespan correlates with TROC in birds and mammals (Box 2). They suggest that lifespan and regulation of telomere length may have coevolved so that the replicative lifespan of individual somatic cell lineages increases with lifespan (Vleck *et al.* 2003). There are two major evolutionary theories of ageing: the antagonistic pleiotropy model and the mutation accumulation model (Rose 1991; Kirkwood & Austad 2000; Partridge & Gems 2002). The former model predicts that ageing has evolved as a result of pleiotropic alleles which are beneficial for survival and reproduction early in life, but that are detrimental for survival and reproduction later in life, being able to accumulate in populations. The latter model predicts that ageing has evolved as a result of alleles with detrimental effects being able to accumulate in populations if they are expressed only later in life when selection pressure is weak. Both theories predict that if the extrinsic rate of mortality increases, selection against senescence will be weak and biological ageing will accelerate (so decreasing lifespan). It is possible that TROC may be, in part, a molecular mechanism that contributes to the rate of ageing and thus of lifespan, as suggested by Hausmann *et al.* (2003b). Examination of TROC in more species will help to clarify this hypothesis.

Differences in TROC among individuals of the same species have so far had very little attention. If telomere length were seen as an age estimator, TROC could be used to estimate the rate of ageing, which may be compared with other life history traits. Potentially, we could ask interesting questions regarding life history trade-offs — for example, is the rate of telomere shortening higher in hard-working parents than lazy ones (i.e. is there some trade-off between energy expenditure and ageing)? TROC may well be different among populations of the same species living in different environmental circumstances, which may in turn influence the level of reactive oxygen species (ROS) (Fig. 1). Such a scenario would enable the investigation of the effects of pollutants or climate on ageing — for example, is TROC different between populations in and around Chernobyl when compared to other sites?

Recent studies, however, suggest that the rate of telomere shortening changes over time (Frenck *et al.* 1998; Rufer *et al.* 1999; Zeichner *et al.* 1999), which may impose some limitations on studies comparing TROC among individuals. For example, it has been reported in humans that telomere shortening in leucocytes occurs rapidly in young children, followed by an apparently stable period (little shortening) between age 4 years to young adulthood, and by gradual shortening in later life (Frenck *et al.* 1998; but see Rufer *et al.* 1999; Zeichner *et al.* 1999). It is still not clear

whether differences in cell turnover rate or in the regulation of telomere length over different stages of life (or both) contribute to this change over time (Frenck *et al.* 1998; Rufer *et al.* 1999; Zeichner *et al.* 1999; Friedrich *et al.* 2001; Brummendorf *et al.* 2002; Sidorov *et al.* 2003). It is likely that the change in telomere shortening over time differs considerably among species.

Estimating chronological or biological age?

Telomere length as an age estimator is promising, but it is not without problems. There is great variation in telomere length among individuals of the same age in a number of bird species (Hausmann & Vleck 2002; Hausmann *et al.* 2003a,b). Similar variation has been described in some mammalian species [e.g. humans, *Homo sapiens* (Hastie *et al.* 1990; Allsopp *et al.* 1992) and western wild mice, *Mus spretus* (Coviello-McLaughlin & Prowse 1997)]. The variation in telomere length results from a combination of variation in initial telomere length and thereafter in TROC. Telomere length therefore correlates with chronological age but does not always predict it reliably (e.g. a 1-year-old tree swallow had a shorter telomere length than one 6 years old; Fig. 2). Also, the reported variations in TROC suggest further limitations. For example, telomeres lengthen with age, albeit linearly, in Leach's storm-petrels, *Oceanodroma leucorhoa* (Box 2). In some species telomere length may change very little or not at all over an entire lifetime, as observed in the European freshwater turtle *Emys orbicularis* (living more than 100 years in captivity), which showed no significant change in the telomere length between embryos and adults (Girondot & Garcia 1998).

Species suitable for estimating chronological ages are probably those with a large change in telomere length over their lifetime and small variation in initial telomere length. For example, in humans accurate age estimation could be achieved from blood and dental pulp samples for forensic purposes (Tsuji *et al.* 2002; Takasaki *et al.* 2003). If such approaches can satisfy the rigorous requirements of forensic science, then there may be other cases in which chronological age estimation by telomere length may find a role. However, for ecological and evolutionary purposes, telomere length might be better employed as an indicator of 'biological age' than as an estimator of chronological age (Aviv 2002a; Harley *et al.* 1992).

Biological age is expressed by variation in lifespan among species and individuals of the same species. Telomere length reflects genome stability (Blackburn 1991; Greider 1996) and is highly heritable in humans (Slagboom *et al.* 1994; Jeanclous *et al.* 2000). As telomeres shorten at each cell division, telomeres are supposed effectively to record the replicative history of somatic cells in vertebrates (Aviv 2002a,b; Goyns 2002). Moreover, recent work has shown that oxidative damage (i.e. reactive oxygen species, ROS) may

be the main cause of telomere shortening (von Zglinicki *et al.* 2000) (Fig. 1). Oxidative stress is incurred both from the environment and from endogenous metabolic processes (Goyns 2002), so that telomere length reflects deleterious environmental factors to which an organism has been exposed, as well as genetic factors. Variation in longevity in the members of the same species may therefore be explained in part by biological age, as revealed through telomere length. Support for the proposition that this concept of biological age may be more important than chronological age comes again from humans, in which Cawthon *et al.* (2003) demonstrated that individuals who had shorter than average telomeres at the age of 60 or older suffered more from diseases of old age later in their lives.

Telomere assays of biological age as a tool in ecology

The direct use of biological age as a trait in ecological, evolutionary and conservational studies may prove particularly fruitful. For example, in Soay sheep older sheep respond differently to weather and density affects than younger sheep, yet previous population models had not dealt specifically with the issue of age structure (Coulson *et al.* 2001). When chronological age structure was incorporated into population models for Soay sheep the ability of these models to predict population crashes was improved dramatically (Coulson *et al.* 2001). How would this model perform if chronological age was substituted with the biological age of sheep? Our prediction is that chronological age may reflect more an individual's experience and learned behaviour, while biological age may be more representative of some aspect of individual physical condition. While both age concepts will overlap considerably we suspect that, for many of the life history issues molecular ecologists wish to address, biological age may be more meaningful.

The use of telomere length as an indicator of biological age will probably be limited to species whose telomere length changes over their lifetime (for most vertebrate species studied, telomere length shortens over a lifetime). It should also be noted that deleterious physical condition caused by short telomeres (e.g. replicative senescence and age-related diseases) may occur only in particularly long-lived species such as humans (Aviv 2002b). In most wild animals it seems unlikely that telomere function (Box 1) will be lost during an animal's lifetime, despite senescence being a common phenomenon among species; (Kirkwood & Austad 2000; Partridge & Gems 2002), because most animals have a much higher extrinsic mortality rate than humans. None the less, telomere assays may, finally, provide opportunities to enable ecologists to gain data on age specific survival rates and age-specific reproductive rates in species where such data are hard or previously impossible to obtain. We believe that there is great potential for the future application of this technique to open up new possibilities

Box 3. Telomere length as a tool in vertebrate conservation management

Telomere length assays may revolutionize the management of many endangered vertebrate species, where data on age may be lacking, but such knowledge might enhance dramatically predictive models and conservation recovery plans. This is especially true of long-lived k-selected vertebrate species that are grossly over-represented in the IUCN *Red Book* (IUCN 2003). The critically endangered kakapo (*Strigops habroptilus*) is an archetypical representative of this type of species. Kakapo are currently returning from near extinction after the discovery of remnant populations in Fiordland and Stewart Island in the 1970s (Elliott *et al.* 2001). In 2003 the population consisted of 86 individuals. However, none of the reproductively active individuals in the current population are of known ages (Elliott *et al.* 2001). With suggestions that some of the breeding birds may be 60+ and possibly older, and with demonstrably different levels of breeding success among this population (Miller *et al.* 2003), it would be useful to know what the age structure of this population is and how reproductive performance varies across individuals' life spans in order to better predict the future trends of this species. Furthermore, knowledge of age, when used in conjunction with traditional

molecular markers, such as microsatellites, would also enable the elucidation of family groups and pedigrees where these are unknown, greatly improving our understanding of the social organization and dynamics of species that have not been subject to long-term study. Given that blood samples of adequate size (> 50 µl) are available for every kakapo, it is possible to use both TRF and Q-PCR assays to provide the knowledge of age needed to help better manage kakapo recovery.

The application of this technology to other species is no less impressive. For example, new molecular tools for ageing together with other established approaches might see an end to lethal sampling programmes such as the scientific whaling programme (Brownell *et al.* 2000; Aron 2001). In the past a desire to understand the age structure of natural populations might appear to be one of the few legitimate areas where lethal sampling might still be justifiable. However, if validated, telomere-based assays of age would make it possible to gain information on: (i) an individual's age via telomere length changes; (ii) species, sex and populational affinities using mtDNA, sex and microsatellite markers (Baker *et al.* 2000); (iii) physiological state via analysis of stress hormones (Wasser *et al.* 2000); (iv) diet through lipid analyses (Olsen & Grahl-Nielsen 2003); and (v) oceanic movement using trace element analyses (Kelly 2000), without lethal sampling.

and avenues in many areas of research, especially in the field of conservation biology (see Box 3). For example, knowledge of biological age in a population of a locally endangered species will be informative in terms of conservation planning (e.g. selecting individuals for introduction and translocation).

Additionally, in a recent paper Stindl (2004) suggests that telomere length of a species may represent 'species age'. He argues that gradual loss of telomere length over thousands of generations may account for many cases of species extinction when telomere length becomes critically short, causing genome instability, and that this same process may result in evolution of new species by reorganizations of chromosomes which are accompanied by subsequent telomere elongation (Stindl 2004). Phylogenetically controlled comparisons of telomere lengths may help to test the validity of these claims.

Conclusions

Although there are some limitations on the use of telomeres in the fields of ecology, evolution and conservation, telomere length (an age estimator) and telomere shortening (an ageing estimator) are poised to become essential tools in

molecular ecology, providing immediate access to population age structure without long-term longitudinal studies. Once refined, telomere assays will contribute to answering questions regarding the relationship between reproduction and ageing (or survival) and other aspects of life history at among- and within-species levels. Applications of estimating the biological age of endangered vertebrate populations seem especially promising (Box 3). The use of various techniques that measure telomere length, particularly the complementary use of TRF analysis and Q-PCR, should facilitate telomere research in a variety of species. In most species, research on the ends of chromosomes has yet to start or has just begun, and further data will undoubtedly lead to methodological improvements (e.g. STELA) that will make telomere analysis more accessible. Time will prove whether 'age' and 'ageing' estimations by measuring telomere properties will or will not lead to a paradigm shift in molecular ecology, but we think it likely.

Acknowledgements

We are grateful to Tim Birkhead, Jake Gratten, Maggie Hall, Ben Hatchwell, Robert Jehle, Jon Slate, Michelle Simeoni, Frank Sin,

Margaret Will and three anonymous reviewers whose comments improved the paper. S. N. is supported by the Foundation for Research Science and Technology, New Zealand and T.A.B. is supported by a Royal Society Leverhulme Trust Senior Research Fellowship.

References

- Albon SD, Coulson TN, Brown D *et al.* (2000) Temporal changes in key factors and key age groups influencing the population dynamics of female red deer. *Journal of Animal Ecology*, **69**, 1099–1110.
- Allsopp RC, Vaziri H, Patterson C *et al.* (1992) Telomere length predicts replicative capacity of human fibroblasts. *Proceedings of the National Academy of Sciences USA*, **89**, 10114–10118.
- Aron W (2001) Scientific whaling. *Science*, **291**, 253.
- Aviv A (2002a) Chronology versus biology: telomeres, essential hypertension, and vascular aging. *Hypertension*, **40**, 229–232.
- Aviv A (2002b) Telomeres, sex, reactive oxygen species, and human cardiovascular aging. *Journal of Molecular Medicine*, **80**, 689–695.
- Baird DM, Rowson J, Wynford-Thomas D, Kipling D (2003) Extensive allelic variation and ultrashort telomeres in senescent human cells. *Nature Genetics*, **33**, 203–207.
- Baker C, Lento G, Cipriano F, Dalebout M, Palumbi S (2000) Scientific whaling: source of illegal products for market? *Science*, **290**, 1695–1695.
- Barja G, Herrero A (2000) Oxidative damage to mitochondrial DNA is inversely related to maximum life span in the heart and brain of mammals. *FASEB Journal*, **14**, 312–318.
- Benetos A, Okuda K, Lajemi M *et al.* (2001) Telomere length as an indicator of biological aging: the gender effect and relation with pulse pressure and pulse wave velocity. *Hypertension*, **37**, 381–385.
- Blackburn EH (1991) Structure and function of telomeres. *Nature*, **350**, 569–573.
- Blasco MA (2003) Telomeres and cancer: a tale with many endings. *Current Opinions in Genetics and Development*, **13**, 70–76.
- Broccoli D, Smogorzewska A, Chong L, de Lange T (1997) Human telomeres contain two distinct Myb-related proteins, TRF1 and TRF2. *Nature Genetics*, **17**, 231–235.
- Brownell RL Jr, Tillman MF, di Sciarra GN, Berggren P, Read AJ (2000) Further scrutiny of scientific whaling. *Science*, **290**, 1696a.
- Brummendorf TH, Mak J, Sabo KM *et al.* (2002) Longitudinal studies of telomere length in feline blood cells: implications for hematopoietic stem cell turnover *in vivo*. *Experimental Hematology*, **30**, 1147–1152.
- Campisi J (1996) Replicative senescence: an old lives' tale? *Cell*, **84**, 497–500.
- Cawthon RM (2002) Telomere measurement by quantitative PCR. *Nucleic Acids Research*, **30**, E47.
- Cawthon RM, Smith KR, O'Brien E, Sivatchenko A, Kerber RA (2003) Association between telomere length in blood and mortality in people aged 60 years or older. *Lancet*, **361**, 393–395.
- Chang E, Harley CB (1995) Telomere length and replicative aging in human vascular tissues. *Proceedings of the National Academy of Sciences USA*, **92**, 11190–11194.
- Charnov E (1991) Evolution of life history variation among female mammals. *Proceedings of the National Academy of Sciences USA*, **88**, 1134–1137.
- Cimino-Reale G, Pascale E, Battiloro E *et al.* (2001) The length of telomeric G-rich strand 3'-overhang measured by oligonucleotide ligation assay. *Nucleic Acids Research*, **29**, E35.
- Coulson T, Catchpole EA, Albon SD *et al.* (2001) Age, sex, density, winter weather, and population crashes in Soay sheep. *Science*, **292**, 1528–1531.
- Coviello-McLaughlin GM, Prowse KR (1997) Telomere length regulation during postnatal development and ageing in *Mus spretus*. *Nucleic Acids Research*, **25**, 3051–3058.
- Ellegren H, Sheldon BC (1997) New tools for sex identification and the study of sex allocation in birds. *Trends in Ecology*, **12**, 255–259.
- Elliott GP, Merton DV, Jansen PW (2001) Intensive management of a critically endangered species: the kakapo. *Biological Conservation*, **99**, 121–133.
- Finkel T, Holbrook NJ (2000) Oxidants, oxidative stress and the biology of ageing. *Nature*, **408**, 239–247.
- Frenck RW Jr, Blackburn EH, Shannon KM (1998) The rate of telomere sequence loss in human leukocytes varies with age. *Proceedings of the National Academy of Sciences USA*, **95**, 5607–5610.
- Friedrich U, Schwab M, Griese EU, Fritz P, Klotz U (2001) Telomeres in neonates: new insights in fetal hematopoiesis. *Pediatric Research*, **49**, 252–256.
- Gan YB, Engelke KJ, Brown CA, Au JLS (2001) Telomere amount and length assay. *Pharmaceutical Research*, **18**, 1655–1659.
- Gemmell NJ, Majluf P (1997) Projectile biopsy sampling of fur seals. *Marine Mammal Science*, **13**, 512–516.
- Girondot M, Garcia J (1998) Senescence and longevity in turtles. What telomeres tell us. In: *Current Studies in Herpetology* (eds Miaud DC, Guyétant R), pp. 133–137. Societa Europaea Herpetologica, Le Bouget du Lac, France.
- Goyns MH (2002) Genes, telomeres and mammalian ageing. *Mechanisms of Ageing and Development*, **123**, 791–799.
- Grant JD, Broccoli D, Muquit M *et al.* (2001) Telometric: a tool providing simplified, reproducible measurements of telomeric DNA from constant field agarose gels. *Biotechniques*, **31**, 1318.
- Greider CW (1996) Telomere length regulation. *Annual Review of Biochemistry*, **65**, 337–365.
- Harley CB, Futcher AB, Greider CW (1990) Telomeres shorten during ageing of human fibroblasts. *Nature*, **345**, 458–460.
- Harley CB, Kim NW, Prowse KR *et al.* (1994) Telomerase, cell immortality, and cancer. *Cold Spring Harbour Symposia on Quantitative Biology*, **59**, 307–315.
- Harley CG, Vaziri HMCC, Allsopp RC (1992) The telomere hypothesis of cellular aging. *Experimental Gerontology*, **27**, 375–382.
- Hastie ND, Dempster M, Dunlop MG *et al.* (1990) Telomere reduction in human colorectal carcinoma and with ageing. *Nature*, **346**, 866–868.
- Hausmann MF, Vleck CM (2002) Telomere length provides a new technique for aging animals. *Oecologia*, **130**, 325–328.
- Hausmann MF, Vleck CM, Nisbet IC (2003a) Calibrating the telomere clock in common terns, *Sterna hirundo*. *Experimental Gerontology*, **38**, 787–789.
- Hausmann MF, Winkler DW, O'Reilly KM *et al.* (2003b) Telomeres shorten more slowly in long-lived birds and mammals than in short-lived ones. *Proceedings of the Royal Society of London B: Biological Science*, **270**, 1387–1392.
- Hemann MT, Strong MA, Hao LY, Greider CW (2001) The shortest telomere, not average telomere length, is critical for cell viability and chromosome stability. *Cell*, **107**, 67–77.

- Henson JD, Neumann AA, Yeager TR, Reddel RR (2002) Alternative lengthening of telomeres in mammalian cells. *Oncogene*, **21**, 598–610.
- Huffman KE, Levene SD, Tesmer VM, Shay JW, Wright WE (2000) Telomere shortening is proportional to the size of the G-rich telomeric-3'-overhang. *Journal of Biological Chemistry*, **275**, 19719–19722.
- IUCN (2003) *2003 IUCN Red List of Threatened Species*. IUCN Gland, Switzerland. [http://www.redlist.org].
- Jeanclous E, Schork NJ, Kyvik KO *et al.* (2000) Telomere length inversely correlates with pulse pressures and is highly familial. *Hypertension*, **36**, 196–200.
- Jones AG, Ardren WR (2003) Methods of parentage analysis in natural populations. *Molecular Ecology*, **12**, 2511–2523.
- Karlseder J, Smogorzewska A, de Lange T (2002) Senescence induced altered telomere state, not telomere loss. *Science*, **295**, 2446–2449.
- Kelly JF (2000) Stable isotopes of carbon and nitrogen in the study of avian and mammalian trophic ecology. *Canadian Journal of Zoology*, **78**, 1–27.
- Kern S, Ackermann M, Stearns SC, Kawecki TJ (2001) Decline in offspring viability as a manifestation of aging in *Drosophila melanogaster*. *Evolution*, **55**, 1822–1831.
- Kirkwood TB, Austad SN (2000) Why do we age? *Nature*, **408**, 233–238.
- Lambertsen RH (1987) A biopsy system for large whales and its use for cytogenetics. *Journal of Mammalogy*, **68**, 443–445.
- Lee WW, Nam KH, Terao K, Yoshikawa Y (2002) Age-related telomere length dynamics in peripheral blood mononuclear cells of healthy cynomolgus monkeys measured by flow FISH. *Immunology*, **105**, 458–465.
- Lindstrom J (1999) Early development and fitness in birds and mammals. *Trends in Ecology*, **14**, 343–348.
- Lingner J, Hughes TR, Shevchenko A *et al.* (1997) Reverse transcriptase motifs in the catalytic subunit of telomerase. *Science*, **276**, 561–567.
- Meyne J, Ratliff RL, Moyzis RK (1989) Conservation of the human telomere sequence (TTAGGG)_n among vertebrates. *Proceedings of the National Academy of Sciences USA*, **86**, 7049–7053.
- Miller HC, Lambert DM, Millar CD, Robertson BC, Minot EO (2003) Minisatellite DNA profiling detects lineages and parentage in the endangered kakapo (*Strigops habroptilus*) despite low microsatellite DNA variation. *Conservation Genetics*, **4**, 265–274.
- Miyashita N, Shiga K, Yonai M *et al.* (2002) Remarkable differences in telomere lengths among cloned cattle derived from different cell types. *Biological Reproduction*, **66**, 1649–1655.
- Ogburn CE, Carlberg K, Ottinger MA *et al.* (2001) Exceptional cellular resistance to oxidative damage in long-lived birds requires active gene expression. *Journal of Gerontology A, Biological Sciences and Medical Sciences*, **56**, B468–B474.
- Olsen E, Grahl-Nielsen O (2003) Blubber fatty acids of minke whales: stratification, population identification and relation to diet. *Marine Biology*, **142**, 13–24.
- Partridge L, Gems D (2002) Mechanisms of ageing: public or private? *Nature Review Genetics*, **3**, 165–175.
- Perls TT, Wilmoth J, Levenson R *et al.* (2002) Life-long sustained mortality advantage of siblings of centenarians. *Proceedings of the National Academy of Sciences USA*, **99**, 8442–8447.
- Prowse KR, Greider CW (1995) Developmental and tissue-specific regulation of mouse telomerase and telomere length. *Proceedings of the National Academy of Sciences USA*, **92**, 4818–4822.
- Ricklefs RE (1977) Evolution of reproductive strategies in birds – reproductive effort. *American Naturalist*, **111**, 453–478.
- Ricklefs RE, Wikelski M (2002) The physiology/life-history nexus. *Trends in Ecology and Evolution*, **17**, 462–468.
- Rose MR (1991) *The Evolutionary Biology of Aging*. Oxford University Press, Oxford.
- Rufer N, Brummendorf TH, Kolvraa S *et al.* (1999) Telomere fluorescence measurements in granulocytes and T lymphocyte subsets point to a high turnover of hematopoietic stem cells and memory T cells in early childhood. *Journal of Experimental Medicine*, **190**, 157–167.
- Rufer N, Dragowska W, Thornbury G, Roosnek E, Lansdorp PM (1998) Telomere length dynamics in human lymphocyte subpopulations measured by flow cytometry. *Nature Biotechnology*, **16**, 743–747.
- Sedivy JM, Shippen DE, Shakirov EV (2003) Surprise ending. *Nature Genetics*, **33**, 114–116.
- Sidorov IA, Gee D, Dimitrov DS (2003) A kinetic model of telomere shortening in infants and adults. *Journal of Theoretical Biology*, **226**, 169–175.
- Slagboom PE, Droog S, Boomsma DI (1994) Genetic determination of telomere size in humans: a twin study of three age groups. *American Journal of Human Genetics*, **55**, 876–882.
- Stearns SC (1989) Trade-offs in life-history evolution. *Functional Ecology*, **3**, 259–268.
- Stearns SC, Koella JC (1986) The evolution of phenotypic plasticity in life-history traits – predictions of reaction norms for age and size at maturity. *Evolution*, **40**, 893–913.
- van Steensel B, de Lange T (1997) Control of telomere length by the human telomeric protein TRF1. *Nature*, **385**, 740–743.
- Stewart SA, Ben-Porath I, Carey VJ *et al.* (2003) Erosion of the telomeric single-strand overhang at replicative senescence. *Nature Genetics*, **33**, 492–496.
- Stindl R (2004) Is telomere erosion a mechanism of species extinction? *Journal of Experimental Zoology (Molecular and Developmental Evolution)*, **302B**, 111–120.
- Takasaki T, Tsuji A, Ikeda N, Ohishi M (2003) Age estimation in dental pulp DNA based on human telomere shortening. *International Journal of Legal Medicine*, **117**, 232–234.
- Tsuji A, Ishiko A, Takasaki T, Ikeda N (2002) Estimating age of humans based on telomere shortening. *Forensic Science International*, **126**, 197–199.
- Turker M, Martin G (1998) Genetics of human disease, longevity, and aging. In: *Principles of Geriatric Medicine and Gerontology* (eds Hazzard W, Blass J, Ettinger JW), pp. 21–44. McGraw-Hill, New York.
- Vaziri H, Dragowska W, Allsopp RC *et al.* (1994) Evidence for a mitotic clock in human hematopoietic stem cells: loss of telomeric DNA with age. *Proceedings of the National Academy of Sciences USA*, **91**, 9857–9860.
- Venkatesan RN, Price C (1998) Telomerase expression in chickens: constitutive activity in somatic tissues and down-regulation in culture. *Proceedings of the National Academy of Sciences USA*, **95**, 14763–14768.
- Vleck CM, Haussmann MF, Vleck D (2003) The natural history of telomeres: tools for aging animals and exploring the aging process. *Experimental Gerontology*, **38**, 791–795.
- Wasser SK, Hunt KE, Brown JL *et al.* (2000) A generalized fecal glucocorticoid assay for use in a diverse array of nondomestic mammalian and avian species. *General and Comparative Endocrinology*, **120**, 260–275.

- Watson J (1972) Origin of concatameric T4 DNA. *Nature*, **239**, 197–201.
- Zeichner SL, Palumbo P, Feng Y *et al.* (1999) Rapid telomere shortening in children. *Blood*, **93**, 2824–2830.
- von Zglinicki T, Pilger R, Sitté N (2000) Accumulation of single-strand breaks is the major cause of telomere shortening in human fibroblasts. *Free Radical Biology and Medicine*, **28**, 64–74.
- Zijlmans JM, Martens UM, Poon SS *et al.* (1997) Telomeres in the mouse have large inter-chromosomal variations in the number of T2AG3 repeats. *Proceedings of the National Academy of Sciences USA*, **94**, 7423–7428.

Shinichi Nakagawa is a PhD student at the University of Sheffield, studying the molecular and behavioural ecology of a house sparrow population on Lundy Island, UK. His research interests include developing and applying the tools (e.g. molecular, biochemical, statistical) that can assist the study of animals. Neil Gemmell heads the molecular ecology laboratory at the University of Canterbury, where an overarching theme has been the development and application of new technologies to ecological and evolutionary questions. Terry Burke heads a research group at Sheffield that has for several years used molecular approaches to address ecological questions.
