



UNIVERSITY OF LEEDS

This is a repository copy of *Triangulated neighborhoods in even-hole-free graphs*.

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/74353/>

Article:

da Silva, MV and Vuskovic, K (2007) Triangulated neighborhoods in even-hole-free graphs. *Discrete Mathematics*, 307 (9-10). 1065 - 1073 . ISSN 0012-365X

<https://doi.org/10.1016/j.disc.2006.07.027>

Reuse

See Attached

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Triangulated Neighborhoods in Even-hole-free Graphs

Murilo V. G. da Silva ^{*} and Kristina Vušković [†]

July 2005; revised July 2006

Abstract

An even-hole-free graph is a graph that does not contain, as an induced subgraph, a chordless cycle of even length. A graph is triangulated if it does not contain any chordless cycle of length greater than three, as an induced subgraph. We prove that every even-hole-free graph has a node whose neighborhood is triangulated. This implies that in an even-hole-free graph, with n nodes and m edges, there are at most $n + 2m$ maximal cliques. It also yields an $O(n^2m)$ algorithm that generates all maximal cliques of an even-hole-free graph. In fact these results are obtained for a larger class of graphs that contains even-hole-free graphs.

Keywords: even-hole-free graphs, triangulated graphs, structural characterization, generating all maximal cliques.

1 Introduction

We say that a graph G *contains* a graph H , if H is isomorphic to an induced subgraph of G . A graph G is *H-free* if it does not contain H . A *hole* is a chordless cycle of length at least four. A hole is *even* (resp. *odd*) if it contains even (resp. odd) number of nodes. An *n-hole* is a hole of length n . A graph is said to be *triangulated* if it does not contain any hole.

We *sign* a graph by assigning 0, 1 weights to its edges in such a way that, for every triangle in the graph, the sum of the weights of its edges is odd. A graph G is *odd-signable* if there is a signing of its edges so that, for every hole in G , the sum of the weights of its edges is odd. Clearly every even-hole-free graph is odd-signable, since we can get a correct signing by assigning a weight of 1 to every edge of the graph.

The graphs that are odd-signable and do not contain a 4-hole are studied in [7], where a decomposition theorem is proved for them. This decomposition theorem is used in [8] to obtain a polynomial time recognition algorithm for even-hole-free graphs.

For $x \in V(G)$, $N(x)$ denotes the set of nodes of G that are adjacent to x , and $N[x] = N(x) \cup \{x\}$. For $V' \subseteq V(G)$, $G[V']$ denotes the subgraph of G induced by V' . For $x \in V(G)$, the graph $G[N(x)]$ is called the *neighborhood* of x .

The main result of this paper is the following structural characterization of odd-signable graphs that do not contain a 4-hole.

^{*}School of Computing, University of Leeds, Leeds LS2 9JT, UK. murilo@comp.leeds.ac.uk.

[†]School of Computing, University of Leeds, Leeds LS2 9JT, UK. vuskovi@comp.leeds.ac.uk. This work was supported in part by EPSRC grant EP/C518225/1.

Theorem 1.1 *Every 4-hole-free odd-signable graph has a node whose neighborhood is triangulated.*

Exactly the same characterization of 4-hole-free Berge graphs (i.e. graphs that do not contain a 4-hole nor an odd hole) is obtained by Parfenoff, Roussel and Rusu in [15]. Note that 4-hole-free graphs in general need not have this property, see Figure 1.




Figure 1: A 4-hole-free graph that has no vertex whose neighborhood is triangulated.

A graph is *Berge* if it does not contain an odd hole nor the complement of an odd hole. A *square-3PC*(\cdot, \cdot) is a graph that consists of three paths between two nodes such that any two of the paths induce a hole, and at least two of the paths are of length 2. A graph G is *even-signable* if there is a signing of its edges so that for every hole in G , the sum of the weights of its edges is even. In [13] Maffray, Trotignon and Vušković show that every square-3PC(\cdot, \cdot)-free even-signable graph has a node whose neighborhood does not contain a long hole (where a *long hole* is a hole of length greater than 4). This result is used in [13] to obtain a combinatorial algorithm of complexity $\mathcal{O}(n^7)$ for finding a clique of maximum weight in square-3PC(\cdot, \cdot)-free Berge graphs. Note that this class of graphs generalizes both 4-hole-free Berge graphs and claw-free Berge graphs (where a *claw* is a graph on nodes x, a, b, c with three edges xa, xb, xc). We show in this paper that key ideas from [13] extend to 4-hole-free odd-signable graphs.

Using Theorem 1.1 one can obtain an efficient algorithm for generating all the maximal cliques in 4-hole-free odd-signable graphs (and in particular even-hole-free graphs). This we describe in Section 2. Theorem 1.1 is proved in Section 3.

Recently Addario-Berry, Chudnovsky, Havet, Reed and Seymour [1] have proved a stronger property of even-hole-free graphs, namely that every even-hole-free graph has a bisimplicial vertex (i.e. a vertex whose neighborhood partitions into two cliques). This characterization immediately yields that for an even-hole-free graph G , $\chi(G) \leq 2\omega(G) - 1$, where $\chi(G)$ is the chromatic number of G and $\omega(G)$ is the size of the largest clique in G (observe that if v is a bisimplicial vertex of G , then its degree is at most $2\omega(G) - 2$, and hence G can be colored with at most $2\omega(G) - 1$ colors). The two characterizations of even-hole-free graphs were discovered independently and at about the same time. The proof of the characterization in [1] is over 40 pages long. Our weaker characterization is enough to obtain an

efficient algorithm for generating all maximal cliques of even-hole-free graphs, and its proof is very short.

2 Generating all the maximal cliques of a 4-hole-free odd-signable graph

For a graph G let k denote the number of maximal cliques in G , n the number of nodes in G and m the number of edges of G . Farber [10] shows that there are $\mathcal{O}(n^2)$ maximal cliques in any 4-hole-free graph. Tsukiyama, Ide, Ariyoshi and Shirakawa [19] give an $\mathcal{O}(nmk)$ algorithm for generating all maximal cliques of a graph, and Chiba and Nishizeki [2] improve this complexity to $\mathcal{O}(m^{1.5}k)$. The complexity is further improved for dense graphs by the $\mathcal{O}(M(n)k)$ algorithm of Makino and Uno [14], where $M(n)$ denotes the time needed to multiply two $n \times n$ matrices. Note that Coppersmith and Winograd show that matrix multiplication can be done in $\mathcal{O}(n^{2.376})$ time [9]. So one can generate all the maximal cliques of a 4-hole-free graph in time $\mathcal{O}(m^{1.5}n^2)$ or $\mathcal{O}(n^{4.376})$.

We now show that Theorem 1.1 implies that there are at most $n + 2m$ maximal cliques in a 4-hole-free odd-signable graph, and it yields an algorithm that generates all the maximal cliques of a 4-hole-free odd-signable graph in time $\mathcal{O}(n^2m)$. In particular, in a weighted graph, a maximum weight clique can be found in time $\mathcal{O}(n^2m)$.

Let \mathcal{C} be any class of graphs closed under taking induced subgraphs, such that for every G in \mathcal{C} , G has a node whose neighborhood is triangulated. Consider the following algorithm for generating all maximal cliques of graphs in \mathcal{C} .

Find a node x_1 of G whose neighborhood is triangulated (if no such node exists, G is not in \mathcal{C} , or in particular, G is not 4-hole-free odd-signable graph by Theorem 1.1). Let $G_1 = G[N[x_1]]$ and $G^1 = G \setminus \{x_1\}$. Every maximal clique of G belongs to G_1 or G^1 . Recursively construct triangulated graphs G_1, \dots, G_n as follows. For $i \geq 2$, find a node x_i of G^{i-1} whose neighborhood is triangulated and let $G_i = G[N_{G^{i-1}}[x_i]]$ and $G^i = G^{i-1} \setminus \{x_i\} = G \setminus \{x_1, \dots, x_i\}$.

Clearly every maximal clique of G belongs to exactly one of the graphs G_1, \dots, G_n . A triangulated graph on n vertices has at most n maximal cliques [11]. So for $i = 1, \dots, n$, graph G_i has at most $1 + d(x_i)$ maximal cliques (where $d(x)$ denotes the degree of vertex x). It follows that the number of maximal cliques of G is at most $\sum_{i=1}^n (1 + d(x_i)) = n + 2m$.

Checking whether a graph is triangulated can be done in time $\mathcal{O}(n + m)$ (using lexicographic breadth-first search [16]). So finding a vertex with triangulated neighborhood can be done in time $\mathcal{O}(\sum_{x \in V(G)} (d(x) + m)) = \mathcal{O}(nm)$. Hence constructing the graphs G_1, \dots, G_n takes time $\mathcal{O}(n^2m)$.

Generating all maximal cliques in a triangulated graph can be done in time $\mathcal{O}(n + m)$ (see, for example, [12]). Hence the overall complexity of generating all maximal cliques in a 4-hole-free odd-signable graph is dominated by the construction of the sequence G_1, \dots, G_n , i.e. it is $\mathcal{O}(n^2m)$.

Note that this algorithm is *robust* in Spinrad's sense [17]: given any graph G , the algorithm either verifies that G is not in \mathcal{C} (or in particular that G is not a 4-hole-free odd-signable graph) or it generates all the maximal cliques of G . Note that, when G is not in \mathcal{C} , the algorithm might still generate all the maximal cliques of G .

3 Proof of Theorem 1.1

For a graph G , let $V(G)$ denote its node set. For simplicity of notation we will sometimes write G instead of $V(G)$, when it is clear from the context that we want to refer to the node set of G . Also a singleton set $\{x\}$ will sometimes be denoted with just x . For example, instead of “ $u \in V(G) \setminus \{x\}$ ”, we will write “ $u \in G \setminus x$ ”.

Let x, y be two distinct nodes of G . A $3PC(x, y)$ is a graph induced by three chordless x, y -paths, such that any two of them induce a hole. We say that a graph G contains a $3PC(\cdot, \cdot)$ if it contains a $3PC(x, y)$ for some $x, y \in V(G)$. $3PC(\cdot, \cdot)$'s are also known as *thetas* (for example in [5]).

Let $x_1, x_2, x_3, y_1, y_2, y_3$ be six distinct nodes of G such that $\{x_1, x_2, x_3\}$ and $\{y_1, y_2, y_3\}$ induce triangles. A $3PC(x_1x_2x_3, y_1y_2y_3)$ is a graph induced by three chordless paths $P_1 = x_1, \dots, y_1$, $P_2 = x_2, \dots, y_2$ and $P_3 = x_3, \dots, y_3$, such that any two of them induce a hole. We say that a graph G contains a $3PC(\Delta, \Delta)$ if it contains a $3PC(x_1x_2x_3, y_1y_2y_3)$ for some $x_1, x_2, x_3, y_1, y_2, y_3 \in V(G)$. $3PC(\Delta, \Delta)$'s are also known as *prisms* (for example in [4]).

A *wheel*, denoted by (H, x) , is a graph induced by a hole H and a node $x \notin V(H)$ having at least three neighbors in H , say x_1, \dots, x_n . Node x is the *center* of the wheel. We say that the wheel (H, x) is *even* when n is even.

It is easy to see that even wheels, $3PC(\cdot, \cdot)$'s and $3PC(\Delta, \Delta)$'s cannot be contained in even-hole-free graphs. In fact they cannot be contained in odd-signable graphs. The following characterization of odd-signable graphs, given in [6], states that the converse is also true. It is in fact an easy consequence of a theorem of Truemper [18].

Theorem 3.1 *A graph is odd-signable if and only if it does not contain an even wheel, a $3PC(\cdot, \cdot)$ nor a $3PC(\Delta, \Delta)$.*

The fact that odd-signable graphs do not contain even wheels, $3PC(\cdot, \cdot)$'s and $3PC(\Delta, \Delta)$'s will be used throughout the rest of the paper.

In the next three lemmas we assume that G is a 4-hole-free odd-signable graph, x a node of G that is not adjacent to every other node of G , C_1 a connected component of $G \setminus N[x]$, and H a hole of $N(x)$. Note that H is an odd hole, else (H, x) is an even wheel.

Lemma 3.2 *If node u of C_1 has a neighbor in H then u is one of the following two types:*

- *Type 1: u has exactly one neighbor in H .*
- *Type 2: u has exactly two neighbors in H , and they are adjacent.*

Proof: If u has two nonadjacent neighbors a and b in H , then $\{a, b, u, x\}$ induces a 4-hole. \square

Let T^3 be a graph on 3 nodes that has exactly one edge.

Let x_1, x_2, x_3, y be four distinct nodes of G such that x_1, x_2, x_3 induce a triangle. A $3PC(x_1x_2x_3, y)$ is a graph induced by three chordless paths $P_1 = x_1, \dots, y$, $P_2 = x_2, \dots, y$ and $P_3 = x_3, \dots, y$, such that any two of them induce a hole. We say that a graph G contains a $3PC(\Delta, \cdot)$ if it contains a $3PC(x_1x_2x_3, y)$ for some $x_1, x_2, x_3, y \in V(G)$. $3PC(\Delta, \cdot)$'s are also known as *pyramids* (for example in [3]).

Lemma 3.3 *If H contains a T^3 all of whose nodes have neighbors in C_1 , then C_1 contains a path P , of length greater than 0, such that $P \cup H$ induces a $3PC(\Delta, \cdot)$, and the nodes of H that have a neighbor in P induce a T^3 .*

Proof: Let C be a smallest subset of C_1 such that $G[C]$ is connected and $H = h_1, \dots, h_n, h_1$ contains a T^3 all of whose nodes have neighbors in C . W.l.o.g. h_1, h_2 and $h_i, 3 < i < n$, have neighbors in C . Let $P = p_1, \dots, p_k$ be a shortest path of C such that p_1 is adjacent to h_1 and p_k is adjacent to h_2 . Note that no intermediate node of P is adjacent to h_1 or h_2 . Also possibly $k = 1$.

Claim 1: No node of $\{h_4, \dots, h_{n-1}\}$ has a neighbor in P .

Proof of Claim 1: Suppose not. Then by minimality of C , h_i has a neighbor in P and w.l.o.g. no node of $\{h_{i+1}, \dots, h_{n-1}\}$ has a neighbor in P . By Lemma 3.2, $p_1, p_k \notin N(h_i) \cap P$. In particular $k > 1$.

First suppose $N(h_n) \cap P \neq \emptyset$. By Lemma 3.2, $h_n p_k$ is not an edge. If $N(h_n) \cap P = p_1$ then $\{x, h_n, h_2, h_1\} \cup P$ induces an even wheel with center h_1 . So h_n has a neighbor in $P \setminus \{p_1, p_k\}$. If $h_i h_n$ is not an edge, then since all of h_1, h_n, h_i have neighbors in $P \setminus p_k$, the minimality of C is contradicted. So $h_i h_n$ is an edge of G . But then all of h_i, h_n, h_2 have neighbors in $P \setminus p_1$ and the minimality of C is contradicted. So $N(h_n) \cap P = \emptyset$.

Let p_r be the node of P with highest index adjacent to h_i . Let H' be the hole induced by $\{h_i, \dots, h_n, h_1, h_2, p_k, \dots, p_r\}$. Since (H', x) cannot be an even wheel, it follows that $h_i, \dots, h_n, h_1, h_2$ is an even subpath of H . Let p_s be the node of P with lowest index adjacent to h_i . Then $\{x, h_i, \dots, h_n, h_1, p_1, \dots, p_s\}$ induces an even wheel with center x . This completes the proof of Claim 1.

By Claim 1, h_i is not adjacent to a node of P . But h_i has a neighbor in C , and since C is connected, let $Q = q_1, \dots, q_l$ be a chordless path in C such that q_1 is adjacent to h_i and q_l has a neighbor in P .

Claim 2: No node of $\{h_4, \dots, h_{n-1}\}$ has a neighbor in $(P \cup Q) \setminus q_1$.

Proof of Claim 2: Suppose that some $h_j \in \{h_4, \dots, h_{n-1}\}$ has a neighbor in $(P \cup Q) \setminus q_1$. Then all of h_1, h_2, h_j have neighbors in $(P \cup Q) \setminus q_1$, contradicting the minimality of C . This completes the proof of Claim 2.

Claim 3: q_1 is of type 1 w.r.t. H .

Proof of Claim 3: By Lemma 3.2 q_1 is of type 1 or type 2. Suppose q_1 is of type 2. We now prove that $N(q_1) \cap H$ is either $\{h_3, h_4\}$ or $\{h_{n-1}, h_n\}$. Assume not. Then q_1 is adjacent to neither h_3 nor h_n . W.l.o.g. $N(q_1) \cap H = \{h_i, h_{i-1}\}$ and $i \neq 4$. If $N(q_l) \cap P \neq p_1$, then $(P \cup Q) \setminus p_1$ is connected and all of h_i, h_{i-1}, h_2 have neighbors in it, contradicting the minimality of C . So $N(q_l) \cap P = p_1$. If $k > 1$, then all of h_i, h_{i-1}, h_1 have neighbors in $(P \cup Q) \setminus p_k$, contradicting the minimality of C . So $k = 1$, and hence by Lemma 3.2, $N(p_1) \cap H = \{h_1, h_2\}$. Since H is odd, the two subpaths of H , h_2, \dots, h_{i-1} and h_i, \dots, h_n, h_1 have different parities.

W.l.o.g. h_2, \dots, h_{i-1} is odd, i.e. i is even. By Claim 2, no node of $\{h_4, \dots, h_{n-1}\}$ has a neighbor in $(P \cup Q) \setminus q_1$. If h_3 has no neighbor in Q then $Q \cup P \cup \{h_2, \dots, h_{i-1}, x\}$ contains an even wheel with center x . So h_3 must have a neighbor in Q . But then h_i, h_{i-1}, h_3 all have neighbors in Q (note that $h_3 h_{i-1}$ is not an edge since $i-1$ is odd greater than 3) contradicting the minimality of C . So $N(q_1) \cap H$ is either $\{h_3, h_4\}$ or $\{h_{n-1}, h_n\}$.

W.l.o.g. $N(q_1) \cap H = \{h_3, h_4\}$. If $N(q_l) \cap P \neq p_k$, then since all of h_1, h_3, h_4 have neighbors in $(P \cup Q) \setminus p_k$, the minimality of C is contradicted. So $N(q_l) \cap P = p_k$.

If $N(h_1) \cap Q \neq \emptyset$, then since all of h_1, h_3, h_4 have neighbors in Q , the minimality of C is contradicted. So $N(h_1) \cap Q = \emptyset$.

Now suppose that $N(h_n) \cap Q \neq \emptyset$. If $k > 1$, then since all of h_2, h_3, h_n have neighbors in $(P \cup Q) \setminus p_1$, the minimality of C is contradicted. So $k = 1$. Let q_r be the neighbor of h_n with highest index. If h_2 does not have neighbor in q_r, q_{r+1}, \dots, q_l , then $\{q_r, q_{r+1}, \dots, q_l, p_1, h_1, h_2, h_n, x\}$ induces an even wheel with center h_1 . So $N(h_2) \cap Q \neq \emptyset$. But then since h_2, h_3, h_n have neighbors in Q , the minimality of C is contradicted. Therefore, $N(h_n) \cap Q = \emptyset$. So, by Claim 2, no node of h_5, \dots, h_n, h_1 has a neighbor in Q .

Suppose $N(h_2) \cap Q \neq \emptyset$. Let q_r be the neighbor of h_2 in Q with lowest index. Then $(H \setminus h_3) \cup \{x, q_1, \dots, q_r\}$ induces an even wheel with center x . Therefore, $N(h_2) \cap Q = \emptyset$. If $k > 1$, then $Q \cup (H \setminus h_3) \cup \{p_k, x\}$ induces an even wheel with center x . So $k = 1$. Let q_s be the node of Q with highest index adjacent to h_3 . Then $\{p_1, q_s, \dots, q_l, h_1, h_2, h_3, x\}$ induces an even wheel with center h_2 . This completes the proof of Claim 3.

Claim 4: $N(q_l) \cap P = p_1$ or p_k .

Proof of Claim 4: Assume not. Then $k > 1$, and both $(P \cup Q) \setminus p_1$ and $(P \cup Q) \setminus p_k$ are connected. $N(h_1) \cap Q = \emptyset$, else all of h_1, h_2, h_i have neighbors in $(P \cup Q) \setminus p_1$, contradicting the minimality of C . Similarly, $N(h_2) \cap Q = \emptyset$.

We now show that h_3 has no neighbor in $P \cup Q$. Suppose it does. Then by Lemma 3.2, h_3 has a neighbor in $(P \cup Q) \setminus p_1$. If $i \neq 4$, then since all h_2, h_3, h_i have neighbors in $(P \cup Q) \setminus p_1$, the minimality of C is contradicted. So $i = 4$. If $N(h_3) \cap (P \cup Q) \neq p_k$, then all of h_1, h_3, h_4 have neighbors in $(P \cup Q) \setminus p_k$, contradicting the minimality of C . So $N(h_3) \cap (P \cup Q) = p_k$. But then $P \cup Q \cup \{h_2, h_3, h_4, x\}$ contains an even wheel with center h_3 . Therefore, h_3 has no neighbor in $P \cup Q$, and similarly neither does h_n .

By minimality of C , $N(q_l) \cap P$ is either a single vertex or two adjacent vertices of P . If $N(q_l) \cap P = \{a, b\}$, where $ab \in E(G)$, then $P \cup Q \cup \{x, h_1, h_2, h_i\}$ induces a $3PC(q_l ab, x h_1 h_2)$. If $N(q_l) \cap P = \{a\}$, then $P \cup Q \cup \{h_1, h_2, \dots, h_i\}$ induces a $3PC(a, h_2)$. This completes the proof of Claim 4.

By Claim 4, w.l.o.g. $N(q_l) \cap P = p_k$.

Claim 5: h_1 does not have a neighbor in $(P \cup Q) \setminus p_1$.

Proof of Claim 5: If $k > 1$, the claim follows from the minimality of C . Now suppose $k = 1$ and $N(h_1) \cap Q \neq \emptyset$. If h_2 has a neighbor in Q , then all of h_1, h_2, h_i have a neighbor in Q , contradicting the minimality of C . So h_2 does not have a neighbor in Q .

Suppose h_n has a neighbor in Q . Note that by Claim 3, such a neighbor is in $Q \setminus q_1$. Then

h_3 cannot have a neighbor in Q , else all of h_n, h_1, h_3 have neighbors in Q , contradicting the minimality of C . But then $(Q \setminus q_1) \cup (H \setminus h_1) \cup \{x, p_1\}$ contains an even wheel with center x . So h_n does not have a neighbor in Q .

Suppose h_3 has a neighbor in Q . By Claim 3, such a neighbor is in $Q \setminus q_1$. Then $(Q \setminus q_1) \cup (H \setminus h_2) \cup x$ contains an even wheel with center x . So h_3 does not have a neighbor in Q .

Let H' be the hole induced by $\{p_1, h_2, \dots, h_i\} \cup Q$, and H'' the hole induced by $\{x, p_1, h_2, h_i\} \cup Q$. Then either (H', h_1) or (H'', h_1) is an even wheel. This completes the proof of Claim 5.

Claim 6: $N(h_n) \cap (P \cup Q) = \emptyset$.

Proof of Claim 6: Assume not. If h_3 has a neighbor in $P \cup Q$ then, by Claim 3, all of h_2, h_3, h_n have a neighbor in $(P \cup Q) \setminus q_1$, contradicting the minimality of C . So $N(h_3) \cap (P \cup Q) = \emptyset$. Let R be a shortest path from h_2 to h_n in the graph induced by $P \cup (Q \setminus q_1) \cup \{h_2, h_n\}$. Then by Claims 2 and 3, $R \cup (H \setminus h_1) \cup x$ induces an even wheel with center x . This completes the proof of Claim 6.

Claim 7: $N(h_3) \cap (P \cup Q) = \emptyset$.

Proof of Claim 7: Assume not. Let R be a shortest path from h_1 to h_3 in the graph induced by $(P \cup Q) \setminus q_1$. Then $R \cup (H \setminus h_2) \cup x$ induces an even wheel with center x . This completes the proof of Claim 7.

If $k > 1$ then the graph induced by $H \cup Q \cup p_k$ contains a $3PC(h_2, h_i)$. So $k = 1$. By symmetry and Claim 5, h_2 does not have a neighbor in Q , and hence $P \cup Q \cup H$ induces a $3PC(\Delta, \cdot)$. \square

Lemma 3.4 *There exists a node of H that has no neighbor in C_1 .*

Proof: Let $H = h_1, \dots, h_n, h_1$ and suppose that every node of H has a neighbor in C_1 . Recall that since (H, x) cannot be an even wheel, H is of odd length. So H contains a T^3 all of whose nodes have neighbors in C_1 . By Lemma 3.3, C_1 contains a path $P = p_1, \dots, p_k$, $k > 1$, such that $P \cup H$ induces w.l.o.g. a $3PC(h_1 h_2 p_k, h_i)$, $3 < i < n$. If i is odd, then $\{x, h_2, \dots, h_i\} \cup P$ induces an even wheel with center x . So i is even.

Let $Q = q_1, \dots, q_l$ be a path in C_1 defined as follows: q_1 is adjacent to $h_j \in H \setminus \{h_1, h_2, h_i\}$ where j is odd, q_l is adjacent to a node of P and no proper subpath of Q has this property. We may assume that P and Q are chosen so that $|P \cup Q|$ is minimized.

By the choice of P and Q , $N(q_l) \cap P$ is either one single vertex or two adjacent vertices of P , and h_j has no neighbor in $Q \setminus q_1$. Note that since n is odd, the two subpaths of H , h_2, \dots, h_i and h_i, \dots, h_n, h_1 are both of even length, so we may assume w.l.o.g. that $2 < j < i$.

Claim 1: At least one node of $\{h_2, \dots, h_{j-1}\}$ (resp. $\{h_{j+1}, \dots, h_n\}$) has a neighbor in Q .

Proof of Claim 1: First suppose that no node of $H \setminus \{h_1, h_j\}$ has a neighbor in Q . Let p_s be the node of P with highest index adjacent to q_l . If $j > 3$, then $\{x, h_2, \dots, h_j, p_s, \dots, p_k\} \cup Q$ induces

an even wheel with center x . So $j = 3$. If $N(h_1) \cap Q = \emptyset$ then $\{x, h_1, h_2, h_3, p_s, \dots, p_k\} \cup Q$ induces an even wheel with center h_2 . So $N(h_1) \cap Q \neq \emptyset$. Let q_r be the node of Q with lowest index adjacent to h_1 . Then $(H \setminus h_2) \cup \{x, q_1, \dots, q_r\}$ induces an even wheel with center x . So at least one node of $H \setminus \{h_1, h_j\}$ has a neighbor in Q .

Next suppose that no node of $\{h_2, \dots, h_{j-1}\}$ has a neighbor in Q . Let p_s be the node of P with highest index adjacent to q_l . If $j > 3$ then $\{x, h_2, \dots, h_j, p_s, \dots, p_k\} \cup Q$ induces an even wheel with center x . So $j = 3$. Let $h_{j'}$ be the node of $\{h_{j+1}, \dots, h_n\}$ with lowest index adjacent to a node of Q . By definition of Q and Lemma 3.2, j' is even. Let q_r be the node of Q with lowest index adjacent to $h_{j'}$. If $j' > 4$ then $\{x, h_j, \dots, h_{j'}, q_1, \dots, q_r\}$ induces an even wheel with center x . So $j' = 4$. If $N(h_1) \cap Q = \emptyset$ then $\{x, h_1, h_2, h_3, p_s, \dots, p_k\} \cup Q$ induces an even wheel with center h_2 . So $N(h_1) \cap Q \neq \emptyset$. In fact, by Lemma 3.2, $N(h_1) \cap (Q \setminus q_1) \neq \emptyset$. Suppose $N(h_4) \cap Q \neq q_1$. Let R be a shortest path from h_4 to h_1 in the graph induced by $(Q \setminus q_1) \cup \{h_1, h_4\}$. Then $\{x, h_1, \dots, h_4\} \cup R$ induces an even wheel with center x . So $N(h_4) \cap Q = q_1$. Suppose $N(q_l) \cap P \neq p_1$ or $i > 4$. Then $\{x, h_2, h_3, h_4, p_s, \dots, p_k\} \cup Q$ induces an even wheel with center h_3 . So $N(q_l) \cap P = p_1$ and $i = 4$. Let R be a shortest path from p_1 to h_1 in the graph induced by $Q \cup \{p_1, h_1\}$. Then $P \cup R \cup \{h_1, h_4, x\}$ induces a $3PC(p_1, h_1)$. Therefore at least one node of $\{h_2, \dots, h_{j-1}\}$ has a neighbor in Q .

Finally suppose that no node of $\{h_{j+1}, \dots, h_n\}$ has a neighbor in Q . Let $h_{j'}$ be a node of h_2, \dots, h_{j-1} such that $N(h_{j'}) \cap Q \neq \emptyset$ and the path from $h_{j'}$ to h_i in the graph induced by $P \cup Q \cup \{h_i, h_{j'}\}$ is minimized. By definition of Q and Lemma 3.2, j' is even. Suppose $N(h_1) \cap Q \neq \emptyset$. Let R be a shortest path from h_j to h_1 in the graph induced by $Q \cup \{h_1, h_j\}$. Then $(H \setminus \{h_2, \dots, h_{j-1}\}) \cup R \cup x$ induces an even wheel with center x . So $N(h_1) \cap Q = \emptyset$. Suppose $N(q_l) \cap P \neq p_k$. Let R be a shortest path from h_i to $h_{j'}$ in the graph induced by $P \cup Q \cup \{h_i, h_{j'}\}$. Note that by definition of Q and $h_{j'}$ and by Lemma 3.2, no node of $\{h_2, \dots, h_{j-1}\}$ has a neighbor in R . Then $(H \setminus \{h_{j'+1}, \dots, h_{i-1}\}) \cup R \cup x$ induces an even wheel with center x . So $N(q_l) \cap P = p_k$. But then $(H \setminus \{h_2, \dots, h_{j-1}\}) \cup P \cup Q$ induces a $3PC(p_k, h_i)$. This completes the proof of Claim 1.

By Claim 1 at least two nodes, say $h_{j'}$ and $h_{j''}$, of $H \setminus \{h_1, h_j\}$ have a neighbor in Q . Note that by definition of Q and Lemma 3.2, j' and j'' are both even. W.l.o.g. $j' < j < j''$. Let $R = r_1, \dots, r_t$ be a shortest path in the graph induced by Q where $N(h_{j'}) \cap R = r_1$ and $N(h_{j''}) \cap R = r_t$. W.l.o.g. and by Lemma 3.2 no other node from $H \setminus \{h_1, h_j\}$ has a neighbor in R .

If $N(h_1) \cap R = \emptyset$, then $(H \setminus \{h_{j'+1}, \dots, h_{j''-1}\}) \cup R \cup x$ induces an even wheel with center x . So $N(h_1) \cap R \neq \emptyset$. Suppose $j' \neq 2$. Let R' be a shortest path from h_1 to $h_{j'}$ in the graph induced by $R \cup \{h_1, h_{j'}\}$. Then $\{x, h_1, \dots, h_{j'}\} \cup R'$ induces an even wheel with center x . Therefore $j' = 2$.

Suppose that $N(h_1) \cap R = r_1$. Then by Lemma 3.2, $N(r_1) \cap H = \{h_1, h_2\}$. If $r_t = q_1$, then by Lemma 3.2, $N(r_t) \cap H = \{h_j, h_{j+1}\}$, and hence $H \cup R$ induces a $3PC(h_1 h_2 r_1, h_{j+1} h_j r_t)$. So $r_t \neq q_1$, and hence $N(r_t) \cap H = \{h_{j''}\}$. Therefore $H \cup R$ induces a $3PC(h_1 h_2 r_1, h_{j''})$. Let R' be a shortest path from q_1 to a node of R in the graph induced by Q . Since $|R \cup R'| < |P \cup Q|$, the choice of P and Q is contradicted.

So $N(h_1) \cap (R \setminus r_1) \neq \emptyset$. Let r_s be the node of R with highest index adjacent to h_1 . If h_j has no neighbor in r_s, \dots, r_t , then $\{x, h_1, \dots, h_{j''}, r_s, \dots, r_t\}$ induces an even wheel with center x . So h_j does have a neighbor in r_s, \dots, r_t , i.e. $r_t = q_1$. By Lemma 3.2, $N(r_t) \cap H = \{h_j, h_{j''}\}$,

where $j'' = j+1$. Note that $i \geq j+1$ and $r_s \neq q_l$. But then $(H \setminus \{h_2, \dots, h_j\}) \cup P \cup \{r_s, \dots, r_t\}$ induces a $3PC(h_1, h_i)$. \square

Note that the above lemma does not work if we allow 4-holes. Consider the odd-signable graph in Figure 2 (one can see that this graph is odd-signable by assigning 0 to the three bold edges and 1 to all the other edges). Let H be the 5-hole induced by the neighborhood of node x . Then every node of H has a neighbor in the unique connected component obtained by removing $N(x) \cup x$.




Figure 2: An odd-signable graph for which Lemma 3.4 does not work.

Let \mathcal{F} be a class of graphs. We say that a graph G is \mathcal{F} -free if G does not contain (as an induced subgraph) any of the graphs from \mathcal{F} .

A class \mathcal{F} of graphs satisfies *property (*) w.r.t. a graph G* if the following holds: for every node x of G such that $G \setminus N[x] \neq \emptyset$, and for every connected component C of $G \setminus N[x]$, if $F \in \mathcal{F}$ is contained in $G[N(x)]$, then there exists a node of F that has no neighbor in C .

The following theorem is proved in [13]. For completeness we include its proof here.

Theorem 3.5 (Maffray, Trotignon and Vušković [13]) *Let \mathcal{F} be a class of graphs such that for every $F \in \mathcal{F}$, no node of F is adjacent to all the other nodes of F . If \mathcal{F} satisfies property (*) w.r.t. a graph G , then G has a node whose neighborhood is \mathcal{F} -free.*

Proof: Let \mathcal{F} be a class of graphs such that for every $F \in \mathcal{F}$, no node of F is adjacent to all the other nodes of F . Assume that \mathcal{F} satisfies property (*) w.r.t. G , and suppose that for every $x \in V(G)$, $G[N(x)]$ is not \mathcal{F} -free. Then G is not a clique (since every graph of \mathcal{F} contains nonadjacent nodes) and hence it contains a node x that is not adjacent to all other nodes of G . Let C_1, \dots, C_k be the connected components of $G \setminus N[x]$, with $|C_1| \geq \dots \geq |C_k|$. Choose x so that for every $y \in V(G)$ the following holds: if C_1^y, \dots, C_l^y are the connected components of $G \setminus N[y]$ with $|C_1^y| \geq \dots \geq |C_l^y|$, then

- $|C_1| > |C_1^y|$, or
- $|C_1| = |C_1^y|$ and $|C_2| > |C_2^y|$, or

- ...
- $|C_1| = |C_1^y|, \dots, |C_{k-1}| = |C_{k-1}^y|$ and $|C_k| > |C_k^y|$, or
- for $i = 1, \dots, k$, $|C_i| = |C_i^y|$ and $k = l$.

Let $N = N(x)$ and $C = C_1 \cup \dots \cup C_k$. For $i = 1, \dots, k$, let N_i be the set of nodes of N that have a neighbor in C_i .

Claim 1: $N_1 \subseteq N_2 \subseteq \dots \subseteq N_k$ and for every $i = 1, \dots, k - 1$, every node of $(N \setminus N_i) \cup (C_{i+1} \cup \dots \cup C_k)$ is adjacent to every node of N_i .

Proof of Claim 1: We argue by induction. First we show that every node of $(N \setminus N_1) \cup (C_2 \cup \dots \cup C_k)$ is adjacent to every node of N_1 . Assume not and let $y \in (N \setminus N_1) \cup (C_2 \cup \dots \cup C_k)$ be such that it is not adjacent to $z \in N_1$. Clearly y has no neighbor in C_1 , but z does. So $G \setminus N[y]$ contains a connected component that contains $C_1 \cup z$, contradicting the choice of x .

Now let $i > 1$ and assume that $N_1 \subseteq \dots \subseteq N_{i-1}$ and every node of $(N \setminus N_{i-1}) \cup (C_i \cup \dots \cup C_k)$ is adjacent to every node of N_{i-1} . Since every node of C_i is adjacent to every node of N_{i-1} , it follows that $N_{i-1} \subseteq N_i$. Suppose that there exists a node $y \in (N \setminus N_i) \cup (C_{i+1} \cup \dots \cup C_k)$ that is not adjacent to a node $z \in N_i$. Then $z \in N_i \setminus N_{i-1}$ and z has a neighbor in C_i . Also y is adjacent to all nodes in N_{i-1} and no node of $C_1 \cup \dots \cup C_i$. So there exist connected components of $G \setminus N[y]$, C_1^y, \dots, C_l^y such that $C_1 = C_1^y, \dots, C_{i-1} = C_{i-1}^y$ and $C_i \cup z$ is contained in C_i^y . This contradicts the choice of x . This completes the proof of Claim 1.

Since $G[N]$ is not \mathcal{F} -free, it contains $F \in \mathcal{F}$. By property (*), a node y of F has no neighbor in C_k . By Claim 1, y is adjacent to every node of N_k , and no node of $N \setminus N_k$ has a neighbor in C . So (since every node of F has a non-neighbor in F) F must contain another node $z \in N \setminus N_k$, nonadjacent to y . But then C_1, \dots, C_k are connected components of $G \setminus N[y]$ and z is contained in $(G \setminus N[y]) \setminus C$, so y contradicts the choice of x . \square

Proof of Theorem 1.1: Let G be a 4-hole-free odd-signable graph. Let \mathcal{F} be the set of all holes. By Lemma 3.4, \mathcal{F} satisfies property (*) w.r.t. G . So by Theorem 3.5, G has a node whose neighborhood is \mathcal{F} -free, i.e. triangulated. \square

4 Final remarks

In a graph G , for any node x , let C_1, \dots, C_k be the connected components of $G \setminus N[x]$, with $|C_1| \geq \dots \geq |C_k|$, and let the numerical vector $(|C_1|, \dots, |C_k|)$ be associated with x . The nodes of G can thus be ordered according to the lexicographic ordering of the numerical vectors associated with them. Say that a node x is *lex-maximal* if the associated numerical vector is lexicographically maximal over all nodes of G . Theorem 3.5 actually shows that for a lex-maximal node x , $N(x)$ is \mathcal{F} -free. This implies the following.

Theorem 4.1 *Let G be a 4-hole-free odd-signable graph, and let x be a lex-maximal node of G . Then the neighborhood of x is triangulated.*

Possibly a more efficient algorithm for listing all maximal cliques can be constructed by searching for a lex-maximal node.

Lemma 3.4 also proves the following decomposition theorem. (H, x) is a *universal wheel* if x is adjacent to all the nodes of H . A node set S is a *star cutset* of a connected graph G if for some $x \in S$, $S \subseteq N[x]$ and $G \setminus S$ is disconnected.

Theorem 4.2 *Let G be a 4-hole-free odd-signable graph. If G contains a universal wheel, then G has a star cutset.*

Proof: Let (H, x) be a universal wheel of G . If $G = N[x]$, then for any two nonadjacent nodes a and b of H , $N[x] \setminus \{a, b\}$ is a star cutset of G . So assume $G \setminus N[x]$ contains a connected component C_1 . By Lemma 3.4, a node $a \in H$ has no neighbor in C_1 . But then $N[x] \setminus a$ is a star cutset of G that separates a from C_1 . \square

In [7] universal wheels in 4-hole-free odd-signable graphs are decomposed with triple star cutsets, i.e. node cutsets S such that for some triangle $\{x_1, x_2, x_3\} \subseteq S$, $S \subseteq N(x_1) \cup N(x_2) \cup N(x_3)$.

References

- [1] L. Addario-Berry, M. Chudnovsky, F. Havet, B. Reed and P. Seymour, *Bisimplicial vertices in even-hole-free graphs*, preprint (2006).
- [2] N. Chiba and T. Nishizeki, *Arboricity and subgraph listing algorithms*, SIAM J. Comput. 14 (1985) 210-223.
- [3] M. Chudnovsky, G. Cornuéjols, X. Liu, P. Seymour and K. Vušković, *Recognizing Berge graphs*, Combinatorica 25 (2005) 143-187.
- [4] M. Chudnovsky, N. Robertson, P. Seymour and R. Thomas, *The strong perfect graph theorem*, to appear in Annals of Mathematics.
- [5] M. Chudnovsky and P. Seymour, *Excluding induced subgraphs*, preprint (2006).
- [6] M. Conforti, G. Cornuéjols, A. Kapoor and K. Vušković, *Even and odd holes in cap-free graphs*, Journal of Graph Theory 30 (1999) 289-308.
- [7] M. Conforti, G. Cornuéjols, A. Kapoor and K. Vušković, *Even-hole-free graphs, Part I: Decomposition theorem*, Journal of Graph Theory 39 (2002) 6-49.
- [8] M. Conforti, G. Cornuéjols, A. Kapoor and K. Vušković, *Even-hole-free graphs, Part II: Recognition algorithm*, Journal of Graph Theory 40 (2002) 238-266.
- [9] D. Coppersmith and S. Winograd, *Matrix multiplication via arithmetic progression*, Journal of Symbolic Computation 9 (1990) 251-280.
- [10] M. Farber, *On diameters and radii of bridged graphs*, Discrete Mathematics 73 (1989) 249-260.

- [11] D.R. Fulkerson and O.A. Gross, *Incidence matrices and interval graphs*, Pacific J. Math. 15 (1965) 835-855.
- [12] M.C. Golumbic, *Algorithmic Graph Theory and Perfect Graphs*, Second Edition, Annals of Discrete Mathematics 57, Elsevier, 2004.
- [13] F. Maffray, N. Trotignon and K. Vušković, *Algorithms for square-3PC(\cdot, \cdot)-free Berge graphs*, preprint (2005), submitted.
- [14] K. Makino and T. Uno, *New algorithm for enumerating all maximal cliques*, Algorithm Theory - SWAT 2004, edited by T. Hagerup and J. Katajainen, LNCS 3111 (2004) 260-272.
- [15] I. Parfenoff, F. Roussel and I. Rusu, *Triangulated neighborhoods in C_4 -free Berge graphs*, Proceedings of WG'99, LNCS 1665 (1999) 402-412.
- [16] D.J. Rose, R.E. Tarjan and G.S. Leuker, *Algorithmic aspects of vertex elimination on graphs*, SIAM J. Comput. 5 (1976) 266-283.
- [17] J. Spinrad, *Efficient Graph Representations*, Field Institute Monographs 19, American Mathematical Society, Providence (2003).
- [18] K. Truemper, *Alpha-balanced graphs and matrices and $GF(3)$ -representability of matroids*, Journal of Combinatorial Theory B 32 (1982) 112-139.
- [19] S. Tsukiyama, M. Ide, H. Ariyoshi and I. Shirakawa, *A new algorithm for generating all the maximal independent sets*, SIAM J. Comput. 6 (1977) 505-517.