

promoting access to White Rose research papers



Universities of Leeds, Sheffield and York
<http://eprints.whiterose.ac.uk/>

This is an author produced version of a paper published in **Proceedings of the National Academy of Sciences of USA.**

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/76303/>

Paper:

De Marcos Lousa, C, van Roermund, CW, Postis, VL, Dietrich, D, Kerr, ID, Wanders, RJ, Baldwin, SA, Baker, A and Theodoulou, FL (2013) *Intrinsic acyl-CoA thioesterase activity of a peroxisomal ATP binding cassette transporter is required for transport and metabolism of fatty acids*. Proceedings of the National Academy of Sciences of USA, 110 (4). 1279 – 1284.

<http://dx.doi.org/10.1073/pnas.1218034110>

Intrinsic acyl-CoA thioesterase activity of a peroxisomal ABC transporter is required for transport and metabolism of fatty acids.

Carine De Marcos Lousa^{1,7}, Carlo W. van Roermund^{2,7}, Vincent L.G.Postis³, Daniela Dietrich^{4*}, Ian D. Kerr⁵, Ronald J.A. Wanders², Stephen A. Baldwin³, Alison Baker^{1,6} and Frederica L. Theodoulou⁴.

1. Centre for Plant Sciences, School of Biology, University of Leeds, Leeds, LS2 9JT, UK 2. Laboratory of Genetic Metabolic Diseases, Departments of Pediatrics and Clinical Chemistry, Academic Medical Centre, University of Amsterdam, The Netherlands. 3. Astbury Centre for Structural Molecular Biology, School of Biomedical Sciences, University of Leeds, Leeds, LS2 9JT, UK 4. Biological Chemistry and Crop Protection Department, Rothamsted Research, Harpenden, AL5 2JQ, UK 5. School of Biomedical Sciences, Queen's Medical Centre, University of Nottingham, Nottingham 6. To whom correspondence should be addressed. Email: a.baker@leeds.ac.uk 7. These authors made an equal contribution * Current address: Centre for Plant Integrative Biology, Sutton Bonington Campus, University of Nottingham, Loughborough, LE12 5RD, UK

Submitted to Proceedings of the National Academy of Sciences of the United States of America

Peroxisomes are organelles that perform diverse metabolic functions in different organisms, but a common function is β -oxidation of a variety of long chain aliphatic, branched and aromatic carboxylic acids. Import of substrates into peroxisomes for β -oxidation is mediated by ABC transporter proteins of subfamily D, which includes the human ALDP protein defective in X-linked adrenoleukodystrophy. Whether substrates are transported as CoA esters or free acids has been a matter of debate. Using COMATOSE (CTS), a plant representative of the ABCD family, we demonstrate that there is a functional and physical interaction between the ABC transporter and the peroxisomal long chain acyl-CoA synthetases LACS6 and 7. We expressed recombinant CTS in insect cells and showed that membranes from infected cells possess fatty acyl-CoA thioesterase activity which is stimulated by ATP. A mutant, S810N, which is defective in fatty acid degradation in vivo retains ATPase activity but has strongly reduced thioesterase activity, providing strong evidence for the biological relevance of this activity. Thus CTS, and most likely the other ABCD family members, represent rare examples of polytopic membrane proteins with an intrinsic additional enzymatic function that may regulate the entry of substrates into the β -oxidation pathway. The cleavage of CoA raises questions about the side of the membrane where this occurs and this is discussed in the context of the peroxisomal CoA budget.

ABC transporter | beta oxidation | COMATOSE | peroxisomes | thioesterase

Introduction

The peroxisome is the sole site of β -oxidation of fatty acids and related molecules in plants and fungi and is essential in metabolism of very long chain fatty acids and bioactive lipid molecules in mammals. ATP Binding Cassette (ABC) proteins of subfamily D are required for the transport of these substrates across the peroxisome membrane (1). These are Pxa1p/Pxa2p in yeast, ALDP (ABCD1), ALDPR (ABCD2) and PMP70 (ABCD3) in mammals and Comatose (CTS; also known as PED3, PXA1 and AtABCD1) in plants. Defects in ALDP result in X-linked adrenoleukodystrophy, a neurological disorder in which very long chain fatty acids accumulate (2). Similarly, *cts* mutants are defective in germination and mobilisation of stored triacylglycerol (3).

Activation by formation of a Coenzyme A (CoA) thioester is a pre-requisite for entry of substrates into β -oxidation (4) but whether ABCD proteins accept free fatty acids or acyl-CoAs has been contentious. Arabidopsis mutants lacking CTS accumulate acyl-CoAs (5) and the basal ATPase activity of the protein is stimulated by acyl-CoAs rather than free fatty acids (6). Studies with yeast cells and trypanosomes suggest that Pxa1p/Pxa2p and the trypanosome ABCD protein GAT1 transport acyl-CoAs (7,8)

and human ALDP, ALDR and Arabidopsis CTS are all able to complement the *Saccharomyces cerevisiae* *pxa1/pxa2* Δ mutant (6,9,10). Conversely, in Arabidopsis, genetic evidence indicates that CTS and the peroxisomal long chain acyl-CoA synthetases LACS6 and 7 act in the same pathway, suggesting that free fatty acids are transported (11). Moreover, knock down of the Arabidopsis peroxisomal adenine nucleotide translocators inhibits fatty acid degradation, consistent with a need for intra peroxisomal ATP for fatty acid activation (12).

To reconcile these differences, a model was proposed in which CTS transports fatty acyl-CoA but the CoA moiety is removed either by the transporter or by a peroxisomal thioesterase (11). Recently, complementation of the *S. cerevisiae* *pxa1/pxa2* Δ mutant by human ALDP was shown to be dependent upon the presence of the peroxisomal fatty acyl-CoA synthetase, Faa2p (13), indicating that CoA is cleaved during or directly after transport. CoA cleavage was demonstrated by isotopic labelling but the protein responsible for the acyl-CoA thioesterase activity was not identified (13). Peroxisomes of mammals, yeast and plants contain acyl-CoA thioesterases which catalyse the hydrolysis of fatty acyl-CoAs to free fatty acids and CoASH but their physiological roles are not well characterised. It has been suggested that these enzymes are important for maintaining CoASH at optimal levels during periods of increased fatty acid oxidation (14).

In this study we demonstrate that CTS requires Faa2p or the targeting of the Arabidopsis synthetases LACS6 or LACS7 to yeast peroxisomes to support β -oxidation of oleate in *S. cerevisiae*. This functional interaction between transporter and synthetase is also reflected in a physical interaction *in planta*. We expressed CTS in insect cells and demonstrated that the protein has intrinsic thioesterase activity which is stimulated by ATP and is essential for fatty acid degradation in yeast and *in planta*. This resolves the long standing debate about the transport substrate for ABCD transporters and provides a rare example of a transporter which also possesses an intrinsic accessory enzyme activity.

Reserved for Publication Footnotes

137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204




Fig. 1. Peroxisomal acyl-CoA synthetase is required for CTS-dependent transport The *Saccharomyces cerevisiae* *pxa1/pxa2Δ* and *pxa1/pxa2/faa2Δ* mutants were transformed as indicated. A, B: yeast cells grown on oleate medium were incubated with [14 C] oleic acid (C18:1) and β -oxidation rates were measured. Genotypes are shown to the left of the graphs. The *fox1Δ* mutant lacks acyl-CoA oxidase (the first enzyme of β -oxidation) and serves as a negative control. The activity in wild type cells was taken as a reference (100 %) and values are means \pm SD of 3 independent experiments. C-E: 17,000.g pellets of oleate-grown *pxa1/pxa2/faa2Δ* cells expressing LACS6 (C), LACS7 (D) or mis-targeted LACS7 (LACS7-PTS; E) were subjected to Nycodenz equilibrium density gradient centrifugation and C8:0-CoA synthetase activity was measured in the different fractions. 3-hydroxy-CoA dehydrogenase (3-HAD) activity was used as a peroxisomal marker. Activity in each fraction is expressed as a percentage of the total activity measured throughout the gradient.




Fig. 2. CTS forms a complex with peroxisomal acyl-CoA synthetase in planta Membrane proteins from *Arabidopsis thaliana* suspension cultured cells were solubilised with β -dodecyl maltoside. A: solubilised membrane proteins were subjected to blue-native PAGE (in first dimension) followed by denaturing SDS-PAGE and immunoblotting for CTS. B: solubilised membrane proteins were separated in a sucrose step gradient. Fractions were collected from the top of the gradient and subjected to immunoblotting with a range of antibodies as indicated. T: total membranes. Positions of molecular weight markers (sizes in kDa) are shown above (first dimension or sucrose gradient) and to the left (second dimension) of panels A and B. C: Solubilised membrane proteins were subjected to immunoprecipitation using anti-CTS beads, separated by SDS PAGE and immunoblotted with anti-LACS and anti-CTS antibody. Lane 2, immunoprecipitate; lane 1, mock-immunoprecipitation without protein extract; lane 3, immunoblot of protein extract probed with anti-LACS and anti-CTS antibody, to show positions of LACS and CTS. Figures to the left of the panel indicate positions of molecular weight markers (sizes in kDa).

Results

Peroxisomal acyl-CoA synthetase is essential for CTS-dependent fatty acid β -oxidation

Expression of CTS in the *S. cerevisiae* *pxa1/pxa2Δ* mutant permitted this strain to metabolise oleate (C18:1) at approximately 90% of the wild type level (Fig. 1A) and (6). A mutation, S810N, that compromises CTS-dependent fatty acid β -oxidation in plants (15,16) also resulted in an inability to support oleate β -oxidation in yeast (Fig 1A), despite the protein being correctly targeted to peroxisomes (Fig. S1). Expression of wild type CTS in the *pxa1/pxa2/faa2Δ* triple mutant resulted in only 12% of the wild type activity (Fig. 1B), in marked contrast to the β -oxidation

activity of cells where CTS was expressed in the *pxa1/pxa2Δ* background (Fig 1A), showing dependence of CTS on Faa2p.

In *Arabidopsis* two peroxisomal acyl-CoA synthetases with a broad substrate range are involved in β -oxidation of fatty acids (11,17). When either of these synthetases was co-expressed with CTS in the triple mutant background, β -oxidation activity was restored to approximately 40% (LACS6) and 80% (LACS7) of the wild type level. Expression of either synthetase in the absence of CTS gave a much smaller rescue of oleate β -oxidation showing that the effect of CTS plus LACS is additive (Fig. 1B). This reflects the ability of yeast to import free fatty acids via an ABCD-independent route that requires Faa2p and which accounts for

205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272




Fig. 3. Recombinant CTS has fatty acyl-CoA thioesterase activity Membranes were isolated from *sf9* cells expressing wild-type and mutant forms of CTS. A: ATPase activity in membranes from control (non-infected) and CTS-expressing cells, in the presence and absence of AIF_x. B: acyl-CoA thioesterase activity of membranes expressing wild type CTS with different acyl-CoA substrates (20 μM) in the presence and absence of ATP. Activity was absent from membranes from control (non-infected) cells. C: 18:0-CoA thioesterase activity as a function of substrate concentration (in the presence of ATP). Values represent mean ± SD (n=4). D: relative 18:0-CoA thioesterase activities of wild type CTS and S810N mutant in the presence and absence of ATP. E: relative ATPase activities of CTS and S810N mutant. F: Ponceau-stained protein profile of membranes expressing wild type CTS and S810N mutant, and corresponding immunoblot (α-CTS) stained with anti-CTS antibody.




Fig. 4. Models showing role of fatty acyl-CoA cleavage in peroxisomal import of substrates for β-oxidation in Arabidopsis Fatty acids (FA) are activated to CoA esters extra-peroxisomally by as-yet unidentified acyl-CoA synthetases. The ABC transporter, CTS, binds fatty acyl-CoAs on the cytosolic side of the peroxisomal membrane, the CoA moiety is cleaved by CTS and free FA are imported into the peroxisome. Transport is energised by ATP hydrolysis. In model A, CoA is released in the peroxisome lumen, where it is re-esterified to acyl-CoAs by the action of peroxisomal acyl-CoA synthetases LACS6 and LACS7. FA-CoAs then enter the β-oxidation pathway. The peroxisomal ATP pool is supplied by peroxisomal adenine nucleotide translocators, PNC1 and PNC2 (not shown for clarity). Alternatively, in model B, CoA is released on the cytosolic side of the peroxisomal membrane and must enter peroxisomes via the PXCN carrier, probably in exchange for AMP. In both cases, a proportion of FA may cross the peroxisomal membrane by a CTS-independent route and is activated by LACS6/7 (indicated by dashed line).

the residual growth of the *pxa1/pxa2Δ* mutant on oleate (Fig. 1A (9,18).

Complementation was dependent on the correct peroxisomal targeting of the synthetase. The localisation and function of LACS6 and 7 were assessed by measuring octanoyl-CoA (C8:0) synthetase activity in different subcellular fractions, since this activity is absent from the cytosol of yeast (18). In *pxa1/pxa2/faa2Δ* cells transformed with LACS6 or 7, C8:0-CoA synthetase activity was detectable in cell lysates, organelle pellets, and purified peroxisomes but was absent from cells transformed with vector lacking insert (Fig. 1C-E; S2A-B). Removal of the PTS1 and 2 peroxisome targeting signals from LACS7 (LACS7-PTS) resulted in loss of peroxisomal C8:0-CoA synthetase activity and inability to restore β-oxidation of oleate (Fig. 1B, E). The peak peroxisomal fraction (#3) from the gradients shown in Fig. 1C and D was assayed for C18:1-CoA synthetase activity. This revealed that LACS7 has a much higher ratio of C18:1- to C8:0-CoA synthetase activity (Fig. S2C) as also reported previously (17),

thereby explaining the more efficient rescue of oleate β-oxidation by LACS7 compared to LACS6 (Fig. 1B).

CTS is present in a complex with LACS6/7 in planta

These results indicate a functional interaction between CTS and the peroxisomal acyl-CoA synthetases LACS6, LACS7 and Faa2p, therefore evidence for a physical interaction was sought. Membranes were prepared from *Arabidopsis thaliana* suspension culture cells, solubilised with β-dodecylmaltoside (β-DDM), separated on a blue native gel followed by a denaturing (SDS-PAGE) second dimension and then immunoblotted with an antibody against CTS. The transporter (MW 150 kDa) was found predominantly in a complex of ca. 700 kDa (Fig 2A). The experiment was repeated using membranes from seedlings that express a CTS-GFP fusion from the native CTS promoter. This construct is functional, as judged by complementation of the *cts-1* mutant (Fig. S3A). On a 2D gel, a similar complex of ca. 700 kDa was observed when probed with anti-GFP antibody, although the apparent size of the denatured CTS-GFP species was approximately 170kDa, as expected from the addition of GFP (28 kDa) (Fig. S3B). The complex is quite stable, since it was also detected in the presence of Triton X-100 (Fig. S3C). When this complex was probed with both anti-CTS and anti-LACS6/7 antibodies (19), two pools of LACS immuno-reactivity were seen (Fig. S3C). The most intense migrated at around 150 kDa, the expected size for the dimeric LACS proteins, but a significant immunoreactivity was also detectable in the higher molecular weight fractions between 700 and 1100 kDa that also contained CTS (Fig. S3C).

Detergent solubilised membranes were fractionated on a sucrose gradient followed by SDS-PAGE and immunoblotting. CTS sedimented in a broad peak with maximum intensity in fractions 6-9, corresponding to complexes >700 kDa (Fig. 2B). The same fractions were probed with antibodies against LACS6/7 and the β-oxidation enzymes acyl-CoA oxidase 1 (ACX1), multifunctional protein 2 (MFP2) and 3-ketoacyl-CoA thiolase (KAT2). MFP2 and KAT2 sedimented in the lightest fractions of the gradient, and showed no co-migration with CTS, as would be expected for soluble proteins. However both LACS and ACX1 showed a distinct minor peak in fraction 8 which also contained CTS.

While co-migration of LACS proteins with CTS is suggestive of their being part of the same complex, this does not provide evidence for a physical interaction between the two. Therefore solubilised membranes were immunoprecipitated with anti-CTS antibodies, and the immunoprecipitates probed with anti-LACS6/7 and anti-CTS antibodies (Fig. 2C). Since the antibody chains from the immunoprecipitation (IP) react with the secondary antibody in the blot, a mock IP was carried out without extract but with anti-CTS antibodies (lane 1) and processed alongside the extract

containing sample (lane 2). Bands present in lane 2 but not lane 1 represent proteins specifically immunoprecipitated by the anti-CTS antibody. The ca. 70 kDa band that is absent from control lane 1 but is present in lane 2 co-migrates with native LACS as detected by immunoblotting of the extract without prior IP (Fig. 2C, lane 3).

CTS exhibits intrinsic thioesterase activity

CTS was expressed in insect cells to investigate its biochemical properties. Expression of the active, full-length protein was demonstrated by the presence in membrane preparations of immunoreactive protein of MW ca. 150 kDa and detection of ATPase activity that was much higher than uninfected controls (Fig. 3A,F), and was inhibited by AIF_x as previously described (6). Insect cell membranes containing CTS exhibited thioesterase activity towards 18:0-CoA which was absent from membranes prepared from uninfected cells (Fig. 3B). ATP stimulated C18:0-CoA thioesterase activity >2 fold. C14:0-CoA thioesterase exhibited about 60% of the activity of C18:0-CoA (Fig. 3B). In the presence of ATP, acyl-CoA thioesterase activity exhibited Michaelis-Menten behaviour; the K_m for C18:0 CoA was $1.05 \pm 0.16 \mu\text{M}$, V_{max} $38.15 \pm 7.27 \text{ nmol mg CTS}^{-1} \text{ s}^{-1}$ and quantification of CTS levels by immunoblotting permitted estimation of k_{cat} at 5.7 s^{-1} (Fig. 3C).

A mutant, *ped3-4*, has been described which is a *cts* missense mutation, S810N. *In planta* this mutant germinates but cannot break down fatty acids and establish (15,16) and it failed to complement the *S. cerevisiae pxa1/pxa2Δ* mutant for oleate β-oxidation (Fig. 1A). Membranes from insect cells expressing CTS_{S810N} exhibited ca. 70% of WT CTS-dependent ATPase activity but only 40% thioesterase activity (Fig. 3D,E). No thioesterase activity was detected in the absence of ATP, indicating that this activity retained nucleotide sensitivity, similar to the wild type (Fig. 3D).

Discussion

Previous studies have provided indirect evidence in favour of both acyl-CoAs and free fatty acids as transport substrates of ABCD proteins. It was proposed that the solution to this paradox is that the CoA moiety is cleaved during transport (11), and indirect evidence based on isotope labelling patterns of yeast cells is consistent with this (13). The current study provides direct evidence that CTS has an ATP-stimulated thioesterase activity although the protein has no obvious homology with known thioesterases. Previously described acyl CoA thioesterases are not stimulated by ATP, thus this appears to be a novel type of activity. By analogy with thioesterases of known structure and mechanism, catalytic residues could potentially include either conserved acidic, hydroxyl, histidine and amide side chains, as in the "hot-dog fold" thioesterases, or the triad of a nucleophilic residue, histidine and acidic residue found in α/β hydrolase-type thioesterases (20). Members of the clade of plant, yeast and metazoan ABCD transporters known to be involved in peroxisomal fatty acid import (21) contain highly conserved residues of these types at a number of positions. In human ALDP some of these have been implicated in X-linked adrenoleukodystrophy, attesting to their functional importance. Future work will systematically explore the role of these residues in CTS, in the context of our structural model of a lumen-facing conformation of the protein (16).

A functional and physical interaction between the peroxisomal ABC transporter CTS and the peroxisomal acyl-CoA synthetases LACS6 and LACS7 could also be demonstrated. Thus CTS (and probably the other peroxisomal ABCD transporters) cleaves acyl-CoA as part of the transport cycle, with the free fatty acid requiring subsequent reactivation by peroxisomal acyl-CoA synthetases (Fig. 4). Vectorial acylation is a common feature of fatty acid transport across membranes. In mammals, fatty acids are transported into mitochondria by conversion to acyl

carnitine by carnitine acyl transferases and re-esterification to CoA in the matrix, thus the mitochondrion maintains a separate CoA pool from the cytosol. Uptake of fatty acids into yeast cells and adipocytes requires co-operation of a long chain acyl CoA synthetase and a fatty acyl transport protein (FATP) that may itself also possess acyl CoA synthetase activity (22). Recently *S. cerevisiae* Fat1p was shown to be dually targeted to both the plasma membrane and the peroxisomal membrane, and to form a functional interaction with Pxa1p/Pxa2p that is required for the stability of the former (13). The activity of Fat1p explains the ability of *faa2Δ* cells to metabolise LCFA in the presence of Pxa1p/Pxa2p (Fig. 1B). CTS cannot substitute for Pxa1p/Pxa2p when Faa2p is also absent, suggesting that it is unable to interact functionally with Fat1p (Fig. 1B). HsABCD1 and HsABCD2 are similarly unable to rescue the *pxa1/pxa2/faa2* triple mutant (13).

The S810N mutant has markedly reduced thioesterase activity and is unable to support oleate β-oxidation when expressed in the *S. cerevisiae pxa1/pxa2Δ* mutant, demonstrating that the thioesterase activity is important for biological function *in vivo*. Ser810 is predicted to be located in the second transmembrane (TM) helix of the second TM domain (i.e. TM helix 8 of the complete CTS molecule, which is a fused heterodimer) and the equivalent residue in ALDP (Ala141) is mutated in some X-ALD patients (www.x-ald.nl), supporting its functional importance. In plants, the S810N mutant (originally described as *ped3-4*; ref. 15) germinates slowly but cannot degrade its oil bodies or establish (15,16). The germination defect of mutants with a severe block in β-oxidation has been ascribed to an inability to metabolise the jasmonic acid precursor, 12-oxophytodienoic acid (OPDA; 23), a plant hormone which is thought to be transported into the peroxisome by CTS for β-oxidation (24). It is possible that cleavage of activated OPDA is less severely affected by the S810N mutation compared to fatty acyl-CoAs and that the residual thioesterase activity in *ped3-4* may be sufficient to reduce OPDA concentrations below the physiological threshold for germination, or alternatively that free OPDA can be accepted by the transporter.

CoA cleavage by CTS might appear to be an energetically wasteful process, since an additional two high energy phosphate bonds are expended for every fatty acid converted to a CoA thioester following import into the peroxisome. However energy expenditure is frequently utilised as a means of regulation or control, e.g. GTP-dependent proof reading in protein synthesis. Which side of the membrane CoA is released is unknown but has important implications for the CoA budget. Release on the cytosolic side would prevent the peroxisome acting as a sink for CoA, especially during periods of high flux through β-oxidation, but would imply an independent uptake route for this cofactor (Fig. 4). The Arabidopsis peroxisome membrane protein PXN was originally identified as an NAD⁺ transporter (25) but has recently been shown to be able to transport CoA (26). Surprisingly, *pxn* knockout mutants were still able to degrade fatty acids, albeit at a reduced rate (25), suggesting the existence of more than one route for CoA import. Thus it is possible that the thioesterase activity of CTS releases the CoA moiety on the luminal side of the peroxisome membrane (Fig. 4). This scenario implies that a mechanism for CoA removal exists, such as degradation via peroxisomal nudix hydrolases (27).

Sucrose gradients and native gels revealed that the majority of CTS resides in a complex of >700 kDa. Because of the effect of the detergent on protein migration it is impossible to size accurately but is indicative of an oligomer of three or more CTS molecules. This is consistent with other ABC proteins including PMP70 and ALDP, which form homo-oligomers (28). A sub-population of LACS and ACX1 proteins was associated with the CTS complex, which is in accordance with the previously reported membrane association of LACS6/7 (19). Although the

functional significance of a physical interaction between ALDP and very long chain acyl-CoA synthetase has been disputed (1,4), here we provide strong evidence for a physical and functional interaction between CTS and LACS6/7 in Arabidopsis. Since CTS, as a broad specificity transporter, appears to be a gateway through which the bulk of β -oxidation substrates must pass (1,6), the cleavage and re-esterification of transported molecules could allow the channelling or prioritising of substrates for metabolism via induction, activation or even specific recruitment of distinct acyl-CoA synthetases. Plants, in particular, contain a large family of acyl activating enzymes, several of which are targeted to the peroxisome (29)

There are a number of examples of transporters which interact functionally with enzymes to create membrane transport metabolons involved in substrate channelling and metabolic regulation (30) and several transporters, particularly in prokaryotes, have acquired related secondary enzyme activities by fusion of additional domains (31). However CTS, and perhaps the other ABCD proteins, appears to be a rare example of a polytopic membrane transporter with an intrinsic additional enzymatic function that controls entry of diverse substrates to a metabolic pathway.

Methods

Plasmids

Details of plasmid construction, mutagenesis and primers are given in Supplementary data.

Yeast expression

Yeast strains are given in Table S1. Yeast growth, β -oxidation, subcellular fractionation and assays for 3-hydroxy-CoA dehydrogenase and acyl-CoA synthetase activity were performed as described in (6, 13) and Supplementary material.

Plant membrane extraction

Arabidopsis suspension culture cells were collected by centrifugation (1,000.g, 15 min), washed twice in cold PBS and once in extraction buffer (EB) [50 mM Tris-HCl pH 8.2, 20% glycerol, 2 mM EDTA, 1 mM PMSF, 5% plant protease inhibitor (Sigma)]. The cell pellet was ground in liquid N_2 and the frozen powder resuspended on ice at 1 mg/ml in EB. Following filtration through Miracloth, the extract was clarified by centrifugation (3000.g, 5 min at 4°C) and membranes were collected by centrifugation at 25,000.g for 20 min at 4°C. The pellet was resuspended in EB.

2D gels

Membranes were resuspended at 1 mg/ml in solubilisation buffer [50 mM Tris-HCl pH 8.2, 200 mM NaCl, 20% glycerol, 1.5 mM $MgCl_2$, 3% DDM, 5% plant protease inhibitor (Sigma), 2 mg/ml Pefabloc] and incubated on ice for 1 h. Insoluble material was removed by centrifugation at 100,000.g for 30 min. Solubilised material was loaded on a 4-16% Blue Native gel (Invitrogen) following manufacturer's recommendations. A single lane was applied to a 10% SDS-PAGE gel and sealed by addition of 6% acrylamide mix. Following electrophoresis, proteins were transferred to PVDF and detected with either CTS or GFP antibody as described in (6).

Sucrose gradients

Solubilised membrane proteins (200 μ l) were layered onto a 25-60% (w/v) sucrose step gradient (step volume of 250 μ l; step increment, 5%). Gradients were centrifuged overnight at 214,000.g at 4°C. Fractions (180 μ l) were collected and precipitated with 17% (w/v) TCA for 20 min on ice. The precipitated proteins were collected by centrifugation at 17,000.g for 30 min at 4°C. Pellets were resuspended in 25 μ l 1M Tris-HCl, pH 9.4. Following

immunoblotting for CTS, membranes were stripped and probed with antisera raised to LACS6/7 (19), ACX1 (32), thiolase (33) and MFP2 (34). The absence of any residual signal was confirmed followed each stripping step.

Immunoprecipitation

Solubilised membrane proteins were diluted at 1 mg/ml in IP buffer [10 mM Tris-HCl pH 7.4, 150 mM NaCl, 0.3% (w/v) DDM] and incubated with pre-equilibrated protein A beads (Thermo Scientific Pierce) for 1 h at 4°C. The beads were removed by centrifugation for 1 min at 1,000.g and the supernatant was diluted twice in IP buffer. 10 μ l of affinity-purified CTS antibody was added and the mix was incubated at 4°C overnight. 50 μ l of protein A beads were then added and incubated for a further 3h at 4°C. The beads were pelleted at 1,000.g for 1 min and washed four times with IP buffer. The immunoprecipitated proteins were eluted with SDS PAGE sample buffer.

Insect cell expression

Recombinant bacmids encoding CTS or its S810N mutant were prepared using the Bac-to-Bac system (Invitrogen), according to the manufacturer's instructions. *Sf9* insect cells (*Spodoptera frugiperda*) were infected with virus (MOI = 3) and cultured with serum-free SF900-SFM medium (Invitrogen) in a shaking incubator at 100 rpm, 28°C for 72 h. The cells were collected by centrifugation at 1,000.g for 10 min and incubated in a hypo-osmotic buffer [5 mM Tris-HCl pH 7.4, containing Mini complete protease inhibitor-EDTA (Roche)] for 15 min at 4°C. 3 mM DTT was then added and cells were further disrupted using 30 strokes of a tight-fitting Dounce homogeniser. The extract was diluted in homogenisation buffer (5 mM Tris-HCl pH 7.4, 0.5 mM $MgCl_2$, 5 mM EGTA, 1 mM PMSF) and centrifuged at 2,000.g for 20 min at 4°C. Membranes were pelleted at 100,000g for 1h at 4°C. The pellet was then resuspended in resuspension buffer [20 mM Tris pH 7.4, 200 mM NaCl, 1.5 mM $MgCl_2$, 20% (v/v) glycerol; 2 mM DTT, plant protease inhibitor mix (Sigma), 2 mg/ml Pefabloc], homogenised in a small Dounce homogeniser and aliquots were flash frozen in liquid nitrogen before storing at -80°C. CTS expression levels were estimated by quantitative immunoblotting.

ATPase assays

ATPase activity was assayed according to (35). Briefly, insect cell membranes expressing wild-type CTS or S810N mutant were mixed with buffer containing 10 mM Tris-HCl pH 7.4, 0.1 mM EGTA, 1 mM DTT, 15 mM $MgSO_4$, 2 mM ouabain and 10 mM sodium azide (final concentrations) in the presence of 10 mM AlF_4^- when indicated and incubated for 3 min at 37°C (total volume, 50 μ l). The reaction was started by the addition of ATP and incubated for 30 min at 37°C.

Thioesterase assays

Acyl-CoA thioesterase activity of CTS-expressing insect cell membranes was measured spectrophotometrically using 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB). Membranes were mixed at a final concentration of 0.3 mg/ml with 200 μ M DTNB, 4 μ M BSA, 10 mM ATP in 50 mM potassium phosphate buffer, pH 8.0 in the presence of various concentrations of acyl-CoAs (Avanti Polar Lipids, Inc.). The reaction was started by the addition of membranes and specific activities were determined by measuring the change in absorbance at 412 nm for 10 min. $\Sigma = 17780 M^{-1} cm^{-1}$.

Acknowledgements.

We thank Ian Graham, Greg Howe, Luigi DeBellis and Doug Muench for antibodies, Amelia Lesiuk and Vincent Agboh for assistance with insect cell culture and Tom Lanyon-Hogg for advice on thioesterase assays. This work was funded by the Biotechnology and Biological Sciences Research Council (BBSRC) (grants BB/F007108/1 to FLT and BB/F007299/1 to AB/SAB). Rothamsted Research receives grant-aided support from the BBSRC of the UK. **Author contributions** CDML, CVR, VP and FLT performed research, CDML, CVR, VP, IDK, RJAW, SAB, FLT and AB designed research and analysed data, DD and IDK contributed new reagents, CDML, AB, SAB and FLT wrote the manuscript. All authors read and approved the manuscript before submission.

functions as a homodimer and accepts acyl-CoA esters. *FASEB J* 22:4201-4208.

- Theodoulou FL, Holdsworth M, Baker A (2006) Peroxisomal ABC transporters. *FEBS Lett* 580:1139-1155.
- Kemp S, Theodoulou FL, Wanders RJ (2011) Mammalian peroxisomal ABC transporters: from endogenous substrates to pathology and clinical significance. *Br J Pharmacol* 164:1753-1766
- Theodoulou FL, Eastmond PJ. (2012) Seed storage oil catabolism: a story of give and take. *Curr Opin Plant Biol* 15:322-328.
- Watkins, PA, Ellis JM (2012) Peroxisomal acyl-CoA synthetases. *Biochim Biophys Acta* 1822:1411-1420
- Footitt S, et al. (2002) Control of germination and lipid mobilization by COMATOSE, the Arabidopsis homologue of human ALDP. *EMBO J* 21:2912-2922.
- Nyathi Y, et al. (2010) The Arabidopsis peroxisomal ABC transporter, comatose, complements the *Saccharomyces cerevisiae* *pxa1 pxa2* mutant for metabolism of long-chain fatty acids and exhibits fatty acyl-CoA-stimulated ATPase activity. *J Biol Chem* 285:29892-29902.
- Verleur N, Hettrema EH, van Roermund CW, Tabak HF, Wanders RJ. (1997) Transport of activated fatty acids by the peroxisomal ATP-binding-cassette transporter Pxa2 in a semi-intact yeast cell system. *Eur J Biochem* 249:657-661.
- Igoillo-Esteve M, et al. (2011) Glycosomal ABC transporters of *Trypanosoma brucei*: characterisation of their expression, topology and substrate specificity. *Int J Parasitol* 41:429-438
- van Roermund CW, et al. (2008) The human peroxisomal ABC half transporter ALDP
- van Roermund CW, Visser WF, Ijlst L, Waterham HR, Wanders RJ. (2011) Differential substrate specificities of human ABCD1 and ABCD2 in peroxisomal fatty acid β -oxidation. *Biochim Biophys Acta* 1811:148-152.
- Fulda M, Schnurr J, Abbadi A, Heinz E, Browse J (2004) Peroxisomal Acyl-CoA synthetase activity is essential for seedling development in *Arabidopsis thaliana*. *Plant Cell* 16:394-405.
- Linka N, et al. (2008) Peroxisomal ATP import is essential for seedling development in *Arabidopsis thaliana*. *Plant Cell* 20:3241-3257.
- van Roermund CW, et al. (2012) Peroxisomal Fatty Acid Uptake Mechanism in *Saccharomyces cerevisiae*. *J Biol Chem* 287:20144-20153.
- Kirky B, Roman N, Kobe B, Kellie S, Forwood JK (2010) Functional and structural properties of mammalian acyl-coenzyme A thioesterases. *Prog Lipid Res* 49:366-377.
- Hayashi M, et al. (2002) Ped3p is a peroxisomal ATP-binding cassette transporter that might supply substrates for fatty acid beta-oxidation. *Plant Cell Physiol* 43:1-11.
- Dietrich D, et al. (2009) Mutations in the Arabidopsis peroxisomal ABC transporter COMATOSE allow differentiation between multiple functions in *planta*: insights from an allelic series. *Mol Biol Cell* 20:530-543.
- Fulda M, Shockey J, Werber M, Wolter FR, Heinz E (2002) Two long-chain acyl-CoA synthetases from Arabidopsis thaliana involved in peroxisomal fatty acid beta-oxidation. *Plant J* 32:93-103.

681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748

18. Hettema EH, et al. (1996) The ABC transporter proteins Pat1 and Pat2 are required for import of long-chain fatty acids into peroxisomes of *Saccharomyces cerevisiae*. *EMBO J* 15:3813-3822.

19. Hayashi H, et al. (2002) Molecular characterization of an Arabidopsis acyl-coenzyme A synthetase localized on glyoxysomal membranes. *Plant Physiol* 130:2019-2026.

20. Cantu DC, Chen Y, Reilly PJ (2010) Thioesterases: a new perspective based on their primary and tertiary structures. *Protein Sci* 19: 1281-1295.

21. Nyathi Y, et al. (2012) Pseudo half-molecules of the ABC transporter, COMATOSE, bind Pex19 and target to peroxisomes independently but both are required for activity. *FEBS Lett* 586: 2280-2286.

22. Kazantzis M, Stahl A (2012) Fatty acid transport proteins; implications in physiology and disease. *Biochim Biophys Acta* 821: 852-857.

23. Dave A, et al. (2011) 12-oxo-phytyldienoic acid accumulation during seed development represses seed germination in Arabidopsis. *Plant Cell* 23:583-599.

24. Theodoulou FL, et al. (2005) Jasmonic acid levels are reduced in COMATOSE ATP-binding cassette transporter mutants. Implications for transport of jasmonate precursors into peroxisomes. *Plant Physiol* 137:835-840

25. Bernhardt K, Wilkinson S, Weber AP, Linka N (2012) A peroxisomal carrier delivers NAD⁺ and contributes to optimal fatty acid degradation during storage oil mobilization. *Plant J* 69:1-13.

26. Agrimi G, Russo A, Pierri CL, Palmieri F (2012) The peroxisomal NAD⁺ carrier of *Arabidopsis thaliana* transports coenzyme A and its derivatives. *J Bioenerg Biomembr* 44:333-340.

27. Antonenkov VD, Hiltunen JK. (2012) Transfer of metabolites across the peroxisomal membrane. *Biochim Biophys Acta* 1822:1374-1386.

28. Guimarães CP, et al. (2004) Mouse liver PMP70 and ALDP: homomeric interactions prevail *in vivo*. *Biochim Biophys Acta* 1689:235-243.

29. Shockey J, Browse J (2011) Genome-level and biochemical diversity of the acyl-activating enzyme superfamily in plants. *Plant J*. 66:143-160.

30. Moraes TF, Reithmeier RAF (2012) Membrane transport metabolons. *Biochim Biophys Acta* 1818:2687-2706.

31. Reddy VS, Shlykov MA, Castillo R, Sun EI, Saier MH Jr (2012) The major facilitator superfamily (MFS) revisited. *FEBS J* 279:2022-2035.

32. Schillmiller AL, Koo AJK, Howe GA (2007) Functional diversification of acyl-coenzyme oxidases in jasmonic acid biosynthesis and action. *Plant Physiol* 143:812-824.

33. Germain V, et al. (2001) Requirement for 3-ketoacyl-CoA thiolase-2 in peroxisome development, fatty acid beta-oxidation and breakdown of triacylglycerol in lipid bodies of Arabidopsis seedlings. *Plant J* 28:1-12.

34. Chuong SD, Mullen RT, Muench DG (2002) Identification of a rice RNA- and microtubule-binding protein as the multifunctional protein, a peroxisomal enzyme involved in the beta-oxidation of fatty acids. *J Biol Chem* 277:2419-2429.

35. Chifflet S, Torriglia A, Chiesa R, Tolosa S (1988) A method for the determination of inorganic phosphate in the presence of labile organic phosphate and high concentrations of protein: Application to lens ATPases. *Anal Biochem* 168:1-4.

Conflict of Interest
None declared

Submission PDF

749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816