

promoting access to White Rose research papers



Universities of Leeds, Sheffield and York
<http://eprints.whiterose.ac.uk/>

This is an author produced version of a paper published in **Hydrological Processes**

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/id/eprint/77261>

Paper:

Brown, LE, Cooper, L, Holden, J and Ramchunder, SJ (2010) *A comparison of stream water temperature regimes from open and afforested moorland, Yorkshire Dales, northern England*. *Hydrological Processes*, 24 (22). 3206 - 3218.

<http://dx.doi.org/10.1002/hyp.7746>

A comparison of stream water temperature regimes from open and afforested moorland, Yorkshire Dales, northern England

Journal:	<i>Hydrological Processes</i>
Manuscript ID:	HYP-09-0459.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	
Complete List of Authors:	Brown, Lee; University of Leeds, Geography Cooper, Luke; University of Leeds, Geography Holden, Joseph; University of Leeds, Geography Ramchunder, Sorain; University of Leeds, Geography
Keywords:	peatland, thermal regime, coniferous plantation, catchment-scale, river, afforestation



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 **A comparison of stream water temperature regimes from**
2 **open and afforested moorland, Yorkshire Dales, northern England**

3
4 Brown, L.E.*, Cooper, L., Holden, J. & Ramchunder, S.

5
6 Affiliation of all authors: School of Geography, University of Leeds, Woodhouse Lane, Leeds,
7 LS2 9JT, UK

8 * Author for correspondence

9 E: l.brown@leeds.ac.uk

10 T: ++44 (0)113 343 3302

11 F: ++44 (0)113 343 3308

12
13 **Submitted to *Hydrological Processes*: 15 March 2010**

14
15
16 **Running head:** Stream water temperature, Yorkshire Dales

17 **Keywords:** afforestation, catchment-scale, coniferous plantation, peatland, river, thermal regime

Abstract

Despite the known importance of water temperature for river ecosystems, the thermal regime of streams and rivers can be heavily modified by afforestation. Although the nature of the heat budget affecting streams in forested catchments shows high variability in space and time, most studies of stream temperature response to afforestation have lacked replication among streams. This study examined the impacts of coniferous forest plantations on stream water temperature at six sites (three forested and three open moorland) in the Yorkshire Dales, northern England. Our aim was to test the hypothesis that afforestation would alter the thermal regime of streams, leading to reduced year-round thermal variability, and cooler summer/warmer winter water temperatures, relative to streams flowing across open moorland. Data collected from April 2007 to March 2009 showed similar thermal dynamics among all six streams over the study period, although variability in forested streams was markedly lower as expected. Mean and maximum daily water temperature were significantly higher in open moorland streams for much of the year but while some forested streams were warmer than individual moorland streams during winter months (November to February), there was considerable overlap in water temperature between moorland and forest streams. Most stream temperature records showed evidence of low/no winter flow and freezing. These results contrast with many previous studies which have reported warmer temperatures in forested versus open moorland streams during winter, a finding which most likely reflects site-specific hydrological, geomorphological and climatological influences on water temperature in addition to afforestation. This study demonstrates the need for replication of hydrological monitoring when examining the effects of basin-scale management practices and provides further evidence for changes in stream thermal regime following afforestation, a practice which is likely to increase in future due to growing demands for increased forest cover in the UK uplands.

1 INTRODUCTION

2 The components that make up the energy budget of rivers and streams are complex and vary both
3 temporally and spatially (Webb and Zhang, 1999; Hannah et al., 2004; Caissie, 2006). Energy
4 inputs to streams may occur through incident short-wave (solar) and long-wave (downward
5 atmospheric) radiation, condensation, friction at the channel beds and banks, and chemical and
6 biological processes. Losses may include reflection of solar radiation, emission of long-wave
7 (back) radiation and evaporation. Sensible heat and water column-bed energy transfers may
8 cause gains or losses. In addition to these exchanges, energy may be advected by in/out flowing
9 stream discharge, evaporated water, groundwater up/downwelling, tributary inflows and
10 precipitation. Improvements in understanding the natural dynamics of, and human influences on,
11 these processes are increasingly being made due to a recent upturn in the availability of accurate
12 and reliable temperature datalogging technology (Webb et al., 2008).

13 Water temperature is widely recognised as one of the most important water quality parameters.
14 For example, it plays a major role influencing the biological composition of streams and rivers
15 and functional processes such as organic matter decomposition and nutrient cycling (Vannote
16 and Sweeney, 1980; Weatherley and Ormerod, 1990a; Poole and Berman, 2001; Acuña et al.,
17 2008). Water temperature influences the chemical characteristics of running waters by affecting
18 the solubility of oxygen and trace metals, and influencing pH (Berner and Berner, 1996). Despite
19 the known importance of water temperature for river ecosystems, the thermal regimes of many
20 streams and rivers have been modified by humans through direct point sources of thermal
21 pollution (e.g. power stations and industry), river regulation, and afforestation/deforestation
22 (Webb et al., 2008).

23
24

1
2 1 The presence of a forest canopy has long been known to modify the amount of solar radiation
3
4 2 and other meteorological factors influencing stream temperatures (Pluhowski, 1972; Moore et
5
6 3 al., 2005a) but the nature of the heat budget affecting streams in forested catchments typically
7
8 4 shows high variability in space and time. Compared to open environments, forests provide a
9
10 5 microclimate of lower wind speeds, higher humidity and less variable air temperature (Hannah et
11
12 6 al., 2008). Dense canopies can reduce solar radiation by as much as 90%, effectively isolating
13
14 7 streams from their main source of energy (i.e. incoming shortwave radiation: Sinokrot and
15
16 8 Stefan, 1993; Poole and Berman, 2001). In North America, the effects of forest harvesting on
17
18 9 stream temperature have been of concern for 50+ years due to the potential for increased water
19
20 10 temperature to negatively affect fish populations (e.g. Johnson and Jones, 2000; Moore et al.,
21
22 11 2005b). Several studies were undertaken in the 1990s in north Wales which generally showed
23
24 12 lower stream water temperatures under coniferous plantations (Weatherley and Ormerod, 1990b;
25
26 13 Webb and Crisp, 2006). However, the effect of afforestation on stream water temperature has
27
28 14 received only minimal attention in other parts of the UK, with most studies being from Scotland
29
30 15 (Malcolm et al., 2004; Webb and Crisp, 2006; Hannah et al., 2008).

31
32
33 16 Some recent studies have provided insights into the likely effects of upland afforestation on
34
35 17 water temperature. Mean stream water temperatures were found to be reduced by ~0.4 to 0.5°C
36
37 18 in coniferous forests in north Wales and southwest Scotland, respectively (Crisp, 1997; Webb
38
39 19 and Crisp, 2006) while mixed forestry in the Cairngorms, Scotland was found to moderate
40
41 20 temperature extremes (Malcolm et al., 2004). A study by Hannah et al. (2008) in mixed
42
43 21 temperate forests in northeast Scotland suggested cooler, less variable water temperatures in
44
45 22 forested reaches due to reduced net radiation but higher sensible and latent heat fluxes. However,
46
47 23 all of these studies reviewed above examined the thermal regime of only one forested stream or
48
49
50
51
52
53
54
55
56
57
58
59
60

1 one open canopy stream despite the known high spatial variability of energy budgets in forests.
2
3
4 2 Only Weatherley & Ormerod (1990b) have considered the thermal regimes of replicate UK
5
6
7 3 streams under afforested conditions, showing mean daily temperatures to be consistently lower
8
9
10 4 in three forested streams compared to three grazed moorlands during spring and summer. The
11
12 5 lack of replicated studies into forest effects on stream temperature means that questions remain
13
14 6 about the transferability of findings related to stream temperature responses to afforestation.
15
16
17 7

18
19 8 In the UK there has been a substantial increase in upland tree cover over the past 50 years
20
21 9 through coniferous plantations. Upland tree cover will also increase in the next 30 years as
22
23 10 economic drivers may incentivise more coniferous plantation, and Scottish and English agencies
24
25 11 aim to meet new targets for mixed broadleaf tree cover in the uplands (Natural England, 2009).
26
27 12 Forests currently cover some 2.8M ha of the UK (~12% of the UK land surface) with
28
29 13 approximately 58% of this cover being coniferous plantation. Much of the upland coniferous
30
31 14 afforestation has occurred in areas where organic and organo-mineral soils dominate. Of
32
33 15 particular concern is afforestation on blanket peats which cover approximately 15 % of the
34
35 16 British Isles (Tallis, 1998). Blanket peatlands strongly influence stream hydrology, tending to
36
37 17 encourage flashy regimes due to limited buffering of rainfall on near-saturated slopes (Holden
38
39 18 and Burt, 2003b). Coniferous afforestation leads to alterations in upland blanket peatland
40
41 19 hydrological processes with associated water quality changes, in particular acidification
42
43 20 (Reynolds et al., 1988; Neal et al., 1992a; Reynolds et al., 1992). Despite the concerns about
44
45 21 afforestation damage to peatland environments, the decreasing profitability of open moorland,
46
47 22 concerns about the need to reduce downstream flood risk and a desire to mitigate climate change
48
49 23 may lead to upland landowners turning over more of their land to plantations in the near future
50
51 24 (Holden et al., 2007; Natural England, 2009).
52
53
54
55
56
57
58
59
60

1
2
3 1
4 2 This study examined the impacts of coniferous afforestation on the thermal regime of six streams
5
6 3 in the Yorkshire Dales, northern England, where there has been recent conversion of open
7
8 4 blanket peat moorland to coniferous plantation. No studies of afforestation effects on stream
9
10 5 thermal regimes have been undertaken in this region. Water temperature measurements were
11
12 6 made for replicate open moorland and afforested study basins, so that spatial variability in the
13
14 7 effects of afforestation could be examined. We hypothesised that coniferous forest plantations
15
16 8 would alter the thermal regime of blanket peatland streams, leading to reduced year-round
17
18 9 thermal variability with cooler summer and warmer winter temperatures relative to open
19
20 10 moorland streams. The study ran for two annual cycles and involved high resolution (15 min)
21
22 11 water temperature and meteorological measurements enabling accurate characterisation of
23
24 12 diurnal dynamics and thermal extremes.
25
26
27
28

29 13 30 14 **METHODS**

31 15 *Study area*

32
33
34 16 Field observations were made in the Green Field Beck and Oughtershaw Beck river basins,
35
36 17 Upper Wharfedale (Figure 1), east of Pen-y-Gent in the Yorkshire Dales National Park, northern
37
38 18 England (54°2'N; 2°2'W) between 1 April 2007 (day 91) and 31 March 2009 (day 90).
39
40 19 Hereafter, dates are referred to using the calendar day of the year and time is quoted as
41
42 20 Greenwich Mean Time (GMT). The Oughtershaw basin is predominantly open blanket peat
43
44 21 moorland (Wallage et al., 2006) whereas the Green Field basin is predominantly covered by
45
46 22 Spruce forest (Lane et al., 2008). The Oughtershaw basin lies directly to the north of the Green
47
48 23 Field basin and therefore the two share similar physical characteristics and drain from west to
49
50 24 east. Both basins lie on Carboniferous-age Great Scar Limestone, Yoredale Series and Millstone
51
52 25 Grit overlain with a thin layer of glacial boulder clay which provides an impermeable substrate
53
54
55
56
57
58
59
60

1 over which blanket peat has developed. Oughtershaw is dominated by *Eriophorum* species with
2 some *Sphagnum* species present. The Green Field basin covers an area of 12.4 km² and
3 Oughtershaw 13.8 km². Altitude ranges from 364 m to 602 m above sea level in the Green Field
4 basin, and from 364 m to 668 m above sea level in the Oughtershaw basin. At the village of
5 Beckermonds, the two streams converge forming the headwaters of the River Wharfe. The
6 annual average precipitation, 1981-2008 inclusive, was 1817 mm ranging from 1383 mm in 1996
7 to 2457 mm in 2008.

8
9 Afforestation of the Green Field basin first occurred between 1970 and 1975 by Green Field
10 Forestry Trust (*G. Hay, United Paper Mills [UPM] TilHill, personal communication*) and trees
11 currently cover an area of 10.9 km². Sitka spruce (*Picea sitchensis*) is the dominant tree species
12 covering over 64% of the plantation, with the remainder being Norwegian Spruce (*Picea abies*),
13 and a few mixed broadleaf species. The forest is currently managed by UPM TilHill with areas
14 intended to be felled and restocked on a 25 year rotation. However, adequate road access for
15 logging vehicles has yet to be installed and the commencement of deforestation remains
16 uncertain.

17 ***Field methods***

18 Three tributaries were chosen for detailed study in both study basins (Table I). All were second
19 order streams, approximately 2-3 m wide and study locations were established approximately 50
20 m upstream from the confluence with the main beck. Site M1 was an exception as the study site
21 was established 50 m upstream of a small road which crosses the stream approximately 200 m
22 upstream from Oughtershaw Beck. At each site, water column temperature was monitored using
23 a Gemini Tinytag temperature datalogger housed in a radiation shield. Water temperature was
24 recorded every 15 minutes, and dataloggers downloaded at quarterly intervals and internal clocks

1
2 1 synchronised. All sub-zero water temperature records were removed from the time-series
3
4 2 because rivers do not typically reach temperatures below 0°C unless super cooled (Mohseni and
5
6 3 Stefan, 1999) and therefore all such records most likely indicated exposure of the sensor due to
7
8 4 low/no flow. All dataloggers were cross-calibrated prior to field deployment to ensure
9
10 5 comparable output data within the certified logger error of $\pm 0.2^\circ\text{C}$.
11
12
13

14 6
15 7 Air temperature was monitored in the two study basins using a Gemini Tinytag temperature
16
17 8 datalogger recording at 15 minute intervals from day 313 (November 9th), 2007. Prior to this
18
19 9 date, air temperature was monitored adjacent to the two river gauges. However, the two
20
21 10 dataloggers were damaged during overbank floods and all data were lost. Subsequently, we
22
23 11 obtained a complete meteorological data set (30 min resolution) for April 2007 to March 2009
24
25 12 from a Davis weather monitor II automatic weather station located at 240m altitude ~20km
26
27 13 northeast ($54^\circ 22' 25''\text{N}$, $2^\circ 2' 24''\text{W}$), and established two new air temperature monitoring sites
28
29 14 (Figure 1). Daily precipitation totals were obtained from the British Atmospheric Data Centre for
30
31 15 the gauge at Beckermonds. To provide contextual information on periods of peak runoff, river
32
33 16 stage data were obtained from the Environment Agency from rated cross-sections located in
34
35 17 bedrock channels for the Oughtershaw Beck and Green Field Beck gauges. Discharge
36
37 18 measurements were made for a range of stages using an electromagnetic current meter and stage-
38
39 19 discharge rating curves were constructed. Flow records for Oughtershaw Beck were available
40
41 20 only to calendar day 2, 2009 following flood damage to the stage recorder. The stage-discharge
42
43 21 curve is uncertain for flows greater than $8 \text{ m}^3 \text{ s}^{-1}$ at Green Field Beck but these high flow events
44
45 22 occurred for <1% of the time.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 1 *Data analysis*
3

4 2 Due to flashy stream flow conditions, temperature records for Sites F2 and F3 were interrupted
5
6 3 by loggers being displaced onto the bank and then recording air temperature instead of stream
7
8 4 water temperature. These erroneous data (which amounted to only 4.8% of the total number of
9
10 5 recordings), and those recorded concurrently at all other sites, were therefore omitted from
11
12 6 subsequent statistical analyses to ensure data sets of the same length. For all remaining data,
13
14 7 Pearson's correlation coefficients were calculated as a measure of association between air-water
15
16 8 temperatures, using 15 minute interval data from day 313, 2007 to day 90, 2009. Cross-
17
18 9 correlation functions were computed to assess lags and leads in the maximum correlation
19
20 10 between air and water temperature up to ± 24 h. Mean water and air temperature statistics were
21
22 11 calculated from 15 minute values rather than daily maxima and minima methods employed
23
24 12 elsewhere (Holden, 2007). T-tests were used to compare monthly mean temperatures of the three
25
26 13 open moorland and three forested streams. All statistical tests were undertaken with SPSS 15.0
27
28 14 and results were considered significant for $P < 0.05$. Temperature-duration curves were
29
30 15 constructed for 15 min water column temperatures at all sites. The form of these curves depicts
31
32 16 the nature of thermal characteristics, as steeper gradient curves reflect higher variability (Brown
33
34 17 et al., 2005; Brown et al., 2006).

35
36
37 18
38 19 To provide a detailed insight into thermal differences between forest and moorland streams over
39
40 20 diurnal timescales using high resolution (15 min) datasets, eight 24 hour time periods were
41
42 21 selected as follows: (i) diurnal water and air temperature records were examined for the two
43
44 22 summer (June 21, 2007/2008) and two winter (December 21, 2007/2008) equinoxes to consider
45
46 23 seasonal 'extremes'; (ii) two days were examined to consider the warmest and coldest days
47
48 24 across the two-year record based on the highest and lowest mean daily air temperatures, and; (iii)
49
50
51
52
53
54
55
56
57
58
59
60

1 the day in winter months (DJF) and the day in summer months (JJA) with the highest daily
2 rainfall total were examined to consider responses to high magnitude precipitation events.

3 4 **RESULTS**

5 Precipitation events were frequent throughout the monitoring period (Figure 2a), although the
6 highest magnitude event (56 mm) was observed in December 2007. The year 2008 was the
7 wettest since monitoring began at Beckermonds in 1981. Mean daily air temperature at
8 Swaledale followed an approximate sinusoidal pattern, peaking on July 27, 2008 (Figure 2;
9 20.4°C). However, peak instantaneous air temperature reached >25°C during both summer 2007
10 and 2008 at Swaledale, and >28°C at Oughtershaw on July 25, 2008 (Table II). Comparison of
11 the shorter air temperature time-series recorded at Oughtershaw and Green Field (November
12 2007 to March 2008) revealed that Swaledale air temperature was on average warmer and had a
13 smaller range than at Oughtershaw Beck and Green Field Beck (Table II). However, Swaledale
14 records were significantly and strongly correlated with both Oughtershaw (R=0.98) and Green
15 Field (R=0.99) indicating highly similar air temperature patterns over time. Both Oughtershaw
16 Beck and Green Field Beck had similar flow regimes, with low baseflow and flashy responses to
17 precipitation events (Figure 3), characteristics which were observed while in the field at the six
18 tributaries used for water temperature monitoring. Despite similar catchment size and relief,
19 runoff from Green Field Beck was typically higher than Oughtershaw Beck during storms.

20 Water column temperatures followed approximate sinusoidal patterns similar to the air
21 temperature records (Figure 4). On average, water column temperatures for the forested sites in
22 Green Field were cooler, less variable and with lower maxima than the moorland streams
23 draining the Oughtershaw Beck basin (Table III). All six water column temperature monitoring
24 sites had highly significant ($P<0.001$) correlations with local air temperature, although

1
2 1 coefficients for the forested streams were all higher than those for the moorland streams (Table
3
4 2 IV). Water temperature lagged air temperature by between 0.75 and 2.5 hrs, although there were
5
6 3 no marked differences in lags between forested and moorland streams.
7
8

9
10 4 The pronounced annual temperature cycles at all forest and open moorland stream sites were
11
12 5 further evident from monthly mean, maximum and minimum water temperatures (Figure 5).
13
14 6 Open moorland streams had significantly higher mean temperatures for 12 of the 24 months
15
16 7 studied but no significant differences were found from November 2007 to February 2008, or
17
18 8 from July 2008 to February 2009. Maximum temperatures were significantly higher in moorland
19
20 9 streams during late spring and summer months of each annual cycle (April – September) whereas
21
22 10 there were no significant differences in monthly temperature minima. The difference in water
23
24 11 column temperature maxima between forested and open moorland streams was further evident
25
26 12 from temperature duration curves (Figure 6). Compared with the forested streams of the Green
27
28 13 Field basin, the moorland streams in the Oughtershaw basin had steeper curves with higher
29
30 14 temperature maxima. Minimum water temperatures were similar across all streams.
31
32 15
33
34
35

36 16 All three moorland sites showed highly similar temporal water column temperature patterns,
37
38 17 particularly during warm time periods (e.g. day 209, 2008; Figure 4). Temporal patterns were
39
40 18 similar to open moorland streams at the forested sites although somewhat muted (e.g. the warm
41
42 19 water temperatures of day 170, 2007, and the low water column temperatures of day 355, 2007).
43
44 20 Clear differences in diurnal temperature cycles were observed as a function of both season and
45
46 21 forest cover (Figure 7). On June 21, 2007, average water temperature was up to 2.3°C higher
47
48 22 (Table V), with markedly higher temperature maxima in moorland streams but similar minima
49
50 23 across all sites (Figure 7a). On December 21, 2007, air temperature remained below freezing for
51
52 24 the entire day, although temperature range at Green Field Beck was markedly lower than at
53
54
55
56
57
58
59
60

1
2 1 Oughtershaw Beck (Figure 7b). Two of the moorland streams (M1 and M3) had sub-zero
3
4 2 temperatures for the entire day implying exposure of the sensor due to low/no streamflow, while
5
6 3 temperatures at M2 and the three forested streams remained approximately constant throughout
7
8 4 the day. June 21, 2008 had lower average, maximum and minimum air temperature than the
9
10 5 preceding year's summer equinox, with an exceptionally large air temperature range for summer
11
12 6 due to overnight sub-zero temperatures (Table V; Figure 7c). Water temperature maxima were
13
14 7 high for moorland sites similar to June 21, 2007, although the magnitude of difference between
15
16 8 moorland and forest streams was lower in 2008. December 21, 2008 was relatively warm for
17
18 9 winter with higher air and water temperatures at all sites compared with December 21, 2007
19
20 10 (Figure 7d), and 7 mm of precipitation was recorded. Air temperature was marginally higher at
21
22 11 Green Field Beck but average, maximum and minimum water temperatures were similar across
23
24 12 all streams with only minimal variation during the day (Table V).

25
26 13
27 14 The warmest mean daily air temperatures at Oughtershaw and Green Field Beck were recorded
28
29 15 on July 25, 2008 (17.6 and 17.3°C, respectively; Table V). The diurnal air temperature cycle was
30
31 16 strongly mirrored in the open moorland streams whereas thermal dynamics were relatively muted
32
33 17 in the forest streams (Figure 8a). The coldest mean daily air temperatures were recorded on
34
35 18 December 31, 2008. Air temperatures were sub-zero for the entire day, although a strong diurnal
36
37 19 cycle was evident with temperatures at Oughtershaw Beck ranging from -13.0 to -0.9°C (Table
38
39 20 V). However, water temperature records for two of the six streams (M1, F2) indicated exposure
40
41 21 of the sensor due to low or no flow, whilst three streams (M2, M3, F3) had near freezing
42
43 22 temperatures for the entire day (Figure 8b). Only stream F1 had water temperature above
44
45 23 freezing with an average water temperature of 1.0°C and a minimum of 0.7°C (Table V).

46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 1 On December 29, 2007, a total of 56mm of precipitation was recorded at Beckermonds and mean
3
4 2 daily air temperatures were 2.7°C (Green Field) and 3.0 °C (Oughtershaw). Whilst the exact
5
6 3 timing of the rainfall event at Beckermonds is unknown, 30 min records from Swaledale
7
8 4 indicated two rainfall peaks, the first at 03:00GMT and the second at 14:00GMT. Air
9
10 5 temperature records from Green Field and Oughtershaw dipped sharply at these times mirroring
11
12 6 records from Swaledale but stream temperature records varied both among and within land use
13
14 7 types, showing several increases and decreases throughout the day (Figure 8c). Temperature
15
16 8 range was marginally higher for moorland than forested streams (Table V) but there was no clear
17
18 9 distinction between the thermal regime streams of Oughtershaw or Green Field. The largest daily
19
20 10 rainfall total recorded during summer months was 41.3mm on June 19, 2008, with 30 min
21
22 11 records from Swaledale indicating a rainfall peak at 02:00GMT. Forest stream temperature
23
24 12 varied less across the day than moorland streams (Figure 8d; Table V) but similar to the winter
25
26 13 rainfall event there were clear overlaps in the thermal regimes of the six streams throughout the
27
28 14 24 hour period.

15 16 **DISCUSSION**

17 This study has demonstrated clear differences in the thermal regime of open and afforested
18 moorland streams in the Yorkshire Dales, northern England. High resolution (15 min) sampling
19 enabled the effects of forest plantations on water temperature diurnal cycles and extremes
20 (maximum/minimum) to be explored in detail. Additionally, our study has provided insights into
21 between-stream variability of thermal regime in forested catchments, whereas in many previous
22 studies, only one stream has been used to characterise water temperature for either forest or open
23 moorland (e.g. Greene, 1950; Neal et al., 1992b; Malcolm et al., 2004; Webb and Crisp, 2006).
24 All open moorland streams typically had higher maximum water temperatures, despite similar

1 climatological (air temperature, rainfall) and hydrological (river runoff) conditions to the
2 forested study areas. In a manner similar to Malcolm et al. (2004), hydroclimatological datasets
3 were used as a context for inferring the physical processes influencing thermal differences
4 between the Oughtershaw Beck open moorland streams and the forested tributaries in Green
5 Field.

6 Our observation that water temperature in streams lacking forestry ranged widely (average range
7 for the three streams was 27.3°C), with higher temperature maxima compared to forested
8 streams, is supported by findings from the River Twyi, Wales (Weatherley and Ormerod,
9 1990b), and Girnock Burn (Malcolm et al., 2004) and Loch Grannoch (Webb and Crisp, 2006) in
10 Scotland. Similar results have been seen in studies from North America (Johnson, 2004; Moore
11 et al., 2005b) and New Zealand (Rowe and Pearce, 1994) where forest streams have been
12 compared to open canopy streams following deforestation. Open moorland streams in the UK
13 flow through environments with only short riparian dwarf shrubs and grasses, meaning channels
14 are typically openly exposed to the atmosphere. While air-water temperature relationships for
15 tributary streams in the Oughtershaw Beck basin were marginally weaker than those at Green
16 Field Beck, the open moorland streams were generally more responsive to temporal (diurnal to
17 inter-annual) meteorological dynamics than forest streams. Incoming shortwave radiation,
18 sensible heat flux and wind speed/humidity (thus latent heat fluxes) are major components of
19 moorland stream energy budgets (Hannah et al., 2004; Hannah et al., 2008) and the relatively
20 low thermal capacity of small streams such as those studied herein also contributes to the high
21 magnitude of temperature fluctuations (Webb et al., 2003). Interestingly, there were no clear
22 between-catchment differences in stream temperature response to high magnitude precipitation
23 events in contrast to other studies that have shown changes of up to $\pm 10^{\circ}\text{C}$ (Chutter, 1970; Smith
24 and Lavis, 1975; Brown and Hannah, 2007). This may be a reflection of the small stream size

1 thus, minimal direct interception of precipitation. Additionally, the dominance of saturation-
2 excess overland flow pathways in peatland systems (Holden and Burt, 2003b) means there are
3 limited rainfall-induced contributions from other water sources during storms (e.g. deep
4 groundwater) that could otherwise significantly alter the stream thermal regime. Furthermore, the
5 timing of the rainfall events on the two days examined herein could account for the minimal
6 stream temperature change (Brown and Hannah, 2007) because streams would be relatively cool
7 during the night and perhaps less likely to show a large precipitation induced temperature
8 response.

9
10 Malcolm et al. (2004) noted subtle differences in thermal regimes between five open moorland
11 stream reaches as a consequence of reach-scale factors such as aspect and width:depth ratios.
12 Results from Oughtershaw Beck supported these findings of between-stream thermal variability
13 but did not fully support the hypothesised effect of aspect as average temperature in the north-
14 facing site M2 was only 0.2°C lower (i.e. within datalogger error range) than the highest
15 recorded mean moorland temperature at south-facing site M3. However, the south-facing site M3
16 did have the highest maximum temperature of any of the six sites which could be due to more
17 direct receipt of incoming solar radiation. More replication of sites is required to properly
18 elucidate the role of aspect and to consider the interactive role of other site-specific variables
19 such as stream discharge dynamics and width:depth ratios.

20
21 Temperature records for sites M1 and M3 had the longest periods of interruption due to low/no
22 flow during winter, whereas M2 flowed year-round. The very small hydraulic conductivity of
23 blanket peats in all but the uppermost few centimetres of the peat mass (Holden and Burt, 2003a)
24 means that these soils are poor regulators of baseflow. Often small peatland streams will dry up
25 at any time of the year if there has not been rainfall, or if the upper layers of the peat remain

1 frozen. Even though there is unfrozen water held in most of the peat in winter, the very low
2 hydraulic conductivity at depth combined with a frozen upper layer means that water is held in
3 situ and free drainage to the stream is inhibited. It may well be that freezing of the peat surface
4 was more common on the exposed Oughtershaw Moss site than within the forested Green Field
5 Beck catchment. Furthermore, forestry tends to increase the hydraulic conductivity of the peat
6 through the action of tree roots, litter and forest drainage systems (Holden et al., 2004) and so
7 very low flows might be better maintained (rather than no flow) in tributaries of Green Field
8 Beck than in tributaries of Oughtershaw Beck.

9
10 In the forested tributaries of Green Field Beck, diurnal dynamics were typically dampened and
11 water temperature was on average 0.8°C cooler than the moorland tributaries of Oughtershaw
12 Beck. This difference was slightly greater than the 0.15°C recorded from the Cairngorms
13 (Hannah et al., 2008), 0.4°C reported from North Wales by Crisp (1997), 0.5°C from southern
14 Scotland by Webb & Crisp (2006) and 0.6°C for a tributary of the Upper River Severn, Wales
15 (Stott and Marks, 2000). These differences are likely due to a combination of differences in
16 climatic conditions during each of the aforementioned studies, as well as more general stream
17 hydrogeomorphological characteristics (i.e. length, aspect, width:depth, discharge: Brown and
18 Hannah, 2008). Forestry characteristics such as tree species and density (Brown, 1969) also vary
19 between study locations, and they can contribute to thermal heterogeneity within individual
20 streams (e.g. Moore et al., 2005b).

21
22 In our study, forest cover had a significant effect on maximum stream temperature, with
23 reductions of up to $\sim 7^{\circ}\text{C}$ as observed in several other studies (Kirby et al., 1991; Neal et al.,
24 1992b; Webb and Crisp, 2006). One of the key reasons for discrepancies between the maximum
25 water temperature of open canopy and forested streams is that shading leads to a significant

1
2 1 reduction in net radiation during summer (Hannah et al., 2008). As net radiation is typically the
3
4 2 major component of river energy budgets, water temperature dynamics are consequently
5
6 3 dampened. A similar magnitude of maximum stream water temperature change between forested
7
8 4 and clearfelled areas has been observed in North American studies (Johnson and Jones, 2000;
9
10 5 Moore et al., 2005b). Analysis of diurnal temperature cycles for the three forested streams herein
11
12 6 showed that site F3 was typically warmer on average compared with F1 and F2, and had with
13
14 7 higher maximum and minimum temperatures. F3 has several small treeless clearings upstream of
15
16 8 the monitoring site which allow incoming radiation to reach the stream and thus heat the water in
17
18 9 a manner not dissimilar to situations seen following forest harvesting (Swift and Messer, 1971;
19
20 10 Johnson and Jones, 2000; Moore et al., 2005b). Inter-site differences in thermal regimes could
21
22 11 also potentially cause differences in hydrological flow paths and groundwater inputs between
23
24 12 individual study basins (Moore et al., 2005a) but further hydrological research is necessary at the
25
26 13 study sites to elucidate the importance of these factors.
27
28
29
30
31

32
33 14 While forest stream temperatures are typically cooler than those of moorland streams in summer
34
35 15 due to shading effects, some previous studies have found forest stream temperatures to be higher
36
37 16 than those of open moorland streams during winter (Smith, 1980; Webb and Crisp, 2006;
38
39 17 Hannah et al., 2008). It has been suggested that during winter, the canopy (in coniferous
40
41 18 plantations) acts to 'insulate' the stream by maintaining warmer air temperatures and reducing
42
43 19 long-wave radiation losses (Webb and Zhang, 2004; Webb and Crisp, 2006). Our results partially
44
45 20 support the suggestion of warmer mean temperatures in forest streams during some winter
46
47 21 months (i.e. Figure 5a; December and January of both years), but our consideration of three
48
49 22 streams showed that these minor differences were not statistically significant. If our study had
50
51 23 focused on one site per land use as per previous studies (for example choosing F1 v M1), mean
52
53
54
55
56
57
58
59
60

1
2 1 January 2007 water temperature for example would have been reported as being 0.5°C higher in
3
4 2 the forested site. In contrast, comparison of F2 v M2 would have shown the moorland site as
5
6 3 being 0.3°C warmer at this time. While we acknowledge that three streams per land-use
7
8 4 represents only a minimal level of replication providing low statistical power, these results
9
10 5 nevertheless highlight the need for replication of future stream temperature monitoring studies to
11
12 6 avoid potential erroneous conclusions that can arise from inter-site thermal differences under the
13
14 7 same land-use.
15
16
17

18
19 8
20 9 Coniferous plantations are often seen as having negative effects on peatland soils, runoff, river
21
22 10 water quality and terrestrial biodiversity (Charman, 2002) yet their moderating effect on stream
23
24 11 temperature extremes may be considered advantageous to poikilothermic stream dwelling
25
26 12 organisms such as salmonids and some invertebrates (Hawkins et al., 1997; Malcolm et al.,
27
28 13 2004). Maximum water temperatures in these Wharfedale forested study streams reached only
29
30 14 17.2°C compared with temperatures of up to 23.8°C in open moorland streams. The afforested
31
32 15 streams in this study therefore offered thermal conditions well below the upper temperature limit
33
34 16 of 22°C for juvenile Atlantic Salmon growth, and less than the 19.5°C limit at which adult
35
36 17 Brown Trout typically stop feeding (e.g. Crisp, 1993; Elliott and Hurley, 1997). Nevertheless,
37
38 18 temperature minima in both forested and open moorland streams were close to or below lethal
39
40 19 limits for all stages of salmonid life cycles (Armstrong et al., 2003) which probably contributes
41
42 20 to the low abundance of salmonids found in these Upper Wharfedale streams (Tetley, 1998). In
43
44 21 contrast to large maximum water temperature differences between forest and open moorland
45
46 22 streams, overall mean water temperature only reached up to 1.2°C higher in the open canopy
47
48 23 Oughtershaw Beck tributaries. Nevertheless, even these seemingly minor thermal differences
49
50 24 may be large enough to induce significant change in UK upland stream ecosystems. For
51
52
53
54
55
56
57
58
59
60

1
2 1 example, a +1°C change to some Welsh headwater streams due to climate warming is considered
3
4 2 likely to reduce spring abundances of benthic macroinvertebrates, which are a key food source
5
6 3 for fish and other predators, by as much as 21% (Durance and Ormerod, 2007).
7
8

9
10 4 Afforestation has been the main cause of net moorland habitat loss over the past century (Holden
11
12 5 et al., 2007) with an estimated 9% of upland UK peatland changed (Cannell et al., 1993). As the
13
14 6 majority of this land use change has involved the planting of commercial coniferous trees
15
16 7 (Forestry Commission, 2007), our study provides an insight into some of the likely wider effects
17
18 8 of this land management practice on stream thermal regimes. Growing demands for increased
19
20 9 tree cover in the UK uplands (Gimingham, 2002; Natural England, 2009), albeit with more
21
22 10 mixed plantations as well as coniferous crops, mean that it is likely that there will be increased
23
24 11 planting in future. Efforts to model ecosystem responses to upland afforestation to inform
25
26 12 decisions about the spatial extent of planting (Nisbet and Broadmeadow, 2003) should clearly
27
28 13 consider the effects on stream thermal regimes as exemplified herein.
29
30
31
32
33

34 15 **ACKNOWLEDGEMENTS**

35
36
37 17 Joseph Holden's contribution was supported by a Philip Leverhulme Prize. Sorain Ramchunder
38
39 18 was supported by a NERC studentship (NER/S/A/2006/14151). We are grateful to George Hay
40
41 19 (UPM TilHill), John Hart (Swarthghyll Farm, Oughtershaw) and Holly Rogerson (High Green
42
43 20 Field) for their kind permission to access the streams used in this study. Emma Appleford, Finn
44
45 21 Barlow-Duncan, Rachel Berman, Rachel Cleator, Matthew Cooper, Edward Higgins and Marc
46
47 22 Holland provided field assistance. The British Atmospheric Data Centre provided daily
48
49 23 precipitation data for Beckermonds, and the Environment Agency provided discharge data for
50
51 24 Oughtershaw Beck and Green Field Beck. Air temperature records for Swaledale were obtained
52
53 25 from Swaledale Weather (www.swaledaleyorkshire.com).
54
55
56
57
58
59
60

1 **Table I.** Water temperature monitoring site characteristics

Site	Green Field Beck basin			Oughtershaw Beck basin		
	F1	F2	F3	M1	M2	M3
Stream name	-	-	Green Field Beck	Mireing Gill	Long Gill	Blea Gill
Co-ordinates	54.22°N 2.20°W	54.21°N 2.23°W	54.21°N 2.26°W	54.24°N 2.21°W	54.23°N 2.21°W	54.24°N 2.23°W
Elevation (m)	325	361	388	458	376	393
Basin Area (km ²)	0.25	0.16	0.43	0.36	0.39	0.98
Slope (m per km)	173	150	94	161	198	153
Aspect	N	N	NE	SW	N	S

1 **Table II.** Air temperature statistics for the entire study period

	Swaledale air	Swaledale air ^a	F air ^a	M air ^a	F air ^b	M air ^b
Mean	9.2	7.7	5.5	5.4	5.5	5.5
Max	25.7	25.7	22.6	28.3	22.6	28.3
Min	-6.8	-6.8	-11.4	-17.2	-11.5	-17.8
Range	32.5	32.5	34.0	45.5	34.1	46.1
St. Dev	5.4	5.3	5.1	5.8	5.1	5.8

2 ^a Statistics for calendar day 313, 2007 to calendar day 90, 2008 for direct comparison of
3 Swaledale, Oughtershaw (M) and Green Field (F) records, based on 30min data

4 ^b Statistics for calendar day 313, 2007 to calendar day 90, 2009 based on 15min data

1 **Table III.** Descriptive statistics calculated from 15 min water column temperature data

Site	F1	F2	F3	M1	M2	M3
Mean	6.9	6.9	7.0	7.7	7.8	8.1
Max	17.2	16.9	17.1	23.0	22.9	23.8
Min	0.0	0.0	0.1	0.0	0.0	0.0
Range	17.2	16.9	17.0	23.0	22.9	23.8
St. Dev	3.2	3.4	3.8	4.0	4.4	4.5

2

For Peer Review

Table IV. Air-water correlation coefficients (R), cross correlation coefficients (X-R) and lag times (hr) based on 15 min water column and air temperature records [All correlations were significant at $P < 0.001$]

Site	R	X-R	Lag (hr)
F1	0.912	0.922	1.0
F2	0.927	0.941	0.75
F3	0.906	0.929	2.5
M1	0.891	0.914	0.75
M2	0.875	0.900	1.5
M3	0.864	0.900	2.25

Table V. Descriptive statistics for water and air temperature records (based on 15 min data) for the eight selected diurnal records. [- denotes no data]

		F1	F2	F3	M1	M2	M3	F air	M air
June 21, 2007	Mean	10.4	10.5	11.6	11.0	12.1	12.7	-	-
	Maximum	11.0	11.8	12.9	14.8	16.3	16.2	-	-
	Minimum	9.8	9.9	10.5	9.1	9.7	10.5	-	-
	Range	1.2	1.9	2.4	5.7	6.6	5.7	-	-
	St. Dev.	0.4	0.5	0.8	1.6	2.0	1.7	-	-
Dec. 21, 2007	Mean	2.0	0.9	0.9	-	0.2	-	-2.8	-4.2
	Maximum	2.6	1.6	1.2	-	0.5	-	-0.7	-1.1
	Minimum	1.0	0.0	0.2	-	0.0	-	-6.8	-11.2
	Range	1.6	1.6	1.1	-	0.5	-	6.1	10.1
	St. Dev.	0.3	0.5	0.3	-	0.1	-	1.6	2.7
June 21, 2008	Mean	8.0	8.0	8.7	8.2	8.9	9.6	7.9	7.7
	Maximum	9.2	9.1	9.4	9.3	10.3	10.5	13.7	14.1
	Minimum	6.2	7.3	8.1	6.8	7.2	8.6	1.9	-2.3
	Range	3.0	1.8	1.4	2.4	3.2	1.9	11.8	16.5
	St. Dev.	0.9	0.5	0.3	0.8	1.0	0.6	2.9	4.3
Dec. 21, 2008	Mean	6.1	6.2	5.9	5.9	6.1	6.4	7.8	7.5
	Maximum	6.3	6.4	6.3	6.3	6.4	6.9	8.7	8.4
	Minimum	5.8	5.8	5.4	5.6	5.7	5.9	6.8	7.0
	Range	0.6	0.6	0.9	0.7	0.8	1.0	1.9	1.5
	St. Dev.	0.2	0.1	0.3	0.2	0.2	0.3	0.5	0.4
Jul. 25, 2008	Mean	11.6	12.7	13.8	15.1	16.6	16.6	17.3	17.6
	Maximum	12.3	13.8	15.6	19.0	21.2	20.8	22.6	25.6
	Minimum	10.9	11.5	12.3	11.6	12.4	13.0	12.3	8.4
	Range	1.4	2.3	3.4	7.4	8.7	7.8	10.4	17.3
	St. Dev.	0.5	0.8	1.2	2.6	3.0	2.8	3.6	5.0
Dec. 31, 2008	Mean	1.0	-	0.2	-	0.0	0.1	-6.5	-8.1
	Maximum	1.8	-	0.4	-	0.1	0.1	-4.0	-0.9
	Minimum	0.7	-	0.1	-	0.0	0.1	-8.6	-13.0
	Range	1.0	-	0.2	-	0.1	0.1	4.7	12.1
	St. Dev.	0.3	-	0.1	-	0.0	0.0	1.2	3.5
Dec. 29, 2007	Mean	5.0	4.5	4.5	4.2	4.6	4.5	3.0	2.7
	Maximum	5.8	5.5	5.7	5.6	5.9	5.8	4.3	4.3
	Minimum	4.5	3.9	4.1	3.6	4.1	3.7	0.5	0.1
	Range	1.3	1.7	1.6	2.0	1.8	2.1	3.8	4.2
	St. Dev.	0.3	0.3	0.4	0.5	0.5	0.5	0.9	1.1
Jun. 19, 2008	Mean	9.0	8.9	9.6	8.8	9.6	10.3	10.1	10.2
	Maximum	9.5	9.1	10.3	10.1	10.8	12.0	12.4	13.2
	Minimum	8.6	8.6	9.0	8.1	8.8	9.0	8.5	6.4
	Range	0.8	0.6	1.3	1.9	2.1	3.1	4.0	6.7
	St. Dev.	0.3	0.2	0.4	0.6	0.6	0.9	1.1	1.4

3

1
2 **1 Figure captions**
3

4
5 **2 Figure 1.** Map of the Oughtershaw Beck and Green Field Beck study basins showing locations
6
7
8 **3** of water column temperature, air temperature, river gauge and precipitation gauge
9

10
11 **4 Figure 2.** (a) Total daily rainfall measured at Beckermonds gauge at the confluence of
12
13 Oughtershaw Beck and Green Field Beck, and; mean daily air temperatures based on 30 min data
14
15
16 **6** measured at (b) Swaledale, (c) Green Field Beck, and (d) Oughtershaw Beck
17

18
19 **7 Figure 3.** Mean daily discharge for Oughtershaw Beck and Green Field Beck.
20

21
22 **8 Figure 4.** Mean daily stream water temperatures at (a) F1, (b) F2, (c) F3, (d) M1, (e) M2, and (f)
23
24
25 **9** M3
26

27
28 **10 Figure 5.** Monthly (a) mean, (b) maximum and (c) minimum water temperatures for forest and
29
30 moorland streams [error bars = 1 standard deviation of the mean and asterisks denote significant
31
32
33 **12** differences at $P < 0.05$]
34

35
36 **13 Figure 6.** Water temperature duration curves for forest and moorland streams
37

38
39 **14 Figure 7.** Diurnal temperature records for equinoxes: (a) June 21, 2007; (b) December 21, 2007;
40
41
42 **15** (c) June 21, 2008, and; (d) December 21, 2008. Note different y-axis scales for June and
43
44 **16** December figures
45

46
47 **17 Figure 8.** Diurnal temperature records for (a) July 25, 2008 [highest mean daily air temperature];
48
49
50 **18** (b) December 31, 2008 [lowest mean daily air temperature]; (c) December 29, 2007 [highest
51
52
53 **19** winter daily precipitation total = 56mm], and; (d) June 19, 2008 [highest summer daily
54
55 **20** precipitation total = 41.3mm]. Note different y-axis scales
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Fig 1.

For Peer Review




Fig. 2.





Fig. 3.




1

2

3

Fig. 4.



1

2 **Fig. 5.**

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60






Fig. 6.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 **Fig. 7.**



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 **Fig. 8.**




References

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
- Acuña, V., Wolf, A. and Tockner, K., 2008. Temperature dependence of stream benthic respiration in an Alpine river network with relevance to global warming. *Freshwater Biology*, 53: 2076-2088.
- Armstrong, J.D., Kemp, P.S., Kennedy, G.J.A., Ladle, M. and Milner, N.J., 2003. Habitat requirements of Atlantic salmon and brown trout in rivers and streams. *Fisheries Research*, 62: 143-170.
- Berner, E.K.B. and Berner, R.A., 1996. *Global Environment. Water, Air and Geochemical Cycles*. Prentice Hall.
- Brown, G.W., 1969. Predicting temperatures of small streams. *Water Resources Research*, 5: 68-75.
- Brown, L.E. and Hannah, D.M., 2007. Alpine stream temperature response to storm events. *Journal of Hydrometeorology*, 8: 952-967.
- Brown, L.E. and Hannah, D.M., 2008. Spatial heterogeneity of water temperature across an alpine river basin. *Hydrological Processes*, 22: 954-967.
- Brown, L.E., Hannah, D.M. and Milner, A.M., 2005. Spatial and temporal water column and streambed temperature dynamics within an alpine catchment: Implications for benthic communities. *Hydrological Processes*, 19: 1585-1610.
- Brown, L.E., Hannah, D.M. and Milner, A.M., 2006. Hydroclimatological influences upon water column and streambed temperature dynamics in an alpine river system. *Journal of Hydrology*, 325: 1-20.
- Caissie, D., 2006. The thermal regime of rivers: a review. *Freshwater Biology*, 51: 1389-1406.
- Cannell, M.G.R., Dewar, R.C. and Pyatt, D.G., 1993. Conifer plantations on drained peatlands in Britain: a net gain or loss of carbon? *Forestry*, 66: 353-369.
- Charman, D., 2002. *Peatlands and Environmental Change*. John Wiley, Chichester.
- Chutter, F.M., 1970. Hydrobiological studies in the catchment of Vaal Dam, South Africa. Part 1. River zonation and the benthic fauna. *International Review of Hydrobiology*, 55: 445-494.
- Crisp, D.T., 1993. The environmental requirements of salmon and trout in fresh water. *Freshwater Forum*, 3: 176-202.
- Crisp, D.T., 1997. Water temperature of Plynlimon streams. *Hydrology and Earth System Sciences*, 1: 535-540.
- Durance, I. and Ormerod, S.J., 2007. Climate change effects on upland stream macroinvertebrates over a 25-year period. *Global Change Biology*, 13(5): 942-957.
- Elliott, J.M. and Hurley, M.A., 1997. A functional model for maximum growth of Atlantic salmon parr, *Salmo salar*, from two populations in northwest England. *Functional Ecology*, 11: 592-603.
- Forestry Commission, 2007. *Forestry facts and figures 2007*. Forestry Commission, Edinburgh, 14 pp.
- Gimingham, C.H., 2002. Foreword. A vision for the uplands. In: T.P. Burt, D.B.A. Thompson and J. Warburton (Editors), *The British Uplands: Dynamics of Change*. JNCC Report 319. Joint Nature Conservancy Committee, Durham, pp. 9-11.
- Greene, G., 1950. Land Use and Trout Streams. *Journal of Soil and Water Conservations*, 5: 125-126.
- Hannah, D.M., Malcolm, I.A., Soulsby, C. and Youngson, A.F., 2004. Heat exchanges and temperatures within a salmon spawning stream in the Cairngorms, Scotland: Seasonal and sub-seasonal dynamics. *River Research and Applications*, 20: 635-652.
- Hannah, D.M., Malcolm, I.A., Soulsby, C. and Youngson, A.F., 2008. A comparison of forest and moorland stream microclimate, heat exchanges and thermal dynamics. *Hydrological Processes*, 22: 919 - 940.
- Hawkins, C.P., Hogue, J.N., Decker, L.M. and Feminella, J.W., 1997. Channel morphology, water temperature, and assemblage structure of stream insects. *Journal of the North American Benthological Society*, 16(4): 728-749.
- Holden, J., 2007. A plea for more careful presentation of near-surface temperature data in geomorphology. *Earth Surface Processes and Landforms*, 32: 1433-1436.
- Holden, J. and Burt, T.P., 2003a. Hydraulic conductivity in upland blanket peat; measurement and variability. *Hydrological Processes*, 17: 1227-1237.
- Holden, J. and Burt, T.P., 2003b. Runoff production in blanket peat covered catchments. *Water Resources Research*, 39: 1191, DOI: 10.1029/2003WR002067.

- 1
2 1 Holden, J., Chapman, P.J. and Labadz, J.C., 2004. Artificial drainage of peatlands: hydrological and
3 2 hydrochemical process and wetland restoration. *Progress in Physical Geography*, 28: 95-123.
- 4 3 Holden, J., Shotbolt, L., Bonn, A., Burt, T.P., Chapman, P.J., Dougill, A.J., Fraser, E.D.G., Hubacek, K.,
5 4 Irvine, B., Kirkby, M.J., Reed, M.S., Prell, C., Stagl, S., Stringer, L.C., Turner, A. and Worrall, F.,
6 5 2007. Environmental change in moorland landscapes. *Earth-Science Reviews*, 82: 75-100.
- 7 6 Johnson, S.L., 2004. Factors influencing stream temperatures in small streams: substrate effects and a
8 7 shading experiment. *Canadian Journal of Fisheries and Aquatic Sciences*, 61: 913-923.
- 9 8 Johnson, S.L. and Jones, J.A., 2000. Stream temperature response to forest harvest and debris flows in
10 9 western Cascades, Oregon. *Canadian Journal of Fisheries and Aquatic Sciences*, 57 (Supp. 2): 30-39.
- 11 10 Kirby, C., Newson, M.D. and Gilman, K., 1991. Plynlimon research: the first two decades. Institute of
12 11 Hydrology Report, 109, 188 pp.
- 13 12 Lane, S.N., Reid, S.C., Tayefia, V., Yu, D. and Hardy, R.J., 2008. Reconceptualising coarse sediment
14 13 delivery problems in rivers as catchment-scale and diffuse Geomorphology, 98: 227-249.
- 15 14 Malcolm, I.A., Hannah, D.M., Donaghy, M.J., Soulsby, C. and Youngson, A.F., 2004. The influence of
16 15 riparian woodland on the spatial and temporal variability of stream water temperatures in an upland
17 16 salmon stream. *Hydrology and Earth System Sciences*, 8: 449-459.
- 18 17 Mohseni, O. and Stefan, H.G., 1999. Stream temperature/air temperature relationship: a physical
19 18 interpretation. *Journal of Hydrology*, 218: 128-141.
- 20 19 Moore, R.D., Spittlehouse, D.L. and Story, A., 2005a. Riparian microclimate and stream temperature
21 20 response to forest harvesting: a review. *Journal of the American Water Resources Association*, 41: 813
22 21 - 834.
- 23 22 Moore, R.D., Sutherland, P., Gomi, T. and Dhakal, A., 2005b. Thermal regime of a headwater stream
24 23 within a clear-cut, coastal British Columbia, Canada. *Hydrological Processes*, 19: 2591-2608.
- 25 24 Natural England, 2009. *Vital Uplands: A 2060 vision for England's upland environment*. Crown
26 25 Copyright, 8 pp.
- 27 26 Neal, C., Reynolds, B., Smith, C.J., Hill, S., Neal, M., Conway, T., Ryland, G.P., Jeffery, H.A., Robson,
28 27 A.J. and Fisher, R., 1992a. The impact of conifer harvesting on stream water pH, alkalinity and
29 28 aluminium concentrations for the British uplands: an example for an acidic and acid-sensitive
30 29 catchment in mid-Wales. *Science of the Total Environment*, 126: 75-87.
- 31 30 Neal, C., Smith, C.J. and Hill, S., 1992b. Forestry impact on upland water quality. Institute of Hydrology
32 31 Report, 119.
- 33 32 Nisbet, T.R. and Broadmeadow, S., 2003. *Opportunity Mapping for Trees and Floods*. Forest Research,
34 33 Farnham.
- 35 34 Pluhowski, E.J., 1972. Unusual temperature variations in two small streams in Northern Virginia.
36 35 *Geological Survey Research; US Geological Survey Professional Paper 800-B: B255-B258*.
- 37 36 Poole, G.C. and Berman, C.H., 2001. An ecological perspective on in-stream temperature: Natural heat
38 37 dynamics and mechanisms of human-caused thermal degradation. *Environmental Management*, 27(6):
39 38 787-802.
- 40 39 Reynolds, B., Neal, C., Hornung, M., Hughes, S. and Stevens, P.A., 1988. Impacts of afforestation on the
41 40 soil solution of a stagnopodzol in mid-Wales. *Water, Air, Soil Pollution*, 38: 55-70.
- 42 41 Reynolds, B., Stevens, P.A., Adamson, J.K., Hughes, S. and Roberts, J.D., 1992. Effects of clear felling
43 42 on stream and soil water aluminium chemistry in three UK forests. *Environmental Pollution*, 77: 157-
44 43 165.
- 45 44 Rowe, L.K. and Pearce, C.H., 1994. Hydrology and related changes after harvesting native forest
46 45 catchments and establishing *Pinus radiata* plantations. Part 3. Stream temperatures. *Hydrological*
47 46 *Processes*, 8: 299-310.
- 48 47 Sinokrot, B.A. and Stefan, H.G., 1993. Stream temperature dynamics - Measurements and modelling.
49 48 *Water Resources Research*, 29(7): 2299-2312.
- 50 49 Smith, B.D., 1980. The effects of afforestation on the trout of a small stream in southern Scotland.
51 50 *Fisheries Management and Ecology*, 11 39-58.
- 52 51 Smith, K. and Lavis, M.E., 1975. Environmental influences on the temperature of a small upland stream.
53 52 *Oikos*, 26: 228-236.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 1 Stott, T. and Marks, S., 2000. Effects of plantation forest clearfelling on stream temperatures in the
2 Plynlimon experimental catchments, mid-Wales. *Hydrology and Earth System Sciences*, B(1): 95-104.
3 Swift, L.W. and Messer, J.B., 1971. Forest cuttings raise temperatures of small streams in the southern
4 Appalachians. *Journal of Soil and Water Conservation*, 26: 111-116.
5 Tallis, J.H., 1998. Growth and degradation of British and Irish blanket mires. *Environmental Reviews*, 6:
6 81 - 122.
7 Tetley, L., 1998. *Yorkshire Dales Anglers' Guide*. Cicerone Press Limited, Milnthorpe, Cumbria, 223 pp.
8 Vannote, R.L. and Sweeney, B.W., 1980. Geographic analysis of thermal equilibria: a conceptual model
9 for evaluating the effect of natural and modified thermal regimes on aquatic insect communities. *The*
10 *American Naturalist*, 115: 667-695.
11 Wallage, Z.E., Holden, J. and McDonald, A.T., 2006. Drain blocking: An effective treatment for reducing
12 dissolved organic carbon loss and water discolouration in a drained peatland. *Science of the Total*
13 *Environment*, 367(2-3): 811-821.
14 Weatherley, N.S. and Ormerod, S.J., 1990a. The constancy of invertebrate assemblages in soft-water
15 streams - Implications for the prediction and detection of environmental-change. *Journal of Applied*
16 *Ecology*, 27(3): 952-964.
17 Weatherley, N.S. and Ormerod, S.J., 1990b. Forests and the temperature of upland streams in Wales: a
18 modelling exploration of the biological effects. *Freshwater Biology*, 24: 109-122.
19 Webb, B.W., Clack, P.D. and Walling, D.E., 2003. Water-air temperature relationships in a Devon river
20 system and the role of flow. *Hydrological Processes*, 17(15): 3069-3084.
21 Webb, B.W. and Crisp, D.T., 2006. Afforestation and stream temperature in a temperate maritime
22 environment. *Hydrological Processes*, 20: 51-66.
23 Webb, B.W., Hannah, D.M., Moore, R.D., Brown, L.E. and Nobilis, F., 2008. Recent advances in stream
24 and river temperature research. *Hydrological Processes*, 22: 902-918.
25 Webb, B.W. and Zhang, Y., 1999. Water temperatures and heat budgets in Dorset chalk water courses.
26 *Hydrological Processes*, 13(3): 309-321.
27 Webb, B.W. and Zhang, Y., 2004. Inter-annual variability in the non-advective heat energy budget of
28 Devon streams and rivers. *Hydrological Processes*(11): 2117-2146.
29
30
31

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



1