



UNIVERSITY OF LEEDS

This is a repository copy of *Salt stress causes peroxisome proliferation, but inducing peroxisome proliferation does not improve NaCl tolerance in Arabidopsis thaliana*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/77548/>

Version: Published Version

---

**Article:**

Mitsuya, S, El-Shami, M, Sparkes, IA et al. (4 more authors) (2010) Salt stress causes peroxisome proliferation, but inducing peroxisome proliferation does not improve NaCl tolerance in Arabidopsis thaliana. PLoS One, 5 (2). e9408. 1 - 10. ISSN 1932-6203

<https://doi.org/10.1371/journal.pone.0009408>

---

**Reuse**

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# Salt Stress Causes Peroxisome Proliferation, but Inducing Peroxisome Proliferation Does Not Improve NaCl Tolerance in *Arabidopsis thaliana*

Shiro Mitsuya<sup>‡a</sup>, Mahmoud El-Shami<sup>‡b</sup>, Imogen A. Sparkes<sup>‡c</sup>, Wayne L. Charlton, Carine De Marcos Lousa, Barbara Johnson, Alison Baker\*

Centre for Plant Sciences, University of Leeds, Leeds, United Kingdom

## Abstract

The PEX11 family of peroxisome membrane proteins have been shown to be involved in regulation of peroxisome size and number in plant, animals, and yeast cells. We and others have previously suggested that peroxisome proliferation as a result of abiotic stress may be important in plant stress responses, and recently it was reported that several rice PEX11 genes were up regulated in response to abiotic stress. We sought to test the hypothesis that promoting peroxisome proliferation in *Arabidopsis thaliana* by over expression of one PEX11 family member, PEX11e, would give increased resistance to salt stress. We could demonstrate up regulation of PEX11e by salt stress and increased peroxisome number by both PEX11e over expression and salt stress, however our experiments failed to find a correlation between PEX11e over expression and increased peroxisome metabolic activity or resistance to salt stress. This suggests that although peroxisome proliferation may be a consequence of salt stress, it does not affect the ability of *Arabidopsis* plants to tolerate saline conditions.

**Citation:** Mitsuya S, El-Shami M, Sparkes IA, Charlton WL, De Marcos Lousa C, et al. (2010) Salt Stress Causes Peroxisome Proliferation, but Inducing Peroxisome Proliferation Does Not Improve NaCl Tolerance in *Arabidopsis thaliana*. PLoS ONE 5(2): e9408. doi:10.1371/journal.pone.0009408

**Editor:** Michael V. Mickelbart, Purdue University, United States of America

**Received:** October 13, 2009; **Accepted:** February 3, 2010; **Published:** February 24, 2010

**Copyright:** © 2010 Mitsuya et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** The authors acknowledge financial support from the Biotechnology and Biological Sciences Research Council (United Kingdom) (www.bbsrc.ac.uk), grant number 24/P13265, and Japan partnering award JPA1811. SM acknowledges the support of the grant 'KAKENHI (20780011)' from the Japan Society for Promotion of Science (http://www.jsps.go.jp/). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: a.baker@leeds.ac.uk

‡a Current address: Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya, Japan

‡b Current address: Faculty of Agriculture, Department of Agronomy, Al-Azhar University, Cairo, Egypt

‡c Current address: School of Life Sciences, Oxford Brookes University, Oxford, United Kingdom

## Introduction

Peroxisomes are eukaryotic organelles which are highly dynamic and pleiomorphic. They possess a single boundary membrane and lack DNA so their protein complement is acquired by post-translational uptake of cytosolically synthesized proteins which possess a suitable peroxisome targeting signal. A few peroxisomal proteins lack a targeting signal of their own but are imported via association with another protein that does contain a targeting signal, so called 'piggy-back' import [1]. In mammalian and yeast cells, peroxisomes have been shown to be capable of arising *de novo* from the endoplasmic reticulum [2,3,4], and, although formal proof is lacking, this seems likely to be the case in plant cells as well [5].

Despite the ability to be synthesised *de novo*, peroxisomes possess division machinery [6], which is partially shared with mitochondria [7]. The capacity to divide and be segregated to daughter cells may reflect the requirement for peroxisomes to increase their number and volume rapidly in response to particular developmental and environmental signals. For example peroxisome proliferators such as clofibrate cause peroxisome proliferation in rodents [8]. Yeast peroxisomes are induced to proliferate when the organism is switched to growth on a carbon source that requires peroxisomal metabolism, such as fatty acids [9] or methanol [10].

Plant peroxisomes proliferate during post germinative growth, when peroxisomal  $\beta$ -oxidation plays an essential role in the mobilization of storage lipid [11], and also upon transition of dark grown seedlings into the light [12].

The molecular machinery involved in peroxisome proliferation and division has begun to be characterized (reviewed in [13,14]). The first protein identified as a regulator of peroxisome size and number was PEX11 from *Saccharomyces cerevisiae*. Disruption of the gene encoding this protein resulted in the formation of a few giant peroxisomes per cell and over expression resulted in many small peroxisomes [15,16]. Subsequently other genes have been discovered which also act to regulate peroxisome size and abundance; Pex25p, Pex27p [17,18], and Pex31p and Pex32p [19,20]. Mammals and plants do not have obvious homologues of these latter genes but have expanded members of the PEX11 family. Phylogenetic analysis has suggested that this expansion has taken place since the divergence of the eukaryotic lineage [21]. There are 3 mammalian isoforms: PEX11 $\alpha$ ,  $\beta$ , and  $\gamma$  which differ in expression pattern and the phenotypic consequences of gene knockout [22,23]. The  $\alpha$ -isoform stimulates peroxisome proliferation in response to metabolic cues, whereas  $\beta$  appears to play a more constitutive role [24].

In *Arabidopsis* there are 5 isoforms, designated PEX11a-e [25], which fall into 3 subfamilies on the basis of sequence similarity and

show different expression profiles [21]. All 5 isoforms were targeted to peroxisomes and resulted in peroxisome elongation and/or proliferation [21,25], although some differences were observed between cultured cells [25] and the transgenic plants [21]. Additionally Orth et al were able to show that PEX11e could partially complement the *S. cerevisiae*  $\Delta pex11$  mutant, thereby demonstrating (partial) conservation of function [21].

In mammals PEX11 was shown to interact with DLP1 (dynamin like protein) [26] and more recently a complex containing DLP1, Pex11 $\beta$  and FIS1 was characterized from mammalian cells [27]. Recent studies also point to the involvement of dynamin related proteins playing a role in peroxisome division in plants. The *apm1* mutant of Arabidopsis [28] has a lesion in dynamin-related protein DRP3A which results in elongated peroxisomes and mitochondria, suggestive of a failure to complete division. The *fis1a* and *fis1b* mutants, homologues of mammalian FIS1 which acts to tether DRPs to peroxisomal and mitochondrial membranes, also show a similar cellular phenotype [29,30]. Arabidopsis PEX11 isoforms d and e were shown to interact with Fis1b using Bimolecular Fluorescence Complementation (BiFC), and to be required for targeting of Fis1b to peroxisomes [31]. Thus PEX11 appears to play a role in recruitment of the molecular machinery for peroxisome division and it is tempting to speculate that the more complex PEX11 families of multicellular organisms reflects an increased requirement to modulate and integrate peroxisomal activities in response to endogenous and environmental stimuli. The demonstration that one specific Arabidopsis PEX11 isoform Pex11b mediates light-induced peroxisome proliferation is consistent with this hypothesis [12].

Peroxisomes compartmentalize many metabolic pathways, but a common function is metabolism of reactive oxygen species. Many peroxisomal activities generate superoxide or hydrogen peroxide [32]. The presence of catalase and other anti-oxidative enzymes such as superoxide dismutase, ascorbate peroxidase, dihydro- and monohydro- ascorbate reductase, glutathione peroxidase and glutathione reductase act to scavenge ROS produced within peroxisomes. As well as causing damage to many cellular macromolecules ROS also play a role in a wide range of signal transduction processes that include developmental, hormone and stress related responses. Thus maintaining an appropriate temporal and spatial balance of ROS underpins many plant responses. Whilst the precise contribution of peroxisomal ROS metabolism to stress responses is difficult to quantify, given the presence of multiple isoenzymes in different compartments and separate pools of anti-oxidants such as glutathione, ascorbate and  $\alpha$ -tocopherol, a number of studies have provided evidence to link peroxisome proliferation with stress conditions. High light intensity [33] ozone, [34,35] metal stress [36,37], salt stress [38], and treatment with the herbicide isoproturon [39] or with the hypolipidemic drug clofibrate [40] have all been reported to increase peroxisome number or modify activity of peroxisome enzymes involved in ROS metabolism (reviewed in [41]).

Many stresses including salt and drought result in abscisic acid (ABA) accumulation, which results in various protective and adaptive responses such as stomatal closure to limit water loss and changes in transcriptional and post transcriptional gene regulation [42]. Microarray studies in Arabidopsis demonstrated a large overlap between salt, drought and ABA induced genes [43]. ABA signaling operates through a variety of second messengers including phospholipid derived molecules, cyclic ADPrbose and hydrogen peroxide [44] and with extensive cross-talk with other phytohormone pathways such as ethylene and jasmonic acid (JA) [45]. Hydrogen peroxide transiently inactivates ABI1 and ABI2,

protein phosphatases which act as negative regulators of ABA signaling [44]. Among the targets of ABA are many transcription factors such as ABI3 [46] and ABI4 [47], which control specific physiological and developmental responses.

The observation that stress increases peroxisome number and activity begs the question is peroxisome proliferation simply a consequence of stress conditions or does peroxisome proliferation result in an accompanying increase in peroxisomal activities and can this provide a protective effect against oxidative stress? In this study we have investigated whether manipulating peroxisome number via over expression of the PEX11e isoform can improve tolerance of Arabidopsis plants to NaCl.

## Results and Discussion


### PEX11e Is an Integral Peroxisomal Membrane Protein and Forms a High Molecular Weight Complex

Pex11e was selected for this study as it is constitutively expressed at a reasonably high level in most Arabidopsis tissues [www.geneinvestigator.ethz.ch](http://www.geneinvestigator.ethz.ch) [48]. AtPEX11e was fused at the C terminus of eYFP and expressed by agroinfiltration in the leaf epidermal cells of tobacco plants which stably express the peroxisome targeted reporter CFP-SKL [49]. Figure 1 panel A shows eYFP fluorescence in ring like-structures. Panel B shows the image from the CFP channel which shows that the same structures contain CFP-SKL and the merged image panel C shows that the fluorescence from the eYFP-PEX11 fusion protein surrounds the matrix marker CFP-SKL as would be expected for a peroxisome membrane protein [49]. Occasionally ring type structures can also be seen with matrix markers (an example is visible in panel B). As peroxisomes sometimes contain matrix inclusions (such as catalase crystals) we hypothesise that such observations represent exclusion of the fluorescent protein from such inclusions. Our results corroborate the findings that N terminally tagged forms of PEX11e target to peroxisomes in transiently transformed BY2 cells [25] and in stably transformed Arabidopsis plants [21] respectively.


Immunofluorescence microscopy was carried out on Arabidopsis suspension culture cells using an antibody raised to a common sequence in PEX11c, d and e. These suspension cultured cells contain very small peroxisomes [49] that contain isocitrate lyase [50]. In Figure 1 panel D punctate structures are detected by the anti-PEX11 antibody that co-localise with the signal obtained with anti-isocitrate lyase (ICL) antibodies (panel E). The merged image panel F shows that these structures are one and the same. As ICL is a peroxisome marker we conclude that native PEX11 is a peroxisomal protein in Arabidopsis as previously reported [21].

Membranes from Arabidopsis cell culture were treated with alkaline sodium carbonate to remove soluble and peripheral membrane proteins and subjected to SDS PAGE and immunoblotting with anti-PEX11c/d/e antibody. PEX11 was recovered quantitatively in the pellet fraction (Fig. 2A) supporting the notion that it is an integral membrane protein in Arabidopsis [21] [25]. It has been reported that PEX11 in *S. cerevisiae* is peripheral and forms a redox sensitive dimer [51]. We could find no evidence of Arabidopsis PEX11 forming dimers in the absence of DTT (Figure 2A right lanes –DTT).

To explore further whether PEX11 is oligomeric, a peroxisome-enriched organelle fraction from Arabidopsis suspension culture cells was incubated with different concentrations of the detergent dodecyl maltoside, or with 1% Triton X-100 and the soluble fraction (100,000 $\times$ g supernatant) separated on a native 3–8% polyacrylamide gel. Western blotting of the gel with anti-PEX11c/d/e antibodies revealed that PEX11 forms a complex of



**Figure 1. Localisation of PEX11e to peroxisomes.** eYFP-PEX11e (panel A, magenta) was transiently co-expressed with the peroxisomal marker CFP-SKL (panel B, green) in tobacco epidermal cells. The two fluorescent markers co-localise in small motile structures 1–2 micrometers in size, typical of peroxisomes (panel C). Arabidopsis suspension cells were immuno-labelled with anti-PEX11 antibody (D, magenta) and anti isocitrate lyase antibody (E, green) a peroxisomal (glyoxysomal) marker protein. The merged figure (F) shows PEX11 in punctate structures containing the glyoxysomal enzyme ICL. Scale bar 2  $\mu$ m.  
doi:10.1371/journal.pone.0009408.g001




**Figure 2. PEX11 is an integral membrane protein that forms a high molecular weight complex.** A. Duplicate 100 microgramme mixed light membrane fractions (10,000 g pellet) from Arabidopsis suspension cells was subjected to extraction by 0.1 M sodium carbonate. Soluble and insoluble fractions were separated by SDS-PAGE (with or without DTT) and immunoblotted with PEX11 affinity purified antibody. B. 100 microgrammes of protein from a peroxisome enriched organelle fraction was solubilised in the indicated concentration of detergent and the solubilised proteins were separated on a 3–8% native gel and immunoblotted with anti-PEX11 affinity purified antiserum.  
doi:10.1371/journal.pone.0009408.g002

approximately 800 kDa in molecular weight in dodecyl maltoside. This complex is unstable in 1% Triton X-100 (Figure 2B). Since the antibodies recognise more than one PEX11 isoform this could be a homo or hetero oligomeric complex as PEX11c, d and e are all expressed in suspension cultured cells [25] (and [www.genevestigator.ethz.ch](http://www.genevestigator.ethz.ch) [48]). Even allowing for the presence of detergent molecules, a complex of 800 kDa, must contain multiple copies of PEX11 possibly in complex with other proteins since the predicted monomeric molecular weight of the various PEX11 isoforms is 25.5–25.9 kDa, and the protein(s) run close to their predicted molecular mass on SDS PAGE (Fig. 2A). PEX11e has been shown to homo-dimerise using an in planta split ubiquitin system [52]. A complex containing FIS1, DLP1 and PEX11 $\beta$  could be detected biochemically in mammalian cells [27]. Recently it was shown that the various Arabidopsis PEX11 isoforms could interact with one another and also with FIS1b which acts as a tethering factor for the mitochondrial and peroxisome associated Dynamin related protein DLP1 in mammals [31]. Thus other proteins involved in peroxisome division are also candidate components of the complex observed in Fig. 2B.

### PEX11e Transcripts and Peroxisome Number Are Increased by Salt Stress

Four members of the rice *PEX11* family have recently been reported to be up regulated by treatment with 200 mM NaCl [53], and we have previously reported the up regulation by salt stress of transcripts for the Arabidopsis peroxisome biogenesis factors *PEX10* and *PEX1* [54] both of which are required for import of peroxisome matrix proteins [55]. Using an identical approach we investigated the changes in steady state levels of *PEX11e* mRNA in response to salt stress in wild type and mutants impaired in abscissic acid and Jasmonate signalling. *PEX11e* transcript was induced two fold after 4 h treatment with 200 mM or 400 mM NaCl (Figure S1). Pools of 13 day old seedlings were treated with 400 mM NaCl or distilled water for 4 h, RNA extracted and transcripts determined by quantitative PCR. All transcripts were normalised relative to Actin2 and the level of expression in the untreated control ecotype set to 100%. In both Landsberg erecta and Columbia ecotypes salt treatment resulted in an approximately 2 fold increase in *PEX11e* transcript (Fig. 3). The increase in expression in response to salt was completely blocked in the *abi1-1* mutant and partially blocked in the *abi3-1* mutant (which is a leaky allele [56]) consistent with a role for abscissic acid in switching on *PEX11e* expression in response to salt. Salt induction still occurred but to a more limited extent in the *abi2-1* and *abi4-103* mutant backgrounds (Fig. 3A). *abi1-1* and *abi2-1* are dominant mutants in type 2C protein phosphatases that are defective in multiple ABA responses [57]. ABI3 and ABI4 are transcription factors that regulate ABA responsive genes. In the *jar1-1* mutant background [58] *PEX11e* is up regulated in the absence of salt and its expression reduced in the presence of salt (Fig. 3), suggesting that salt induction of *PEX11e* may require jasmonate signalling. JAR1 conjugates jasmonic acid to isoleucine to form JA-Ile [59]. JA-Ile promotes association of JAZ transcriptional repressor proteins with SCF<sup>COI1</sup>, the F box subunit of a multi component ubiquitin E3 ligase, leading to their degradation and de-repression of JA-responsive genes [60]. Interestingly, 306 nucleotides 5' to the ATG of the *PEX11e* gene, in the intron within the 5' UTR, there is a perfect match to the sequence CGTCA which mediates methyl jasmonate responsiveness of the Barley *LOX1* gene [61] and within the intron (+166) and promoter (+633) is a match to the binding site for MYC2 (ACGTG/CACGT) [62] the key JA transcriptional activator. MYC2 is itself upregulated by both JA and ABA, although ABA induction is independent of ABI1 and most likely



**Figure 3. Transcriptional response of PEX11e to salt treatment in wild type and signalling mutant backgrounds.** A. Pools of 13 day old seedlings of the indicated genotypes were treated with either 400 mM NaCl solution (black bars) or distilled water (white bars) for 4 h. RNA was extracted and the relative levels of expression of *PEX11e* determined by quantitative real time PCR. Untreated control values were set to 1. Data shown are the means of three independent RNA extractions performed on different sets of material grown at the same time with the PCR reactions run in duplicate. B. Model for control of *PEX11e* expression by ABA and JA. doi:10.1371/journal.pone.0009408.g003


occurs by ABA activation of the JA pathway[45]. A possible model for the control of *PEX11e* expression by ABA and JA is presented in Figure 3B. The response of *PEX11e* to salt in the different mutant backgrounds mirrors that of *PEX10* and the  $\beta$ -oxidation enzyme 3-ketoacyl thiolase. [54].

Since *PEX11e* is up regulated by salt, and up regulation of *PEX11e* has been reported to induce peroxisome proliferation [21,31] we tested whether salt stress resulted in peroxisome

proliferation. Seedlings of the A5 line which expresses a peroxisomal targeted GFP reporter [63] were grown on 0.5x MS with 1% sucrose supplemented with 150 mM NaCl, or 200 mM mannitol or no supplement for 5 days and GFP in the roots imaged by confocal microscopy. In the control roots peroxisomes were visible (Fig. 4A control). In roots from plants grown on 150 mM NaCl peroxisomes appeared far more abundant (Figure 4B 150 mM NaCl). This effect is due to ionic stress rather than dehydration as peroxisome numbers were not increased in mannitol treated roots (Fig. 4C, 200 mM mannitol), indeed peroxisomes appeared less abundant in mannitol treated roots. Interestingly *PEX1* and *PEX10* transcripts increased in response to salt but not sorbitol treatment [54] suggesting that peroxisomes are responding to the ionic component of salt stress.

### Over Expression of PEX11e Increases Peroxisome Number but Does Not Increase Peroxisome Activity or Tolerance to Salt

Since peroxisome proliferation is induced in salt stress conditions as well as by over expressing PEX11e, we tested the hypothesis that increasing peroxisome number by up regulating PEX11e would provide resistance to salt stress. Columbia ecotype was transformed with a *PEX11e* cDNA construct under the control of the 35S promoter. Three independent transformants (plants 5, 9 and 14) were selected based on high levels of PEX11 expression (Figure S2). T3 (in the case of plant 5 descendents) and T4 (in the case of plant 9 and 14 descendents) were tested for homozygosity (based on 100% kanamycin resistance) and PEX11 expression (based on western blotting). An antibody raised against the  $\beta$  subunit of mitochondrial ATP synthase was used as a loading




**Figure 4. NaCl treatment causes an increase in peroxisome number.** Seeds expressing the peroxisome targeted GFP-MFP2 fusion protein (A5 line) were sown on 0.5x MS media supplemented with 1% sucrose and either 150 mM NaCl, 200 mM mannitol or no addition (control). After 5 days the peroxisomes in the roots were imaged by confocal microscopy. Scale bar = 20  $\mu$ m. doi:10.1371/journal.pone.0009408.g004

control. Only one line 5-6-1 consistently showed over expression of PEX11. Five lines were selected for further study, three descended from plant 5 (5-3-1 5-3-4 and 5-6-1) and two each from plants 9 (9-20-2 and 9-20-3) and 14 (14-9-1 and 14-9-3) (Figure 5).


Peroxisome function was tested in the transgenic lines using three different experiments. Firstly the hypocotyl length of 5 day old seedlings grown in the dark in the presence or absence of sucrose was determined. All the lines tested showed a mild reduction in hypocotyl length (50–85%) compared to wild type when grown in the absence of sucrose, which was restored by the inclusion of sucrose in the medium (Fig. 6A). As sucrose dependence can be indicative of impaired  $\beta$ -oxidation, next the growth of the transgenic lines on 2, 4-dichloro-phenoxybutyric acid (2,4-DB) was examined (Fig. 6B). This compound is bioactivated by  $\beta$ -oxidation to the herbicide 2,4D, resulting in inhibition of root growth. All the lines retained sensitivity to 2,4D as expected. Root growth of wild type Columbia was reduced to 20% of the untreated control at 0.3 and 0.4  $\mu$ M 2,4 DB comparable with previously published results [64]. Lines 5-3-1 and 5-3-4 showed no resistance to 2,4DB whereas lines 5-6-1 9-20-2, 9-20-3, 14-9-1 and 14-9-3 showed increased resistance to 0.3  $\mu$ M 2,4DB indicative of a partial block in conversion of this compound to 2,4D. These lines also had the shortest hypocotyls when grown in the absence of sucrose. For comparison a strong  $\beta$ -oxidation mutant such as *ped3-4* is completely resistant to 0.8  $\mu$ M 2,4-DB and, when grown in the absence of sucrose, produces very short hypocotyls that are ca 20% of the length of hypocotyls of plants grown on sucrose [64]. Thus the PEX11e transgenics appear to have a mild  $\beta$ -oxidation deficiency phenotype. This is in contrast to results obtained by Orth et al [21] who saw no sucrose dependence of PEX11 e over expressing lines, although these authors did not report on 2,4DB sensitivity.

Finally, to investigate a  $\beta$ -oxidation independent function, catalase activity was measured in homogenates from leaf material of the transgenic lines but no striking differences were observed between the lines (data not shown). Western blotting was carried out with anti-thiolase antibodies (on dark grown seedlings) and anti-glycolate oxidase antibodies (on green leaf tissue). While lines 5-3-1 and 5-3-4 showed slightly reduced levels of thiolase protein, there were no changes of level of protein correlated with increased expression of PEX11e. Likewise there were no striking alterations of protein level of glycolate oxidase in the transgenic lines (Figure S3).

Plants of lines 5-3-1, 5-6-1, 9-20-2 and 14-9-1 were crossed to the A5 line as male or female parent and F1 selected on kanamycin plus basta plates to select for both transgenes. Leaves were analysed by confocal microscopy after 14 days and the




**Figure 5. Expression level of PEX11e in transgenic Arabidopsis plants.** A crude membrane fraction was prepared from 5 day old seedlings of the lines indicated grown in the dark. Twenty five microgrammes of protein from each line was separated by SDS-PAGE and immunoblotted with anti-PEX11 antibody or anti ATP $\beta$  antibody. doi:10.1371/journal.pone.0009408.g005



**Figure 6. Physiological analysis of PEX11e transgenics.** A. Hypocotyl growth in the presence and absence of sucrose. Seedlings were grown for 5 days in the dark on 0.5xMS medium  $\pm$  1% sucrose and hypocotyl length measured. Results are mean  $\pm$  SE for 3 replicates of 3–5 seedlings per line. B. Root growth in response to 2,4DB. Seedlings were grown for 5 days in the dark on plates containing 0.5xMS, 0.5% sucrose plus the indicated amount of 2,4DB or 2,4D at 0.05  $\mu$ g/mL. Results are mean  $\pm$ SE of 3 replicates of 3–5 seedlings per line normalised to the length of untreated (control) seedlings for each line. doi:10.1371/journal.pone.0009408.g006

number of peroxisomes quantified (Fig. 7). Crosses with the 5-6-1 line as male or female parent resulted in an increase in peroxisome number consistent with previous reports for over expression of PEX11e [21,25]. The other lines when crossed to the A5 line did not result in an increase in peroxisome number compared to the A5 control. Interestingly, the peroxisome number was always higher when the PEX11e transgenic was used as the female parent, compared to the reciprocal cross. A reduction in peroxisome number compared to the A5 line was seen in both reciprocal crosses with the 9-20-2 and 14-9-1 lines. This reduction in peroxisome number along with the loss of PEX11e over expression from descendants of plants selected for high expression may indicate that high levels of PEX11e expression are not well tolerated by Arabidopsis.

To test if any of the PEX11e lines had greater resistance to salt stress seeds were grown on medium supplemented with the




**Figure 7. Over-expression of PEX11e results in increased peroxisome number.** Reciprocal crosses were made between the PEX11e transgenic lines shown in figure 5 and the A5 line. F1 seed were plated on 0.5x MS supplemented with 1% sucrose, basta and hygromycin to select for both transgenes. Control (homozygous A5 plants) were grown without selection. A. After 2 weeks peroxisomes were observed in the leaves by confocal microscopy. B. Peroxisomes were counted in 10 fields ( $146.57 \times 146.57 \mu\text{m}^2$ ) per cross. \* and \*\* represent significant differences from A5 at  $t < 0.01$  and  $t < 0.05$ , respectively (Student's t-test). doi:10.1371/journal.pone.0009408.g007

indicated amounts of NaCl or mannitol for 5 days, the roots were measured and growth was expressed relative to that of the control. None of the lines tested showed any increased resistance to NaCl (Fig. 8A and B). To test the effect of long term salt stress plants were grown in soil and watered with NaCl supplemented water at the indicated concentrations. After 24 days fresh weight was determined (fig. 8). Again there was no effect on fresh weight.

## Conclusions

Arabidopsis PEX11e is targeted to peroxisomes in tobacco epidermal cells when expressed as a fusion protein with eYFP, and native PEX11c,d,e are peroxisomal in Arabidopsis suspension cells. Pex11c,d,e behave as integral membrane proteins and no evidence could be found for a redox sensitive dimer as has been reported for *S. cerevisiae* PEX11[51]. On native gels PEX11 migrates as a large complex. PEX11e is up regulated by NaCl



**Figure 8. Seedlings with increased level of PEX11e do not show increased resistance to NaCl stress.** Seeds from the lines shown in Figure 5 were grown on 0.5x MS medium plus the indicated amount of NaCl or Mannitol. Root length was measured after 5 days and plotted as a % of the unsupplemented (control) value for each line (A and B). After 24 days the fresh weight of the shoots was determined and expressed as a % of the unsupplemented (control) (C). Graphs show the mean and standard error of 3–5 seedlings from each of 3 independent experiments.

doi:10.1371/journal.pone.0009408.g008

stress in an ABI 3- ABI1- and JAR1-dependent manner suggesting ABA and JA signalling act downstream from salt stress to regulate *PEX11e* expression. Ionic but not dehydration stress resulted in an increase in peroxisome number, but increasing peroxisome number by up regulation of *PEX11e* did not provide increased resistance to salt stress in seedlings or older plants. This is in contrast to studies which have shown that over expression of specific peroxisomal enzymes ascorbate peroxidase [65] and betaine aldehyde dehydrogenase [66] gave enhanced stress tolerance. However the studies reported here did not find any evidence that increased *PEX11e* abundance actually increased peroxisome activity. The transgenic lines rather showed a mild  $\beta$ -oxidation phenotype in seedlings and no indication of increased

catalase activity in leaves. Although increased *PEX11e* expression by salt could provide a mechanism for peroxisome proliferation under salt stress conditions, the hypothesis that the quantity of peroxisomes limits plant stress tolerance proved incorrect and increasing peroxisome number by manipulating peroxisome biogenesis related genes is unlikely to be a useful strategy for increasing salt tolerance.

## Materials and Methods

### Construction and Characterisation of Transgenic Plants

The *PEX11e* gene was amplified from an Arabidopsis 2 day seedling cDNA library (a gift from Prof Ian Graham, University of York) with the primers (5' ATGACTACACTAGATTTGACC 3'), and antisense (5' TCTTCAACTTGGGGCGCGATG 3') and the product cloned into pGEM-T easy and fully sequenced. Overexpression constructions were carried out using the GATEWAY system (Invitrogen). The construct was generated by PCR with two primers: attB1-*AtPex11-2* (5'-GGGGACAAGTTTGTACAAAAAAGCAGGCTTCACCATGACTACACTAGATTTGACC-3') and attB2-*AtPex11-2* (5'-GGGGACCACCTTTGTA-CAAGAAAGCTGGGTCTCATGATTTCTTCAACTTGG-3') The PCR products flanked by *attB1* and *attB2* (underlined) were recombined into pDONR-207 (Invitrogen).

The intermediate, *AtPex11e attL1 -attL2*-containing pDONR-207 vector was recombined with the Gateway enabled destination vector pGD625 [67] containing the 35S promoter and nos terminator which we used for overexpression. Inserts containing clones were verified by restriction enzyme digestion and by sequencing. Arabidopsis plants ecotype Columbia were transformed by the floral dip method [68], selected on kanamycin and checked for overexpression of *PEX11e* by western blotting. Three T1 plants (lines 5, 9 and 14) were selected for further analysis (Figure S1). Experiments described in this paper were performed on homozygous T4 (lines beginning 5) or T5 (lines beginning 9 or 14) plants which were selected from parents showing high level of expression of *PEX11e* by western in the previous generation. For confocal analysis reciprocal crosses were made to plants of the A5 line [63] which expresses a peroxisomal targeted GFP-MFP2 fusion protein.

### Antibodies, Polyacrylamide Gel Electrophoresis and Western Blotting

A polyclonal antiserum was raised in rabbit to the peptide VLYLNKAEARDKICRAIQYGSKFLSC corresponding to amino acids 15 to 40 of Arabidopsis *PEX11e* (Genosphere technologies). This region is highly conserved in *PEX11c*, *d* and *e*, therefore the antibody would be expected to cross react with these isoforms as well. The carboxy-terminal C was added to facilitate affinity purification of antibodies by immobilising the peptide antigen to SulfoLink Coupling Gel (Pierce) according to the manufacturer's standard protocol, yielding affinity purified antibodies with protein concentrations of 0.7 and 0.9 mg ml<sup>-1</sup>. Proteins were separated by standard SDS-PAGE and transferred to nitrocellulose membranes (0.45  $\mu$ m, Micron Separations, Schleicher&Schuell). Except where indicated, sample volumes loaded on gels were equivalent. Membranes were blocked overnight at 4°C in TBST (50 mM Tris-HCl, pH 7.4, 200 mM NaCl, and 0.1% v/v Tween 20) containing 5% (w/v) non-fat dry milk, then incubated in TBST plus 5% (w/v) non-fat dry milk (TBST-milk) containing affinity-purified *PEX11e* antibodies at dilutions of 1:2,000. Incubations with primary antibody were carried out 2 h at temperature room. After four 10-min washes in TBST, membranes were incubated with horseradish peroxidase-



conjugated goat-anti-rabbit secondary antibodies (Sigma) for 2 h at 1:3,000 dilution in TBST-milk. After four 10-min washes in TBST, membranes were developed by enhanced chemi-luminescence and the signal was recorded on X-Ray film (Fuji Medical).

### Preparation of Membrane Fractions from Seedlings

Seedlings were ground to a fine powder in liquid N<sub>2</sub>. Homogenisation buffer (50 mM Tris HCl pH 8.2, 2 mM EDTA, 20% v/v glycerol, 1 mM PMSF, 2% protease inhibitor cocktail, 0.5 mM DTT) was added at a ratio of 0.1 mL per 0.1 g fresh weight and ground again to produce a slurry. The homogenate was centrifuged 10 min 800 g 4°C, the supernatant carefully removed and centrifuged for 30 minutes at 4°C 100,000 g, the supernatant was removed and the pellet resuspended in 30 microlitre homogenization buffer plus 5 microlitres 10% SDS.

### Preparation of Membrane Fractions from Cell Culture

Cell culture was pelleted and washed twice in PBS and once in homogenisation buffer 50 mM Tris-HCl pH 8.2, 2 mM EDTA, 20% Glycerol, and protease inhibitor tablets (Sigma). The cell pellet was then ground in liquid nitrogen for 10 min, and resuspended in homogenisation buffer at 1 ml/g ground material. This was finally filtered twice through miracloth (Millipore) and spun for 10 min at 3000 g at 4°C. The supernatant was submitted to a second centrifugation at 10000 g for 30 min at 4°C. The pellet was resuspended in homogenisation buffer and the protein concentration was measured with BCA reagent (Pierce) with bovine serum albumen as standard.

### Carbonate Extraction

100 microgrammes of proteins were aliquoted and either left untreated (T) or extracted by 0.1 M Na<sub>2</sub>CO<sub>3</sub> for 30 min on ice. Aliquots were then spun for 20 min at 4°C. The supernatant was precipitated with 20% TCA for 20 min and pelleted at 14000 g for 20 min. Pellets were resuspended in sample buffer with or without DTT as indicated and loaded on a denaturing gel. As a control thiolase (peroxisome matrix protein) signal was found in the supernatant (not shown) confirming the success of carbonate extraction.

### Native Gel

100 microgrammes of protein from a peroxisome enriched organelle fraction was aliquoted and solubilised in the indicated concentration of detergent (dodecyl maltoside; DDM, or Triton X-100; Triton) for 30 minutes on ice. After a clearing spin of 100,000 g for 20 minutes at 4°C, the samples were separated on a NuPAGE 3–8% native gel (Invitrogen) according to the manufacturer's instructions. After transfer to PVDF the membrane was incubated with anti-PEX11 affinity purified antiserum.

### Salt Stress Experiments

Treatment of seedlings, quantitative real time PCR experiments and analysis of data were carried out exactly as described in Charlton et al., 2005. Primers 11-2F

5'-TGA ATT GCT TGG ACG TAT ATC ACT TT -3' and 11-2R 5'-CAC CAA TCT CGA CTG CAC TTG T-3' were used.

For seedling experiments seeds were grown on 0.5x MS media, supplemented as described in the figure legends. Plates were maintained in a growth room (16 h light 20 °C) for the required number of days before being scanned and root length quantified using the programme Image J. For longer term effects of salt stress seeds were germinated on 0.5x MS agar plates. After 1 week they

were transplanted to soil and watered with tap water for a further week. Plants were then watered with tap water supplemented with 100, 200 or 300 mM NaCl for a further 24 days before measuring the fresh weight.

### 2,4DB Resistance and Sucrose Dependence

Experiments were carried out as described [64].

### Confocal Microscopy and Immunofluorescence

AtPEX11e was amplified using Pfx from the AtPEX11 overexpression vector (see previously) using primers AtPEX11-2A (5' GGGGACAAGTTTGTACAAAAAAGCA GCCTTC-CCGCCAATGACTACACTAGATTTGACCAGAG 3') and AtPEX11-2B (5' GGGGACCACCTTTGTACAAGAAAGCTGGTCTCAAGGTGTCTTCAACTTGG 3'), cloned into pDONOR207 and subsequently recombined into the plant binary destination vector 35S-eYFP-cassette-Nos::pCAMBIA 1300 [49] resulting in eYFP-AtPEX11e. All vectors were sequenced and verified. *Agrobacterium tumefaciens* was transformed according to the freeze thaw procedure [69]. Transient expression in tobacco leaf epidermal cells was carried out according to Sparkes et al [70] where eYFP-AtPEX11e and CFP-SKL were infiltrated at an optical density of 0.1 and 0.04 respectively. Dual expression was imaged using a Zeiss inverted LSM510 using the settings stated in Sparkes et al. [49]. Arabidopsis cell culture was harvested 4 days after subculture, immunofluorescence and subsequent imaging were carried out according to Sparkes et al. [49], where anti-PEX11 antibody (1:40) and anti-ICL (1:1000) with Texas red-conjugated goat anti-rabbit IgG (H+L; Molecular Probes; 1:100) and FITC-conjugated goat anti-rabbit IgG (Sigma Aldrich, 1:40) respective secondary antibodies were used.

To determine the effect of NaCl on the morphology of peroxisomes, A5 seedlings were grown on 0.5x MS with 1% (w/v) sucrose supplemented with 150 mM NaCl, 200 mM mannitol or no supplement for 5 days and the middle part of the roots was used for confocal microscopy. For numerical analysis of peroxisomes in Pex11-e overexpressing plants, the seedlings of reciprocal crosses between Pex11-e introduced plants and A5 line were grown on 0.5x MS with 1% (w/v) sucrose plates containing 50 µg/ml kanamycin and 10 µg/ml glufosinate ammonium (BASTA) for 2 weeks. Then the abaxial epidermis of leaves was used for confocal microscopy and the number of peroxisomes was obtained from confocal microscope images captured from 150 µm × 150 µm.

### Supporting Information

**Figure S1** PEX11e is upregulated by salt stress. Seedlings were treated with 200 mM or 400 mM NaCl for 4.5 h then the steady state levels of PEX11e transcripts determined by QPCR. Found at: doi:10.1371/journal.pone.0009408.s001 (1.76 MB TIF)

**Figure S2** Primary transformants with increased level of expression of PEX11e. Forty microgrammes of membrane protein from 35 day old primary transformant plants number 5, 9 and 14 were separated by SDS PAGE. Panel A immunoblot with anti-PEX11c/d/e antiserum. Panel B Coomassie protein stain of the same membrane protein fractions. Found at: doi:10.1371/journal.pone.0009408.s002 (0.45 MB TIF)

**Figure S3** PEX11e transgenics do not have altered levels of glycolate oxidase or 3-ketoacyl thiolase protein. Western blot analysis of thiolase and glycolate oxidase in total protein extracts (20 µg per lane) from dark grown seedlings (thiolase) and green leaves of 4 week old plants (glycolate oxidase) of the indicated lines.

The Ponceau S stained membrane is shown in each case to verify equal loading of the lanes.

Found at: doi:10.1371/journal.pone.0009408.s003 (6.98 MB TIF)

## Acknowledgments

We thank Mrs F. Moulton for excellent technical assistance.

## References

- Brown L-A, Baker A (2008) Shuttles and cycles; transport of proteins into the peroxisome matrix (review). *Molecular Membrane Biology* 25: 363–375.
- Hoepfner D, Schildknecht D, Braakman I, Philippsen P, Tabak HF (2005) Contribution of the endoplasmic reticulum to peroxisome formation. *Cell* 122: 85–95.
- Kim PK, Mullen RT, Schumann U, Lippincott-Schwartz J (2006) The origin and maintenance of mammalian peroxisomes involves a de novo PEX16-dependent pathway from the ER. *Journal Of Cell Biology* 173: 521–532.
- Tam YYC, Fagarasanu A, Fagarasanu M, Rachubinski RA (2005) Pex3p initiates the formation of a preperoxisomal compartment from a subdomain of the endoplasmic reticulum in *Saccharomyces cerevisiae*. *Journal Of Biological Chemistry* 280: 34933–34939.
- Mullen RT, Trelease RN (2006) The ER-peroxisome connection in plants: Development of the “ER semi-autonomous peroxisome maturation and replication” model for plant peroxisome biogenesis. *Biochimica Et Biophysica Acta-Molecular Cell Research* 1763: 1655–1668.
- Fagarasanu A, Fagarasanu M, Rachubinski RA (2007) Maintaining peroxisome populations: A story of division and inheritance. *Annual Review Of Cell And Developmental Biology* 23: 321–344.
- Schrader M (2006) Shared components of mitochondrial and peroxisomal division. *Biochimica Et Biophysica Acta-Molecular Cell Research* 1763: 531–541.
- Hess R, Staubli W, Riess W (1965) Nature of the hepatomegalic effect produced by ethyl-chlorophenoxy-isobutyrate in the rat. *Nature* 208: 856–858.
- Gurvitz A, Rottensteiner H (2006) The biochemistry of oleate induction: Transcriptional upregulation and peroxisome proliferation. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 1763: 1392.
- van der Klei IJ, Yurimoto H, Sakai Y, Veenhuis M (2006) The significance of peroxisomes in methanol metabolism in methylotrophic yeast. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* 1763: 1453.
- Baker A, Graham I, Holdsworth M, Smith S, Theodoulou F (2006) Chewing the fat:  $\beta$ -oxidation in signalling and development. *Trends in Plant Science* 11: 124–132.
- Desai M, Hu J (2008) Light Induces Peroxisome Proliferation in Arabidopsis Seedlings through the Photoreceptor Phytochrome A, the Transcription Factor HY5 HOMOLOG, and the Peroxisomal Protein PEROXIN11b. *Plant Physiol* 146: 1117–1127.
- Thoms S, Erdmann R (2005) Dynamin-related proteins and Pex11 proteins in peroxisome division and proliferation. *Febs Journal* 272: 5169–5181.
- Hu JP (2007) Plant peroxisome multiplication: Highly regulated and still enigmatic. *Journal Of Integrative Plant Biology* 49: 1112–1118.
- Erdmann R, Blobel G (1995) Giant Peroxisomes In Oleic Acid-Induced *Saccharomyces-Cerevisiae* Lacking The Peroxisomal Membrane-Protein Pmp27p. *Journal Of Cell Biology* 128: 509–523.
- Marshall PA, Krimkevich YI, Lark RH, Dyer JM, Veenhuis M, et al. (1995) Pmp27 Promotes Peroxisomal Proliferation. *Journal Of Cell Biology* 129: 345–355.
- Tam YYC, Torres-Guzman JC, Vizeacoumar FJ, Smith JJ, Marelli M, et al. (2003) Pex11-related proteins in peroxisome dynamics: A role for the novel peroxin Pex27p in controlling peroxisome size and number in *Saccharomyces cerevisiae*. *Molecular Biology of the Cell* 14: 4089–4102.
- Rottensteiner H, Stein K, Sonnenhol E, Erdmann R (2003) Conserved function of Pex11p and the novel Pex25p and Pex27p in peroxisome biogenesis. *Molecular Biology of the Cell* 14: 4316–4328.
- Vizeacoumar FJ, Torres-Guzman JC, Tam YYC, Aitchison JD, Rachubinski RA (2003) YHR150w and YDR479c encode peroxisomal integral membrane proteins involved in the regulation of peroxisome number, size, and distribution in *Saccharomyces cerevisiae*. *Journal of Cell Biology* 161: 321–332.
- Vizeacoumar FJ, Torres-Guzman JC, Bouard D, Aitchison JD, Rachubinski RA (2004) Pex30p, Pex31p, and Pex32p form a family of peroxisomal integral membrane proteins regulating peroxisome size and number in *Saccharomyces cerevisiae*. *Molecular Biology of the Cell* 15: 665–677.
- Orth T, Reumann S, Zhang XC, Fan JL, Wenzel D, et al. (2007) The PEROXIN11 protein family controls peroxisome proliferation in Arabidopsis. *Plant Cell* 19: 333–350.
- Li XL, Baumgart E, Morrell JC, Jimenez-Sanchez G, Valle D, et al. (2002) PEX11 beta deficiency is lethal and impairs neuronal migration but does not abrogate peroxisome function. *Molecular and Cellular Biology* 22: 4358–4365.
- Li XL, Baumgart E, Dong GX, Morrell JC, Jimenez-Sanchez G, et al. (2002) PEX11 alpha is required for peroxisome proliferation in response to 4-phenylbutyrate but is dispensable for peroxisome proliferator-activated receptor alpha-mediated peroxisome proliferation. *Molecular and Cellular Biology* 22: 8226–8240.
- Schrader M, Reuber BE, Morrell JC, Jimenez-Sanchez G, Obie C, et al. (1998) Expression of PEX11 beta mediates peroxisome proliferation in the absence of extracellular stimuli. *Journal of Biological Chemistry* 273: 29607–29614.
- Lingard MJ, Trelease RN (2006) Five Arabidopsis peroxin 11 homologs individually promote peroxisome elongation, duplication or aggregation. *Journal Of Cell Science* 119: 1961–1972.
- Li XL, Gould SJ (2003) The dynamin-like GTPase DLP1 is essential for peroxisome division and is recruited to peroxisomes in part by PEX11. *Journal of Biological Chemistry* 278: 17012–17020.
- Kobayashi S, Tanaka A, Fujiki Y (2007) Fis1, DLP1, and Pex11p coordinately regulate peroxisome morphogenesis. *Experimental Cell Research* 313: 1675.
- Mano S, Nakamori C, Kondo M, Hayashi M, Nishimura M (2004) An Arabidopsis dynamin-related protein, DRP3A, controls both peroxisomal and mitochondrial division. *Plant Journal* 38: 487–498.
- Zhang XC, Hu JP (2009) Two small protein families, DYNAMIN-RELATED PROTEIN3 and FISSION1, are required for peroxisome fission in Arabidopsis. *Plant Journal* 57: 146–159.
- Zhang XC, Hu JP (2008) FISSION1A and FISSION1B Proteins Mediate the Fission of Peroxisomes and Mitochondria in Arabidopsis. *Molecular Plant* 1: 1036–1047.
- Lingard MJ, Gidda SK, Bingham S, Rothstein SJ, Mullen RT, et al. (2008) Arabidopsis PEROXIN11c-e, FISSION1b, and DYNAMIN-RELATED PROTEIN3A Cooperate in Cell Cycle-Associated Replication of Peroxisomes. *Plant Cell* 20: 1567–1585.
- Nyathi Y, Baker A (2006) Plant peroxisomes as a source of signalling molecules. *Biochimica Et Biophysica Acta-Molecular Cell Research* 1763: 1478–1495.
- Ferreira RMB, Bird B, Davies DD (1989) The Effect of Light on the Structure and Organization of Lemna Peroxisomes. *J Exp Bot* 40: 1029–1035.
- Morré DJ, Selldén G, Ojanperä K, Sandelius AS, Egger A, et al. (1990) Peroxisome proliferation in Norway spruce induced by ozone. *Protoplasma* 155: 58.
- Oksanen E, Haikio E, Sober J, Karnosky DF (2004) Ozone-Induced H<sub>2</sub>O<sub>2</sub> Accumulation in Field-Grown Aspen and Birch Is Linked to Foliar Ultrastructure and Peroxisomal Activity. *New Phytologist* 161: 791.
- Palma JM, Gomez M, Yanez J, Del Rio LA (1987) Increased Levels of Peroxisomal Active Oxygen-Related Enzymes in Copper-Tolerant Pea Plants. *Plant Physiol* 85: 570–574.
- McCarthy I, Romero-Puertas MC, Palma JM, Sandalio LM, Corpas FJ, et al. (2001) Cadmium induces senescence symptoms in leaf peroxisomes of pea plants. *Plant, Cell & Environment* 24: 1065–1073.
- Corpas FJ, Gomez M, Hernandez JA, del Rio LA (1993) Metabolism of activated oxygen in peroxisomes from two *Pisum sativum* L. cultivars with different sensitivity to sodium chloride. *Journal of Plant Physiology* 141.
- de Felipe MR, Lucas MM, Pozuelo JM (1988) *J Plant Physiol* 132: 67–73.
- Palma JM, Garrido M, Rodriguez-Garcia MI, Del Rio LA (1991) Peroxisome proliferation and oxidative stress mediated by activated oxygen species in plant peroxisomes. *Archives of Biochemistry and Biophysics* 287: 68–74.
- del Rio LA, Sandalio LM, Corpas FJ, Palma JM, Barroso JB (2006) Reactive Oxygen Species and Reactive Nitrogen Species in Peroxisomes. Production, Scavenging, and Role in Cell Signaling. *Plant Physiol* 141: 330–335.
- Zhu JK (2002) Salt and drought stress signal transduction in plants. *Annual Review of Plant Biology* 53: 247–273.
- Seki M, Narusaka M, Ishida J, Nanjo T, Fujita M, et al. (2002) Monitoring the expression profiles of 7000 Arabidopsis genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *Plant Journal* 31: 279–292.
- Himmelbach A, Yang Y, Grill E (2003) Relay and control of abscisic acid signaling. *Current Opinion in Plant Biology* 6: 470–479.
- Lorenzo O, Chico JM, Sanchez-Serrano JJ, Solano R (2004) JASMONATE-INSENSITIVE1 Encodes a MYC Transcription Factor Essential to Discriminate between Different Jasmonate-Regulated Defense Responses in Arabidopsis. *Plant Cell* 16: 1938–1950.
- Giraudat J, Hauge BM, Valon C, Smalle J, Parcy F, et al. (1992) Isolation of the Arabidopsis ABI3 Gene by Positional Cloning. *Plant Cell* 4: 1251–1261.
- Finkelstein RR, Wang ML, Lynch TJ, Rao S, Goodman HM (1998) The Arabidopsis Abscisic Acid Response Locus ABI4 Encodes an APETALA 2 Domain Protein. *Plant Cell* 10: 1043–1054.
- Zimmermann P, Hirsch-Hoffmann M, Hennig L, Gruissem W (2004) GENEVESTIGATOR. Arabidopsis Microarray Database and Analysis Toolbox. *Plant Physiol* 136: 2621–2632.

## Author Contributions

Conceived and designed the experiments: SM MES IAS WLC CDML AB. Performed the experiments: SM MES IAS WLC CDML BJ. Analyzed the data: SM IAS WLC CDML AB. Contributed reagents/materials/analysis tools: MES BJ. Wrote the paper: AB. Performed essential supporting work: BJ.

49. Sparkes IA, Hawes C, Baker A (2005) AtPEX2 and AtPEX10 are targeted to peroxisomes independently of known endoplasmic reticulum trafficking routes. *Plant Physiology* 139: 690–700.
50. Tugal HB, Pool M, Baker A (1999) Arabidopsis 22-kilodalton peroxisomal membrane protein. Nucleotide sequence analysis and biochemical characterization. *Plant Physiology* 120: 309–320.
51. Marshall PA, Dyer JM, Quick ME, Goodman JM (1996) Redox-sensitive homodimerization of Pex11p: A proposed mechanism to regulate peroxisomal division. *Journal of Cell Biology* 135: 123–137.
52. Rahim G, Bischof S, Kessler F, Agne B (2009) In vivo interaction between atToc33 and atToc159 GTP-binding domains demonstrated in a plant split-ubiquitin system. *J Exp Bot* 60: 257–267.
53. Nayidu NK, Wang L, Xie WB, Zhang CJ, Fan CZ, et al. (2008) Comprehensive sequence and expression profile analysis of PEX11 gene family in rice. *Gene* 412: 59–70.
54. Charlton WL, Matsui K, Johnson B, Graham I, Ohme-Takagi M, et al. (2005) Salt-induced expression of peroxisome-associated genes requires components of the ethylene, jasmonate and abscisic acid signalling pathways. *Plant Cell Environ* 28: 513–524.
55. Nito K, Kamigaki A, Kondo M, Hayashi M, Nishimura M (2007) Functional classification of Arabidopsis peroxisome biogenesis factors proposed from analyses of knockdown mutants. *Plant And Cell Physiology* 48: 763–774.
56. Nambara E, Naito S, McCourt P (1992) A mutant of Arabidopsis which is defective in seed development and storage protein accumulation is a new *abi3* allele. *The Plant Journal* 2: 435–441.
57. Leung J, Merlot S, Giraudat J (1997) The Arabidopsis ABSCISIC ACID-INSENSITIVE2 (*ABI2*) and *ABI1* Genes Encode Homologous Protein Phosphatases 2C Involved in Abscisic Acid Signal Transduction. *Plant Cell* 9: 759–771.
58. Staswick PE, Tiryaki I, Rowe ML (2002) Jasmonate Response Locus *JAR1* and Several Related Arabidopsis Genes Encode Enzymes of the Firefly Luciferase Superfamily That Show Activity on Jasmonic, Salicylic, and Indole-3-Acetic Acids in an Assay for Adenylation. *Plant Cell* 14: 1405–1415.
59. Staswick PE, Tiryaki I (2004) The Oxylinin Signal Jasmonic Acid Is Activated by an Enzyme That Conjugates It to Isoleucine in Arabidopsis. *Plant Cell* 16: 2117–2127.
60. Thines B, Katsir L, Melotto M, Niu Y, Mandaokar A, et al. (2007) JAZ repressor proteins are targets of the SCFCO11 complex during jasmonate signalling. *Nature* 448: 661.
61. Rouster J, Leah R, Mundy J, Cameron-Mills V (1997) Identification of a methyl jasmonate-responsive region in the promoter of a lipoxygenase 1 gene expressed in barley grain. *The Plant Journal* 11: 513–523.
62. Chini A, Fonseca S, Fernandez G, Adie B, Chico JM, et al. (2007) The JAZ family of repressors is the missing link in jasmonate signalling. *Nature* 448: 666.
63. Cutler SR, Ehrhardt DW, Griffitts JS, Somerville CR (2000) Random GFP: cDNA fusions enable visualization of subcellular structures in cells of Arabidopsis at a high frequency. *Proceedings of the National Academy of Sciences of the United States of America* 97: 3718–3723.
64. Dietrich D, Schmuths H, Lousa CD, Baldwin JM, Baldwin SA, et al. (2009) Mutations in the Arabidopsis Peroxisomal ABC Transporter COMATOSE Allow Differentiation between Multiple Functions In Planta: Insights from an Allelic Series. *Molecular Biology Of The Cell* 20: 530–543.
65. Li YJ, Hai RL, Du XH, Jiang XN, Lu H (2009) Over-expression of a *Populus* peroxisomal ascorbate peroxidase (*PpAPX*) gene in tobacco plants enhances stress tolerance. *Plant Breeding* 128: 404–410.
66. Kishitani S, Takanami T, Suzuki M, Oikawa M, Yokoi S, et al. (2000) Compatibility of glycinebetaine in rice plants: evaluation using transgenic rice plants with a gene for peroxisomal betaine aldehyde dehydrogenase from barley. *Plant Cell Environ* 23: 107–114.
67. Favaro R, Pinyopich A, Battaglia R, Kooiker M, Borghi L, et al. (2003) MADS-Box Protein Complexes Control Carpel and Ovule Development in Arabidopsis. *Plant Cell* 15: 2603–2611.
68. Clough SJ, Bent AF (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *The Plant Journal* 16: 735–743.
69. Hofgen R, Willmitzer L (1988) Storage of competent cells for *Agrobacterium* transformation. *Nucleic Acids Research* 16: 9877.
70. Sparkes I, Runions J, Kearns A, Hawes C (2006) Rapid, transient expression of fluorescent fusion proteins in tobacco plants and generation of stably transformed plants. *Nature Protocols* 1: 2019–2025.