



This is a repository copy of *Functional genomics reveals that Clostridium difficile Spo0A coordinates sporulation, virulence and metabolism.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/78494/>

Version: Supplemental Material

---

**Article:**

Pettit, L.J., Browne, H.P., Yu, L. et al. (10 more authors) (2014) Functional genomics reveals that Clostridium difficile Spo0A coordinates sporulation, virulence and metabolism. BMC Genomics, 15. 160. ISSN 1471-2164

<https://doi.org/10.1186/1471-2164-15-160>

---

**Reuse**


Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>



Additional file 1: Identification of differentially expressed genes in *C. difficile* 630 *spo0A* mutant by transcriptional profiling. Scatter plot of the log2 fold changes against the normalised mean read abundance per gene (calculated at the base level). Red dots represent genes considered to be significantly differentially expressed ( $P = \leq 0.01$ ). Black dots signify genes not deemed to be significantly differentially expressed according to these criteria.