

This is a repository copy of *Targeting subgroups to realise value:the cost-effectiveness of omalizumab in severe allergic asthma.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/80776/>

Version: Submitted Version

Conference or Workshop Item:

Faria, Rita orcid.org/0000-0003-3410-1435, McKenna, Claire orcid.org/0000-0002-7759-4084 and Palmer, Stephen John orcid.org/0000-0002-7268-2560 Targeting subgroups to realise value:the cost-effectiveness of omalizumab in severe allergic asthma. In: UNSPECIFIED. (Unpublished)

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Targeting subgroups to realise value: The cost-effectiveness of omalizumab in severe allergic asthma

Rita Faria, Claire McKenna and Steve Palmer
Centre for Health Economics, University of York, UK


Background

- Patients with uncontrolled severe persistent allergic asthma balance asthma control with adverse effects from treatment.
- Omalizumab offers an effective alternative to stepping up treatment but at a substantial cost.
- Although omalizumab is unlikely to be cost-effective for the overall patient population, it may be good value for money for severe subgroups.

Population and subgroups

- Population reflects marketing authorisation
 - Patients uncontrolled at step 4
 - Patients controlled at step 5
 - Two age groups: 6-11 and ≥ 12
- Population subgroups defined by indicators of severity:
 - Hospitalisation for asthma in past year
 - Maintenance OCS use (step 5)
 - ≥ 3 exacerbations in the past year


The inputs



Objectives

- Evaluate cost-effectiveness of omalizumab for severe persistent asthma for the overall patient population covered by the marketing authorisation and relevant subgroups under the UK NHS perspective.
- Omalizumab 75mg or 150mg solution for injection + optimised therapy vs optimised therapy alone (step 4 or step 5)
- Assessment for response to treatment at 16 weeks:
 - Non-responders should discontinue omalizumab


The model



Results

Population	Age	ICER, /QALY
Overall	6-11 years	98,072€
	≥ 12 years	91,276€
Hospitalisation	6-11 years	54,324€
	≥ 12 years	51,646€
Maintenance OCS	≥ 12 years	58,712€
≥ 3 Exacerbations	6-11 years	91,106€
	≥ 12 years	89,094€

ICERs > conventional thresholds of cost-effectiveness used in the UK NHS of 23,400€ (£20,000) and 35,100€ (£30,000)/QALY



Conclusions

- Omalizumab improves the health outcomes of patients but at a high cost to the UK NHS.
- ICERs were lower in more severe subgroup but still above conventional cost-effectiveness thresholds.
- Asthma-related mortality is one of the key cost-effectiveness drivers:
 - ↑ asthma-related mortality → ↓ ICER

References and Acknowledgements

This project was funded by the National Institute for Health Research Health Technology Assessment (NIHR HTA) Programme on behalf of the National Institute for Health and Clinical Excellence. The views expressed are those of the authors and not necessarily those of the NIHR HTA Programme. Any errors are the responsibility of the authors.

Electronic Medicines Compendium. *Xolair 150mg powder and solvent for solution for injection. Summary of product characteristics*. 2013. [cited 30/04/2013]. Available from: <http://www.medicines.org.uk/emc/medicine/17029#INDICATIONS>.

Bousquet J, Siergiejko Z, Swiebocka E, Humbert M, Rabe KF, Smith N, et al. Persistency of response to omalizumab therapy in severe allergic (IgE-mediated) asthma. *ALLERGY* 2011;**66**:671-8.

Humbert M, Beasley R, Ayres J, Slavin R, Hebert J, Bousquet J, et al. Benefits of omalizumab as add-on therapy in patients with severe persistent asthma who are inadequately controlled despite best available therapy (GINA 2002 step 4 treatment): INNOVATE. *Allergy* 2005;**60**:309-16.

Lanier B, Bridges T, Kulus M, Taylor AF, Berhane I, Vidaurre CF. Omalizumab for the treatment of exacerbations in children with inadequately controlled allergic (IgE-mediated) asthma. *Journal of Allergy and Clinical Immunology* 2009;**124**:1210-6.

de Vries F, Setakis E, Zhang B, van Staa TP. Long-acting beta2-agonists in adult asthma and the pattern of risk of death and severe asthma outcomes: a study using the GPRD. *Eur Respir J* 2010;**36**:494-502.

Novartis Pharmaceuticals UK Ltd. *Xolair® (omalizumab) for the Treatment of Severe Persistent Allergic Asthma (review of TA133 and TA201). Manufacturer Submission of Evidence* 2012 12/01/2012. Available from: <http://guidance.nice.org.uk/TA/WaveR110/Consultation/EvaluationReport/ManufacturerSubmissions>