



UNIVERSITY OF LEEDS

This is a repository copy of *Characterisation of 11 β -hydroxysteroid dehydrogenase 1 in human orbital adipose tissue: a comparison with subcutaneous and omental fat.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/81076/>

Version: Published Version

Article:

Bujalska, IJ, Durrani, OM, Abbott, J et al. (8 more authors) (2007) Characterisation of 11 β -hydroxysteroid dehydrogenase 1 in human orbital adipose tissue: a comparison with subcutaneous and omental fat. *Journal of Endocrinology*, 192 (2). 279 - 288. ISSN 0022-0795

<https://doi.org/10.1677/JOE-06-0042>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Characterisation of 11 β -hydroxysteroid dehydrogenase 1 in human orbital adipose tissue: a comparison with subcutaneous and omental fat

Iwona J Bujalska¹, Omar M Durrani², Joseph Abbott², Claire U Onyimba^{1,2}, Pamela Khosla^{1,2}, Areeb H Moosavi³, Tristan T Q Reuser³, Paul M Stewart¹, Jeremy W Tomlinson¹, Elizabeth A Walker^{1,*} and Saaeha Rauz^{2,*}

¹Department of Endocrinology, Division of Medical Sciences, ²Academic Unit of Ophthalmology, Division of Immunity and Infection, University of Birmingham, ³Birmingham and Midland Eye Centre, Dudley Road, Birmingham B18 7QU, UK

(Requests for offprints should be addressed to S Rauz Academic; Email: s.rauz@bham.ac.uk)

*(E A Walker and S Rauz share the role of senior author)

Abstract

Glucocorticoids (GCs) have a profound effect on adipose biology increasing tissue mass causing central obesity. The pre-receptor regulation of GCs by 11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1) that activates cortisol from cortisone has been postulated as a fundamental mechanism underlying the metabolic syndrome mediating adipocyte hyperplasia and hypertrophy in the omental (OM) depot. Orbital adipose tissue (OF) is the site of intense inflammation and tissue remodelling in several orbital inflammatory disease states. In this study, we describe features of the GC metabolic pathways in normal human OF depot and compare it with subcutaneous (SC) and OM depots. Using an automated histological characterisation technique, OF adipocytes were found to be significantly smaller (parameters: area, maximum diameter and perimeter) than OM and SC adipocytes ($P < 0.001$). Although immunohistochemical analyses demonstrated resident CD68⁺ cells in all three

whole tissue adipose depots, OF CD68 mRNA and protein expression exceeded that of OM and SC (mRNA, $P < 0.05$; protein, $P < 0.001$). In addition, there was higher expression of glucocorticoid receptor (GR) α mRNA in the OF whole tissue depot ($P < 0.05$). Conversely, 11 β -HSD1 mRNA together with the markers of late adipocyte differentiation (FABP4 and G3PDH) were significantly lower in OF. Primary cultures of OF preadipocytes demonstrated predominant 11 β -HSD1 oxo-reductase activity with minimal dehydrogenase activity. Orbital adipocytes are smaller, less differentiated, and express low levels of 11 β -HSD1 but abundant GR α compared with SC and OM. OF harbours a large CD68⁺ population. These characteristics define an orbital microenvironment that has the potential to respond to sight-threatening orbital inflammatory disease.

Journal of Endocrinology (2007) **192**, 279–288

Introduction

Adipose tissue is a highly active endocrine organ regulated not only by a range of external factors, such as the autonomic nervous system and circulating hormones, but also has the ability to modulate its own metabolic activity. One important group of regulating factors are the adrenocorticosteroids. Of these, glucocorticoids (GCs) have a profound effect on adipose biology increasing tissue mass giving rise to obesity in conditions such as Cushing's disease or following the use of exogenous glucocorticoids. Their role in simple obesity is, however, controversial. Regulation of adipose tissue mass involves integration of a variety of complex mechanisms, including the enzyme 11 β -hydroxysteroid dehydrogenase type 1 (11 β -HSD1). 11 β -HSD1 is an NADPH-dependent short-chain alcohol dehydrogenase, localised in the lumen of the endoplasmic reticulum, that interconverts active and

inactive GCs (Tomlinson *et al.* 2004). Although it has bidirectional capacity, *in vivo* it functions as an oxo-reductase activating cortisol from cortisone due to the provision of a cofactor, NADPH, by a second regulatory enzyme, hexose-6-phosphate dehydrogenase (H6PDH), thereby mediating glucocorticoid receptor (GR) responses (Draper *et al.* 2003, Bujalska *et al.* 2005). 11 β -HSD1 is involved in a number of disease processes, including insulin resistance (Kotelevtsev *et al.* 1997), osteoporosis (Cooper *et al.* 2000, Tomlinson *et al.* 2000), glaucoma (Rauz *et al.* 2003), ocular surface renewal (Onyimba *et al.* 2006), inflammation (Thieringer *et al.* 2001, Zhang *et al.* 2005) and the metabolic syndrome – a cluster of cardiovascular risk factors, including hypertension, obesity, hyperlipidaemia and insulin resistance (Masuzaki *et al.* 2003).

11 β -HSD1 is highly expressed in human omental (OM) and subcutaneous (SC) tissue, although expression and activity are higher in OM preadipocytes, the fibroblast-like

precursors of mature adipocytes, when compared with SC adipose tissue (Bujalska *et al.* 1997, 1999). The enzyme is induced upon adipocyte differentiation in human adipose tissue cultures (preadipocytes to adipocytes) and is thought to be related to a 'switch' in enzyme set point from dehydrogenase (preadipocytes) to oxo-reductase (adipocytes) without any significant change in 11 β -HSD1 mRNA levels (Bujalska *et al.* 2002). This observation is explained by the induction of H6PDH across differentiation (Bujalska *et al.* 2005). With the known effect of glucocorticoids on adipose tissue function and distribution, it has been postulated that the enhanced conversion of cortisone to cortisol within omental adipose tissue plays an important role in the pathogenesis of central obesity (Bujalska *et al.* 2002, Tomlinson *et al.* 2002).

Cortisol generation after exogenous introduction of cortisone, however, is reduced with increasing body mass index (BMI), and a reduction in activity and expression of 11 β -HSD1 in OM preadipocytes derived from samples taken from obese volunteers, is seen. These data suggest that hepatic 11 β -HSD1 protects against the metabolic effects of simple obesity (Tomlinson *et al.* 2004). Nevertheless, central obesity is a characteristic feature of mice overexpressing 11 β -HSD1 within adipocytes (Masuzaki *et al.* 2001). The mechanisms underlying obesity are complex, but recent findings suggest a further role for chronic inflammation that may activate adipose tissue. Increased accumulation of CD68⁺ macrophages is seen in SC adipose depots in patients with high BMI and obese phenotypes, and increased expression of monocyte chemoattractant protein (MCP)-1, macrophage inflammatory protein (MIP)-1 α and MAC-1 is evident in rodent white adipose tissue after diet-induced obesity (Weisberg *et al.* 2003, Wellen & Hotamisligil 2003, Xu *et al.* 2003).

Orbital fat (OF) represents a highly specialised adipose tissue depot that occupies the space behind the eyeball (known as the orbit). OF fills most of the orbital cavity, and surrounds the globe, extraocular muscles, nerves and vessels providing support to these delicate structures and protecting against sight-threatening mechanical and inflammatory trauma (Wolfram-Gabel & Kahn 2002). Two types of orbital adipose tissue have been defined: anterior OF consisting chiefly of small adipocytes that are separated by thick conjunctival septa providing support to intraorbital structures, and posterior OF consisting of larger adipocytes separated by thin conjunctival septa enabling structural movements within the orbit (Bremond-Gignac *et al.* 2004). Over 60% of orbital disorders are inflammatory in origin (Rootman 2002). OF is the site for intense inflammation in several disease states of the orbit including local conditions, such as orbital cellulitis, orbital myositis, idiopathic orbital inflammatory disease, as well as systemic metabolic disorders (thyroid-associated ophthalmopathy) and generalised vasculitides (Wegener's granulomatosis). Orbital fibroblasts and other resident orbital cells (macrophages and dendritic cells) are known to participate in the initiation and perpetuation of many of these disease processes secreting mediators that are crucial to orchestrating orbital immune and inflammatory responses and

tissue matrix remodelling (Burnstine *et al.* 1998, Heufelder 2000). OF, like SC and OM depots, expresses GR (Heufelder *et al.* 1992, Burnstine *et al.* 1998, Muhlberg *et al.* 2000). Exogenous GCs are known to modulate the autocrine and paracrine interactions between OF and resident or infiltrating inflammatory cells (lymphocytes and macrophages), leading to the secretion of chemokines and cytokines (Burnstine *et al.* 1998, Valyasevi *et al.* 2001, Kumar & Bahn 2003, Chen *et al.* 2005), mediating OF proliferation (Heufelder & Bahn 1994) and production of matrix proteins (Kaback & Smith 1999, Han & Smith 2005). To date, no studies have characterised the role of 11 β -HSD1 and the pre-receptor regulation of endogenous glucocorticoids in OF. In this study, we describe features of the GC metabolic pathways in normal human OF depot and compare it with SC and OM depots.

Materials and Methods

Collection of human adipose tissue

Paired OM and SC fat was obtained from 50 non-obese (BMI < 25 kg/m²) patients undergoing elective abdominal surgery for non-malignant conditions. Surgically acquired normal orbital adipose tissue specimens (avoiding the anterior and nasal orbital adipose tissue) were obtained from 40 non-obese (BMI < 25 kg/m²) patients undergoing orbital or eyelid reconstructive procedures carried out at a regional tertiary referral centre (Birmingham and Midland Eye Centre, Birmingham, UK). Patients with orbital tumours, those on exogenous corticosteroids, or with underlying endocrine disease were excluded from the study. Each fat specimen was fixed in formalin, stored in RNAlater at -20 °C for RNA extraction or placed in a sterile universal container for immediate processing where sufficient starting tissue was available.

Characterisation of whole tissue adipocyte morphology

Formalin fixed paraffin embedded 5 μ m sections of OF ($n=17$), SC ($n=8$) and OM tissue ($n=8$) were stained with haematoxylin and eosin and photographed in triplicate using an Olympus-BH-2-RFC microscope (Olympus UK Ltd, Southall, Middx, UK) and MicroFireTM-S99808 camera (Indigo Scientific, Baldock, Herts, UK). The images were processed using the green channel to enhance edge prominence using Image-Pro Plus (v4.0, Media Cybernetics, Silver Spring, MD, USA). Software parameters were optimised in order to identify the visual edge of the cell membrane and generate an outline of the cell profiles for analysis. The scaling factor was calculated by photographing a micro-grid counting area of 500 \times 584 μ m capturing a minimum of 100 cells (the exact number varying according to the size of the adipocytes). Images were taken in triplicate for each specimen, i.e. approximately 300 adipocytes were evaluated for each histological section examined. The following descriptive variables were measured: cell diameter,

length of longest line joining two points and passing through centroid; roundness, $((\text{perimeter})^2/4\pi\text{area})$; radius ratio, largest radius/smallest radius; feret length, calliper longest length; cell area (μm^2); heterogeneity, fraction of pixels that deviate more than 10% from average intensity and margination, relative distribution of object intensity between centre and margin (larger value represents brighter centre and 0.33 represents homogeneity).

Immunohistochemistry

Immunohistochemical analyses were performed on 5 μm sections OF, SC and OM tissues mounted on charged slides.

CD68 immunohistochemistry Dewaxed and rehydrated slides of OF, SC and OM sections were blocked with hydrogen peroxide for 15 min before antigen retrieval using Tris/EDTA tris-buffered saline (TBS) buffer at pH 7.8 and a microwave at full power (900 W) for 15 min. All sections were allowed to cool to room temperature before being rinsed in water. All incubations and washes were performed using the Shandon sequenza. A biotin block was performed on the sections for 10 min using the Zymed avidin/biotin blocking kit (Invitrogen Ltd) before incubating with primary antibody (CD68 clone PG-M1 (Dako, Carpinteria, CA, USA) 1:100 dilution) in 10% (v/v) normal swine serum in TBS buffer for 60 min. Negative control sections were incubated in TBS buffer without primary antibody. The positive control sections used were human tonsil tissue. The sections were then washed in TBS buffer before incubating in Link reagent (Dakocytomation (yellow)) for 10 min. The sections were then rinsed again in TBS buffer (pH 7.8), incubated with streptavidin reagent from Dakocytomation (red) for 10 min, rinsed in TBS buffer (pH 7.8) before incubating with CheMate liquid 3,3'-diaminobenzidine (DAB; TBS buffer, 1:50) for approximately 10 min. Finally, sections were counterstained with Meyer's haematoxylin for 45 s and processed 100% (v/v) ethanol and xylene. CD68 positive cells were counted in triplicate by masked observers in three identical fields of view per slide for each adipose depot.

11 β -HSD immunohistochemistry Immunoperoxidase studies were performed on OF sections using antisera raised in sheep against human 11 β -HSD1 (amino acid residues 18–33) and 11 β -HSD2 (amino acid residues 137–160 and 334–358; The Binding Site, Birmingham, UK). Antibody dilutions were in the ratio of 1:100 for 11 β -HSD1 and 1:100 for 11 β -HSD2. Control sections included the omission of primary antibody and use of antibody pre-treated with the immunising peptides. Secondary antibodies comprised donkey anti-sheep peroxidase conjugate (1:100; Binding Site). Sections were visualised with the peroxidase substrate DAB. Human

liver and kidney sections were used as positive control tissues for 11 β -HSD1 and 11 β -HSD2 respectively.

Orbital preadipocyte isolation and primary cultures

Orbital ($n=12$), SC ($n=34$) and OM ($n=34$) preadipocytes were isolated using methodology previously reported (Bujalska *et al.* 2002). The OF protocol was adapted to maximise yield from the small sample size. Briefly, orbital adipose tissue biopsies were washed in PBS containing 50 000 U/l penicillin and 50 000 $\mu\text{g/l}$ streptomycin (Life Technologies, Inc.). The tissue was minced with sterile scissors in a Class II culture hood and digested with collagenase class I (2 mg/ml; Worthington Biochemical Corp., Reading, UK) in 1 \times Hanks' Balanced Salt Solution (Gibco Invitrogen Corporation) for 45 min at 37 °C. Samples were centrifuged at 500 g for 1 min, and the supernatant containing connective tissue debris, collagenase and lipid was removed leaving a pellet containing preadipocytes. The cellular pellet was washed with Dulbecco's modified Eagle's medium DMEM/Nutrient Mixture F-12 (Gibco) containing 15% (v/v) foetal calf serum (Gibco) and seeded on 48-well plates. Cells were left overnight to attach and all unattached cells (including red blood cells) were washed the following day with 1 \times Hanks' Balanced Salt Solution. Enzyme assays were performed in duplicate when preadipocytes were confluent at approximately 10 days of culture (see below). The identity of cells were confirmed by differentiating preadipocytes from one well of the 48-well plate into mature lipid containing adipocytes (>90% cultured cells) using chemically defined media.

11 β -HSD enzyme assays in cultured orbital preadipocytes

In confluent OF preadipocytes, the culture medium was replaced with serum-free medium 2 h before incubation. Dehydrogenase activity (cortisol to cortisone conversion) was assessed using 100 nM unlabelled cortisol (Sigma Chemical Co) diluted in serum-free medium and tracer amounts (1.5 nM) of [^3H]cortisol (specific activity 74.0 Ci/mmol; NEN, Boston, MA, USA) at 37 °C for 24 h. Conversion of cortisone to cortisol (oxo-reductase) was analysed by incubating cells with 100 nM cortisone and tracer amounts of [^3H]cortisone (50 000 c.p.m.) synthesised in-house (Stewart *et al.* 1994). After 24-h incubation, steroids were extracted from the medium with ten volumes of dichloromethane, separated by thin-layer chromatography with chloroform:ethanol (92:8) as a mobile phase and the fractional conversion of steroids was calculated after scanning analysis using a Bioscan 2000 radioimaging detector (Bioscan, Washington, DC, USA). Following enzyme assay, cell monolayers were lysed in 1 ml water for subsequent protein assays. Total protein in each well was determined using a standard protein assay reagent (Bio-Rad), and enzyme activities were expressed as pmol/h per mg. All assays were carried out in duplicate for each preadipocyte primary culture derived from individual patient samples.

Whole human adipose tissue RNA extraction and reverse transcription

Total RNA was extracted from whole OF ($n=8$), SC ($n=8$) and OM ($n=7$) tissue stored in RNAlater at -20°C using a single-step extraction method (Tri-Reagent; Sigma-Aldrich). RNA integrity was assessed by electrophoresis on 1% (w/v) agarose gels, and quantity was determined spectrophotometrically at OD₂₆₀. One microgram of total RNA was reverse transcribed in 50 μl total volume using TaqMan Reverse Transcription Reagents (Applied Biosystems, Warrington, Cheshire, UK) according to the manufacturer's guidelines. The reaction was terminated by heating the cDNA to 95°C for 5 min and stored at -20°C until required.

Real-time reverse transcriptase-PCR

11 β -HSD1, 11 β -HSD2, H6PDH, GR α , FABP4, G3PDH and CD68 mRNA levels were measured using real-time PCR using an ABI 7500 system (Perkin-Elmer, Biosystems, Warrington, UK). PCR was performed in 25 μl reactions on 96-well plates in reaction buffer containing TaqMan universal PCR mastermix (Applied Biosystems), 3 mM Mn(Oac)₂, 200 μM deoxy-NTPs, 1.25 U AmpliTaq Gold polymerase (Perkin-Elmer), 1.25 U AmpErase UNG (Perkin-Elmer), 900 nmol primers, 100–200 nmol TaqMan probe and 25–50 ng cDNA. All reactions were multiplexed with primers specific for 18S rRNA (provided as a pre-optimised mix; Perkin-Elmer, Beaconsfield, Bucks, UK) as an internal reference and were carried out in duplicate. Data were obtained as Ct values (the cycle number at which logarithmic PCR plots cross a calculated threshold line) and used to determine dCt values (dCt = Ct of the target gene minus Ct of the internal reference, 18S). All target gene probes were labelled with the fluorescent label 6-carboxyfluorescein, and the housekeeping gene was labelled with the fluorescent label VIC. Reactions were as follows: 50°C for 2 min, 95°C for 10 min, and then 40 cycles of 95°C for 15 s and 60°C for 1 min. All expression data are presented as fold-change when compared with OF using the equation $2^{-\Delta\Delta\text{Ct}}$ ($\Delta\Delta\text{Ct}$ is the difference between dCt values in SC or OM minus dCt in OF). Sequences of oligonucleotide primers and probes were as follows: (i) 11 β -HSD1 antisense primer 5'-AGGAAAGCTCATGGGAGGACTAG-3', sense primer 5'-ATGGTGAA-TATCATCATGAAAAAGATTC-3', probe 5'-CATGCT-CATTCTCAACCACATCACCAACA-3'; (ii) GR α antisense primer 5'-GCGATGGTCTCAGAAACCAAAC-3', sense primer 5'-GAGATTACAGAGGAAGTTATCCTCTGC-3', probe 5'-TGCAGTGAAGGTTGCTGAGGCTCTGA-3'; (iii) H6PDH antisense primer 5'-CAGGTGTCTAGTGCACATTGAC-3', sense primer 5'-GTAGCCCACTCTCTC-GTCCAA-3', probe 5'-AAGGCACGCCCTCCCAGCG-3'; (iv) G3PDH antisense primer 5'-CCATCAGTTCATCGG-CAAGAT-3', sense primer 5'-TCGTCTACCCCCTTAA-TAAGAGATATG-3', probe 5'-AGGCCATCTGAAGGC

AAACGCC-3'. Primers and probes for CD68 and FABP4 were from Expression Assays Applied Biosystems.

Ethics and statistical methods

The collection of human tissues for *in vitro* investigation was approved by the local ethics committee. Informed consent was obtained from all patient volunteers.

Statistical analysis was performed using the software package SPSS for Windows Version 11.5.1 (SPSS, Inc., Chicago, IL, USA, 2002), Minitab Release 14.12 (Minitab Ltd, Coventry, UK) and Prism for Windows Version 4.03c (GraphPad Software, Inc., San Diego, CA, USA). Histological characterisation comparisons between the three fat depots were made using a nested general linear model regression analysis for replicated measures of data (to minimise loss of data by averaging individual readings), followed by integrated pairwise comparisons. Data are presented as means and 95% confidence intervals. Enzyme assay data are presented as means with standard errors of the separate experiments. Real-time PCR analyses were performed with dCt values to exclude potential bias due to averaging and transforming data through the equation $2^{-\Delta\Delta\text{Ct}}$ to give a fold-change in gene expression. Statistical analyses were performed using a non-parametric Mann-Whitney test for non-normal data, or *t*-test/ANOVA for normally distributed data. *P* values <0.05 were accepted as statistically significant.

Results*Characterisation of whole tissue adipocyte morphology*

Adipocyte diameter, perimeter and area (Fig. 1A–C respectively) were identified as optimal parameters differentiating adipocyte morphology from the different depots. Mean cell diameter, perimeter, adipocyte area and feret length defined similar trends (OF $<$ OM $<$ SC; $P < 0.001$) across the three adipocyte depots. OF adipocytes were the smallest adipocyte with mean cell diameter (50.7 μm , 95%CI 49.8–51.6), shortest perimeter (149 μm , 95%CI 146.7–152.7) and smallest area (2470 μm^2 , 95% CI 2382–2560). SC adipocytes were the largest according to each of these parameters. Radius ratio, cell roundness and margination followed similar trends to each other across the groups but these trends were not significant (OF $>$ OM $<$ SC; $P = \text{NS}$, data not shown), and cell count per area of interest did not differentiate significantly between depots. Representative haematoxylin and eosin-stained photomicrographs detailing histomorphometric characterisation of the three adipose depots are shown in Fig. 1D–F.

Evaluation of CD68⁺ protein and mRNA expression in human adipose depots

The largest resident CD68⁺ population was found in the OF depot (median 7.6 (range 3.8–19.6) cell counts/unit area)




Figure 1 Adipocyte morphology. Diameter of the individual adipocyte (A), perimeter cell diameter (B) and area (C), were identified as optimal characteristics defining adipocyte morphology. Smallest adipocytes were found in the orbital fat (OF) depot (D), while SC adipocytes were the largest cell (E). Omental (OM) adipocytes were statistically larger than OF but smaller than SC (F) (OF, $n=17$; SC, $n=8$; OM, $n=8$; results are expressed as mean and 95% confidence intervals; $*P<0.001$).

when compared with SC (median 0.7 (range 0.1–6.5) cell counts/unit area, $P<0.001$) and OM (median 2.9 (range 0.6–5.2) cell counts/unit area, $P<0.001$; Fig. 2A). These findings were endorsed by a similar trend in CD68 mRNA expression for each depot (Fig. 2B (OF=1; SC=0.21; $P<0.05$; OM 0.38 < 0.05 (fold-change when compared with OF))). Representative photomicrographs showing CD68⁺ macrophages in each whole adipose tissue depot (OF, SC and OM) are shown in Fig. 2C–E.

Characterisation of 11 β -HSD1 expression and activity in orbital fat

Immunoreactivity was seen for 11 β -HSD1 antibody in OF sections (Fig. 3A) and around the central vein of human liver (positive control; Fig. 3C). Minimal staining was detected with 11 β -HSD1 pre-adsorbed antibody (data not shown). No staining was seen with 11 β -HSD2 antibody in OF tissue (Fig. 3B), whereas good staining was visualised in human distal collecting duct epithelium in the kidney (positive control, Fig. 3D).

Primary stromal cultures of OF preadipocytes, confirmed 11 β -HSD1 oxo-reductase activity (0.69 ± 0.5 pmol/h per mg (mean \pm S.E.M.)) when incubated for 24 h with 100 nM cortisone and minimal dehydrogenase activity (0.11 ± 0.08 pmol/h per mg (mean \pm S.E.M.); $P<0.05$; Fig. 4A–C). Oxo-reductase activity was greatest in OM preadipocytes when compared with SC (43.7 ± 5.2 vs 17.6 ± 3.6 pmol/mg per h (mean \pm S.E.M.); $P<0.001$) and OF preadipocytes (OF versus SC; $P<0.01$; OF versus OM; $P<0.001$; Fig. 4D).

Comparison of 11 β -HSD1 and glucocorticoid-related target gene expression in whole adipose tissue depots

11 β -HSD1 mRNA was expressed within OF tissue, although quantitative real-time PCR analyses across the three whole adipose tissue depots revealed lowest levels of expression in OF when compared with SC and OM adipose depots (OF=1; SC=8.1, $P<0.001$; OM=10 < 0.001 (fold-change when compared with OF); Fig. 5A). Nevertheless, these data reflect our 11 β -HSD1 oxo-reductase activity data detailed earlier.

Whole tissue analyses indicated the highest expression of GR α mRNA in OF when compared with SC and OM adipose tissue (Fig. 5A). Conversely, H6PDH mRNA (NADPH provider for 11 β -HSD1 oxo-reductase activity; Fig. 5A) and markers of late adipocyte differentiation (FABP4, G3PDH) were significantly lower in OF (Fig. 5B).

Discussion

In this paper, we have described a novel reproducible automated method for the histomorphometric analysis of adipose tissue from differing human depots. This has been performed in the context of gene expression profiling and has identified important depot specific differences in adipocyte biology and GC metabolism.

In comparison with OM and SC depots, orbital adipocytes were significantly smaller, expressed lower levels of adipocyte differentiation markers (FABP4 and G3PDH), but revealed a higher CD68⁺ cellular infiltrate. Examination of the key players of the GC metabolic pathways, confirmed that all




Figure 2 Immunohistochemistry using human anti-CD68 for macrophages confirmed a large resident population in the OF depot when compared with SC and OM (A). Results are expressed as median and full range of counts. These findings were endorsed by real-time RT-PCR CD68 mRNA expression in each of the adipose depots (B). Results are expressed as fold-change when compared with orbital fat; dCt are shown in table below (* $P < 0.05$; † $P < 0.001$). Representative sections of OF, SC and OM adipose depots with identified macrophages (M ϕ) are shown in C, D and E respectively.

three depots expressed GR α mRNA in keeping with earlier published studies (Heufelder *et al.* 1992, Burnstine *et al.* 1998, Muhlberg *et al.* 2000, Boullu-Ciocca *et al.* 2003), but expression in the OF depot was significantly higher. In addition, our data has defined 11 β -HSD1 oxo-reductase activity and mRNA expression in OF together with its regulatory enzyme, H6PDH, although these were at a lower level than that seen in SC and OM tissues.

These observations highlight the importance of adipose tissue, depot-specific analyses. Adipose tissue is regarded as an 'endocrine organ' and depot specificity of gene expression between SC and OM depots is well described (Montague *et al.* 1997). Furthermore, these changes are functionally important with differing propensities for adipocyte differentiation, proliferation, apoptosis and lipid metabolism between these depots (Vikman *et al.* 1995, Niesler *et al.* 1998). Our data show that orbital adipose tissue is clearly distinct from both SC and OM depots. The fundamental explanation for these differences lies, in addition to genetic and environmental factors, in the embryological origins of adipose connective tissue. Whereas SC and OM 'peripheral' adipose depots and fibroblasts are derived from mesoderm, orbital

fibroblasts originate from neuroectoderm (Smith 2004). Studies have shown that orbital and peripheral fibroblasts have both distinguishing morphological phenotypic features (Smith *et al.* 1995), but also at a molecular level, orbital fibroblasts have distinct defining characteristics, including expression of surface receptors, cellular proteins, responses to cytokines and the ability to secrete and remodel matrix proteins (Prabhakar *et al.* 2003, Ajan & Weetman 2004, Smith 2004). As orbital fibroblasts are neural crest in origin, they possess considerable plasticity – under appropriate *in vitro* conditions, almost 50% may be differentiated into lipid containing adipocytes (Smith *et al.* 2002). Hence, it has been proposed that the unique nature of the orbital molecular environment accounts for its underlying susceptibility to autoimmune diseases (Kaminski *et al.* 2003).

The impact of the macrophage infiltrate in orbital adipose tissue may also be important particularly as cytokines have potent effects upon adipocyte biology. Recent studies have shown that chronic inflammation activates adipose tissue in simple obesity defined by an increased accumulation of CD68⁺ macrophages, increased adipocyte size and increased TNF α in patients with high BMIs (Weisberg *et al.* 2003,




Figure 3 11 β -HSD immunohistochemistry confirmed 11 β -HSD1 expression in human orbital fat (A) and not 11 β -HSD2 (B). Human liver and kidney were used as positive control tissues for 11 β -HSD1 (C) and 11 β -HSD2 (D). No staining was seen in the negative control sections ($n=5$).




Figure 4 Primary cultures of OF preadipocytes confirmed 11 β -HSD1 oxo-reductase activity and minimal dehydrogenase activity when incubated for 24 h with 100 nM cortisone and cortisol respectively ($n=12$) (A). Representative radiochromatograph traces are shown for dehydrogenase (B) and oxo-reductase (C) activity. (D) Oxo-reductase activity was greatest in OM preadipocytes followed by SC and OF adipocytes (* $P<0.05$; † $P<0.01$; ‡ $P<0.001$).




Figure 5 Whole adipose tissue real time RT-PCR analyses showed significantly higher expression of 11 β -HSD1 and H6PDH mRNA in SC and OM when compared with OF. Conversely, highest expression of GR α mRNA in OF when compared with SC and OM is seen (A). FABP4, G3PDH mRNA expression (markers of late adipocyte differentiation) were significantly lower in OF (B). Results are expressed as fold-change when compared with orbital fat; dCt are shown in tables below (* $P < 0.05$; $^{\dagger}P < 0.001$).

Wellen & Hotamisligil 2003, Neels & Olefsky 2006, Pietiläinen *et al.* 2006). There is considerable heterogeneity in monocyte-derived lineages (e.g. macrophages) and the different monocyte subsets reflect the developmental stages with distinct physiological roles, such as recruitment into inflammatory lesions or entry to normal tissues (Gordon & Taylor 2005). It is thought that with the expansion of adipose tissue during weight gain, induction of a range of signalling pathways activates the adipose vascular fraction allowing monocytes to extravasate through the endothelial cell layer into the adipose tissue where they differentiate into macrophages localising predominantly around dead adipocytes (Neels & Olefsky 2006). Crosstalk between adipocytes, macrophages and endothelial cells aggravates the inflammatory state, resulting in increased secretion of pro-inflammatory cytokines/chemokines, adipokines and angiogenic factors. Furthermore, the interaction of cytokines and GC metabolism may be an additional contributing factor. TNF α , IL-6 and IL-1 β are potent inducers of 11 β -HSD1 expression and activity in adipose tissue (Tomlinson *et al.* 2001). Macrophage-derived cytokine production *in vivo* may serve to augment cortisol generation through induction of 11 β -HSD1. TNF α specifically, in addition to modulating 11 β -HSD1 activity, inhibits adipocyte differentiation, proliferation and promotes apoptosis and de-differentiation (Petruschke & Hauner 1993, Prins *et al.* 1997). This process may be of considerable relevance to the orbital fat microenvironment. The orbital fat samples analysed in this study were all from healthy controls undergoing elective orbital or lid surgery for non-inflammatory disease states, and despite this, normal OF harboured a large CD68 $^{+}$ population

and adipocytes were less differentiated when compared with SC and OM. These data reflect an orbital molecular environment that has an increased potential and plasticity that is vital to combat inflammatory, immune-mediated or infective insult.

The role of GCs in the regulation of adipocyte biology is exemplified by patients with Cushing's syndrome who develop florid, but reversible central obesity. GCs are essential for adipocyte differentiation (Hauner *et al.* 1987), inhibit omental preadipocyte proliferation (Tomlinson *et al.* 2002), but interestingly, promote proliferation in the SC depot (Bader *et al.* 2002). 11 β -HSD1 overexpression in adipose tissues in rodent models induces obesity and the metabolic syndrome (Masuzaki *et al.* 2001, 2003) and selective 11 β -HSD1 inhibitors have been suggested as a novel therapeutic strategy (Alberts *et al.* 2002, 2003, Hermanowski-Vosatka *et al.* 2005). The identification of 11 β -HSD1 in human orbital adipose tissue is an important finding and raises the possibility that the local generation of GC within this depot may be functionally important. Although levels were lower than those observed in SC and OM depots, the increased expression of GR α is likely to compensate for this. Low levels of oxo-reductase activity have been documented in other tissue depots, notably bone (Cooper *et al.* 2002). In this tissue, low basal levels are believed to be physiologically significant, as induction of oxo-reduction activity increasing GC bioavailability is seen when osteoblasts are challenged with pro-inflammatory cytokines (Cooper *et al.* 2001). Critically, unlike the SC and OM depots, the orbital fat depot lies within a confined space surrounding tissues susceptible to compression. Therefore, rapid expansion,

perhaps as a consequence of enhanced GC autocrine generation through 11 β -HSD1, may have important sight-threatening clinical implications. The possibility of dysregulation of 11 β -HSD1 in disease states (particularly those with underlying inflammatory and immune-mediated aetiologies) requires further investigation, and a detailed characterisation of expression and functional importance in these conditions is now warranted.

To conclude, orbital adipose tissue is a biologically distinct adipose tissue depot. Adipocytes are smaller and less differentiated than SC and OM. The identification of 11 β -HSD1 and H6PDH expression and activity within the depot, together with an abundant GR α and a large CD68⁺ population define a microenvironment that has the potential to respond to sight-threatening orbital inflammatory disease. Our data raise the exciting possibility that the autocrine generation of cortisol may be functionally important in regulating orbital adipocyte biology and that this may be amenable to therapeutic modulation.

Acknowledgements

We would like to thank Charles Shaikh and Susan Hughes for their technical support in the laboratory, and Timothy Marshall for his advice on the statistical analyses. *Grant Support:* Wellcome Trust Research, UK; SWBH NHS Trust R&D Major Award 2006. The authors have no proprietary interest in the products described in this article. The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

References

Ajjan RA & Weetman AP 2004 New understanding of the role of cytokines in the pathogenesis of Graves' ophthalmopathy. *Journal of Endocrinological Investigation* **27** 237–245.

Alberts P, Engblom L, Edling N, Forsgren M, Klingström G, Larsson C, Rönquist-Nii Y, Öhman B & Abrahamson L 2002 Selective inhibition of 11 β -hydroxysteroid dehydrogenase type 1 decreases blood glucose concentrations in hyperglycaemic mice. *Diabetologia* **45** 1528–1532.

Alberts P, Nilsson C, Selen G, Engblom L, Edling N, Norling S, Klingstrom G, Larsson C, Forsgren M, Ashkzari M *et al.* 2003 Selective inhibition of 11 β -hydroxysteroid dehydrogenase type 1 improves hepatic insulin sensitivity in hyperglycaemic mice strains. *Endocrinology* **144** 4755–4762.

Bader T, Zoumakis E, Friedberg M, Hiroi N, Chrousos G & Hochberg Z 2002 Human adipose tissue under *in vitro* inhibition of 11beta-hydroxysteroid dehydrogenase type 1: differentiation and metabolism changes. *Hormone and Metabolic Research* **34** 752–757.

Boullu-Ciocca S, Paulmyer-Lacroix O, Fina F, Ouafik LH, Alessi M-C, Oliver C & Grino M 2003 Expression of the mRNAs coding for the glucocorticoid receptor isoforms in obesity. *Obesity Research* **11** 925–929.

Bremond-Gignac D, Copin H, Cussenot O, Lassau J-P & Henin D 2004 Anatomical, histological and mesoscopic study of the adipose tissue of the orbit. *Surgical and Radiologic Anatomy* **26** 297–302.

Bujalska IJ, Kumar S & Stewart PM 1997 Does central obesity reflect 'Cushing's disease of the omentum'? *Lancet* **349** 1210–1213.

Bujalska IJ, Kumar S, Hewison M & Stewart PM 1999 Differentiation of adipose stromal cells: the roles of glucocorticoids and 11{beta}-hydroxysteroid dehydrogenase. *Endocrinology* **140** 3188–3196.

Bujalska IJ, Walker EA, Hewison M & Stewart PM 2002 A switch in dehydrogenase to reductase activity of 11 β -hydroxysteroid dehydrogenase type 1 upon differentiation of human omental adipose stromal cells. *Journal of Clinical Endocrinology and Metabolism* **87** 1205–1210.

Bujalska IJ, Draper N, Michialidou Z, Tomlinson JW, White PC, Chapman KE, Walker EA & Stewart PM 2005 Hexose-6-phosphate dehydrogenase confers oxo-reductase activity upon 11 β -hydroxysteroid dehydrogenase type 1. *Journal of Molecular Endocrinology* **35** 675–684.

Burnstine MA, Elner SG & Elner VM 1998 Orbital fibroblast chemokine modulation: effects of dexamethasone and cyclosporin A. *British Journal of Ophthalmology* **82** 318–322.

Chen B, Tsui S & Smith TJ 2005 IL-1{beta} induces IL-6 expression in human orbital fibroblasts: identification of an anatomic-site specific phenotypic attribute relevant to thyroid-associated ophthalmopathy. *Journal of Immunology* **175** 1310–1319.

Cooper MS, Walker EA, Bland R, Fraser WD, Hewison M & Stewart PM 2000 Expression and functional consequences of 11 β -hydroxysteroid dehydrogenase activity in human bone. *Bone* **27** 375–381.

Cooper MS, Bujalska I, Rabbitt EA, Walker EA, Bland R, Sheppard MC, Hewison M & Stewart PM 2001 Modulation of 11 β -hydroxysteroid dehydrogenase isozymes by proinflammatory cytokines in osteoblasts: an autocrine switch from glucocorticoid inactivation to activation. *Journal of Bone and Mineral Research* **16** 1037–1044.

Cooper MS, Rabbitt EH, Goddard PE, Bartlett WA, Hewison M & Stewart PM 2002 Osteoblastic 11 β -hydroxysteroid dehydrogenase type 1 activity increases with age and glucocorticoid exposure. *Journal of Bone and Mineral Research* **17** 979–986.

Draper N, Walker EA, Bujalska IJ, Tomlinson JW, Chalder SM, Arlt W, Lavery GG, Bedendo O, Ray DW, Laing I *et al.* 2003 Mutations in the genes encoding 11beta-hydroxysteroid dehydrogenase type 1 and hexose-6-phosphate dehydrogenase interact to cause cortisone reductase deficiency. *Nature Genetics* **34** 434–439.

Gordon S & Taylor PR 2005 Monocyte and macrophage heterogeneity. *Nature Reviews, Immunology* **5** 953–964.

Han R & Smith TJ 2005 Induction by IL-1 β of tissue inhibitor of metalloproteinase-1 in human orbital fibroblasts: modulation of gene promoter activity by IL-4 and IFN- γ . *Journal of Immunology* **174** 3072–3079.

Hauner H, Schmid P & Pfeiffer EF 1987 Glucocorticoids and insulin promote the differentiation of human adipocyte precursor cells into fat cells. *Journal of Clinical Endocrinology and Metabolism* **64** 832–835.

Hermanowski-Vosatka A, Balkovec JM, Cheng K, Chen HY, Hernandez M, Koo GC, Le Grand CB, Li Z, Metzger JM, Mundt SS *et al.* 2005 11{beta}-HSD1 inhibition ameliorates metabolic syndrome and prevents progression of atherosclerosis in mice. *Journal of Experimental Medicine* **202** 517–527.

Heufelder AE 2000 Pathogenesis of ophthalmopathy in autoimmune thyroid eye disease. *Reviews in Endocrine and Metabolic Disorders* **1** 87–95.

Heufelder AE & Bahn RS 1994 Modulation of Graves' orbital fibroblast proliferation by cytokines and glucocorticoid receptor agonists. *Investigative Ophthalmology and Visual Science* **35** 120–127.

Heufelder AE, Bahn RS & Smith TJ 1992 Regulation by glucocorticoids of interferon gamma-induced HLA-DR antigen expression in cultured human orbital fibroblasts. *Clinical Endocrinology* **37** 59–63.

Kaback LA & Smith TJ 1999 Expression of hyaluronan synthase messenger ribonucleic acids and their induction by interleukin-1{beta} in human orbital fibroblasts: potential insight into the molecular pathogenesis of thyroid-associated ophthalmopathy. *Journal of Clinical Endocrinology and Metabolism* **84** 4079–4084.

Kaminski HJ, Li Z, Richmonds C, Ruff RL & Kusner L 2003 Susceptibility of ocular tissues to autoimmune diseases. *Annals of the New York Academy of Sciences* **998** 362–374.

Kotelvtsev Y, Holmes MC, Burchell A, Houston PM, Schmol D, Jamieson P, Best R, Brown R, Edwards CRW, Seckl JR *et al.* 1997 11 β -Hydroxysteroid dehydrogenase type 1 knockout mice show attenuated glucocorticoid-inducible responses and resist hyperglycaemia on obesity and stress. *PNAS* **94** 14924–14929.

Kumar S & Bahn RS 2003 Relative overexpression of macrophage-derived cytokines in orbital adipose tissue from patients with Graves' ophthalmopathy. *Journal of Clinical Endocrinology and Metabolism* **88** 4246–4250.

- Masuzaki H, Paterson J, Shinyama H, Morton NM, Mullins JJ, Seckl JR & Flier JS 2001 A transgenic model of visceral obesity and the metabolic syndrome. *Science* **294** 2166–2170.
- Masuzaki H, Yamamoto H, Kenyon CJ, Elmquist JK, Morton NM, Paterson JM, Shinyama H, Sharp MGF, Fleming S, Mullins JJ *et al.* 2003 Transgenic amplification of glucocorticoid action in adipose tissue causes high blood pressure in mice. *Journal of Clinical Investigation* **112** 83–90.
- Montague CT, Prins JB, Sanders L, Digby JE & O'Rahilly S 1997 Depot- and sex-specific differences in human leptin mRNA expression: implications for the control of regional fat distribution. *Diabetes* **46** 342–347.
- Muhlberg T, Joba W, Spitzweg C, Schworm H-D, Heberling H-J & Heufelder AE 2000 Interleukin-1 receptor antagonist ribonucleic acid and protein expression by cultured Graves' and normal orbital fibroblasts is differentially modulated by dexamethasone and irradiation. *Journal of Clinical Endocrinology and Metabolism* **85** 734–742.
- Neels JG & Olefsky JM 2006 Inflamed fat: what starts the fire? *Journal of Clinical Investigation* **116** 33–35.
- Niesler CU, Siddle K & Prins JB 1998 Human preadipocytes display a depot-specific susceptibility to apoptosis. *Diabetes* **47** 1365–1368.
- Onyimba CU, Vijapurapu N, Curnow SJ, Khosla P, Stewart PM, Murray PI, Walker EA & Rauz S 2006 Characterisation of the prereceptor regulation of glucocorticoids in the anterior segment of the rabbit eye. *Journal of Endocrinology* **190** 483–493.
- Petruschke T & Hauner H 1993 Tumor necrosis factor- α prevents the differentiation of human adipocyte precursor cells and causes delipidation of newly developed fat cells. *Journal of Clinical Endocrinology and Metabolism* **76** 742–747.
- Pietiläinen KH, Kannisto K, Korshennikova E, Rissanen A, Kaprio J, Ehrenborg E, Hamsten A & Yki-Järvinen H 2006 Acquired obesity increases CD68 and TNF- α and decreases adiponectin gene expression in adipose tissue – a study in monozygotic twins. *Journal of Clinical Endocrinology and Metabolism* **91** 2776–2781.
- Prabhakar BS, Bahn RS & Smith TJ 2003 Current perspective on the pathogenesis of Graves' disease and ophthalmopathy. *Endocrine Reviews* **24** 802–835.
- Prins JB, Niesler CU, Winterford CM, Bright NA, Siddle K, O'Rahilly S, Walker NI & Cameron DP 1997 Tumor necrosis factor- α induces apoptosis of human adipose cells. *Diabetes* **46** 1939–1944.
- Rauz S, Cheung CMG, Wood PJ, Coca-Prados M, Walker EA, Murray PI & Stewart PM 2003 Inhibition of 11 β -hydroxysteroid dehydrogenase type 1 lowers intraocular pressure in patients with ocular hypertension. *Quarterly Journal of Medicine* **96** 481–490.
- Rootman J 2002 Pathophysiologic patterns of orbital disease. In *Diseases of the Orbit: a Multidisciplinary Approach*, 2nd edn, Ed. J Rootman. Philadelphia, PA: Lippincott Williams & Wilkins.
- Smith TJ 2004 Novel aspects of orbital fibroblast pathology. *Journal of Endocrinological Investigation* **27** 246–253.
- Smith TJ, Sempowski GD, Wang HS, Del Vecchio PJ, Lippe SD & Phipps RP 1995 Evidence for cellular heterogeneity in primary cultures of human orbital fibroblasts. *Journal of Clinical Endocrinology and Metabolism* **80** 2620–2625.
- Smith TJ, Koumas L, Gagnon A, Bell A, Sempowski GD, Phipps RP & Sorisky A 2002 Orbital fibroblast heterogeneity may determine the clinical presentation of thyroid-associated ophthalmopathy. *Journal of Clinical Endocrinology and Metabolism* **87** 385.
- Stewart PM, Murry BA & Mason JI 1994 Human kidney 11 β -hydroxysteroid dehydrogenase is a high affinity nicotinamide adenine dinucleotide-dependent enzyme and differs from the cloned type I isoform. *Journal of Clinical Endocrinology and Metabolism* **79** 480–484.
- Thieringer R, Grande CBL, Carbin L, Cai T, Wong B, Wright SD & Hermanowski-Vosatka A 2001 11 β -Hydroxysteroid dehydrogenase type 1 is induced in human monocytes upon differentiation to macrophages. *Journal of Immunology* **167** 30–35.
- Tomlinson JW, Bujalska I, Stewart PM & Cooper MS 2000 The role of 11 β -hydroxysteroid dehydrogenase in central obesity and osteoporosis. *Endocrine Research* **26** 711–722.
- Tomlinson JW, Moore J, Cooper MS, Bujalska I, Shahmanesh M, Burt C, Strain A, Hewison M & Stewart PM 2001 Regulation of expression of 11 β -hydroxysteroid dehydrogenase type 1 in adipose tissue: tissue specific induction by cytokines. *Endocrinology* **142** 1982–1989.
- Tomlinson JW, Sinha B, Bujalska I, Hewison M & Stewart PM 2002 Expression of 11 β -hydroxysteroid dehydrogenase type 1 in adipose tissue is not increased in human obesity. *Journal of Clinical Endocrinology and Metabolism* **87** 5630–5635.
- Tomlinson JW, Walker EA, Bujalska IJ, Draper N, Lavery GG, Cooper MS, Hewison M & Stewart PM 2004 11 β -Hydroxysteroid dehydrogenase type 1: a tissue-specific regulator of glucocorticoid response. *Endocrine Reviews* **25** 831–866.
- Valyasevi RW, Jyonougi SC, Dutton CM, Munsakul N & Bahn RS 2001 Effect of tumour necrosis factor- α , interferon- γ , and expression of thymotropin receptor in human orbital preadipocyte fibroblasts. *Journal of Clinical Endocrinology and Metabolism* **86** 903–908.
- Vikman HL, Hreniuk SP, Kauffman GF, LaNoue KF, Martin LF & Ohisalo JJ 1995 Cyclic AMP and glycerol concentrations in freeze-clamped human omental and subcutaneous adipose tissues. *International Journal of Obesity and Related Metabolic Disorders* **19** 388–391.
- Weisberg SP, McCann DM, Desai M, Rosenbaum M, Leibel RL & Ferrante AW 2003 Obesity is associated with macrophage accumulation in adipose tissue. *Journal of Clinical Investigation* **112** 1796–1808.
- Wellen KE & Hotamisligil GS 2003 Obesity-induced inflammatory changes in adipose tissue. *Journal of Clinical Investigation* **112** 1785–1788.
- Wolfram-Gabel R & Kahn JL 2002 Adipose body of the orbit. *Clinical Anatomy* **15** 186–192.
- Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS, Tartaglia LA *et al.* 2003 Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *Journal of Clinical Investigation* **112** 1821–1830.
- Zhang TY, Ding X & Daynes RA 2005 The expression of 11 β -hydroxysteroid dehydrogenase type I by lymphocytes provides a novel means for intracrine regulation of glucocorticoid activities. *Journal of Immunology* **174** 879–889.

Received in final form 3 November 2006

Accepted 6 November 2006

Made available online as an Accepted Preprint

13 November 2006