



UNIVERSITY OF LEEDS

This is a repository copy of *Applications of Laplace-Carleson embeddings to admissibility and controllability*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/81897/>

Version: Accepted Version

Article:

Jacob, B, Partington, JR orcid.org/0000-0002-6738-3216 and Pott, S (2014) Applications of Laplace-Carleson embeddings to admissibility and controllability. *SIAM Journal on Control and Optimization*, 52 (2). pp. 1299-1313. ISSN 0363-0129

<https://doi.org/10.1137/120894750>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Applications of Laplace–Carleson embeddings to admissibility and controllability

Birgit Jacob* Jonathan R. Partington[†] Sandra Pott[‡]

January 31, 2014

Abstract

It is shown how results on Carleson embeddings induced by the Laplace transform can be used to derive new and more general results concerning the weighted (infinite-time) admissibility of control and observation operators for linear semigroup systems with q -Riesz bases of eigenvectors. As an example, the heat equation is considered. Next, a new Carleson embedding result is proved, which gives further results on weighted admissibility for analytic semigroups. Finally, controllability by smoother inputs is characterised by means of a new result about weighted interpolation.

Keywords. Semigroup system, controllability, admissibility, Hardy space, weighted Bergman space, Interpolation, Carleson measure.

2000 Subject Classification. 30D55, 30E05, 47A57, 47D06, 93B05, 93B28.

1 Introduction

The main purpose of this note is to show how recent results on Carleson embeddings, mostly derived in [14], may be applied to the theory of well-posed linear systems. We shall discuss concepts such as admissibility, controllability and observability, for which some standard references are the books [23, 20] and the survey [9]. Our basic tools will be the theory of a class of function spaces known as Zen spaces, which include the standard Hardy and Bergman spaces.

The structure of the paper is as follows. In Section 2, we review the basics of the theory of admissibility for diagonal semigroups, introduce the key embedding results for Zen spaces, and derive new admissibility results in this context. Section 3 treats the case of analytic semigroups (corresponding to measures supported in a sector), where we are able to move away from the Hilbertian

*Fachbereich C - Mathematik und Naturwissenschaften, Bergische Universität Wuppertal, Gaußstraße 20, 42097 Wuppertal, Germany, jacob@math.uni-wuppertal.de

[†]School of Mathematics, University of Leeds, Leeds LS2 9JT, U.K. J.R.Partington@leeds.ac.uk

[‡]Centre for Mathematics, Faculty of Science, Lund University, Sölvegatan 18, 22100 Lund, Sweden, Sandra.Pott@math.lu.se

(L^2) context and study general L^p spaces. We also derive a new embedding theorem and apply it to weighted admissibility. Finally, in Section 4 we consider notions of controllability, which are linked with interpolation questions.

Although the norms most commonly used in physical applications are unweighted L^1 and L^2 norms, it is also useful to consider Sobolev spaces (control by smoother inputs) and weighted L^p spaces (when the size of the input may depend on time), see [4, 24] and the references therein.

2 Admissibility for diagonal semigroups

Let A be the infinitesimal generator of a C_0 -semigroup $(T(t))_{t \geq 0}$ defined on a Hilbert space H , and consider the system

$$\frac{dx(t)}{dt} = Ax(t) + Bu(t), \quad x(0) = x_0, \quad t \geq 0, \quad (1)$$

where $u(t) \in \mathbb{C}$ is the input at time t , and $B : \mathbb{C} \rightarrow D(A^*)'$ is the control operator. Here $D(A^*)'$ is the completion of H with respect to the norm

$$\|x\|_{D(A^*)'} = \|(\beta - A)^{-1}x\|_H,$$

for any $\beta \in \rho(A)$. To ensure that the state $x(t)$ lies in H it is sufficient that $B \in \mathcal{L}(\mathbb{C}, D(A^*)')$, that $\int_0^\infty T(t)Bu(t) dt$ lies in H for each input $u \in L^2(0, \infty)$, and

$$\left\| \int_0^\infty T(t)Bu(t) dt \right\|_H \leq m_0 \|u\|_{L^2(0, \infty)}$$

for some $m_0 > 0$ (the *infinite-time admissibility* condition for B). We note that the C_0 -semigroup $(T(t))_{t \geq 0}$ has an extension to $D(A^*)'$.

Dually, we may consider the system

$$\frac{dx(t)}{dt} = Ax(t), \quad y(t) = Cx(t),$$

with $x(0) = x_0$, say. Here $C : D(A) \rightarrow \mathbb{C}$ is an A -bounded observation operator mapping; i.e., for some $m_1, m_2 > 0$,

$$\|Cz\| \leq m_1 \|z\| + m_2 \|Az\|.$$

C is *infinite-time admissible*, if there is an $m_0 > 0$ such that $y \in L^2(0, \infty)$ and $\|y\|_2 \leq m_0 \|x_0\|$ for all $x_0 \in D(A)$. Note that $y(t) = CT(t)x_0$ for every $x_0 \in D(A)$.

In this paper we shall refer simply to *admissibility* throughout.

The duality here is that B is an admissible control operator for $(T(t))_{t \geq 0}$ if and only if B^* is an admissible observation operator for the dual semigroup $(T(t)^*)_{t \geq 0}$.

Admissibility is an important concept in the theory of well-posed systems, and we refer to the survey [9] and the book [23] for the basic background to the subject. For diagonal semigroups, admissibility is linked with the theory of Carleson measures as in [8, 25]; namely, supposing that A has a Riesz basis of eigenvectors, with eigenvalues (λ_k) lying in the open left half-plane \mathbb{C}_- , then a scalar control operator corresponding to a sequence (b_k) is admissible if and only if the measure

$$\mu := \sum_k |b_k|^2 \delta_{-\lambda_k}$$

is a Carleson measure for the Hardy space $H^2(\mathbb{C}_+)$ on the right half-plane: this means that the canonical embedding $H^2(\mathbb{C}_+) \rightarrow L^2(\mathbb{C}_+, \mu)$ is bounded. In fact, the extension to normal semigroups has also been made [26].

Generalizations to α -admissibility, in which u must lie in $L^2(0, \infty; t^\alpha dt)$ for $-1 < \alpha < 0$, were studied by Wynn [28]. The key fact here is that under the Laplace transform the space $L^2(0, \infty; t^\alpha dt)$ is mapped to a weighted Bergman space, and for these there are analogues of the Carleson measure theorem available.

The results below enable us to take this generalization further and consider admissibility in the sense of the input lying in much more general spaces $L_w^2(0, \infty) = L^2(0, \infty; w(t) dt)$.

In order to state the link between admissibility and embeddings in the greatest generality possible, we assume now that $1 \leq q < \infty$ and the semigroup $(T(t))_{t \geq 0}$ acts on a Banach space X with a q -Riesz basis of eigenvectors (ϕ_k) ; that is, $T(t)\phi_k = e^{\lambda_k t} \phi_k$ for each k , with each λ_k lying in \mathbb{C}_- , and (ϕ_k) is a Schauder basis of X such that for some $C_1, C_2 > 0$ we have

$$C_1 \sum |a_k|^q \leq \left\| \sum a_k \phi_k \right\|^q \leq C_2 \sum |a_k|^q$$

for all sequences (a_k) in ℓ^q . In practice, this will mean that without loss of generality we can assume that $X = \ell^q$ and that the eigenvectors of the generator of $(T(t))_{t \geq 0}$, denoted by A , are the canonical basis of ℓ^q . We suppose also that we have a Banach space Z of functions on $(0, \infty)$, which may be an L^p space or a weighted space $L^2(0, \infty; w(t) dt)$, whose dual space Z^* can be regarded, respectively, as either $L^{p'}(0, \infty)$ or $L^2(0, \infty; w(t)^{-1} dt)$ in a natural way. Here, and throughout the paper, p' denotes the conjugate exponent to p , i.e., $p' = p/(p-1)$.

The following result links admissibility and Laplace–Carleson embeddings (that is, Carleson embeddings induced by the Laplace transform).

Theorem 2.1 *Let B be a linear bounded map from \mathbb{C} to $D(A^*)'$ corresponding to the sequence (b_k) . The control operator B is Z -admissible for $(T(t))_{t \geq 0}$, that is, there is a constant $m_0 > 0$ such that*

$$\left\| \int_0^\infty T(t) B u(t) dt \right\|_X \leq m_0 \|u\|_Z, \quad u \in Z,$$

if and only if the Laplace transform induces a continuous mapping from Z into $L^q(\mathbb{C}_+, d\mu)$, where μ is the measure $\sum |b_k|^q \delta_{-\lambda_k}$.

Proof: Clearly we may suppose without loss of generality that $X = \ell^q$ and that $\phi_k = e_k$, the standard basis of X . We have that

$$\begin{aligned} \left\| \int_0^\infty T(t)Bu(t) dt \right\|_X &= \left\| \int_0^\infty \sum_k e^{\lambda_k t} b_k e_k u(t) dt \right\|_X \\ &= \left(\sum_k |\hat{u}(-\lambda_k)|^q |b_k|^q \right)^{1/q}, \end{aligned}$$

from which the result follows easily. ■

A duality argument gives the corresponding result for observation operators.

Theorem 2.2 *Let C be a linear bounded map from $D(A)$ to \mathbb{C} . The observation operator C is Z -admissible for $(T(t))_{t \geq 0}$, that is, there is a constant $m_0 > 0$ such that*

$$\|CT(\cdot)x\|_Z \leq m_0 \|x\|_X$$

for all $x \in D(A)$, if and only if the Laplace transform induces a continuous mapping from Z^* into $L^{q'}(\mathbb{C}_+, d\mu)$, where μ is the measure $\sum |c_k|^{q'} \delta_{-\lambda_k}$ and $c_k := C\phi_k$.

Proof: Again we may suppose without loss of generality that $X = \ell^q$ and that $\phi_k = e_k$, the standard basis of X . Z -admissibility is equivalent to the condition that

$$\sup_{f,x} \left| \int_0^\infty CT(t)xf(t) dt \right| < \infty,$$

where we take $f \in Z^*$ and $x = (x_k) \in D(A)$ both of norm 1. Calculating $CT(t)x$, we see that this is equivalent to the condition that

$$\sup_{\|f\|=\|x\|=1} \left| \sum_k c_k \hat{f}(-\lambda_k) x_k \right| < \infty,$$

and, taking the supremum over the set of x of norm 1 in $D(A)$, which is dense, we obtain

$$\sup_{\|f\|=1} \|(c_k \hat{f}(-\lambda_k))\|_{q'} < \infty,$$

which is easily seen to be equivalent to the boundedness of the Laplace–Carleson embedding from Z^* into $L^{q'}(\mathbb{C}_+, d\mu)$. ■

In general we shall state our results in terms of control operators, leaving the interested reader to deduce the corresponding results for observation operators.

Now let $\tilde{\nu}$ be a positive regular Borel measure on $[0, \infty)$ satisfying the following (Δ_2) -condition:

$$R := \sup_{r>0} \frac{\tilde{\nu}[0, 2r)}{\tilde{\nu}[0, r)} < \infty. \tag{\Delta_2}$$

Let ν be the positive regular Borel measure on $\overline{\mathbb{C}_+} = [0, \infty) \times \mathbb{R}$ given by $d\nu = d\tilde{\nu} \otimes d\lambda$, where λ denotes Lebesgue measure. In this case, for $1 \leq p < \infty$, we call

$$A_\nu^p = \left\{ f : \mathbb{C}_+ \rightarrow \mathbb{C} \text{ analytic} : \sup_{\varepsilon > 0} \int_{\overline{\mathbb{C}_+}} |f(z + \varepsilon)|^p d\nu(z) < \infty \right\}$$

a Zen space on \mathbb{C}_+ . If $\tilde{\nu}(\{0\}) > 0$, then by standard Hardy space theory, f has a well-defined boundary function $\tilde{f} \in L^p(i\mathbb{R})$, and we can give meaning to the expression $\int_{\overline{\mathbb{C}_+}} |f(z)|^p d\nu(z)$. Therefore, we write

$$\|f\|_{A_\nu^p} = \left(\int_{\overline{\mathbb{C}_+}} |f(z)|^p d\nu(z) \right)^{1/p},$$

a notation that makes sense whether or not $\tilde{\nu}(\{0\}) = 0$. Clearly the space A_ν^2 is a Hilbert space.

Well-known examples of Zen spaces are Hardy spaces $H^p(\mathbb{C}_+)$, where $\tilde{\nu}$ is the Dirac measure at 0, or the standard weighted Bergman spaces A_α^p , where $d\tilde{\nu}(r) = r^\alpha dr$, $\alpha > -1$. Some further examples constructed from Hardy spaces on shifted half planes were given by Zen Harper in [6, 7].

The following proposition, given in [14], is elementary and appears for special cases in [6, 7]. Partial results are also given in [2, 3].

Proposition 2.3 (*Proposition 2.3 in [14]*) *Let A_ν^2 be a Zen space, and let $w : (0, \infty) \rightarrow \mathbb{R}_+$ be given by*

$$w(t) = 2\pi \int_0^\infty e^{-2rt} d\tilde{\nu}(r) \quad (t > 0).$$

Then the Laplace transform defines an isometric map $\mathcal{L} : L_w^2(0, \infty) \rightarrow A_\nu^2$.

Note that the existence of the integral is guaranteed by the (Δ_2) -condition.

We shall require the following Laplace–Carleson Embedding Theorem from [14].

Theorem 2.4 (*Theorem 2.4 in [14]*) *Let A_ν^2 be a Zen space, $\nu = \tilde{\nu} \otimes \lambda$, and let $w : (0, \infty) \rightarrow \mathbb{R}_+$ be given by*

$$w(t) = 2\pi \int_0^\infty e^{-2rt} d\tilde{\nu}(r) \quad (t > 0). \quad (2)$$

Let μ be a positive regular Borel measure on \mathbb{C}_+ . Then the following are equivalent:

1. *The Laplace transform \mathcal{L} given by $\mathcal{L}f(z) = \int_0^\infty e^{-tz} f(t) dt$ defines a bounded linear map*

$$\mathcal{L} : L_w^2(0, \infty) \rightarrow L^2(\mathbb{C}_+, \mu).$$

2. *For a sufficiently large $N \in \mathbb{N}$, there exists a constant $\kappa > 0$ such that*

$$\int_{\mathbb{C}_+} \left| (\mathcal{L}t^{N-1}e^{-\lambda t})(z) \right|^2 d\mu(z) \leq \kappa \int_0^\infty |t^{N-1}e^{-\lambda t}|^2 w(t) dt \text{ for each } \lambda \in \mathbb{C}_+.$$

3. There exists a constant $\kappa > 0$ such that

$$\mu(Q_I) \leq \kappa \nu(Q_I) \text{ for each Carleson square } Q_I.$$

where Q_I denotes the Carleson square $Q_I = \{z = x + iy \in \mathbb{C}_+ : iy \in I, 0 < x < |I|\}$.

From this we may deduce a result characterizing admissibility for normal semigroups in the sense of $L_w^2(0, \infty)$.

Theorem 2.5 *Suppose that $A : D(A) \subset H \rightarrow H$ has a Riesz basis (ϕ_k) of eigenvectors with eigenvalues (λ_k) satisfying $\operatorname{Re} \lambda_k < 0$ and let B be a linear bounded map from \mathbb{C} to $D(A^*)'$ given by the sequence (b_k) . Let A_ν^2 be a Zen space and let w be given by (2). Then the following statements are equivalent.*

1. B is an admissible control operator with respect to $L_w^2(0, \infty)$, that is, there exists a constant $m_0 > 0$ such that

$$\left\| \int_0^\infty T(t)Bu(t) dt \right\|_H \leq m_0 \|u\|_{L_w^2(0, \infty)}$$

for every $u \in L_w^2(0, \infty)$.

2. For a sufficiently large $N \in \mathbb{N}$, there exists a constant $\kappa > 0$ such that

$$\|(\lambda - A)^{-N}B\|^2 \leq \kappa \int_0^\infty |t^{N-1}e^{-\lambda t}|^2 w(t) dt, \quad (\lambda \in \mathbb{C}_+).$$

3. There exists a constant $\kappa > 0$ such that

$$\mu(Q_I) \leq \kappa \nu(Q_I) \text{ for each Carleson square } Q_I,$$

where $\mu = \sum_k |b_k|^2 \delta_{-\lambda_k}$.

Proof: This follows from Theorem 2.1 and Theorem 2.4, taking $q = 2$ and $Z = L_w^2(0, \infty)$. Note that the resolvent condition follows because

$$\|(\lambda - A)^{-N}B\|^2 = \sum_k |\lambda - \lambda_k|^{-2N} |b_k|^2 = \int_{\mathbb{C}_+} \frac{d\mu(z)}{|\lambda + z|^{2N}},$$

and the Laplace transform of $t^{N-1}e^{-\lambda t}$ is a constant multiple of $(\lambda + z)^{-N}$. ■

Remark 2.6 *Theorem 2.5 in the case that $\tilde{\nu}$ equals the Dirac measure in 0, that is $A_\nu^2 = H^2(\mathbb{C}_+)$, is due to Ho and Russell [8], and Weiss [25]. In the case $d\tilde{\nu} = r^\alpha dr$, $\alpha \in (-1, 0)$ the result is due to Wynn [28]. Using Theorem 2.5 any $\alpha > -1$ can now be considered.*

In [5], Haak applied Carleson measure theory to find conditions for admissibility of the system (1) above, with A generating a diagonal semigroup defined on ℓ^q , and inputs lying in the space $L^p(0, \infty)$. Using the Hausdorff–Young inequality, it is possible to characterize admissibility in the situation $p \leq 2$ and $1 < p' \leq q < \infty$.

Theorem 2.7 *Let $p \leq 2$ and $1 < p' \leq q < \infty$. Suppose that $A : D(A) \subset \ell^q \rightarrow \ell^q$ is a diagonal operator with eigenvalues (λ_k) satisfying $\operatorname{Re} \lambda_k < 0$, and let B be a linear bounded map from \mathbb{C} to $D(A^*)'$ corresponding to the sequence (b_k) . Then the following statements are equivalent.*

1. *B is an admissible control operator with respect to $L^p(0, \infty)$, that is, there exists a constant $m_0 > 0$ such that*

$$\left\| \int_0^\infty T(t)Bu(t) dt \right\|_{\ell^q} \leq m_0 \|u\|_{L^p(0, \infty)}$$

for every $u \in L^p(0, \infty)$.

2. *There exists a constant $\kappa > 0$ such that*

$$\mu(Q_I) \leq \kappa |I|^{q/p'}$$

for all intervals in $I \subset i\mathbb{R}$, where $\mu = \sum_k |b_k|^q \delta_{-\lambda_k}$.

3. *There exists a constant $\kappa > 0$ such that*

$$\|(\lambda - A)^{-1}B\|_{\ell^q} \leq \kappa |\operatorname{Re} \lambda|^{-1/p}$$

for all $\lambda \in \mathbb{C}_+$.

The theorem is a corollary of Theorem 2.1 combined with [14, Thm. 3.2]. Moreover the equivalence of Part 1 and 2 can be found in [5].

Example 2.8 *We study the one-dimensional Schrödinger equation on the interval $[0, 1]$, which is given by*

$$\begin{aligned} \frac{\partial z}{\partial t}(\zeta, t) &= i \frac{\partial^2 z}{\partial \zeta^2}(\zeta, t) - z(\zeta, t), & \zeta \in (0, 1), t \geq 0, \\ \frac{\partial z}{\partial \zeta}(0, t) &= 0, \quad \frac{\partial z}{\partial \zeta}(1, t) = u(t), & t \geq 0, \\ z(\zeta, 0) &= z_0(\zeta), & \zeta \in (0, 1). \end{aligned}$$

The PDE can be written equivalently in the form (1) with $X = \ell^q$, $Ae_n = (-in^2\pi^2 - 1)e_n$, and $b_n = i\sqrt{2}$ for each n . Let $q \geq p'$ and $p \leq 2$. By Theorem 2.7, the operator B is an admissible control operator with respect to $L^p(0, \infty)$.

3 Admissibility for analytic semigroups

There is no general characterization of admissibility of the system (1) above, with A generating a diagonal semigroup defined on ℓ^q , and inputs lying in the space $L^p(0, \infty)$, $p > 2$, as there is no known full characterization of boundedness of Laplace–Carleson embeddings

$$L^p(0, \infty) \rightarrow L^q(\mathbb{C}_+, \mu), \quad f \mapsto \mathcal{L}f = \int_0^\infty e^{-t} f(t) dt.$$

However, characterizations are possible in some cases with additional information on the support on the measure.

If the measure μ is supported on a sector $S(\theta) = \{z \in \mathbb{C}_+ : |\arg z| < \theta\}$ for some $0 < \theta < \frac{\pi}{2}$, then the oscillatory part of the Laplace transform can be discounted, and a full characterization of boundedness can be achieved (see also [5], Theorem 3.2 for an alternative characterization by means of a different measure).

Theorem 3.1 (Theorem 3.3 in [14]) *Let μ be a positive regular Borel measure supported in a sector $S(\theta) \subset \mathbb{C}_+$, $0 < \theta < \frac{\pi}{2}$, and let $q \geq p > 1$. Then the following are equivalent:*

1. *The Laplace–Carleson embedding*

$$\mathcal{L} : L^p(0, \infty) \rightarrow L^q(\mathbb{C}_+, \mu), \quad f \mapsto \mathcal{L}f,$$

is well-defined and bounded.

2. *There exists a constant $\kappa > 0$ such that $\mu(Q_I) \leq \kappa|I|^{q/p'}$ for all intervals in $I \subset i\mathbb{R}$ which are symmetric about 0.*
3. *There exists a constant $\kappa > 0$ such that $\|\mathcal{L}e^{-z}\|_{L^q_\mu} \leq \kappa\|e^{-z}\|_{L^p}$ for all $z \in \mathbb{R}_+$.*

From this we may deduce a result characterizing admissibility for analytic semigroups with respect to $L^p(0, \infty)$.

Theorem 3.2 *Let $1 < p \leq q < \infty$. Suppose that $A : D(A) \subset \ell^q \rightarrow \ell^q$ is a diagonal operator with eigenvalues (λ_k) satisfying $\operatorname{Re} \lambda_k < 0$ and $(-\lambda_k) \subset S(\theta)$ for some $\theta \in (0, \frac{\pi}{2})$, and let B be a linear bounded map from \mathbb{C} to $D(A^*)'$ given by the sequence (b_k) . Then the following statements are equivalent.*

1. *B is an admissible control operator with respect to $L^p(0, \infty)$, that is, there exists a constant $m_0 > 0$ such that*

$$\left\| \int_0^\infty T(t)Bu(t) dt \right\|_{\ell^q} \leq m_0 \|u\|_{L^p(0, \infty)}$$

for every $u \in L^p(0, \infty)$.

2. *There exists a constant $\kappa > 0$ such that*

$$\mu(Q_I) \leq \kappa \nu |I|^{q/p'}$$

for all intervals in $I \subset i\mathbb{R}$ which are symmetric about 0, where $\mu = \sum_k |b_k|^q \delta_{-\lambda_k}$.

3. *There exists a constant $\kappa > 0$ such that*

$$\|(z - A)^{-1}B\|_{\ell^q} \leq \kappa z^{-1/p}$$

for all $z \in \mathbb{R}_+$.

Remark 3.3 Let μ , θ , p and q be as in Theorem 3.2. In [5], Theorem 3.2, the equivalence of the following statements is shown:

1. B is an admissible control operator with respect to $L^p(0, \infty)$, that is, there exists a constant $m_0 > 0$ such that

$$\left\| \int_0^\infty T(t)Bu(t) dt \right\|_{\ell^q} \leq m_0 \|u\|_{L^p(0, \infty)}$$

for every $u \in L^p(0, \infty)$.

2. There exists a constant $\kappa > 0$ such that $\tilde{\mu}(Q_I) \leq \kappa |I|^{q/p}$ for all intervals in $I \subset i\mathbb{R}$ which are symmetric about 0, where $d\tilde{\mu}(z) = |z|^q d\mu(\frac{1}{z})$.

Now let us consider the case $p > q$ for sectorial measures μ . In [14] a condition in terms of the balayage S_μ of μ has been obtained. Recall that the balayage S_μ of a positive Borel measure μ on \mathbb{C}_+ is given by $S_\mu(t) = \int_{\mathbb{C}_+} p_z(t) d\mu(z)$, where p_z denotes the Poisson kernel of the right half plane. Let $S_n := \{z \in \mathbb{C} \mid 2^{n-1} < \operatorname{Re} z \leq 2^n\}$.

Theorem 3.4 (Theorem 3.5 in [14]) Let μ be a positive regular Borel measure supported in a sector $S(\theta) \subset \mathbb{C}_+$, $0 < \theta < \frac{\pi}{2}$ and let $1 \leq q < p < \infty$. Then the following are equivalent:

1. The embedding

$$\mathcal{L} : L^p(\mathbb{R}_+) \rightarrow L^q(\mathbb{C}_+, \mu), \quad f \mapsto \mathcal{L}f,$$

is well-defined and bounded.

2. The sequence $(2^{-nq/p'} \mu(S_n))$ is in $\ell^{p/(p-q)}(\mathbb{Z})$.
3. The sequence $(2^{n/p} \|\mathcal{L}e^{-2^n}\|_{L_\mu^q})$ is in $\ell^{qp/(p-q)}(\mathbb{Z})$.

If $p' < q$, then the above is also equivalent to

4. $t^{q(2-p)/p} S_\mu \in L^{p/(p-q)}(\mathbb{R})$.

As a corollary we obtain the following result.

Theorem 3.5 Let $1 \leq q < p < \infty$. Suppose that $A : D(A) \subset \ell^q \rightarrow \ell^q$ is a diagonal operator with eigenvalues (λ_k) satisfying $\operatorname{Re} \lambda_k < 0$ and $(-\lambda_k) \subset S(\theta)$ for some $\theta \in (0, \frac{\pi}{2})$, and let B be a linear bounded map from \mathbb{C} to $D(A^*)'$ given by the sequence (b_k) . Then the following statements are equivalent.

1. B is an admissible control operator with respect to $L^p(0, \infty)$, that is, there exists a constant $m_0 > 0$ such that

$$\left\| \int_0^\infty T(t)Bu(t) dt \right\|_{\ell^q} \leq m_0 \|u\|_{L^p(0, \infty)}$$

for every $u \in L^p(0, \infty)$.

2. The sequence $(2^{-nq/p'} \mu(S_n))$ is in $\ell^{p/(p-q)}(\mathbb{Z})$.
3. The sequence $(2^{n/p} \|(2^n - A)^{-1} B\|_{\ell^q})$ is in $\ell^{qp/(p-q)}(\mathbb{Z})$.

If $p' < q$, then the above is also equivalent to

4. $t^{q(2-p)/p} S_\mu \in L^{p/(p-q)}(\mathbb{R})$.

Here $\mu = \sum_k |b_k|^q \delta_{-\lambda_k}$.

Example 3.6 We study the one-dimensional heat equation on the interval $[0, 1]$, which is given by

$$\begin{aligned} \frac{\partial z}{\partial t}(\zeta, t) &= \frac{\partial^2 z}{\partial \zeta^2}(\zeta, t), & \zeta \in (0, 1), t \geq 0, \\ \frac{\partial z}{\partial \zeta}(0, t) &= 0, \quad \frac{\partial z}{\partial \zeta}(1, t) = u(t), & t \geq 0, \\ z(\zeta, 0) &= z_0(\zeta), & \zeta \in (0, 1). \end{aligned}$$

This PDE can be written equivalently in the form (1) with $X = \ell^2$, $Ae_n = -n^2 \pi^2 e_n$, and b_n defined by $b_n = 1$ for each n . By Theorem 3.2 (for $1 \leq p \leq 2$) and Theorem 3.5 (for $2 \leq p < \infty$), the operator B is an admissible control operator with respect to $L^p(0, \infty)$ if and only if $p \geq 4/3$.

Control problems involving smoother, Sobolev-space valued, controls are related to embeddings of the form

$$\mathcal{H}_\beta^p(0, \infty) \rightarrow L^q(\mathbb{C}_+, \mu), \quad f \mapsto \mathcal{L}f = \int_0^\infty e^{-t} f(t) dt,$$

given by the Laplace transform \mathcal{L} . Here, for $\beta > 0$ the space $\mathcal{H}_\beta^p(0, \infty)$ is given by

$$\begin{aligned} \mathcal{H}_\beta^p(0, \infty) &= \left\{ f \in L^p(\mathbb{R}_+) : \int_0^\infty \left| \left(\frac{d}{dx} \right)^\beta f(t) \right|^p dt < \infty \right\}, \\ \|f\|_{\mathcal{H}_\beta^p}^p &= \|f\|_p^p + \left\| \left(\frac{d}{dx} \right)^\beta f \right\|_p^p. \end{aligned}$$

Here $(\frac{d}{dx})^\beta f$ is defined as a fractional derivative via the Fourier transform. We call \mathcal{H}_β^p the Sobolev space of index β .

In [14, Corollaries 3.7 and 3.8] the following characterisations of the boundedness of the Laplace–Carleson embedding has been proved.

Proposition 3.7 (Corollary 3.7 in [14]) *Let μ be a positive Borel measure supported in a sector $S(\theta) \subset \mathbb{C}_+$, $0 < \theta < \frac{\pi}{2}$, and let $q \geq p > 1$. Then the following are equivalent:*

1. The Laplace–Carleson embedding

$$\mathcal{L} : \mathcal{H}_\beta^p(0, \infty) \rightarrow L^q(\mathbb{C}_+, \mu), \quad f \mapsto \mathcal{L}f,$$

is well-defined and bounded.

2. There exists a constant $\kappa > 0$ such that $\mu_{q,\beta}(Q_I) \leq \kappa|I|^{q/p'}$ for all intervals in $I \subset i\mathbb{R}$ which are symmetric about 0. Here, $d\mu_{q,\beta}(z) = (1 + \frac{1}{|z|^{q\beta}})d\mu(z)$.
3. There exists a constant $\kappa > 0$ such that $\|\mathcal{L}e^{-\cdot z}\|_{L^q_\mu} \leq \kappa\|e^{-\cdot z}\|_{\mathcal{H}^p_\beta}$ for all $z \in \mathbb{R}_+$.

Proposition 3.8 (Corollary 3.8 in [14]) *Let μ be a positive regular Borel measure supported in a sector $S(\theta) \subset \mathbb{C}_+$, $0 < \theta < \frac{\pi}{2}$ and let $1 \leq q < p$, $\beta \geq 0$. Suppose that $S_{\mu,\beta,q} \in L^{p/(p-q)}$. Then the embedding*

$$\mathcal{L} : \mathcal{H}^p_\beta(0, \infty) \rightarrow L^q(\mathbb{C}_+, \mu), \quad f \mapsto \mathcal{L}f,$$

is well-defined and bounded.

As an application we are able to characterize admissibility with respect to Sobolev space valued control functions.

Theorem 3.9 *Let $1 < p < q < \infty$. Suppose that $A : D(A) \subset \ell^q \rightarrow \ell^q$ is a diagonal operator with eigenvalues (λ_k) satisfying $\operatorname{Re} \lambda_k < 0$ and $(-\lambda_k) \subset S(\theta)$ for some $\theta \in (0, \frac{\pi}{2})$, and let B be a linear bounded map from \mathbb{C} to $D(A^*)'$. Write $\mu = \sum |b_k|^q \delta_{-\lambda_k}$. Then the following statements are equivalent.*

1. B is an admissible control operator with respect to $\mathcal{H}^p_\beta(0, \infty)$, that is, there exists a constant $m_0 > 0$ such that

$$\left\| \int_0^\infty T(t)Bu(t) dt \right\|_{\ell^q} \leq m_0 \|u\|_{\mathcal{H}^p_\beta(0, \infty)},$$

for every $u \in \mathcal{H}^p_\beta(0, \infty)$.

2. There exists a constant $\kappa > 0$ such that $\mu_{q,\beta}(Q_I) \leq \kappa|I|^{q/p'}$ for all intervals in $I \subset i\mathbb{R}$ which are symmetric about 0. Here, $d\mu_{q,\beta}(z) = (1 + \frac{1}{|z|^{q\beta}})d\mu(z)$.
3. There exists a constant $\kappa > 0$ such that $\|(z - A)^{-1}B\|_{\ell^q} \leq \kappa\|e^{-\cdot z}\|_{\mathcal{H}^p_\beta}$ for all $z \in \mathbb{R}_+$.

Theorem 3.10 *Let $1 < q < p < \infty$. Suppose that $A : D(A) \subset \ell^q \rightarrow \ell^q$ is a diagonal operator with eigenvalues (λ_k) satisfying $\operatorname{Re} \lambda_k < 0$ and $(-\lambda_k) \subset S(\theta)$ for some $\theta \in (0, \frac{\pi}{2})$, and let B be a linear bounded map from \mathbb{C} to $D(A^*)'$. Suppose that $S_{\bar{\mu},\beta,q} \in L^{p/(p-q)}$. Then B is an admissible control operator with respect to $\mathcal{H}^p_\beta(0, \infty)$, that is, there exists a constant $m_0 > 0$ such that*

$$\left\| \int_0^\infty T(t)Bu(t) dt \right\|_{\ell^q} \leq m_0 \|u\|_{\mathcal{H}^p_\beta(0, \infty)},$$

for every $u \in \mathcal{H}^p_\beta(0, \infty)$.

In the setting of analytic diagonal semigroups, it is also possible to characterise $L^2((0, \infty), t^\alpha dt)$ -admissibility of control operators and $L^2((0, \infty), t^{-\alpha} dt)$ -admissibility of observation operators in the range $0 < \alpha < 1$ in terms of a Carleson-type condition or a resolvent condition (compare this with the counterexample by Wynn in [27] for a diagonal semigroup).

Generally, the difficulty in the case $\alpha > 0$ stems from the well-known fact that the boundedness of Carleson embeddings on Dirichlet space cannot be tested on reproducing kernels or by means of a simple Carleson-type condition [21]. In the case of sectorial measures, however, such a characterisation is possible, at least for $0 < \alpha < 1$. For an interval $I \subset i\mathbb{R}$, let T_I denote the right half of the Carleson square Q_I .

Theorem 3.11 *Let μ be a positive Borel measure supported in a sector $S(\theta) \subset \mathbb{C}_+$, $0 < \theta < \frac{\pi}{2}$, and let $0 < \alpha < 1$. Then the following are equivalent:*

1. *The Laplace–Carleson embedding*

$$\mathcal{L} : L^2((0, \infty), t^\alpha dt) \rightarrow L^2(\mathbb{C}_+, \mu), \quad f \mapsto \mathcal{L}f,$$

is well-defined and bounded.

2. *There exists a constant $\gamma > 0$ such that*

$$\mu(T_I) \leq \gamma |I|^{1-\alpha}$$

for all intervals in $I \subset i\mathbb{R}$ which are symmetric about 0.

3. *There exists a constant $\kappa > 0$ such that*

$$\|\mathcal{L}t^{-\alpha}e^{-tz}\|_{L^2(\mathbb{C}_+, \mu)} \leq \kappa \|t^{-\alpha}e^{-tz}\|_{L^2(t^\alpha dt)}$$

for all $z \in \mathbb{R}_+$.

Proof: The implication (1) \Rightarrow (3) is immediate. For (3) \Rightarrow (2), let $z = z_I = |I|/2$ denote the centre of the Carleson square Q_I over an interval $I \subset i\mathbb{R}$ which is symmetric about 0. Then the modulus of the function

$$(\mathcal{L}t^{-\alpha}e^{-tz_I})(s) = \frac{\Gamma(1-\alpha)}{(z_I + s)^{-\alpha+1}} \tag{3}$$

is bounded below by $\Gamma(1-\alpha)\frac{1}{(2z_I)^{-\alpha+1}}$ on T_I , and therefore

$$\begin{aligned} \mu(T_I) &\leq \frac{(2z_I)^{2-2\alpha}}{\Gamma(1-\alpha)^2} \int_{\mathbb{C}_+} |(\mathcal{L}t^{-\alpha}e^{-tz_I})(s)|^2 d\mu(s) \\ &\leq \kappa^2 \frac{(2z_I)^{2-2\alpha}}{\Gamma(1-\alpha)^2} \|t^{-\alpha}e^{-tz_I}\|_{L^2(t^\alpha dt)}^2 = \frac{\kappa^2}{\Gamma(1-\alpha)} |I|^{1-\alpha}. \end{aligned}$$

Let us now consider (2) \Rightarrow (1). We use the argument from [14], Thm 3.3. For $n \in \mathbb{Z}$, let

$$T_n = \{x + iy \in \mathbb{C}_+ : 2^{n-1} < x \leq 2^n, -2^{n-1} < y \leq 2^{n-1}\}.$$

That is, T_n is the right half of the Carleson square Q_{I_n} over the interval $I_n = \{y \in i\mathbb{R}, |y| \leq 2^{n-1}\}$. The T_n are obviously pairwise disjoint.

Without loss of generality we assume $0 < \theta < \arctan(\frac{1}{2})$, in which case $S(\theta) \subseteq \bigcup_{n=-\infty}^{\infty} T_n$ (otherwise, we also have to use finitely many translates $T_{n,k}$ of each T_n , for which the same estimates apply).

Now let $z \in T_n$ for some $n \in \mathbb{Z}$. Then we obtain, for $f \in L^2((0, \infty), t^\alpha dt)$,

$$\begin{aligned} |\mathcal{L}f(z)| &\leq \int_0^\infty |e^{-zt}| |f(t)| dt \leq \int_0^\infty |t^{-\alpha/2} e^{-2^{n-1}t}| |f(t) t^{\alpha/2}| dt \\ &\leq C_\alpha 2^{(-n+1)(1-\alpha/2)} (M(t^{\alpha/2} f))(2^{-n+1}), \end{aligned}$$

where $C_\alpha > 0$ is a constant dependent only on the L^1 -integration kernel $\phi_\alpha(t) = \chi_{[0, \infty)}(t+1)(t+1)^{\alpha/2} e^{-t-1}$, and Mf is the Hardy–Littlewood maximal function. We refer to e.g. [22], page 57, equation (16) for a pointwise estimate between the maximal function induced by the kernel ϕ_α , and M . Note that we can easily dominate ϕ_α by a positive, radial, decreasing L^1 function here. Consequently,

$$\begin{aligned} \int_{S(\theta)} |\mathcal{L}f(z)|^2 d\mu(z) &\leq C_\alpha^2 \sum_{n=-\infty}^{\infty} 2^{(-n+1)(2-\alpha)} (M(t^{\alpha/2} f)(2^{-n+1}))^2 \mu(T_n) \\ &\leq \gamma C_\alpha^2 \sum_{n=-\infty}^{\infty} 2^{(-n+1)(2-\alpha)} 2^{n(1-\alpha)} (M(t^{\alpha/2} f)(2^{-n+1}))^2 \\ &= \gamma C_\alpha^2 2^{1-\alpha} \sum_{n=-\infty}^{\infty} 2^{-n+1} M(t^{\alpha/2} f)(2^{-n+1})^2 \\ &\lesssim \|f\|_{L^2(t^\alpha dt)}^2. \end{aligned}$$

■

We therefore have the following corollary of Theorem 3.11 and Theorem 2.1.

Corollary 3.12 *Let $0 < \alpha < 1$. Suppose that $A : D(A) \subset \ell^2 \rightarrow \ell^2$ is a diagonal operator with eigenvalues (λ_k) satisfying $\operatorname{Re} \lambda_k < 0$ and $(-\lambda_k) \subset S(\theta)$ for some $\theta \in (0, \frac{\pi}{2})$, and let B be a linear bounded map from \mathbb{C} to $D(A^*)'$. Write $\mu = \sum |b_k|^2 \delta_{-\lambda_k}$. Then the following statements are equivalent.*

1. *B is an admissible control operator with respect to $L^2((0, \infty), t^\alpha dt)$, that is, there exists a constant $m_0 > 0$ such that*

$$\left\| \int_0^\infty T(t) B u(t) dt \right\|_{\ell^2} \leq m_0 \|u\|_{L^2((0, \infty), t^\alpha dt)},$$

for every $u \in L^2((0, \infty), t^\alpha dt)$.

2. *There exists a constant $\kappa > 0$ such that $\mu(T_I) \leq \kappa |I|^{1-\alpha}$ for all intervals in $I \subset i\mathbb{R}$ which are symmetric about 0.*
3. *There exists a constant $\kappa > 0$ such that*

$$\|\mathcal{L}t^{-\alpha} e^{-\lambda t}\|_{L^2(\mathbb{C}_+, \mu)} \leq \kappa \|t^{-\alpha} e^{-\lambda t}\|_{L^2(t^\alpha dt)}$$

for all $\lambda \in \mathbb{R}_+$.

4. There exists a constant $\kappa > 0$ such that

$$\|(\lambda - A)^{\alpha-1}B\| \leq \kappa\lambda^{(\alpha-1)/2}$$

for all $\lambda \in \mathbb{R}_+$.

Proof: It is enough to check that conditions 3 and 4 are equivalent. Note that for $\lambda \in \mathbb{R}_+$

$$\|(\lambda - A)^{\alpha-1}B\|^2 = \sum_k |b_k|^2 |\lambda - \lambda_k|^{2\alpha-2},$$

which is a constant multiple of $\|\mathcal{L}t^{-\alpha}e^{-\lambda t}\|_{L^2(\mathbb{C}_+, \mu)}^2$ (cf. (3)). Similarly, we have that $\|t^{-\alpha}e^{-\lambda t}\|_{L^2(t^\alpha dt)}^2$ is a constant multiple of $\lambda^{\alpha-1}$. ■

Note that the corollary is also valid in the limiting case $\alpha = 0$.

4 Exact controllability for diagonal systems

We consider the equation

$$\frac{dx(t)}{dt} = Ax(t) + Bu(t), \quad (4)$$

with solution

$$x(t) = T(t)x_0 + \int_0^t T(t-s)Bu(s) ds,$$

suitably interpreted. We now consider an exponentially stable C_0 semigroup $(T(t))_{t \geq 0}$, on a Hilbert space H , i.e.,

$$\|T(t)\| \leq Me^{-\lambda t}, \quad (t \geq 0),$$

for some $M > 0$ and $\lambda > 0$. Suppose first that B is admissible. Then we have a bounded operator $\mathcal{B}_\infty : L^2(0, \infty; U) \rightarrow H$, defined by

$$\mathcal{B}_\infty u = \int_0^\infty T(t)Bu(t) dt.$$

The system is *exactly controllable*, if its range $R(\mathcal{B}_\infty)$ equals H . If B is not admissible, then the operator \mathcal{B}_∞ is commonly defined as a mapping into a larger (extrapolation) space and exact controllability requires that its image contains H .

Although we shall be working with exact controllability in infinite time, it is possible (but much more difficult) to use interpolation and embedding techniques in Paley–Wiener spaces in order to obtain analogous results on finite-time controllability. Some results along these lines can be found in [13].

In [10] exact controllability for diagonal systems with scalar inputs was characterised in terms of Carleson measures (a version for multivariable inputs was

given in [11]). In particular if A has a Riesz basis of eigenvectors, with eigenvalues (λ_n) (as always, assumed to lie in \mathbb{C}_- , but with no separation assumptions made), then with a control operator corresponding to a sequence (b_n) the system is exactly controllable *if and only if*

$$\nu_\lambda := \sum_n \frac{|\operatorname{Re} \lambda_n|^2}{|b_n|^2 \prod_{k \neq n} p(\lambda_n, \lambda_k)^2} \delta_{-\lambda_n}$$

is a Carleson measure. Here $p(\lambda_n, \lambda_k)$ is the pseudo-hyperbolic metric, i.e.,

$$p(\lambda_n, \lambda_k) = \left| \frac{\lambda_n - \lambda_k}{\lambda_n + \bar{\lambda}_k} \right|.$$

Exact controllability by inputs in Sobolev spaces \mathcal{H}_β^2 with $0 < \beta < 1/2$ was characterised in [12, Thm. 3.8]. In [14, Thm. 3.9] the following result was proved, which enable us to dispense with the restriction on β .

Theorem 4.1 *Let μ be a positive Borel measure on the right half plane \mathbb{C}_+ and let $\beta > 0$. Then the following are equivalent:*

1. *The Laplace–Carleson embedding*

$$\mathcal{H}_\beta^2(0, \infty) \rightarrow L^2(\mathbb{C}_+, \mu)$$

is bounded.

2. *The measure $|1 + z|^{-2\beta} d\mu(z)$ is a Carleson measure on \mathbb{C}_+ .*

In [19], [18], the following theorem was proved:

Theorem 4.2 *Let $(g_k)_{k \in \mathbb{N}}$ be a sequence of nonzero complex numbers and let $(z_k)_{k \in \mathbb{N}}$ be a Blaschke sequence in \mathbb{C}_+ . Write $b_{\infty, k} = \prod_{j \neq k} p(z_j, z_k)$. Let $m_2 = \sup_{(a_k) \in \ell^2, \|(a_k)\|_2=1} \inf_{f \in H^2(\mathbb{C}_+), g_k f(z_k) = a_k} \|f\|_{H^2}$. Then*

$$m_2 = \|\mathcal{J}_{\mu_2}\|$$

where

$$\mu_2 = \sum_{k=1}^{\infty} \frac{|2 \operatorname{Re} z_k|^2}{|b_{\infty, k} g_k|^2} \delta_{z_k},$$

and \mathcal{J}_{μ_2} is the Carleson embedding

$$\mathcal{J}_{\mu_2} : H^2(\mathbb{C}_+) \rightarrow L^2(\mathbb{C}_+, \mu_2).$$

Using Theorem 4.1, we obtain

Corollary 4.3 *Let $(g_k)_{k \in \mathbb{N}}$ be a sequence of nonzero complex numbers and let $(z_k)_{k \in \mathbb{N}}$ be a Blaschke sequence in \mathbb{C}_+ . Write $b_{\infty, k} = \prod_{j \neq k} p(z_j, z_k)$. Let*

$$m_\beta = \sup_{(a_k) \in \ell^2, \|(a_k)\|_2=1} \inf_{f \in \mathcal{H}_\beta^2} \{\|f\|_{\mathcal{H}_\beta^2} : g_k \mathcal{L}f(z_k) = a_k\}.$$

Then $m_\beta < \infty$, if and only if there is a constant $\kappa > 0$ such that

$$\sum_{z_k \in Q_I} \frac{|2 \operatorname{Re} z_k|^2 |1 + z_k|^{2\beta}}{|b_{\infty, k}|^2 |g_k|^2} \leq \kappa |I| \text{ for all intervals } I \subset i\mathbb{R}.$$

Finally, Corollary 4.3 enables us to characterize controllability by inputs in Sobolev spaces \mathcal{H}_β^2 .

Theorem 4.4 *Let $\beta > 0$. Suppose that $A : D(A) \subset H \rightarrow H$ has a Riesz basis (ϕ_k) of eigenvectors with eigenvalues (λ_k) satisfying $\operatorname{Re} \lambda_k < 0$ and let B be a linear bounded map from \mathbb{C} to $D(A^*)'$. Write $b_{\infty,k} = \prod_{j \neq k} p(\lambda_j, \lambda_k)$. Then the following statements are equivalent.*

1. System (4) is exactly controllable with respect to \mathcal{H}_β^2 , that is,

$$H \subset \mathcal{B}_\infty(\mathcal{H}_\beta^2).$$

2. there is a constant $\kappa > 0$ such that

$$\sum_{-\lambda_k \in Q_I} \frac{|2 \operatorname{Re} \lambda_k|^2 |1 - \lambda_k|^{2\beta}}{|b_{\infty,k}|^2 |g_k|^2} \leq \kappa |I| \text{ for all intervals } I \subset i\mathbb{R}.$$

Similar results can be proved for null-controllability, along the lines of [10].

5 Conclusion

Other techniques for analysing admissibility and controllability are of course available (see, for example, [1, 16, 17, 20, 23]), and these include Carleman inequalities, multipliers, moment problems and spectral analysis. However, the advantage of the present approach is that it gives a complete answer in terms of necessary and sufficient conditions in the case of diagonal systems and weighted input spaces. As a referee has pointed out, further questions may be asked about non-homogeneous problems, perturbed systems (including non-linear terms), and indeed stabilizability. These would make interesting research projects in their own right, and beyond the scope of this paper, although the issue of perturbed systems is addressed in the context of Carleson measures in the very recent paper [15].

Acknowledgements

This work was supported by EPSRC grant EP/I01621X/1. The third author also acknowledges partial support of this work by a Heisenberg Fellowship of the German Research Foundation (DFG).

References

- [1] A. Bensoussan, G. Da Prato and M. Delfour, *Representation and control of infinite dimensional systems*, Systems & Control: Foundations & Applications, Birkhäuser, Boston, 2007.
- [2] N. Das and J.R. Partington, Little Hankel operators on the half-plane. *Integral Equations Operator Theory* 20 (1994), no. 3, 306–324.

- [3] P. Duren, E.A. Gallardo-Gutiérrez and A. Montes-Rodríguez, A Paley–Wiener theorem for Bergman spaces with application to invariant subspaces. *Bull. Lond. Math. Soc.* 39 (2007), no. 3, 459–466.
- [4] S. Ervedoza and E. Zuazua, A systematic method for building smooth controls for smooth data, *Discrete Contin. Dyn. Syst. Ser. B*, 14 (2010), no. 4, 1375–1401.
- [5] B. H. Haak, On the Carleson measure criterion in linear systems theory. *Complex Anal. Oper. Theory* 4 (2010), no. 2, 281–299.
- [6] Z. Harper, Boundedness of convolution operators and input-output maps between weighted spaces. *Complex Anal. Oper. Theory* 3 (2009), no. 1, 113–146.
- [7] Z. Harper, Laplace transform representations and Paley–Wiener theorems for functions on vertical strips. *Doc. Math.* 15 (2010), 235–254.
- [8] L.F. Ho and D.L. Russell, Admissible input elements for systems in Hilbert space and a Carleson measure criterion, *SIAM J. Control Optim.* 21 (1983), no. 4, 614–640. Erratum, *SIAM J. Control Optim.* 21 (1983), no. 6, 985–986.
- [9] B. Jacob and J.R. Partington, Admissibility of control and observation operators for semigroups: a survey. *Current trends in operator theory and its applications*, 199–221, Oper. Theory Adv. Appl., 149, Birkhäuser, Basel, 2004.
- [10] B. Jacob and J.R. Partington, On controllability of diagonal systems with one-dimensional input space. *Systems Control Lett.* 55 (2006), no. 4, 321–328.
- [11] B. Jacob, J.R. Partington and S. Pott, Interpolation by vector-valued analytic functions, with applications to controllability. *J. Funct. Anal.* 252 (2007), no. 2, 517–549.
- [12] B. Jacob, J.R. Partington and S. Pott, Tangential interpolation in weighted vector-valued H^p spaces, with applications. *Complex Anal. Oper. Theory* 3 (2009), no. 3, 697–727.
- [13] B. Jacob, J.R. Partington and S. Pott, Weighted interpolation in Paley–Wiener spaces and finite-time controllability. *J. Funct. Anal.* 259 (2010), no. 9, 2424–2436.
- [14] B. Jacob, J.R. Partington and S. Pott, On Laplace–Carleson embedding theorems. *J. Funct. Anal.* 264 (2013), no. 3, 783–814.
- [15] B. Jacob, J.R. Partington and S. Pott, Weighted multiple interpolation and the control of perturbed semigroup systems, *J. Evol. Equ.*, 13 (2013), 395–410.

- [16] J.-L. Lions, *Contrôlabilité exacte, perturbations et stabilisation de systèmes distribués. Tome 1*, Recherches en Mathématiques Appliquées [Research in Applied Mathematics], 8, masson, Paris, 1988.
- [17] Z.-H. Luo, B.-Z. Guo O. and Morgul, *Stability and stabilization of infinite dimensional systems with applications*, Communications and Control Engineering Series, Springer-Verlag, London, 1999.
- [18] J.D. McPhail, A weighted interpolation problem for analytic functions. *Studia Math.* 96 (1990), no. 2, 105–116.
- [19] H.S. Shapiro and A.L. Shields, On some interpolation problems for analytic functions. *Amer. J. Math.* 83 (1961), 513–532.
- [20] O. Staffans, *Well-posed Linear Systems*. Encyclopedia of Mathematics and its Applications, 103. Cambridge University Press, Cambridge, 2005.
- [21] D.A. Stegenga, Multipliers of the Dirichlet space. *Illinois J. Math.* 24 (1980), no. 1, 113–139.
- [22] E.M. Stein, *Topics in Harmonic Analysis related to the Littlewood–Paley Theory*. Annals of Mathematics Studies, No. 63 Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1970.
- [23] M. Tucsnak and G. Weiss, *Observation and Control for Operator Semigroups*. Birkhäuser Advanced Texts: Basler Lehrbücher. Birkhäuser Verlag, Basel, 2009.
- [24] M. Tucsnak and G. Weiss, From exact observability to identification of singular sources, submitted.
- [25] G. Weiss, Admissibility of input elements for diagonal semigroups on l^2 , *Systems Control Lett.* 10 (1988), no. 1, 79–82.
- [26] G. Weiss, A powerful generalization of the Carleson measure theorem? *Open Problems in Mathematical Systems and Control Theory*, 267–272, Comm. Control Engrg. Ser., Springer, London, 1999.
- [27] A. Wynn, Counterexamples to the discrete and continuous weighted Weiss conjectures. *SIAM J. Control Optim.* 48 (2009), no. 4, 2620–2635.
- [28] A. Wynn, α -admissibility of observation operators in discrete and continuous time. *Complex Anal. Oper. Theory* 4 (2010), no. 1, 109–131.