



Deposited via The University of Leeds.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/84039/>

Version: Accepted Version

Proceedings Paper:

Darwin, Z (2013) Assessing and responding to maternal stress (ARMS): antenatal psychosocial assessment in research and practice. In: Journal of Reproductive and Infant Psychology. Society for Reproductive and Infant Psychology, 16-18 Sep 2013, Hull, UK. Taylor & Francis (Routledge), E6 - E6. ISSN: 0264-6838. EISSN: 1469-672X.

<https://doi.org/10.1080/02646838.2014.892345>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Society for Reproductive and Infant Psychology
Annual Doctoral Thesis Award

Author: Darwin, Z.
Thesis Title: **Assessing and Responding to Maternal Stress
(ARMS): Antenatal Psychosocial Assessment in
Research and Practice**

Abstract:

Background: Routine Antenatal Psychosocial Assessment (APA) has been introduced into UK antenatal care; however, the ways in which maternity service providers assess and respond to maternal stress are subject of debate.

Methods: Pregnant women (n=191) completed a questionnaire at booking. Details including mental health assessment and referrals were obtained from their health records following delivery. A sub-sample of women (n=22) experiencing high maternal stress were interviewed longitudinally antenatally and postnatally.

Findings: Maternal stress was common. Women reported its significant impact but struggled to self-assess severity. Mental health history and treatment history were under-disclosed, particularly concerning anxiety. Although high depression and high anxiety symptoms frequently co-occurred, measuring either in isolation missed possible cases. Clinical practice used the Whooley depression case finding questions; these identified only half the possible cases identified using the Edinburgh Postnatal Depression Scale in a research context. Incorporating the Arroll 'help' question improved specificity but substantially compromised sensitivity, missing nine in ten possible cases.

Locally, APA was introduced without attention to topics of relevance to women, context of disclosure or adequate resources for consistently responding to identified need. Women experiencing, or at risk of, mild-moderate mental health disorder in pregnancy were thus usually ineligible for further support.

Implications: Care pathways are needed that encompass both assessing and responding to maternal stress, where communication with health professionals, subsequent referral and management are addressed. The development, implementation and evaluation of low-cost resources embedded in such pathways are a priority and the ARMS research offers a foundation on which to build.