



UNIVERSITY OF LEEDS

This is a repository copy of *The influence of slope and peatland vegetation type on riverine dissolved organic carbon and water colour at different scales*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/84984/>

Version: Accepted Version

Article:

Parry, LE, Chapman, PJ, Palmer, SM et al. (3 more authors) (2015) The influence of slope and peatland vegetation type on riverine dissolved organic carbon and water colour at different scales. *Science of the Total Environment*, 527-52. 530 - 539. ISSN 0048-9697

<https://doi.org/10.1016/j.scitotenv.2015.03.036>

© 2015, Elsevier. Licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International
<http://creativecommons.org/licenses/by-nc-nd/4.0/>

Reuse







Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



 N
 Catchment boundary
  Sample point
  Ericaceous
  Bare peat
  Mixed
  Graminoids
  Sedge
  Shadow