



UNIVERSITY OF LEEDS

This is a repository copy of *Population Changes in a Community of Alkaliphilic Iron-Reducing Bacteria Due to Changes in the Electron Acceptor: Implications for Bioremediation at Alkaline Cr(VI)-Contaminated Sites.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/86539/>

Version: Supplemental Material

Article:

Fuller, SJ, Burke, IT, McMillan, DG et al. (2 more authors) (2015) Population Changes in a Community of Alkaliphilic Iron-Reducing Bacteria Due to Changes in the Electron Acceptor: Implications for Bioremediation at Alkaline Cr(VI)-Contaminated Sites. *Water, Air, and Soil Pollution*, 226 (6). 180 - 180. ISSN 0049-6979

<https://doi.org/10.1007/s11270-015-2437-z>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Supplementary Information for:

Population Changes in a Community of Alkaliphilic Iron Reducing Bacteria Due to Changes in the Electron Acceptor: Implications for Bioremediation at Alkaline Cr(VI) Contaminated Sites

Samuel J. Fuller^a, Ian T. Burke^{b}, Duncan G. G. McMillan^c, Weixuan Ding^d, Douglas I. Stewart^{a*}*

^a School of Civil Engineering, University of Leeds, Leeds, LS2 9JT, UK

^b School of Earth and Environment, University of Leeds, Leeds LS2 9JT, UK

^c University Hospital Jena, Friedrich-Schiller University, Jena, 07743, Germany

^d School of Process, Environmental and Materials Engineering, University of Leeds, Leeds, LS2 9JT, UK

* Corresponding Authors: d.i.stewart@leeds.ac.uk and i.t.burke@leeds.ac.uk.

This section consists of 3 pages containing 2 figures.

Further data on the growth of alkaliphilic iron reducing community in FeCr media




Figure S1. Growth of alkaliphilic Fe(III) reducing community in FeCr medium with initial Cr(VI) concentrations of 100, 150, 500 and 4500 $\mu\text{mol.L}^{-1}$. (A) Cr(VI) concentration with time (B) Total Fe(II) concentration with time (C) cell numbers with time (D) pH with time (data shown is complementary to that shown in Figure 3).




Figure S2: Percentage of bacteria sequences recovered from FeCr8500 medium assigned to each OTU.