



UNIVERSITY OF LEEDS

This is a repository copy of *On asymptotic Boronkov-Sakhanenko inequality with unbounded parameter set*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/87558/>

Version: Accepted Version

Article:

Abu-Shanab, R and Veretennikov, AY (2015) On asymptotic Boronkov-Sakhanenko inequality with unbounded parameter set. *Theory of Probability and Mathematical Statistics*, 90. pp. 1-12. ISSN 0094-9000

<https://doi.org/10.1090/tpms/945>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

ON ASYMPTOTIC BOROVKOV–SAKHANENKO INEQUALITY WITH UNBOUNDED PARAMETER SET

UDC 519.226.3+ 519.233.2

R. ABU-SHANAB AND A. YU. VERETENNIKOV

ABSTRACT. Integral analogues of Cramér–Rao’s inequalities for Bayesian parameter estimators proposed initially by Schützenberger (1958) and later by van Trees (1968) were further developed by Borovkov and Sakhanenko (1980). In the paper, new asymptotic versions of such inequalities are established under ultimately relaxed regularity assumptions and under a locally uniform non-vanishing of the prior density and with \mathbf{R}^1 as a parameter set. Optimality of Borovkov–Sakhanenko’s asymptotic lower bound functional is established.

АНОТАЦІЯ. Інтегральні аналоги нерівностей Крамера–Рао для байєсовських параметричних оцінок, що були запропоновані спочатку Шутценберже (1958) і пізніше ван Трісом (1968), набули подальшого розвитку в роботі Боровкова і Саханенка (1980). В даній статті встановлено нові асимптотичні версії таких нерівностей при найбільш послаблених умовах регулярності, умові локально рівномірної асимптотичної невідродженості апіорної щільності і для параметричної множини \mathbf{R}^1 . Встановлено оптимальність нижньої межі, що задається асимптотичним функціоналом Боровкова–Саханенка.

АННОТАЦИЯ. Интегральные аналоги неравенств Крамера–Рао для байесовских параметрических оценок, предложенные первоначально Шутценберже (1958) и позднее ван Трисом (1968), получили дальнейшее развитие в работе Боровкова и Саханенко (1980). В данной статье установлены новые асимптотические версии таких неравенств при наиболее слабых условиях регулярности, условия локально равномерной невырожденности априорной плотности и с параметрическим множеством \mathbf{R}^1 . Установлена оптимальность нижней границы, представленной асимптотическим функционалом Боровкова–Саханенко.

1. INTRODUCTION

The goal of the paper is to present new versions of Borovkov–Sakhanenko’s inequalities [1, 2] also known as Cramér–Rao (CR) type *integral* inequalities; they may be also interpreted as lower bounds for the Bayesian mean square error. For their asymptotic inequality Borovkov and Sakhanenko required Riemann integrability of the prior density; here only Lebesgue integrability is assumed along with some local non-degeneracy (the latter was not necessary in [1, 2]).

More precisely, first of all for a *finite* sample size n an auxiliary version of Borovkov–Sakhanenko’s inequality (3.3) for unbounded parameter sets under certain relaxed conditions in comparison to [1, 2] (see the Proposition 3.1). Namely, regularity of the prior density—denoted in the sequel by q —will be relaxed; instead, new conditions of nondegeneracy type will be assumed. The inequality (3.3) involves some auxiliary function, h_ε , and practically all technical issues relate to the question of how to choose this function in the best way, at least, asymptotically. Note that *if* the ratio q/I (where I is

2010 *Mathematics Subject Classification*. Primary 62F12, 62F15.

Key words and phrases. Cramér–Rao bounds, Borovkov–Sakhanenko bounds, integral information inequalities, asymptotic efficiency.

The paper is based on a chapter in the first author’s PhD Thesis [5]. Both authors are grateful to Professor Sakhanenko for his comments. The second author’s work was partially supported by RFBR grant 13-01-12447 ofi_m2.

Fisher's information) were from the class C^1 then the most natural choice would have been $h_\varepsilon = q/I$; but if this ratio does not belong to this class, then some approximation and smoothing is required.

Secondly, for large samples an *asymptotic* version of the inequality (3.3) is of high interest and this asymptotic inequality is established also under new assumptions in the Theorem 2.1, which is the main result of the paper. One more version of sufficient assumptions is presented in the Theorem 2.2.

Finally, in the Theorem 2.3 the optimal choice of Borovkov–Sakhanenko's asymptotical functional J (inverse Fisher's information integrated with the prior density) is established. This functional is strictly better—i.e., greater—than the Schützenberger–van Trees functional, although, this remains practically unknown in the literature where asymptotic efficiency is traditionally compared with the latter under the name of van Trees alone; some comments about the names could be found in the last section of the paper.

The importance of the asymptotic Borovkov–Sakhanenko's inequality for the theory may be appreciated after its comparison with the standard definition of asymptotic normality and efficiency. For many natural estimators—including Bayesian and under certain conditions maximum likelihood ones – the asymptotic lower bound provided by Borovkov–Sakhanenko's functional is attained due to their asymptotic normality with a correct limiting variance given by inverse Fisher's information. This is of a certain theoretical importance because it shows that the notion of “asymptotically optimal estimator” may be correctly defined in this way.

The paper consists of Introduction and six further sections: 2—Assumptions and main results; 3—Auxiliary results; 4, 5 and 6—Proofs of three main results; 7—Discussion.

2. ASSUMPTIONS AND MAIN RESULTS

Let us consider a family of probability densities ($f(x | \theta), x \in \mathbf{R}^1$) with respect to Lebesgue's measure, with a parameter $\theta \in \Theta$ where Θ is a domain in \mathbf{R}^1 . In the present paper we tackle the case $\Theta = \mathbf{R}^1$; other unbounded cases may be treated similarly. We assume that there is a *prior density* ($q(\theta), \theta \in \Theta$) and denote $f(x, \theta) := f(x|\theta)q(\theta)$; for the sample $X = (X_1, \dots, X_n)$ of size n of i.i.d. random variables from the distribution $f(\cdot | \theta)$ denote by $L(X | \theta)$ the likelihood function, $L(X | \theta) = \prod_{k=1}^n f(X_k | \theta)$.

Let $\theta_n^*(X)$ denote any estimator of θ where n is the sample size. The quality of the estimator is assessed by the complete mean square error, i.e., by the integral

$$\int (E_\theta(\theta_n^*(X) - \theta)^2) q(\theta) d\theta \equiv \mathbf{E}(\theta_n^* - \theta)^2,$$

where E_θ means expectation with respect to the density $L(x_1, \dots, x_n | \theta)$ and the integration $\int \dots d\theta$ is performed over the support of q , i.e., in our case over the whole line \mathbf{R}^1 (see the assumptions below). The notation \mathbf{E} without the lower index θ is used for the “complete expectation”, i.e., with respect both to X and θ .

Clearly, the estimator depends also on the sample size n and in the asymptotical sense one may be interested in a lower bound for the functional

$$\liminf_{n \rightarrow \infty} n \int (E_\theta(\theta_n^*(X) - \theta)^2) q(\theta) d\theta. \quad (2.1)$$

We assume that the derivative function $\partial L(X | \theta)/\partial \theta$ exists in the classical sense for each X and θ (this may be slightly relaxed to a derivative in L_2 in a usual manner) and that the Fisher information function is finite,

$$I(\theta) = E_\theta \left(\frac{\partial \ln f(X_1 | \theta)}{\partial \theta} \right)^2 < \infty.$$

These standard conditions are a part of the setting and will not be repeated in the main assumptions below.

The problem under consideration is lower bounds for the value (2.1). The **main assumptions** of the paper are as follows.

(A1)

$$0 < J := \int_{-\infty}^{\infty} \frac{q(t)}{I(t)} dt < \infty, \quad \int_{-m}^m \sqrt{I(t)} dt < \infty, \quad \forall m > 0.$$

(A2) For every $m > 0$ there exists $C_m > 0$ such that

$$C_m^{-1} \leq \frac{q(t)}{I(t)} \leq C_m, \quad -m < t < m,$$

and

$$\inf_{t \in [-m, m]} I(t) > 0, \quad \forall m > 0.$$

As a consequence of (A2), $\inf_{t \in [-m, m]} q(t) > 0$, for every $m > 0$.

Remark 1. Note that continuity of I is not required in (A2).

Remark 2. We do not discuss more general cases with a prior density q which may vanish at some points. In such a case certain generalizations look also possible; however, auxiliary constructions would be more involved.

Theorem 2.1. *Let the assumptions (A1) and (A2) be satisfied. Then,*

$$\liminf_{n \rightarrow \infty} n \mathbb{E}(\theta^* - \theta)^2 = \liminf_{n \rightarrow \infty} n \int_{-\infty}^{\infty} E_t(\theta^* - t)^2 q(t) dt \geq J. \quad (2.2)$$

Another version of the assumption (A2) will be used in the next result. Note that under continuity and non-degeneracy of I , Riemann integrability of q/I and the same integrability of q are equivalent; the latter was used in [1, 2].

(A2') Function q/I is *Riemann integrable* on every bounded interval in \mathbf{R}^1 and

$$\inf_{|t| \leq m} q(t) = q_m^- > 0$$

for every $m > 0$ and the function I is continuous.

Theorem 2.2. *Let the assumptions (A1) and (A2') be satisfied. Then the inequality (2.2) holds true.*

Theorem 2.3. *Assume $\int_{-\infty}^{\infty} (q/I)(t) dt < \infty$. Then the optimal choice of h in the maximization problem*

$$\frac{\left(\int_{-\infty}^{\infty} h(t) dt \right)^2}{\int_{-\infty}^{\infty} I(t) \frac{h^2(t)}{q(t)} dt} \rightarrow \sup_{h \in L_1(\mathbf{R})} \quad (2.3)$$

is provided by

$$h = c \frac{q}{I}, \quad \text{with any } c > 0, \quad (2.4)$$

and up to such positive constant multiplier the solution of (2.3) is unique.

Remark 3. The meaning of the last Theorem may be seen from the auxiliary inequality (3.3). Indeed, it follows from this inequality that its right hand side with the term $n^{-1} \int \dots$ and with an arbitrary function h (replacement of h_ε) dropped from the denominator may serve as a lower asymptotic bound for the left hand side. Hence, to achieve the best lower bound one should solve the maximization problem (2.3). So, the Theorem 2.3 implies that the asymptotical Borovkov-Sakhanenko's functional J may not be improved by choosing some better auxiliary function h .

On the other hand, it may be noted that due to strict Jensen's inequality, the value J is, generally speaking, strictly greater than the Schützenberger–van Trees' functional $(\int_{\mathbf{R}} I(t)q(t) dt)^{-1}$.

3. AUXILIARY RESULTS

Let us state two technical results.

Proposition 3.1. *Let $h_\varepsilon(t)$ be a C^1 -smooth function satisfying for any $x = (x_1, \dots, x_n)$*

$$\lim_{t \rightarrow \pm\infty} th_\varepsilon(t)L(x | t) = 0, \quad (3.1)$$

and

$$\int_{-\infty}^{\infty} h_\varepsilon(t) dt < \infty, \quad (3.2)$$

and let $\int_{-m}^m \sqrt{I(t)} dt < \infty$ for any $m > 0$. Then,

$$n \int_{-\infty}^{\infty} E_t(\theta^* - t)^2 q(t) dt \geq \frac{\left(\int_{-\infty}^{\infty} h_\varepsilon(t) dt\right)^2}{\int_{-\infty}^{\infty} I(t) \frac{h_\varepsilon^2(t)}{q(t)} dt + \frac{1}{n} \int_{-\infty}^{\infty} \frac{(h'_\varepsilon(t))^2}{q(t)} dt}. \quad (3.3)$$

Remark 4. Practically all papers on the subject contain one or another version of this inequality, see, e.g., [1, Theorem 30.1]. However, the authors did not succeed to find in earlier works the assumption (3.1), which seems to be necessary for a rigorous presentation in the case of unbounded $\Theta = \mathbf{R}^1$.

Proof. First of all, note that it suffices to prove (3.3) assuming that both the left hand side and the denominator in the right hand side are finite.

The basic identity on which the proof is based reads,

$$\mathbb{E} \left((\theta^*(X) - \theta) \frac{(L(X | \theta)h_\varepsilon(\theta))'_\theta}{L(X, \theta)} \right) = \mathbb{E} \frac{h_\varepsilon(\theta)}{q(\theta)} = \int_{-\infty}^{\infty} h_\varepsilon(t) dt. \quad (3.4)$$

Note that the left hand side in (3.4) is finite due to the Cauchy–Bouniakovsky–Schwarz inequality and the earlier assumption. In turn, (3.4) follows from the following. We have,

$$\mathbb{E} \left((\theta^*(X) - \theta) \frac{(L(X | \theta)h_\varepsilon(\theta))'_\theta}{L(X, \theta)} \right) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (\theta^*(x) - t) (L(x | t)h_\varepsilon(t))'_t dt dx_1 \dots dx_n.$$

Here the multiple integral with respect to $dx_1 \dots dx_n$ is denoted by a single integral symbol. Since this multiple integral converges absolutely, let us consider the internal integral

$$\begin{aligned} \int_{-\infty}^{\infty} (\theta^*(x) - t) (L(x | t)h_\varepsilon(t))'_t dt &= \lim_{M, N \rightarrow \infty} \int_{-M}^N (\theta^*(x) - t) (L(x | t)h_\varepsilon(t))'_t dt \\ &= \lim_{M, N \rightarrow \infty} \left((\theta^*(x) - t) (L(x | t)h_\varepsilon(t)) \Big|_{t=-M}^{t=N} - \int_{-M}^N (\theta^*(x) - t)'_t (L(x | t)h_\varepsilon(t)) dt \right) \\ &= \lim_{M, N \rightarrow \infty} \int_{-M}^N L(x | t)h_\varepsilon(t) dt = \int_{-\infty}^{\infty} L(x | t)h_\varepsilon(t) dt. \end{aligned}$$

We have used explicitly (3.1). Integating with respect to $dx_1 \dots dx_n$, we get (3.4) as required. Now the Cauchy–Bouniakovsky–Schwarz inequality applied to (3.4) gives (3.3). The Proposition 3.1 is proved. \square

Lemma 3.2. *Let the assumption (A1) hold true, and let there exist a sequence*

$$0 \leq q_m(t) \uparrow q(t) \quad (\text{a.e.})$$

as $m \rightarrow \infty$, such that for any estimator θ^* ,

$$\liminf_{n \rightarrow \infty} n \int_{-\infty}^{\infty} E_t(\theta_n^*(X) - t)^2 \tilde{q}_m(t) dt \geq \int_{-\infty}^{\infty} \frac{\tilde{q}_m(t)}{I(t)} dt, \quad (3.5)$$

where

$$\tilde{q}_m(t) = \frac{q_m(t)}{\kappa_m}, \quad \text{and} \quad \kappa_m = \int_{-\infty}^{\infty} q_m(\theta) d\theta.$$

Then,

$$\liminf_{n \rightarrow \infty} n \mathbb{E}(\theta_n^*(X) - \theta)^2 = \liminf_{n \rightarrow \infty} n \int_{-\infty}^{\infty} E_t(\theta_n^*(X) - t)^2 q(t) dt \geq J. \quad (3.6)$$

Proof. Note that without loss of generality, we may assume $\kappa_m > 0$ for all m . The proof of the Lemma follows from the monotone convergence theorem. Indeed, $\kappa_m \rightarrow 1$, $m \rightarrow \infty$, due to the assumption $0 \leq q_m(t) \uparrow q(t)$ (a.e.). So, for any m ,

$$\begin{aligned} \liminf_{n \rightarrow \infty} n \int_{-\infty}^{\infty} E_t(\theta_n^*(X) - t)^2 q(t) dt &\geq \kappa_m \liminf_{n \rightarrow \infty} n \int_{-\infty}^{\infty} E_t(\theta_n^*(X) - t)^2 \tilde{q}_m(t) dt \\ &\geq \kappa_m \int_{-\infty}^{\infty} \frac{\tilde{q}_m(t)}{I(t)} dt = \int_{-\infty}^{\infty} \frac{q_m(t)}{I(t)} dt. \end{aligned}$$

Hence,

$$\liminf_{n \rightarrow \infty} n \int_{-\infty}^{\infty} E_t(\theta_n^*(X) - t)^2 q(t) dt \geq \lim_{m \rightarrow \infty} \int_{-\infty}^{\infty} \frac{q_m(t)}{I(t)} dt = \int_{-\infty}^{\infty} \frac{q(t)}{I(t)} dt = J,$$

as required. The Lemma 3.2 is proved. \square

4. PROOF OF THEOREM 2.1

1. The calculus is based on several approximations—smoothing and truncations—of the function

$$h(t) := \frac{q(t)}{I(t)}, \quad t \in \Theta = \mathbf{R}. \quad (4.1)$$

Note that another method of smoothing was suggested in [7] in relation to some other version of Borovkov–Sakhanenko’s lower bound with the analysis of the approximation errors. First of all, we will approximate q by appropriate q_m and apply the Lemma 3.2. Let

$$q_m(t) := q(t) \mathbb{1}_{\{-m+1 < t < m-1\}}, \quad m > 2.$$

Then, $0 \leq q_m(t) \uparrow q(t)$, as $m \uparrow \infty$. Denote

$$\kappa_m := \int_{-m}^m q_m(\theta) d\theta \quad \text{and} \quad \tilde{q}_m(t) := \frac{q_m(t)}{\kappa_m}.$$

To prove the Theorem, it suffices to show that for every m ,

$$\liminf_{n \rightarrow \infty} n \int_{-m}^m E_t(\theta_n^*(X) - t)^2 \tilde{q}_m(t) dt \geq \int_{-m}^m \frac{\tilde{q}_m(t)}{I(t)} dt. \quad (4.2)$$

Denote

$$S_m := \text{supp}(q_m) = [-m + 1, m - 1],$$

and

$$h_{0,m}(t) := \frac{\tilde{q}_m(t)}{I(t)},$$

and consider the following continuous piece-wise linear function $\varphi = \varphi_{\varepsilon, m}$, with $\varepsilon \leq 1$ and $m > 2$,

$$\varphi_{\varepsilon, m}(t) = \begin{cases} (\varepsilon + 1)t + \varepsilon m, & -\infty < t \leq -m + 1, \\ \left(1 - \frac{\varepsilon}{m-1}\right)t, & -m + 1 \leq t \leq m - 1, \\ (\varepsilon + 1)t - \varepsilon m, & m - 1 \leq t < \infty. \end{cases}$$

Notice that

$$\begin{aligned} \varphi_{\varepsilon, m}(-m) &= -m, & \varphi_{\varepsilon, m}(-m + 1) &= -m + 1 + \varepsilon, \\ \varphi_{\varepsilon, m}(m - 1) &= m - 1 - \varepsilon, & \varphi_{\varepsilon, m}(m) &= m, \\ 0 < \frac{1}{2} &\leq \varphi'_{\varepsilon, m} \leq 2; & \sup_{-\infty < t < \infty} |\varphi'_{\varepsilon, m}(t) - 1| &\rightarrow 0, & \varepsilon \rightarrow 0, \\ & & \sup_{-m \leq t \leq m} |\varphi_{\varepsilon, m}(t) - t| &\rightarrow 0, & \varepsilon \rightarrow 0, \end{aligned}$$

and also

$$\tilde{q}_m(-m + 1-) = \tilde{q}_m(m - 1+) = 0.$$

It follows from the construction of the functions $\varphi_{\varepsilon, m}$ that for any $m > 2$,

$$\sup_{-\infty < v < \infty} \left| 1 - \frac{1}{2\varepsilon} \int_{\varphi_{\varepsilon, m}^{-1}(v-\varepsilon)}^{\varphi_{\varepsilon, m}^{-1}(v+\varepsilon)} dt \right| \rightarrow 0, \quad \varepsilon \rightarrow 0. \quad (4.3)$$

Let

$$h_{\varepsilon, m}(t) := \frac{1}{2\varepsilon} \int_{\varphi_{\varepsilon, m}(t)-\varepsilon}^{\varphi_{\varepsilon, m}(t)+\varepsilon} h_{0, m}(v) dv. \quad (4.4)$$

Since $q_m \equiv 0$ outside $[-m + 1, m - 1]$, then

$$h_{\varepsilon, m}(t) = 0 \quad \text{for } \varepsilon \leq 1 \quad \text{and } |t| \geq m.$$

Hence, the functions $h_{\varepsilon, m}$ with $\varepsilon \leq 1$ satisfy the assumption (3.1).

Moreover, the function $h_{\varepsilon, m}(t)$ defined in (4.4) is absolutely continuous and differentiable almost everywhere, with a.e. (in)equalities,

$$\begin{aligned} h'_{\varepsilon, m}(t) &= \frac{1}{2\varepsilon} \{h_{0, m}(\varphi_{\varepsilon, m}(t) + \varepsilon) - h_{0, m}(\varphi_{\varepsilon, m}(t) - \varepsilon)\}, \\ |h'_{\varepsilon, m}(t)| &\leq \frac{1}{2\varepsilon} \{h_{0, m}(\varphi_{\varepsilon, m}(t) + \varepsilon) + h_{0, m}(\varphi_{\varepsilon, m}(t) - \varepsilon)\}. \end{aligned}$$

Since $q_m \leq q$, $\tilde{q}_m(t) = q_m(t)/\kappa_m$, $q(t)/I(t) \leq C$, and $h_{0, m}(t) = \tilde{q}_m(t)/I(t)$, we get,

$$0 \leq h_{0, m}(t) = \frac{\tilde{q}_m(t)}{I(t)} = \frac{q_m(t)}{I(t)\kappa_m} \leq \frac{q(t)}{I(t)\kappa_m} \leq \frac{C}{\kappa_m}.$$

Therefore, there exists C' such that for every ε small enough, and every m large enough,

$$|h'_{\varepsilon, m}(t)| \leq \frac{C'}{\varepsilon}.$$

The function $h_{\varepsilon, m}(t)$ satisfies all conditions of the Proposition 3.1, so,

$$n \int_{-m}^m E_t(\theta^* - t)^2 \tilde{q}_m(t) dt \geq \frac{\left(\int_{-m}^m h_{\varepsilon, m}(t) dt\right)^2}{\int_{-m}^m I(t) \frac{h_{\varepsilon, m}^2(t)}{\tilde{q}_m(t)} dt + \frac{1}{n} \int_{-m}^m \frac{(h'_{\varepsilon, m}(t))^2}{\tilde{q}_m(t)} dt}. \quad (4.5)$$

2. Let us show that

$$\int_{-m}^m h_{\varepsilon, m}(t) dt \rightarrow \int_{-m}^m h_{0, m}(t) dt, \quad (4.6)$$

and

$$\int_{-m}^m I(t) \frac{h_{\varepsilon, m}^2(t)}{\tilde{q}_m(t)} dt \rightarrow \int_{-m}^m \frac{\tilde{q}_m(t)}{I(t)} dt, \quad \varepsilon \rightarrow 0. \quad (4.7)$$

If we manage to choose ε as a function of n so that the term $\frac{1}{n} \int_{-m}^m \frac{(h'_{\varepsilon,m}(t))^2}{\tilde{q}_m(t)} dt$ vanishes in the limit as $n \rightarrow \infty$, then the assertions (4.6)–(4.7) would imply (4.5) with $h_{\varepsilon,m}$ replaced by h_m , which would allow to apply the Lemma 3.2.

3. Recall that $h_{0,m} \equiv 0$ outside the interval $[-m+1, m-1]$, so that $\int_{-m+\varepsilon}^{m-\varepsilon} h_{0,m}(t) dt = \int_{-m}^m h_{0,m}(t) dt$ for any $\varepsilon \leq 1$. Because of this and by virtue of (4.3) and Lebesgue's dominated convergence theorem and assuming that always $\varepsilon \leq 1$ we have,

$$\begin{aligned} \int_{-m}^m h_{\varepsilon,m}(t) dt &= \int_{-m}^m dt \frac{1}{2\varepsilon} \int_{\varphi_{\varepsilon,m}(t)-\varepsilon}^{\varphi_{\varepsilon,m}(t)+\varepsilon} h_{0,m}(v) dv = \int_{-m+\varepsilon}^{m-\varepsilon} h_{0,m}(v) dv \frac{1}{2\varepsilon} \int_{\varphi_{\varepsilon,m}^{-1}(v-\varepsilon)}^{\varphi_{\varepsilon,m}^{-1}(v+\varepsilon)} dt \\ &= \int_{-m}^m \mathbb{1}_{\{-m+\varepsilon < v < m-\varepsilon\}} h_{0,m}(v) dv \frac{1}{2\varepsilon} \int_{\varphi_{\varepsilon,m}^{-1}(v-\varepsilon)}^{\varphi_{\varepsilon,m}^{-1}(v+\varepsilon)} dt \\ &\rightarrow \int_{-m}^m h_{0,m}(v) dv, \quad \varepsilon \rightarrow 0. \end{aligned}$$

So, the convergence (4.6) holds true.

4. To show (4.7), we notice that

$$\begin{aligned} &\int_{-m}^m \frac{I(t)}{\tilde{q}_m(t)} (h_{\varepsilon,m}^2(t) - h_{0,m}^2(t)) dt \\ &= \int_{-m}^m \frac{I(t)}{\tilde{q}_m(t)} (h_{\varepsilon,m}(t) - h_{0,m}(t)) (h_{\varepsilon,m}(t) + h_{0,m}(t)) dt. \end{aligned}$$

Since the terms $I(t)/\tilde{q}_m(t)$ and $(h_{\varepsilon,m}(t) + h_{0,m}(t))$ are uniformly bounded on S_m , it suffices to establish convergence

$$\int_{-m}^m |h_{\varepsilon,m}(t) - h_{0,m}(t)| dt \rightarrow 0, \quad \varepsilon \rightarrow 0. \quad (4.8)$$

Let $\delta > 0$, and, given m , let us approximate the function $h_{0,m}(t)$ in $L_1[-m, m]$ by some continuous function $h_{0,m}^\delta(t)$, which equals identically zero outside the interval $[-m, m]$, so that

$$\int_{-m}^m |h_{0,m}(t) - h_{0,m}^\delta(t)| dt < \delta.$$

This is, clearly, possible, since the space $C[-m, m]$ is a dense subspace in $L_1[-m, m]$. Then, assuming that $\varepsilon > 0$ is so small that the left hand side in (4.3) does not exceed the value 1 (recall that it actually tends to zero with ε) and denoting

$$h_{\varepsilon,m}^\delta(t) := \frac{1}{2\varepsilon} \int_{\varphi_{\varepsilon,m}(t)-\varepsilon}^{\varphi_{\varepsilon,m}(t)+\varepsilon} h_{0,m}^\delta(v) dv, \quad (4.9)$$

we get,

$$\begin{aligned} \int_{-m}^m |h_{\varepsilon,m}(t) - h_{\varepsilon,m}^\delta(t)| dt &= \int_{-m}^m \left| \frac{1}{2\varepsilon} \int_{\varphi_{\varepsilon,m}(t)-\varepsilon}^{\varphi_{\varepsilon,m}(t)+\varepsilon} (h_{0,m}(v) - h_{0,m}^\delta(v)) dv \right| dt \\ &\leq \int_{-m}^m \frac{1}{2\varepsilon} \int_{\varphi_{\varepsilon,m}(t)-\varepsilon}^{\varphi_{\varepsilon,m}(t)+\varepsilon} |h_{0,m}(v) - h_{0,m}^\delta(v)| dv dt \\ &= \int_{-m}^m |h_{0,m}(v) - h_{0,m}^\delta(v)| dv \frac{1}{2\varepsilon} \int_{\varphi_{\varepsilon,m}^{-1}(v-\varepsilon)}^{\varphi_{\varepsilon,m}^{-1}(v+\varepsilon)} dt \leq 2\delta. \end{aligned}$$

Hence, for the values of ε satisfying the above,

$$\begin{aligned} \int_{-m}^m |h_{\varepsilon,m}(t) - h_{0,m}(t)| dt &\leq \int_{-m}^m |h_{\varepsilon,m}(t) - h_{\varepsilon,m}^\delta(t)| dt + \int_{-m}^m |h_{0,m}(t) - h_{0,m}^\delta(t)| dt \\ &\quad + \int_{-m}^m |h_{\varepsilon,m}^\delta(t) - h_{0,m}^\delta(t)| dt \\ &\leq 2\delta + \delta + \int_{-m}^m |h_{\varepsilon,m}^\delta(t) - h_{0,m}^\delta(t)| dt. \end{aligned}$$

For every fixed $\delta > 0$, the latter integral tends to zero as $\varepsilon \rightarrow 0$, because the function $h_{0,m}^\delta(t)$ is uniformly continuous, and due to convergence

$$\sup_t |h_{\varepsilon,m}^\delta(t) - h_{0,m}^\delta(t)| dt \rightarrow 0, \quad \varepsilon \rightarrow 0,$$

by virtue of (4.3) and (4.9). Therefore, for every $\delta > 0$,

$$\limsup_{\varepsilon \rightarrow 0} \int_{-m}^m |h_{\varepsilon,m}(t) - h_{0,m}(t)| dt \leq 3\delta.$$

However, the left hand side here does not depend on δ , hence, (4.8) holds true, which implies (4.7).

5. From (4.5), (4.6) and (4.7) we conclude that

$$\liminf_{n \rightarrow \infty} n \int_{-m}^m E_t(\theta^* - t)^2 \tilde{q}_m(t) dt \geq \frac{\left(\int_{-m}^m h_{0,m}(t) dt \right)^2}{\int_{-m}^m h_{0,m}(t) dt + \limsup_{n \rightarrow \infty} \frac{1}{n} \int_{-m}^m \frac{(h'_{\varepsilon,m}(t))^2}{\tilde{q}_m(t)} dt}.$$

We estimate,

$$\frac{1}{n} \int_{-m}^m \frac{(h'_{\varepsilon,m}(t))^2}{\tilde{q}_m(t)} dt \leq \frac{1}{n} \int_{-m}^m \frac{C'^2}{\varepsilon^2 \tilde{q}_m(t)} dt = \frac{C'^2}{\varepsilon^2 n} \int_{-m}^m \frac{1}{\tilde{q}_m(t)} dt.$$

For any fixed m , we can choose $\varepsilon = \varepsilon(n) = C'n^{-1/5}$, so that $\lim_{n \rightarrow \infty} 1/(\varepsilon^2 n) = 0$. Hence, we obtain,

$$\liminf_{n \rightarrow \infty} n \int_{-m}^m E_t(\theta^* - t)^2 \tilde{q}_m(t) dt \geq \int_{-m}^m h_{0,m}(t) dt.$$

Since

$$h_{0,m}(t) = \frac{\tilde{q}_m(t)}{I(t)},$$

then due to the Lemma 3.2 the desired asymptotic inequality (2.2) follows. The Theorem 2.1 is proved.

5. PROOF OF THEOREM 2.2

1. Let us denote,

$$q_-^m := \inf_{-m \leq t \leq m} q(t) > 0,$$

see the assumption (A2'). As in the proof of the Theorem 2.1, we will approximate q by an appropriate \tilde{q}_m and apply the Lemma 3.2. Let

$$\begin{aligned} q_m(t) &:= q(t) \mathbb{1}_{\{-m+1 < t < m-1\}}, \quad m > 1, \\ \kappa_m &= \int_{-m}^m q_m(\theta) d\theta \quad \text{and} \quad \tilde{q}_m(t) = \frac{q_m(t)}{\kappa_m}. \end{aligned}$$

To prove the Theorem, it suffices to show that for every m ,

$$\liminf_{n \rightarrow \infty} n \int_{-m}^m E_t(\theta_n^*(X) - t)^2 \tilde{q}_m(t) dt \geq \int_{-m}^m \frac{\tilde{q}_m(t)}{I(t)} dt. \quad (5.1)$$

For the function $\int_{-m}^m (h'_\varepsilon(t))^2 / \tilde{q}_m(t) dt$, the following notation will be used,

$$H_m(\varepsilon) := \int_{-m}^m (h'_\varepsilon(t))^2 / \tilde{q}_m(t) dt.$$

Let

$$h_{0,m}(t) := \tilde{q}_m(t) / I(t),$$

$$\bar{h}_{\varepsilon,m}(t) := \min_{|u| \leq \varepsilon} \frac{\tilde{q}_m(t+u)}{I(t+u)}, \quad \tilde{h}_{\varepsilon,m}(t) := \bar{h}_{\varepsilon,m}(t) \wedge \frac{q_m^m}{\varepsilon}, \quad -m \leq t \leq m,$$

and

$$h_{\varepsilon,m}(t) := \frac{1}{2\varepsilon} \int_{t-\varepsilon}^{t+\varepsilon} \tilde{h}_{\varepsilon,m}(v) dv. \quad (5.2)$$

With this definition, we clearly have

$$\tilde{h}_{\varepsilon,m}(t) \leq h_{0,m}(t), \quad \text{and} \quad 0 \leq h_{\varepsilon,m}(t) \leq h_{0,m}(t). \quad (5.3)$$

Now, the function $h_{\varepsilon,m}$ defined in (5.2) is absolutely continuous and differentiable almost everywhere, with

$$|h'_{\varepsilon,m}(t)| \leq \frac{C\tilde{q}_m(t)}{\varepsilon} \wedge \frac{\tilde{q}_m(t)}{I(t)},$$

and $h_{\varepsilon,m}(-m) = h_{\varepsilon,m}(m) = 0$, for any $\varepsilon > 0$. Due to the assumption (A2'), the function $H_m(\varepsilon)$ is finite, and, moreover,

$$H_m(\varepsilon) \leq \frac{C}{\varepsilon^2} \int_{-m}^m \frac{\tilde{q}_m^2(t)}{\tilde{q}_m(t)} dt \leq \frac{C}{\varepsilon^2}.$$

2. Let us show that

$$\tilde{h}_{\varepsilon,m}(t) \rightarrow h_{0,m}(t), \quad \varepsilon \downarrow 0 \quad (\text{a.e.}) \quad (5.4)$$

For that, due to the Lebesgue dominated convergence theorem, it suffices to show that

$$\int_{-m}^m (h_{0,m}(t) - \tilde{h}_{\varepsilon,m}(t)) dt \downarrow 0, \quad \varepsilon \downarrow 0. \quad (5.5)$$

This follows similarly to [1, Proof of Theorem 30.5], where this hint is applied to the function q . We have, by virtue of the Riemann integrability condition and of the theorem about convergence of Darboux' integral sums,

$$\sum_k \bar{h}_{\delta,m}(2k\delta) 2\delta \rightarrow \int_{-m}^m h_{0,m}(t) dt, \quad \delta \rightarrow 0,$$

$$\sum_k \bar{h}_{\delta,m}((2k+1)\delta) 2\delta \rightarrow \int_{-m}^m h_{0,m}(t) dt, \quad \delta \rightarrow 0.$$

Let us estimate the difference,

$$0 \leq \sum_k (\bar{h}_{\delta,m}(2k\delta) - \tilde{h}_{\delta,m}(2k\delta)) 2\delta \leq 2\delta \sum_k \bar{h}_{\delta,m}(2k\delta) \mathbb{1}_{\{\bar{h}_{\delta,m}(2k\delta) > q_m^m / (2\delta)\}}.$$

Since $h_{0,m}$ is Riemann integrable, it must be bounded on $[-m, m]$, and so is $\bar{h}_{\delta,m} \leq h_{0,m}$. Since $\inf_{t \in [-m, m]} \tilde{q}_m(t) > 0$, then it follows from (A2') that $\tilde{h}_{\delta,m} \equiv \bar{h}_{\delta,m}$ as δ is small enough. Then, of course,

$$\mathbb{1}_{\{\bar{h}_{\delta,m}(2k\delta) > q_m^m / (2\delta)\}} = 0.$$

Therefore, the sum $\sum_k \bar{h}_{\delta,m}(2k\delta) \mathbb{1}_{\{\bar{h}_{\delta,m}(2k\delta) > q_m^m / (2\delta)\}}$ equals zero if δ is small enough. So,

$$0 \leq \sum_k (\bar{h}_{\delta,m}(2k\delta) - \tilde{h}_{\delta,m}(2k\delta)) 2\delta \rightarrow 0, \quad \delta \rightarrow 0.$$

Similarly,

$$0 \leq \sum_k (\bar{h}_{\delta,m}((2k+1)\delta) - \tilde{h}_{\delta,m}((2k+1)\delta))2\delta \rightarrow 0, \quad \delta \rightarrow 0.$$

Hence,

$$\begin{aligned} \int_{-m}^m \tilde{h}_{\varepsilon,m}(t) dt &\geq \left(\sum_k \tilde{h}_{2\varepsilon,m}(4k\varepsilon)2\varepsilon + \sum_k \tilde{h}_{2\varepsilon,m}((4k+2)\varepsilon)2\varepsilon \right) \\ &\rightarrow \int_{-m}^m h_{0,m}(t) dt, \quad \varepsilon \rightarrow 0. \end{aligned} \quad (5.6)$$

Since $\int_{-m}^m \tilde{h}_{\varepsilon,m} \leq \int_{-m}^m h_{0,m}$, the latter convergence implies (5.5). Notice that, strictly speaking, so far we have shown *just* convergence

$$\int_{-m}^m (h_{0,m}(t) - \tilde{h}_{\varepsilon,m}(t)) dt \rightarrow 0, \quad \varepsilon \downarrow 0,$$

which may be monotone or not. But, by construction, the function $\tilde{h}_{\varepsilon,m}$ increases with ε decreasing. This implies (5.5). Hence, (5.4) holds true *almost everywhere* on $-m \leq t \leq m$.

3. Notice that $h_{\varepsilon,m}$ satisfies the assumptions of the Proposition 1, being differentiable and since it vanishes at $-m$ and m . So, we get, with $\varepsilon = (Cn)^{-1/3}$,

$$n \int_{-m}^m E_{\theta}(\theta_n^* - \theta)^2 \tilde{q}_m(\theta) d\theta \geq \frac{\left(\int_{-m}^m h_{\varepsilon,m}(t) dt \right)^2}{\int_{-m}^m I(t) h_{\varepsilon,m}(t)^2 / \tilde{q}_m(t) dt + n^{-1/3}}. \quad (5.7)$$

Hence, to complete the proof, it suffices to establish

$$\int_{-m}^m h_{\varepsilon,m}(t) dt \rightarrow \int_{-m}^m h_{0,m}(t) dt, \quad (5.8)$$

and

$$\int_{-m}^m \frac{I(t) h_{\varepsilon,m}(t)^2}{\tilde{q}_m(t)} dt \rightarrow \int_{-m}^m \frac{\tilde{q}_m(t)}{I(t)} dt \equiv \int_{-m}^m \frac{I}{\tilde{q}_m} h_{0,m}^2 dt, \quad \varepsilon \rightarrow 0. \quad (5.9)$$

4. We have,

$$\begin{aligned} 0 &\leq \int_{-m}^m (h_{0,m}(t) - h_{\varepsilon,m}(t)) dt = \int_{-m}^m \left(h_{0,m}(t) - \frac{1}{2\varepsilon} \int_{t-\varepsilon}^{t+\varepsilon} \tilde{h}_{\varepsilon,m}(v) dv \right) dt \\ &= \int_{-m}^m \frac{1}{2\varepsilon} \int_{t-\varepsilon}^{t+\varepsilon} (h_{0,m}(t) - \tilde{h}_{\varepsilon,m}(v)) dv dt \\ &= \int_{-m}^m \frac{1}{2\varepsilon} \int_{t-\varepsilon}^{t+\varepsilon} (h_{0,m}(t) - \tilde{h}_{\varepsilon,m}(t)) dv dt + \int_{-m}^m \frac{1}{2\varepsilon} \int_{t-\varepsilon}^{t+\varepsilon} (\tilde{h}_{\varepsilon,m}(t) - \tilde{h}_{\varepsilon,m}(v)) dv dt. \end{aligned}$$

Here,

$$\int_{-m}^m \frac{1}{2\varepsilon} \int_{t-\varepsilon}^{t+\varepsilon} (h_{0,m}(t) - \tilde{h}_{\varepsilon,m}(t)) dv dt = \int_{-m}^m (h_{0,m}(t) - \tilde{h}_{\varepsilon,m}(t)) dt \rightarrow 0, \quad \varepsilon \rightarrow 0,$$

due to (5.5). On the other hand,

$$\begin{aligned} &\int_{-m}^m \frac{1}{2\varepsilon} \int_{t-\varepsilon}^{t+\varepsilon} (\tilde{h}_{\varepsilon,m}(t) - \tilde{h}_{\varepsilon,m}(v)) dv dt \\ &= \int \tilde{h}_{\varepsilon,m}(t) dt - \int \tilde{h}_{\varepsilon,m}(v) \left(\frac{1}{2\varepsilon} \int_{v-\varepsilon}^{v+\varepsilon} 1 dt \right) dv = 0. \end{aligned}$$

Thus, indeed, (5.8) holds true.

5. Further, by virtue of (5.3) and (5.8), we also have,

$$\begin{aligned} 0 &\leq \int \frac{I(t)}{\tilde{q}_m(t)} (h_{0,m}^2(t) - h_{\varepsilon,m}^2(t)) dt = \int h_{0,m}^{-1} (h_{0,m}(t) - h_{\varepsilon,m}(t)) (h_{0,m}(t) + h_{\varepsilon,m}(t)) dt \\ &\leq \int h_{0,m}^{-1} (h_{0,m}(t) - h_{\varepsilon,m}(t)) 2h_{0,m}(t) dt = 2 \int (h_{0,m}(t) - h_{\varepsilon,m}(t)) dt \rightarrow 0, \quad \varepsilon \rightarrow 0. \end{aligned}$$

This shows (5.9). Now, from (5.7), (5.8) and (5.9) the desired inequality (4.2) follows. By virtue of the Lemma 1, this finally implies (2.2). The Theorem 2.2 is proved.

6. PROOF OF THEOREM 2.3

1. First of all, let us show that considering h which might change sign *may not* give a better bound. To this aim, we note that the maximization problem (2.3), may be presented as an equivalent maximization problem,

$$\max_{h: \int_{-\infty}^{\infty} I(t) \frac{h^2(t)}{q(t)} dt = 1} \int_{-\infty}^{\infty} h(t) dt. \quad (6.1)$$

It is clear that allowing to change sign for h from positive to negative at some points may not increase the value of the integral $\int_{-\infty}^{\infty} h(t) dt$. In other words, the global maximizer h in the problem (6.1) may be, indeed, only non-negative.

2. For any $h \not\equiv 0$ by the Cauchy–Bouniakovsky–Schwarz inequality we have,

$$\frac{\left(\int_{-\infty}^{\infty} h(t) dt \right)^2}{\int_{-\infty}^{\infty} I(t) \frac{h^2(t)}{q(t)} dt} \leq \int_{-\infty}^{\infty} \frac{q(t)}{I(t)} dt, \quad (6.2)$$

since

$$\left(\int_{-\infty}^{\infty} h(t) dt \right)^2 \leq \int_{-\infty}^{\infty} \frac{q}{I}(t) dt \times \int_{-\infty}^{\infty} I(t) \frac{h^2(t)}{q(t)} dt.$$

On the other hand, if we choose $h = cq/I$ (with any $c > 0$), then

$$\frac{\left(\int_{-\infty}^{\infty} h(t) dt \right)^2}{\int_{-\infty}^{\infty} I(t) \frac{h^2(t)}{q(t)} dt} = \frac{c^2 \left(\int q/I(t) dt \right)^2}{\int (I/q) c^2 (q/I)^2(t) dt} = \int_{-\infty}^{\infty} q(t)/I(t) dt.$$

An equality sign in (6.2) is only possible for the choice of h where $I(t) \frac{h^2(t)}{q(t)} = \text{const} \times q/I$, by virtue of the equality part of the Cauchy–Bouniakovsky–Schwarz inequality. The latter equation implies that necessarily $|h| = cq/I$. Since optimal h should be non-negative, this means that the optimal choice is, indeed, provided by (2.4) and the Theorem 2.3 is proved.

7. DISCUSSION

In this little section some less known issues about the Schützenberger–van Trees’ and Borovkov–Sakhanenko’s lower bounds will be discussed. First of all, for a long time it was a common knowledge that integral CR bounds were introduced in [9]. In fact, more than ten years earlier an estimate of this type with a short but rigorous proof was published in [8]. It may be interesting to note that in [8] CR bounds are called Fréchet–Cramér bounds (see [3]). Hence, it would be more than appropriate to call such bounds, as a minimum, by two names, Schützenberger–van Trees. These bounds are in use in the literature in the issues of asymptotical efficiency of estimators (see, for example, [4]). Nevertheless, as was emphasized earlier in the paper more precise asymptotical bounds are provided by Borovkov–Sakhanenko’s inequality.

Next, note that any limiting assertion becomes more useful if some rate of convergence is established. In this sense, Borovkov–Sakhanenko’s results required some complementary bounds of remainder terms. Under certain additional smoothness, such bounds have been established in [1], [2], [6]. However, even without convergence rate, a limiting assertion may be helpful as such since it shows asymptotical properties of estimators under less restrictive conditions in comparison to what is needed for evaluating errors. Here results like the Theorems 2.1 or 2.2 below may be of some help.

Finally, note that in [10] some development of Borovkov–Sakhanenko’s inequalities was presented for a bounded parameter set. As it turns out, unboundedness of this set requires a lot more technical work and additional assumptions so as to tackle this unboundedness.

REFERENCES

1. A. A. Borovkov, *Mathematical Statistics*, Gordon and Breach, Amsterdam, 1998.
2. A. A. Borovkov and A. I. Sakhanenko, *Estimates for averaged quadratic risk*, Probab. Math. Statist. **1** (1980), no. 2, 185–195. (Russian)
3. M. Fréchet, *Sur l’extension de certaines évaluations statistiques au cas de petits échantillons*, Rev. Inst. Internat. Statist. **11** (1943), 182–205.
4. R. D. Gill and B. Y. Levit, *Applications of the van Trees inequality: a Bayesian Cramer–Rao bound*, Bernoulli **1** (1995), 59–79.
5. Reman Abu-Shanab, *Information Inequalities and Parameter Estimation*, PhD Thesis, University of Leeds, UK, 2009.
6. A. E. Shemyakin, *Rao–Cramér type integral inequalities for the estimates of a vector parameter*, Theory Probab. Appl. **33** (1985) no. 3, 426–434.
7. A. E. Shemyakin, *Rao–Cramér type multidimensional integral inequalities for parametric families with singularities*, Sib. Math. J. (1992), 706–715.
8. M. P. Schützenberger, *A propos de l’inégalité de Fréchet–Cramér*, Publ. Inst. Statist. Univ. Paris **7** (1958), no. 3/4, 3–6; <http://igm.univ-mlv.fr/~berstel/Mps/Travaux/A/1958FrechetInstStat.pdf>
9. H. van Trees, *Detection, Estimation and Modulation Theory*, vol. I, Wiley, New York, 1968.
10. A. Yu. Veretennikov, *On asymptotic information integral inequalities*, Theory of Stochastic Processes **13(29)** (2007), no. 1–2, 294–307.

P.O.Box 32038, DEPARTMENT OF MATHEMATICS, COLLEGE OF SCIENCE, UNIVERSITY OF BAHRAIN, KINGDOM OF BAHRAIN

E-mail address: raboshanab@uob.edu.bh

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LS2 9JT, UNITED KINGDOM & INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS, MOSCOW, RUSSIA & NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, MOSCOW, RUSSIA

E-mail address: a.veretennikov@leeds.ac.uk

Received 01/11/2013