

This is a repository copy of *The perils of taxonomic inconsistency in quantitative palaeoecology: experiments with testate amoeba data.*

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/88139/>

Version: Submitted Version

---

**Article:**

Payne, Richard J., Lamentowicz, Mariusz and Mitchell, Edward A. D. (2011) The perils of taxonomic inconsistency in quantitative palaeoecology: experiments with testate amoeba data. *Boreas*. pp. 15-27. ISSN 0300-9483

<https://doi.org/10.1111/j.1502-3885.2010.00174.x>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

The perils of taxonomic inconsistency in quantitative palaeoecology: experiments with testate amoeba data

RICHARD J. PAYNE, MARIUSZ LAMENTOWICZ AND EDWARD A.D. MITCHELL

Payne, R.J., Lamentowicz, M. & Mitchell, E.A.D.: The perils of taxonomic inconsistency in quantitative palaeoecology: experiments with testate amoeba data

A fundamental requirement of quantitative palaeoecology is consistent taxonomy between a modern training set and palaeoecological data. In this study we assess the possible consequences of violation of this requirement by simulating taxonomic errors in testate amoeba data. Combinations of easily-confused taxa were selected and data manipulated to reflect confusion of these taxa, transfer functions based on unmodified data were then applied to these modified data sets. Initially these experiments were carried out one error at a time using four modern training sets, subsequently multiple errors were separately simulated in both four modern training sets and four palaeoecological datasets. Some plausible taxonomic confusions caused major biases in reconstructed values. In the case of two palaeoecological datasets a single consistent taxonomic error was capable of changing the pattern of environmental reconstruction beyond all recognition, totally removing any real palaeoenvironmental signal. The issue of taxonomic consistency is one which many researchers would rather ignore; our results show that the consequences of this may ultimately be severe.

Keywords: Testate amoebae; Palaeoecology; Transfer Functions; Peatlands; Palaeohydrology; Palaeoclimatology

Richard J. Payne (e-mail: [r.payne@mmu.ac.uk](mailto:r.payne@mmu.ac.uk)), Department of Environmental & Geographical Sciences, Manchester Metropolitan University, Chester Street, Manchester, M1 5GD, United Kingdom, and Geography, School of Environment and Development, The University of Manchester, Oxford Road, Manchester, M13 9PL, United Kingdom; Mariusz Lamentowicz, Department of Biogeography and

35 Palaeoecology, Adam Mickiewicz University, Dziegiełowa 27, 61-680 Poznań,  
36 Poland; Edward Mitchell, WSL, Swiss Federal Institute for Forest, Snow and  
37 Landscape Research, Ecosystem Boundaries Research Unit, Wetlands Research  
38 Group, Station 2, CH-1015 Lausanne, Switzerland, École Polytechnique Fédérale de  
39 Lausanne (EPFL), Laboratory of Ecological Systems, Station 2, CH - 1015 Lausanne,  
40 Switzerland, and Laboratory of Soil Biology, University of Neuchâtel, Rue Emile  
41 Argand 11, CH-2009 Neuchâtel, Switzerland

42

43

44

45 Quantitative palaeoecology generally proceeds by modelling the relationship between  
46 species and an environmental variable in modern environments and then applying this  
47 model to palaeoenvironmental data to produce quantitative estimates of environmental  
48 changes through time. Among the basic requirements of this ‘transfer function’  
49 approach is that ‘the fossil data-sets used for reconstruction purposes should be of  
50 comparable taxonomy and nomenclature... as the modern training set’ (Birks 1995)  
51 i.e. that individuals of the same species are identified consistently and called the same  
52 name in both the modern and palaeoecological data (Belyea 2007). However, there  
53 are good reasons to suppose that this assumption is sometimes violated; human error  
54 is inevitable and in some microfossil groups there is considerable uncertainty  
55 regarding the underlying taxonomy. Such a microfossil group is the testate amoebae, a  
56 group of protists which are abundant in many aquatic to terrestrial ecosystems and  
57 whose solid shells (‘tests’) may be preserved long after death (Fig. 1), allowing  
58 community changes to be tracked through time. Testate amoebae are increasingly  
59 used in palaeoecology, in particular as proxies for hydrological change, and therefore  
60 palaeoclimate, in peatlands (Charman 2001; Mitchell *et al.* 2008).

61 The taxonomy of testate amoebae is not straightforward. Difficulties start with  
62 the problem of applying a biological species concept to micro-organisms which, as far  
63 as we know, overwhelmingly reproduce asexually and for which there are little  
64 genetic data (Schlegel & Meisterfeld 2003). Testate amoeba taxonomy is built around  
65 the concept of morphospecies, that consistent morphological forms represent valid  
66 taxonomic units, at least in the absence of any superior approach (Finlay *et al.* 1996;  
67 Finlay 1998). However there are no biometric data for many morphospecies, leaving  
68 considerable room for personal interpretation of what degree of difference justifies the  
69 erection of new morphospecies and what can simply be considered intraspecific  
70 variability (Medioli *et al.* 1987; Odgen & Meisterfeld 1989). Delineation of species is  
71 further complicated by considerable morphological variability in tests (Heal 1963;  
72 Wanner 1999; Bobrov & Mazei 2004). Testate amoebae can show marked phenotypic  
73 plasticity (Lüftnegger *et al.* 1988; Wanner & Meisterfeld 1994; Wanner 1999) and in  
74 some taxa (adaptive) polymorphism (Schönborn 1992). The test morphology of taxa  
75 which build their shells from particles in their environment (xenosomes) depends on  
76 the available material; large particles may obscure the underlying test morphology  
77 (Ogden 1983). It is probable that many described taxa may just represent extreme  
78 forms of this morphological variability. A difference in taxonomies between

79 'lumpers' and 'splitters' is highly apparent in the literature. For instance the  
80 *Centropyxis constricta* of Medioli & Scott (1983) would probably include 20 or more  
81 species and subspecies considered separable by Chardez (1967).

82 Issues with the differentiation of morphospecies are common to other micro-  
83 organisms (e.g. Mann & Droop 1996; Pawlowski *et al.* 2002). However in the case of  
84 testate amoebae these issues are particularly acute due to the inadequacies of the  
85 taxonomic literature. Unlike for instance freshwater diatom analysis, where the floras  
86 of Krammer & Lange-Bertalot (1986, 1988, 1991a, b) are widely used (at least as a  
87 baseline), there is no 'standard text' for testate amoeba taxonomy. The obscurity of  
88 testate amoebae to many biologists, combined with the general decline in  
89 morphological taxonomic research over recent decades (Lee 2000; Wheeler 2004)  
90 have contributed to the poor state of testate amoeba taxonomy. Those attempting to  
91 apply testate amoeba analysis in ecology and palaeoecology are forced to use a  
92 fragmented body of literature, much of which dates back to the early part of the last  
93 century, and much of which is mutually-contradictory. There are no clear rules for  
94 separating many taxa and few taxonomic keys are available (none of which are  
95 comprehensive and few of which are in English, the *de facto* language of modern  
96 science).

97 In environmental studies using testate amoebae these problems are particularly  
98 serious because of the large number of tests which must be counted; typically at least  
99 100 individuals per sample and 40-50 samples (Payne & Mitchell 2009). This number  
100 of tests pragmatically requires that all identification and counting be carried out using  
101 light microscopy under normal (200x to 400x) magnifications. Many fine taxonomic  
102 distinctions rest on very subtle features which are simply not practicable under these  
103 conditions (e.g. in *Euglypha*: Wylezich *et al.* 2002, *Cyphoderia*: Todorov *et al.* 2009;  
104 Heger *et al.* in press, and *Diffflugia*: Ogden 1983). In palaeoecology problems are  
105 compounded by the loss of diagnostic features. The division between taxa with lobose  
106 and filose pseudopodia is the most fundamental in testate amoebae taxonomy but is  
107 not applicable in palaeoecology. Diagnostic features of the test such as spines may be  
108 lost through taphonomic processes or in sample preparation and tests may become  
109 compressed (Charman *et al.* 2000). Taxonomic schemes used in palaeoecology are  
110 therefore a compromise between practical simplicity and loss of palaeoenvironmental  
111 discernment (Charman *et al.* 2000). Given all these problems it would be little  
112 surprise if there were considerable taxonomic differences among researchers. In the

113 absence of a formal inter-comparison exercise it is impossible to know to what extent  
114 different researchers apply the same name to different taxa or different names to the  
115 same taxon. We can however make observations that: i) The taxonomic literature  
116 lacks clarity. ii) There are considerable differences in the taxonomic resolution  
117 adopted by different studies. iii) Inter-comparison exercises for other microfossil  
118 groups used in Quaternary palaeoecology have shown considerable variability  
119 between different analysts and research groups (Munro *et al.* 1990; Pederson &  
120 Moseholm 1993; Kelly *et al.* 2002; Prygiel *et al.* 2002). For instance, in the diatom  
121 inter-comparison exercise of Kelly *et al.* (2002) some taxa were identified correctly  
122 less than 20% of the time. iv) When researchers are learning testate amoeba taxonomy  
123 several mistakes are consistently made.

124 On the basis of these observations we feel it would be naïve to assume that  
125 taxonomies are identical among all researchers. In this study we attempt to gain an  
126 understanding of the possible implications of taxonomic variability for environmental  
127 reconstructing by simulating possible errors in previously established modern and  
128 palaeoecological datasets.

129

## 130 Methods

131

132 Four modern training sets and four palaeoecological datasets were used in our  
133 experiments. The four modern training sets are all derived from *Sphagnum*-  
134 dominated, mostly ombrotrophic mires and span a considerable region from North  
135 America to western Asia (Table 1). They are: i) Poland, from peatlands of Poland  
136 (Lamentowicz *et al.* 2005, 2007, 2008); ii) Jura, from peatlands in the Jura Mountains  
137 of France and Switzerland (Mitchell *et al.* 1999, 2001); iii) Turkey, from the Sürmene  
138 Ağaçaşlı Yaylası peatland in north-eastern Turkey (Payne *et al.* 2008); and iv)  
139 Alaska, from peatlands in south-central Alaska (Payne *et al.* 2006). The final selected  
140 transfer function models were used in our experiments to infer depth to water table  
141 (DWT; Table 1). The four palaeoecological datasets are: 1. 'Site DLB', a peatland in  
142 sub-Arctic Alaska (Payne *et al.* unpublished, but see Payne & Mitchell 2009); 2. Praz-  
143 Rodet, a peatland in Switzerland (Mitchell *et al.* 2001); 3. Tuchola, a peatland in  
144 Poland (Lamentowicz *et al.* 2008), and 4. Jelenia Wyspa, another peatland in Poland  
145 (Lamentowicz *et al.* 2007). All of these palaeoecological datasets have an applicable  
146 transfer function from the same area (i.e. the Alaska, Jura and Poland training sets,

147 Table 1) which was produced by the same analysts. We are as confident as possible  
148 that these palaeoecological datasets and their respective transfer functions have  
149 consistent taxonomic schemes.

150 A first step in our experiments was to select pairs of species which we  
151 considered could be confused (Table 2). Our combinations were based on three  
152 sources of evidence: i) Our assessment of the distinctiveness of the taxon based upon  
153 the literature, in particular where taxa have been considered inseparable by some  
154 authors. ii) Our observations of the mistakes made by undergraduate and postgraduate  
155 students in learning testate amoeba taxonomy. iii) Our own experience of learning  
156 testate amoeba taxonomy. We produced separate lists of taxon combinations for each  
157 of our training sets, reflecting the differing communities encountered in those studies  
158 and the slightly different taxonomic schemes adopted by the analysts. For simplicity  
159 we refer to each of these taxon combinations as an ‘error combination’, however with  
160 some of these pairings we note that the distinction between the taxa may not always  
161 be clear. We would not claim that our taxon combinations reflect all possible errors or  
162 that all of these errors have a high probability. However, we do feel that our taxon  
163 combinations include all of the most common confusions. Three sets of experiments  
164 were conducted:

#### 165 Individual errors

166 The first group of experiments used only the modern training sets and was designed to  
167 quantitatively investigate the impacts of individual errors on transfer function  
168 predictions. We identified three possible ways in which each pair of species could be  
169 confused: 1. All of taxon A could be recorded as taxon B. 2. All of taxon B could be  
170 recorded as taxon A. 3. The taxa could be switched. The training set data were then  
171 transformed to reflect each of these three types of error for each of the taxon pairs  
172 identified. So for instance with the Alaska data we identified 15 taxon pairs (Table 2),  
173 which could each be transformed in three different ways giving a total of 45 possible  
174 individual modifications to the data. We then applied the transfer function derived  
175 from the original, unmodified training set to each of these modified data-sets in turn to  
176 predict depth to water table (DWT). This approach of applying a transfer function  
177 based on a training set to the same training set but with simulated taxonomic errors is  
178 not representative of any real-world situation but is a useful tool to investigate the  
179 impact that these errors might have on transfer function results.

180 Inferred depth to water table values (termed ‘testate amoeba-inferred depth to  
181 water table’: TI-DWT) were compared to predictions based on the unmodified data  
182 set and residuals calculated ( $\text{TI-DWT}_{\text{original}} - \text{TI-DWT}_{\text{modified}}$ ). Differences between  
183 predictions based on the original and modified data were calculated in terms of root  
184 mean square error (RMSE),  $R^2$  and the maximum difference between predictions for  
185 any one sample (Maximum Bias). All transfer function analyses were carried out  
186 using  $C^2$  (Juggins 2003).

#### 187 Multiple errors

188 To investigate the cumulative impact of more than one error we also carried out  
189 experiments simulating multiple errors in our modern training sets. The same taxon  
190 combinations were used as in the individual errors experiments. A random numbers  
191 system was used to select a taxon pair, with each pair assigned an equal probability of  
192 selection. Where more than two taxa could be confused with each other only one  
193 taxon pair could be selected at a time (where more than one pair were selected the  
194 data were kept unchanged). Each taxon pair could be transformed in one of the three  
195 ways described above with each of these three modifications given an equal  
196 probability of being selected. The number of errors in the data was steadily increased  
197 up to the maximum number of possible changes, with fifteen repetitions for each error  
198 total. The transfer function based on the unmodified training set was then applied to  
199 this modified training set and RMSE,  $R^2$  and Maximum Bias calculated as above.

200 A related possible source of bias in inferred values is that taxonomic errors in  
201 a training set lead to selection of a different transfer function model structure which  
202 may, in itself, lead to differences in model output. To investigate the potential  
203 implications of this issue alternative model structures (WA, WA-Tol, WA-PLS, ML)  
204 were tested using the maximum number of simulated errors in each training set and 15  
205 replicates. The best performing model was selected based on  $\text{RMSEP}_{\text{jack}}$  with no  
206 penalty for model complexity.

#### 207 Errors in palaeoecological sequences

208 To see how the simulated errors might affect palaeoenvironmental inference we also  
209 manipulated the four palaeoecological data-sets and then applied transfer functions  
210 based on unmodified training sets. The same taxon combinations were used when  
211 simulating errors in the palaeoecological data-sets as were used in the two  
212 experiments simulating errors in training sets described above. The number of errors  
213 was successively increased from one to ten. Transfer functions based on the



214 unmodified training set data were applied and TI-DWT values calculated for each  
215 modified palaeoecological data-set.

216

217 Results

218

219 Individual errors

220 Results of individual error experiments are shown in Table 2. With all training sets a  
221 few error combinations have a great deal more impact on predictions than most  
222 others. With the Poland data much the most significant error combination is *Diffflugia*  
223 *globulosa/Cyclopyxis arcelloides*, introducing a mean error of up to 2.5 cm (7% of the  
224 total measured DWT range) depending on which of the three permutations is  
225 considered, the next most important error combination is *Arcella vulgaris/Arcella*  
226 *discoides* (RMSE $\leq$ 0.55 cm, 1.5% measured range). With the Jura data the two most  
227 important error combinations are *Cyclopyxis arcelloides/Phryganella acropodia*,  
228 leading to a mean error of up to 1.95 cm (4% measured range) and *Centropyxis*  
229 *aerophila/Centropyxis platystoma*, leading to a mean error of up to 1.1 cm (2%  
230 measured range). With the Turkey data the most important error combination is  
231 *Corythion dubium/Trinema lineare*, leading to a mean error of up to 1.7 cm (2%  
232 measured range). With the Alaska data the most important error combinations are  
233 *Euglypha ciliata/Euglypha strigosa* (RMSE $\leq$ 3.06 cm, 5% measured range), *Nebela*  
234 *tincta/Nebela penardiana* (RMSE $\leq$ 2.78 cm, 4.6% measured range) and *Heleopera*  
235 *petricola/Heleopera sphagni* (RMSE $\leq$ 2.13 cm, 3.5% measured range). Maximum bias  
236 data show that many of these single errors lead to the predicted TI-DWT values of  
237 some samples changing by more than 10 cm, and in some cases more than 20 cm.  
238 These are highly significant changes; 20 cm represents the DWT difference between a  
239 lawn and a low hummock.

240 Multiple errors

241 When multiple errors are simulated there is a steady increase in the deviation of  
242 predictions from those based on the unmodified data (Fig. 2). With the Alaska data  
243 there is an approximately equal division between samples with TI-DWT over- and  
244 under-predicted relative to the original data. However with the other three data-sets  
245 there is a trend in one direction; with the Poland data this is towards under-prediction  
246 of TI-DWT while with the Jura and Turkey data this is towards over-prediction of TI-  
247 DWT. This directional bias is most apparent with the Jura data with the TI-DWT

248 values of the majority of samples being over-predicted relative to the unmodified data.  
249 These directional biases are largely driven by just a few errors, so with the Jura data  
250 the trend is mostly due to the *N. tinctoria*/*N. parvula* combination, with the Poland data  
251 the trend is mostly due to the *C. arcelloides*/*D. globulosa* combination and with the  
252 Turkey data the trend is mostly attributable to the *C. dubium*/*T. lineare* and *H.*  
253 *petricola* /*H. rosea* combinations.

254 If alternative transfer function model structures are tested using the training sets  
255 with simulated errors a different model structure is selected with 93% of replicates  
256 with the Jura data, 60% of replicates with the Poland data, 40% of replicates with the  
257 Turkey data and in no replicates with the Alaska data.

258

259 Errors in palaeoecological sequences

260 The consequences of these errors for palaeoecological reconstruction are shown in  
261 Fig. 3A-D. With the Site DLB data (Fig. 3A) the most distinct features of the  
262 reconstruction based on unmodified data are pronounced wet phases at the base of the  
263 profile, from 52-56 cm and from 25-28 cm. These wet phases generally remain  
264 apparent even when taxonomic errors are introduced, although with increasing  
265 number of errors the phases become less distinct in some experiments. A notable  
266 change with even one error is a period of higher values between 11 and 15 cm due to  
267 counting *Centropyxis ecornis* as *Centropyxis laevigata*. With the Praz Rodet data (Fig.  
268 3B) simulated errors make relatively little difference to reconstructed values. The  
269 maximum deviation is 7.6 cm but in none of these experiments is the TI-DWT  
270 reconstruction different enough to change interpretation of the record. With the  
271 Tuchola data (Fig. 3C) even a single error can drastically change the pattern of the  
272 reconstruction: If *Cyclopyxis arcelloides* is recorded as *Diffflugia globulosa* it  
273 fundamentally changes the reconstruction giving an overall reduction in predicted  
274 values, introducing a period of rapidly fluctuating values between 20 and 120 cm  
275 depth and adding a trough at 360 cm. Interpretation of these data with and without this  
276 error would be utterly different. Increasing error load slightly increases the variability  
277 of predictions, but the overall pattern is largely determined by whether or not *C.*  
278 *arcelloides* and *D. globulosa* are confused.

279 With the Jelenia Wyspa data (Fig. 3D) the difference that even a single error can  
280 make is even more marked. Again the most important error is recording *C. arcelloides*  
281 as *D. globulosa*. This error leads to a general under-prediction of TI-DWT by 5 cm or

282 more and an almost total difference in the pattern of change. Introducing this error  
283 leads to the reconstruction of major TI-DWT peaks at 42, 95 and 110 cm, features  
284 which are totally absent in the reconstruction based on unmodified data. One of the  
285 most distinctive features of the TI-DWT reconstruction based on the unmodified data  
286 is a period of high values between 50 and 65 cm. However in several experiments  
287 with one or more errors this feature is less distinct or not apparent at all. In these  
288 experiments *Centropyxis cassis* has been recorded as either *Centropyxis platystoma* or  
289 *Centropyxis aerophila*. With increasing number of errors there is an increasing  
290 variability in the pattern of reconstructed change, although reconstructions group  
291 around two basic patterns determined by whether *C. arcelloides/D. globulosa* are  
292 confused or not. In some experiments where both *C. arcelloides/D. globulosa*, and *C.*  
293 *cassis* and *C. aerophila* or *C. platystoma* are confused TI-DWT values deviate from  
294 the unmodified data by more than 17 cm.

295

## 296 Discussion

297 All of our experiments make several important assumptions: they assume that  
298 mistakes are made consistently, that these are all possible errors and all have an equal  
299 probability, and they do not account for tests simply over-looked or mistaken for taxa  
300 not included in the transfer function and therefore excluded. While we acknowledge  
301 that our experiments represent a considerable simplification of the real way in which  
302 taxonomic errors may affect transfer function output the results are undeniably  
303 revealing. While many possible errors make very little difference to predicted values  
304 some possible errors can change predicted values drastically, giving reconstructions  
305 which bear little apparent resemblance to those based on full data.

306 The specific errors which produce major impacts in our experiments seem by  
307 no means improbable. For instance the confusion of *C. dubium* with *T. lineare*  
308 (important in the Turkey training set) and *E. ciliata* with *E. strigosa* (important in the  
309 Alaska training set) are both common mistakes among our students. The most  
310 dramatic illustration of the possible impacts of taxonomic errors in our experiments is  
311 provided by the experiments simulating errors in palaeoecological data sets from  
312 Tuchola and Jelenia Wyspa. Major differences in reconstructions are produced by  
313 confusing *D. globulosa* and *C. arcelloides*, two taxa that have a similar overall  
314 morphology and would probably be grouped by Charman *et al.* (2000) or Medioli &  
315 Scott (1983). The drastic impact that this error makes is particularly notable given the

316 relative scarcity of these taxa in the Tuchola data, constituting only 2.7% of total tests  
317 and only exceeding 5% of count in 5 samples. In the Jelenia Wyspa data the taxa are  
318 slightly more abundant, constituting 10.1% of total tests. The difference that this  
319 single change makes to the reconstructions highlights the extent to which the pattern  
320 of palaeoenvironmental reconstruction may be determined by just a few important  
321 taxa. It is worryingly easy to envisage a scenario where somebody, perhaps relatively  
322 new to testate amoebae palaeoecology and using one of the more agglomerative  
323 taxonomies as their main guide, could make such an error to produce an  
324 environmental reconstruction which is substantially biased, or in the worst case  
325 entirely an artefact of taxonomic inconsistency. Taxonomic errors in a training set  
326 may change the transfer function model structure selected, but it is likely that this  
327 change alone would have limited impact on model output (cf. Booth 2007).

328         The large impacts of some of the simulated errors may suggest the need to  
329 group these potentially problematic taxa in our transfer functions. However these taxa  
330 frequently have significantly differing hydrological optima, therefore a corollary of  
331 the impacts of these errors is that if these taxa are grouped considerable ecological  
332 information will be lost. In the worst case grouping may considerably bias  
333 reconstructions. If one of a pair of taxa is well represented in a training set and the  
334 other not, the ecological optima of the group will mostly match that of the first taxon,  
335 however if the second taxon is more abundant in palaeoecological samples then  
336 reconstructed values will be biased.

337         In the absence of any formal taxonomic inter-comparison it is not possible to  
338 make any definitive assessment of how much of a problem taxonomic inconsistency  
339 may be *in praxis*. We would suggest that these errors are far from implausible.  
340 However, whether or not these specific taxonomic errors are very likely, our results  
341 suggest a wider point, that it is possible for taxonomic errors to radically distort  
342 environmental reconstructions. Taxonomic errors will not necessarily make any  
343 significant difference to environmental reconstruction; indeed, most errors will  
344 probably make very little difference. However, there is the potential for a single  
345 taxonomic mistake made consistently to so change an environmental reconstruction  
346 that the real palaeoecological signal is totally masked. Although our experiments only  
347 consider water table reconstruction in peatlands it is likely that similar results would  
348 be found when considering reconstruction of other variables and in other  
349 environments. Problems may be particularly acute in minerotrophic peatlands where

350 there may be a greater abundance of ‘difficult’ taxa (e.g. genera *Diffflugia* and  
351 *Centropyxis*).

352 Taxonomic comparability is critical; what a palynomorph used in  
353 palaeoecology is called matters little as long as the name is used *consistently*. For  
354 instance, non-pollen palynomorphs are commonly referred to as simply a numbered  
355 ‘type’ as the origin of the palynomorph may not be known (van Geel 2001). Given the  
356 taxonomic limitations imposed by palaeoecological counting some authors have  
357 considered it necessary to use a parallel naming system, for instance Joosten & de  
358 Klerk (2002) have suggested the differentiation of fossil pollen from plant species  
359 (and indeed modern pollen) by referring to the former in SMALL CAPITALS. While we  
360 do not feel that such a system is necessarily required for testate amoebae we would  
361 appeal for clarity in the description of taxonomies used in palaeoecological studies of  
362 testate amoebae. Until a revised taxonomic framework with clear identification  
363 criteria and keys is available and consistently used, researchers publishing training  
364 sets should clearly state identification criteria and the taxa included in groupings  
365 where these are not obvious.

366 Extreme caution should be used when applying transfer functions, particularly  
367 when using training sets counted by different analysts. Researchers attempting to use  
368 a transfer function derived by other analysts should work in close cooperation to  
369 ensure the same identification criteria are consistently employed. In our experience  
370 this is best done by close communication during counting, rather than trying to post-  
371 hoc adjust the taxonomy of a palaeoecological data-set to fit the taxonomy of a  
372 transfer function. Comparison of photographs of difficult taxa between analysts is a  
373 useful approach to ensure this consistency. Where there is any doubt at all over the  
374 criteria for differentiating taxa these taxa should be grouped or excluded from the  
375 data-sets. The fact that extremely large reconstruction errors can be introduced by  
376 relatively modest taxonomic errors adds to the case for comparing testate amoeba-  
377 based records with other data in a multi-proxy approach, and ideally replicating  
378 records with multiple cores. All palaeoecological techniques are imperfect, testate  
379 amoeba analysis is no exception.

380 There appears to be a tendency in testate amoeba-based palaeoecological  
381 reconstruction to use boot-strapping to derive estimates of standard errors and  
382 consider any changes which exceed these error bars (or even do not: Hendon &  
383 Charman 2004) to be a palaeoecological ‘signal’. However, these standard errors only

384 provide an estimate of the error inherent in the model, additional errors may well be  
385 introduced if the transfer function does not provide an adequate fit to the  
386 palaeoecological data (cf. Wilmshurst *et al.* 2003) or taxonomic errors are made. In  
387 our experiments even quite minor taxonomic errors produced a bias that significantly  
388 exceeded the boot-strapped standard errors. Boot-strapped standard errors should be  
389 used with caution as other sources of error can produce biases which considerably  
390 exceed these estimates.

391 To ensure taxonomic consistency there is a need for a common standard  
392 taxonomy which can be applied uniformly among analysts given the constraints  
393 imposed by counting large numbers of sub-fossil tests using optical microscopy. The  
394 guide of Charman *et al.* (2000) is the best attempt at this and is widely used (79  
395 citations in ‘Google Scholar’ at the time of writing). However, the taxonomic scheme  
396 set out has not met with uniform acceptance with many authors either not adopting  
397 this scheme or adapting it to varying extents. Major reasons for this lack of consistent  
398 use may include the exclusion of some relatively common peatland taxa (e.g.  
399 *Euglypha cristata*, *Tracheleuglypha dentata*) and the broad ‘types’ adopted for some  
400 groups of taxa (perhaps most notably the ‘*Cyclopyxis arcelloides* type’). The guide of  
401 Charman *et al.* (2000) provides a first attempt at a difficult task and is a very useful  
402 contribution. However we would argue that now, ten years after publication, is the  
403 time for a reconsideration and refinement of the scheme in an attempt to achieve a  
404 broad consensus. A consistent taxonomy is essential given increasing attempts to  
405 compare and combine modern data-sets while the more widespread use of testate  
406 amoebae in palaeoecology means that more environmental reconstructions are being  
407 produced using transfer functions derived by other researchers. Taxonomic  
408 inconsistency is a neglected issue in biological sciences, but its consequences may  
409 ultimately be very severe (Bortolus 2008).

410

#### 411 Conclusions

412 • Errors of taxonomy and enumeration are inevitable in palaeoecology. Testate  
413 amoeba analysis is likely to be particularly susceptible to such errors due to  
414 the inadequacies of the taxonomy.

415 • Our experiments suggest that some likely confusions can produce significant  
416 biases in quantitative environmental reconstructions.

417 • These results call for improvement of the taxonomic baseline. For now,

418 extreme caution should be used when applying transfer functions and especially  
419 interpreting small changes.

420 • There are many possible causes of bias in environmental reconstructions.  
421 Taxonomic inconsistency is but one of these.

422

423

424

425

426 Acknowledgements- RJP was supported by fellowships from the University of  
427 Manchester and British Institute at Ankara. Funding to EM by Swiss NSF projects no.  
428 205321-109709/1, 205321-109709/2 and CCES projects RECORD and BigLink is  
429 kindly acknowledged. ML was supported by the Foundation for Polish Science (FNP)  
430 (Outgoing Fellowship KOLUMB). We thank Pim van der Knaap and John Birks for  
431 constructive comments on a previous version of the manuscript.

432

433 *'Truth is mighty and will prevail. There is nothing the matter with this, except that it*  
434 *ain't so.'* (Mark Twain)

435

436

437

## 438 References

- 439 Belyea, L.R. 2007: Revealing the Emperor's new clothes: niche-based  
440 palaeoenvironmental reconstruction in the light of recent ecological theory. *The*  
441 *Holocene* 17, 683-688.
- 442 Birks, H.J.B. 1995: Quantitative palaeoecological reconstructions. *In* Maddy, D. &  
443 Brew, S. (eds.): Statistical modelling of Quaternary science data, 161-54. Quaternary  
444 Research Association, Cambridge.
- 445 Bobrov, A.A. & Mazei, Y. 2004: Morphological variability of testate amoebae  
446 (Rhizopoda: Testacealobosea) in natural populations. *Acta Protozoologica* 43, 133–  
447 146.
- 448 Booth, R.K. 2007: Testate amoebae as proxies for mean annual water-table depth in  
449 Sphagnum-dominated peatlands of North America. *Journal of Quaternary Science* 23,  
450 43-57.
- 451 Bortolus, A. 2008: Error cascades in the biological sciences: The unwanted  
452 consequences of using bad taxonomy in ecology. *Ambio* 37, 114-118.
- 453 Chardez, D. 1967: Histoire naturelle des Protozoaires Thecamoebiens. *Les*  
454 *Naturalistes Belges* 48, 484-586.
- 455 Charman, D., Hendon, D. & Woodland, W. 2000: *The identification of testate*  
456 *amoebae (Protozoa: Rhizopoda) in peats*. 147pp. Quaternary Research Association  
457 Technical Guide Series, Cambridge.
- 458 Charman, D.J. 2001: Biostratigraphic and palaeoenvironmental applications of testate  
459 amoebae. *Quaternary Science Reviews* 20, 1753-1764.
- 460 Finlay, B.J. 1998: The global diversity of protozoa and other small species.  
461 *International Journal of Parasitology* 28, 29-48.
- 462 Finlay, B.J., Corliss, J.O., Estban, G. & Fenchel, T. 1996: Biodiversity at the  
463 microbial level: the number of free-living ciliates in the biosphere. *Quarterly Review*  
464 *of Biology* 71, 221-237.
- 465 Heal, O.W. 1963: Morphological variation in certain Testacea (Protozoa: Rhizopoda).  
466 *Archiv für Protistenkunde* 106, 351-368.
- 467 Heger, T.J., Mitchell, E.A.D., Todorov, M., Golemansky, V., Lara, E. & Pawlowski,  
468 J. In press: Phylogenies of euglyphid testate amoebae (Cercozoa: Euglyphida) reveal  
469 rare transitions between the marine supralittoral and the freshwater or terrestrial  
470 environments. *Molecular Phylogenetics and Evolution*.



- 471 Hendon, D. & Charman, D.J. 2004: High-resolution peatland water-table changes for  
472 the past 200 years: the influence of climate and implications for management. *The*  
473 *Holocene* 14, 125-134.
- 474 Joosten, H. & de Klerk, P. 2002: What's in a name? Some thoughts on pollen  
475 classification, identification and nomenclature in Quaternary palynology. *Review of*  
476 *Paleobotany and Palynology* 122, 29-45.
- 477 Juggins, S. 2003: C2 user guide. *Software for ecological and palaeoecological data*  
478 *analysis and visualisation*. University of Newcastle, Newcastle Upon Tyne.
- 479 Kelly, M.G., Bayer, M.M., Hürlimann, J. & Telford, R.J. 2002: Human error and  
480 quality assurance in diatom analysis. In: du Buf, H. & Bayer, M.M. (eds.): Automatic  
481 diatom identification, 75-92. World Scientific Publishing, Singapore.
- 482 Krammer, K. & Lange-Bertalot, H. 1986: *Bacillariophyceae. i: Teil Naviculaceae*.  
483 230pp. Gustav Fischer Verlag, Stuttgart.
- 484 Krammer, K. & Lange-Bertalot, H. 1988: *Bacillariophyceae. ii Teil: Bacillariaceae,*  
485 *Epithemiaceae, Surirellaceae*. 596pp. Gustav Fischer Verlag, Stuttgart.
- 486 Krammer, K. & Lange-Bertalot, H. 1991a: *Bacillariophyceae. iii Teil: Centrales,*  
487 *Fragilariaceae, Eunotiaceae*. 576pp. Gustav Fischer Verlag, Stuttgart.
- 488 Krammer, K. & Lange-Bertalot, H. 1991b: *Bacillariophyceae. iv Teil:*  
489 *Achnantheaceae, kritische Ergänzungen zu Navicula (Lineolatae) und Gomphonema*  
490 *Gesamtliteraturverzeichnis*. 437pp. Gustav Fischer Verlag, Stuttgart.
- 491 Lamentowicz, M. & Mitchell, E.A.D. 2005: The ecology of testate amoebae (Protists)  
492 in Sphagnum in north-west Poland in relation to peatland ecology. *Microbial Ecology*  
493 50, 48-63.
- 494 Lamentowicz, M., Tobolski, K. & Mitchell, E.A.D. 2007: Palaeoecological evidence  
495 for anthropogenic acidification of a kettle-hole peatland in northern Poland. *The*  
496 *Holocene* 17, 1185-1196.
- 497 Lamentowicz, M., Obremaska, M. & Mitchell, E.A.D. 2008: Autogenic succession,  
498 land-use change, and climatic influences on the Holocene development of a kettle  
499 hole mire in Northern Poland. *Review of Paleobotany and Palynology* 151, 21-40.
- 500 Lee, M.S.Y. 2000: A worrying systematic decline. *Trends in Ecology and Evolution*  
501 15, 346.
- 502 Lüftnegger, G., Petz, W., Berger, H., Foissner, W. & Adam, H. 1988: Morphologic  
503 and biometric characterization of twenty-four soil testate amoebae (Protozoa,  
504 Rhizopoda). *Archiv für Protistenkunde* 136, 153-189.

- 505 Mann, D.G. & Droop, S.J.M. 1996: Biodiversity, biogeography and conservation of  
506 diatoms. *Hydrobiologia* 336, 19-32.
- 507 Medioli, F.S. & Scott, D.B. 1983: Holocene Arcellacea (Thecamoebians) from  
508 Eastern Canada. 63pp. Cushman Foundation for Foraminiferal Research, Special  
509 Publication 21, Washington DC.
- 510 Medioli, F.S., Scott, D.B. & Abbott, B.H. 1987: A case study of protozoan intraclonal  
511 variability: Taxonomic implications. *Journal of Foraminiferal Research* 17, 28-47.
- 512 Mitchell, E.A.D., Buttler, A.J., Warner, B.G. & Gobat, J.M. 1999: Ecology of testate  
513 amoebae (Protozoa: Rhizopoda) in Sphagnum peatlands in the Jura mountains,  
514 Switzerland and France. *Ecoscience* 6, 565-576.
- 515 Mitchell, E.A.D., van der Knaap, W.O., van Leeuwen, J.F.N., Buttler, A., Warner,  
516 B.G. & Gobat, J.M. 2001: The palaeoecological history of the Praz-Rodet bog (Swiss  
517 Jura) based on pollen, plant macrofossils and testate amoebae (Protozoa). *The*  
518 *Holocene* 11, 65-80.
- 519 Mitchell, E.A.D., Charman, D.J. & Warner, B.G. 2008: Testate amoebae analysis in  
520 ecological and paleoecological studies of wetlands: past, present and future.  
521 *Biodiversity and Conservation* 17, 2115-2137.
- 522 Munro, M.A.R., Kreiser, A.M., Battarbee, R.W., Juggins, S., Stevenson, A.C.,  
523 Anderson, D.S., Anderson, N.J., Berge, F., Birks, H.J.B., Davis, R.B., Flower, R.J.,  
524 Fritz, S.C., Haworth, E.J., Jones, V.J., Kingston, J.C. & Renberg, I. 1990: Diatom  
525 quality control and data handling. *Philosophical Transactions of the Royal Society of*  
526 *London B: Biological Sciences* 327, 257- 261.
- 527 Ogden, C.G. & Meisterfeld, R. 1989: The taxonomy and systematics of some species  
528 of Cucurbitella, Diffflugia and Netzelia (Protozoa: Rhizopoda); with an evaluation of  
529 diagnostic characters. *European Journal of Protistology* 27, 109-128.
- 530 Ogden, C.G. 1983: Observations on the systematics of the genus Diffflugia in Britain  
531 (Rhizopoda, Protozoa). *Bulletin of the British Museum of Natural History (Zoology)*  
532 44, 1-73.
- 533 Pawlowski, J., Fahrni, J.F., Brykczynska, U., Habura, A. & Bowser, S.S. 2002:  
534 Molecular data reveal high taxonomic diversity of allogromiid foraminifera in  
535 Explorers Cove (McMurdo Sound, Antarctica). *Polar Biology* 25, 96-105.
- 536 Payne, R., Kishaba, K., Blackford, J. & Mitchell, E. 2006: The ecology of testate  
537 amoebae in southcentral Alaskan peatlands: Building transfer function models for  
538 palaeoenvironmental inference. *The Holocene* 16, 403-414.


- 539 Payne, R., Charman, D., Matthews, S. & Eastwood, W. 2008: Testate amoebae as  
540 palaeoclimate proxies in Sürmene Ağaçaş Yaylasi peatland (Northeast Turkey).  
541 *Wetlands* 28, 311-323.
- 542 Payne R.J. & Mitchell, E.A.D. 2009: How many is enough? Determining adequate  
543 count totals for ecological and palaeoecological studies of testate amoebae. *Journal of*  
544 *Paleolimnology* 42, 483-495.
- 545 Pedersen, B. & Moseholm, L. 1993: Precision of the daily pollen count. Identifying  
546 sources of variation using variance component models. *Aerobiologia* 9, 15-26.
- 547 Prygiel, J., Carpentier, P., Almeida, S., Coste, M., Druart, J.-C., Ector, L., Guillard,  
548 D., Honore, M-A., Iserentant, R., Ledeganck, P., Lalanne-Cassou, C., Lesniak, C.,  
549 Mercier, I., Moncaut, P., Nazart, M., Nouchet, N., Peres, F., Peeters, V., Rimet, F.,  
550 Rumeau, A., Sabater, S., Straub, F., Torrisi, M., Tudesque, L., Van de Vijver, B.,  
551 Vidal, H., Vizinet, J. & Zydek, N. 2002: Determination of the biological diatom index  
552 (IBD NF T 90–354): Results of an intercomparison exercise. *Journal of Applied*  
553 *Phycology* 14, 27–39.
- 554 Schönborn, W. 1992: Adaptive polymorphism in soil-inhabiting testate amebas  
555 (Rhizopoda) - its importance for delimitation and evolution of asexual species. *Archiv*  
556 *für Protistenkunde* 142, 139-155.
- 557 Schlegel, M. & Meisterfeld, R. 2003: The species problem in protozoa revisited. *European*  
558 *Journal of Protistology* 39, 349-355.
- 559 Todorov, M., Golemanski, V., Mitchell, E.A.D. & Heger, T.J. 2009: Morphology,  
560 biometry and taxonomy of five freshwater and marine interstitial testate amoebae of  
561 the genus *Cyphoderia* (Testaceafilosia: Cyphoderiidae). *Journal of Eukaryotic*  
562 *Microbiology* 56, 279-289.
- 563 Van Geel, B. 2001: Non-pollen palynomorphs, In Smol, J.P., Birks, H.J.B. & Last,  
564 W.M. (eds.): *Tracking Environmental Change Using Lake Sediments. Volume 3:*  
565 *Terrestrial, Algal, and Siliceous Indicators*, 99-119. Kluwer Academic Publishers,  
566 Dordrecht.
- 567 Wanner, M. 1999: A review on the variability of testate amoebae: Methodological  
568 approaches, environmental influences and taxonomical implications. *Acta*  
569 *Protozoologica* 38, 15–29.

- 570 Wanner, M. & Meisterfeld, R. 1994: Effects of some environmental factors on the  
571 shell morphology of testate amoebae (Rhizopoda, Protozoa). *European Journal of*  
572 *Protistology* 30, 191–195.
- 573 Wheeler, Q.D. 2004: Taxonomic triage and the poverty of phylogeny. *Philosophical*  
574 *Transactions of the Royal Society of London B: Biological Sciences* 359, 571-583.
- 575 Wilmshurst, J., Wisser, S. & Charman, D. 2003: Reconstructing Holocene water tables  
576 in New Zealand using testate amoebae: differential preservation of tests and  
577 implications for the use of transfer functions. *The Holocene* 13, 61-72.
- 578 Wylezich, C., Meisterfeld, R., Meisterfeld, S. & Schlegel, M. 2002: Phylogenetic analyses of  
579 small subunit ribosomal RNA coding regions reveal a monophyletic lineage of euglyphid  
580 testate amoebae (order Euglyphida). *Journal of Eukaryotic Microbiology* 49, 108-118.
- 581
- 582

## 583 FIGURES and TABLES

584


585 Figure 1. Illustrations of selected testate amoeba taxa discussed in this paper. A.  
 586 *Nebela tinctoria* var. *major*. B. *N. tinctoria* var. *major* and *N. tinctoria*. C. *N. marginata*. D. *N.*  
 587 *carinata*. E. *N. tinctoria* var. *major*. F. *N. flabellulum*. G. *N. penardiana*. H. *N. militaris*.  
 588 I. *Centropyxis aerophila*. J. *C. aerophila* var. *sphagnicola*. K. *C. ecornis*. L. *C.*  
 589 *laevigata*. M. *C. platystoma*. N. *Phryganella acropodia*. O. *Diffflugia globulosa*. P.  
 590 *Corythion dubium*. Q & R. *Trinema lineare*. S. *Euglypha ciliata*. T. *E. compressa*. U.  
 591 *E. strigosa*. Scale bar is 20µm for P,Q and R, 50µm for others.



592

593


594 Figure 2. Results of multiple error experiments (see Methods) with four modern  
 595 training sets. Plots A-D show residuals ( $TI-DWT_{original} - TI-DWT_{modified}$ ), plots E-H  
 596 show the same data presented as an overall mean TI-DWT deviation. Box plots show  
 597 median (central line), first and third quartiles (grey box), tenth and ninetieth  
 598 percentiles ('whiskers') and fifth and ninety-fifth percentiles (dots).




599

600 Figure 3. Results of errors in palaeoecological sequences experiments (see Methods)  
 601 with palaeoecological data from A) 'Site DLB', Alaska, B) Praz-Rodet, Swiss Jura,  
 602 C) Tuchola, Poland, and D) Jelenia Wyspa, Poland. For each dataset the plot on the  
 603 upper left shows reconstruction based on unmodified data and the adjacent plot shows  
 604 percentage of tests contributed by the taxa which could be confused. Other plots show  
 605 reconstructions for increasing number of errors from 1-10 with fifteen cycles of  
 606 random re-selection for each error total.

607





609  
610  
611  
612  
613  
614

Table 1.

Attributes of the datasets used in this study showing number of samples ( $n$ ), and for modern training sets: transfer function model structure, jack-knifed root mean square error of prediction (RMSEP), Maximum Bias and  $R^2$ . Location given in parentheses after palaeoecological data set name indicates applicable transfer function.

Location	$n$	Model structure	RMSEP <sub>jack</sub> (cm)	Max Bias <sub>jack</sub> (cm)	$R^2_{jack}$	Reference
Modern training sets:						
Poland	84	WA-Tol, Inverse deshrinking	4.6	9.0	0.71	Lamentowicz <i>et al.</i> (2007)*
Jura	37	WA-PLS (2 component)	8.0	21	0.62	Mitchell <i>et al.</i> (1999, 2001) <sup>†</sup>
Turkey	42	ML	7.1	21	0.81	Payne <i>et al.</i> (2008)



Alaska	91	WA-PLS (2 component)	9.7	14	0.55	Payne <i>et al.</i> (2006)
Palaeoecological data sets:						
Site DLB (Alaska)	71					Payne <i>et al.</i> (unpublished)
Praz Rodet (Jura)	57					Mitchell <i>et al.</i> (2001)
Tuchola (Poland)	50					Lamentowicz <i>et al.</i> (2008)
Jelenia Wyspa (Poland)	38					Lamentowicz <i>et al.</i> (2007)

\*Values slightly different from published due to re-calculation of percentages. †Re-calculated using WA-PLS, see Payne and Mitchell (2009).

615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635

Table 2. Results of individual error experiments (Methods section 1) for A) Poland (Lamentowicz *et al.* 2007), B) Jura (Mitchell *et al.* 1999, 2001), C) Turkey (Payne *et al.* 2008), D) Alaska (Payne *et al.* 2006). Showing, taxon pair (A and B), percentage of total tests these taxa represent, number of occurrences of each taxon (N), DWT optima estimated by weighted averaging ('WA Optima') and impact of simulated errors in terms of RMSE, maximum bias and  $R^2$  between TI-DWT based on original and modified datasets. Each taxon pair could be changed in three ways: all of taxon A could be counted as taxon B (A→B), all of taxon B could be counted as taxon A (A←B), and the two taxa could be switched (A↔B).

#### A) Poland

Taxon A	Taxon B	% total		N		WA optima		RMSE		
		A	B	A	B	A	B	A→B	A←B	A↔B
<i>Corythion dubium</i>	<i>Corythion-Trinema</i> type	0.80	0.03	13	4	23.08	20.90	0.03	0.00	0.03
<i>Cyclopyxis arcelloides</i>	<i>Diffugia globulosa</i>	3.63	1.74	33	6	4.36	-0.18	2.33	0.28	2.49
<i>Nebela parvula</i>	<i>Nebela tincta</i>	1.37	2.40	32	33	19.04	21.59	0.04	0.08	0.08
<i>Nebela bohémica</i>	<i>Nebela collaris</i>	2.49	0.12	24	6	11.60	19.72	0.19	0.02	0.20
<i>Nebela militaris</i>	<i>Nebela collaris</i>	1.21	0.12	15	6	25.11	19.72	0.12	0.01	0.11
<i>Heleopera sphagni</i>	<i>Heleopera petricola</i>	0.42	1.56	15	31	13.29	13.02	0.00	0.01	0.01
<i>Heleopera sylvatica</i>	<i>Heleopera petricola</i>	0.16	1.56	5	31	20.10	13.02	0.01	0.05	0.06
<i>Euglypha strigosa</i>	<i>Euglypha compressa</i>	0.25	0.43	10	11	19.75	6.92	0.11	0.06	0.17
<i>Euglypha compressa</i>	<i>Euglypha ciliata</i>	0.43	0.41	11	8	6.92	6.51	0.02	0.02	0.02
<i>Euglypha ciliata</i>	<i>Euglypha strigosa</i>	0.41	0.25	8	10	6.51	19.75	0.40	0.05	0.07
<i>Centropyxis cassis</i>	<i>Centropyxis aerophila</i>	0.27	0.07	5	3	13.98	7.41	0.03	0.03	0.07
<i>Centropyxis aerophila</i>	<i>Centropyxis platystoma</i>	0.07	0.03	3	2	7.41	8.68	0.00	0.00	0.00
<i>Centropyxis cassis</i>	<i>Centropyxis platystoma</i>	0.27	0.03	5	2	13.98	8.68	0.05	0.01	0.05
<i>Amphitrema stenostoma</i>	<i>Amphitrema wrightianum</i>	0.11	0.65	5	5	0.08	0.06	0.01	0.06	0.06
<i>Arcella artocrea</i>	<i>Arcella catinus</i>	0.03	3.05	4	35	11.64	15.08	0.00	0.15	0.15
<i>Arcella discoides</i>	<i>Arcella vulgaris</i>	7.58	2.20	33	17	1.36	3.15	0.43	0.16	0.55
<i>Arcella gibbosa</i>	<i>Arcella hemispherica</i>	0.59	0.59	6	5	0.77	-0.23	0.02	0.02	0.05

636

637

## B) Jura

Taxon A	Taxon B	% total		N		WA optima		RMSE		
		A	B	A	B	A	B	A→B	A←B	A↔B
<i>Arcella artocrea</i>	<i>Arcella catinus</i>	0.10	1.64	7	19	13.16	26.33	0.06	0.88	0.92
<i>Centropyxis aerophila</i>	<i>Centropyxis platystoma</i>	2.10	0.95	17	8	17.17	23.31	1.10	0.50	1.07
<i>Corythion dubium</i>	<i>Trinema</i> type	5.31	3.70	33	20	24.97	26.38	0.49	0.34	0.36
<i>Cyclopyxis arcelloides</i>	<i>Diffugia globulosa</i>	0.55	0.24	7	1	11.12	3.00	0.02	0.01	0.03
<i>Cyclopyxis arcelloides</i>	<i>Phryganella acropodia</i>	0.55	2.99	7	28	11.12	28.25	0.32	1.76	1.95
<i>Diffugia longicollis</i>	<i>Diffugia oblonga</i>	0.37	0.02	3	1	27.35	16.00	0.26	0.01	0.27
<i>Euglypha alveolata</i>	<i>Euglypha tuberculata</i>	0.01	0.01	1	1	41.00	8.00	0.02	0.01	0.03
<i>Euglypha ciliata</i>	<i>Euglypha compressa</i>	2.08	0.29	31	8	21.66	26.25	0.72	0.10	0.69
<i>Euglypha ciliata</i>	<i>Euglypha strigosa</i>	2.08	1.04	31	19	21.66	25.78	0.30	0.15	0.27
<i>Euglypha laevis</i>	<i>Euglypha rounda</i>	1.66	2.62	22	24	24.24	24.75	0.27	0.42	0.47
<i>Euglypha strigosa</i>	<i>Euglypha compressa</i>	1.04	0.29	19	8	25.78	26.25	0.21	0.06	0.22
<i>Heleopera petricola</i>	<i>Heleopera rosea</i>	2.47	2.82	27	22	26.90	26.04	0.29	0.33	0.52
<i>Nebela bohemica</i>	<i>Nebela collaris</i>	0.72	0.23	6	5	20.68	23.20	0.13	0.04	0.09
<i>Nebela carinata</i>	<i>Nebela marginata</i>	0.18	0.91	5	9	8.82	9.59	0.01	0.05	0.05
<i>Nebela militaris</i>	<i>Nebela collaris</i>	6.62	0.23	30	5	27.85	23.20	0.81	0.03	0.83
<i>Nebela parvula</i>	<i>Nebela tincta</i>	0.04	14.68	2	37	29.35	29.29	0.01	5.87	5.86
<i>Nebela penardiana</i>	<i>Nebela tubulosa</i>	0.42	0.69	8	8	19.12	16.41	0.12	0.20	0.23
<i>Phryganella acropodia</i>	<i>Diffugia globulosa</i>	2.99	0.24	28	1	28.25	3.00	1.88	0.15	2.00
<i>Sphenoderia lenta</i>	<i>Tracheleuglypha dentata</i>	0.13	0.81	5	13	17.01	23.01	0.04	0.25	0.21

638

639

## C) Turkey

Species A	Species B	% total		N		WA optima		RMSEP		
		A	B	A	B	A	B	A→B	A←B	A↔B
<i>Phryganella acropodia</i>	<i>Cyclopyxis arcelloides</i>	1.04	0.27	22	3	39.74	9.34	0.03	0.00	0.03
<i>Cyclopyxis eurystoma</i>	<i>Phryganella acropodia</i>	0.84	1.04	8	22	68.28	39.74	0.22	0.35	0.37
<i>Cyclopyxis arcelloides</i>	<i>Cyclopyxis eurystoma</i>	0.27	0.84	3	8	9.34	68.28	0.55	0.17	0.72
<i>Corythion dubium</i>	<i>Trinema lineare</i>	8.24	1.41	31	13	47.40	63.76	1.65	0.35	1.59
<i>Euglypha compressa</i>	<i>Euglypha ciliata</i>	0.12	0.49	5	15	25.39	48.87	0.01	0.13	0.12
<i>Euglypha strigosa</i>	<i>Euglypha compressa</i>	0.07	0.12	4	5	30.29	25.39	0.01	0.01	0.01
<i>Euglypha strigosa</i>	<i>Euglypha ciliata</i>	0.07	0.49	4	15	30.29	48.87	0.00	0.03	0.03
<i>Heleopera rosea</i>	<i>Heleopera petricola</i>	3.45	0.08	27	2	41.03	28.59	0.90	0.01	0.90
<i>Nebela penardiana</i>	<i>Nebela tubulosa</i>	0.03	0.03	2	2	29.63	29.46	0.00	0.00	0.00
<i>Nebela tincta</i>	<i>Nebela penardiana</i>	0.47	0.03	14	2	43.69	29.63	0.01	0.00	0.01
<i>Centropyxis aerophila</i> type	<i>Plagiopyxis cf. callida</i>	2.33	0.06	20	2	57.28	12.62	0.38	0.01	0.38

640

641

642

## D) Alaska

Taxon A	Taxon B	% total		N		WA optima		RMSE		
		A	B	A	B	A	B	A→B	A←B	A↔B
<i>Arcella arenaria</i>	<i>Arcella artocrea</i>	2.02	0.10	58	4	35.79	30.92	0.44	0.02	0.46
<i>Centropyxis ecornis</i>	<i>Centropyxis laevigata</i>	0.76	1.26	19	20	28.35	44.19	0.48	0.80	1.28
<i>Centropyxis aerophila</i>	<i>Centropyxis platystoma</i>	3.05	0.12	38	5	26.43	28.06	0.95	0.04	0.93
<i>Corythion dubium</i>	<i>Trinema</i> spp.	4.81	0.96	48	33	31.44	29.41	1.32	0.26	1.10
<i>Diffugia globulosa</i>	<i>Phryganella acropodia</i> type	0.15	6.89	3	85	19.59	34.72	0.01	0.29	0.29
<i>Euglypha ciliata</i>	<i>Euglypha compressa</i>	4.95	0.83	67	28	35.76	37.60	0.78	0.13	0.75
<i>Euglypha ciliata</i>	<i>Euglypha strigosa</i>	4.95	0.23	67	11	35.76	23.47	3.06	0.14	2.97
<i>Euglypha strigosa</i>	<i>Euglypha compressa</i>	0.23	0.83	11	28	23.47	37.60	0.18	0.64	0.82
<i>Heleopera petricola</i>	<i>Heleopera sylvatica</i>	3.84	0.31	43	12	32.45	33.42	0.57	0.05	0.58
<i>Heleopera petricola</i>	<i>Heleopera sphagni</i>	3.84	3.74	43	33	32.45	24.39	1.17	1.14	2.13
<i>Nebela penardiana</i>	<i>Nebela marginata</i>	0.06	0.33	3	6	18.27	18.35	0.02	0.10	0.09
<i>Nebela tincta</i>	<i>Nebela penardiana</i>	3.25	0.06	60	3	42.25	18.27	2.74	0.05	2.78
<i>Hyalosphenia elegans</i>	<i>Nebela militaris</i>	3.98	1.76	47	40	32.03	46.80	2.59	1.15	2.71
<i>Euglypha rotunda</i>	<i>Tracheleuglypha dentata</i>	1.15	0.03	32	3	31.69	14.52	0.74	0.02	0.73
<i>Tracheleuglypha dentata</i>	<i>Sphenoderia lenta</i>	0.03	0.35	3	12	14.52	20.68	0.00	0.04	0.04

643

644

645

646  
647  
648  
649  
650  
651  
652