



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/88619/>

Version: Accepted Version

Article:

Novara, G., Catto, J.W.F., Wilson, T. et al. (2015) Systematic review and cumulative analysis of perioperative outcomes and complications after robot-assisted radical cystectomy. *European Urology*, 67 (3). 376 - 401. ISSN: 0302-2838

<https://doi.org/10.1016/j.eururo.2014.12.007>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Systematic Review and Cumulative Analysis of Perioperative Outcomes and Complications After Robot-assisted Radical Cystectomy

Giacomo Novara ^{a,*}, James W.F. Catto ^b, Timothy Wilson ^c, Magnus Annerstedt ^d, Kevin Chan ^c, Declan G. Murphy ^e, Alexander Motttrie ^f, James O. Peabody ^g, Eila C. Skinner ^h, Peter N. Wiklund ⁱ, Khurshid A. Guru ^{j,†}, Bertram Yuh ^{c,†}

^a Department of Surgery, Oncology, and Gastroenterology - Urology Clinic

University of Padua, Italy

^b Academic Urology Unit, University of Sheffield, Sheffield, UK

^c City of Hope National Cancer Center Duarte, CA, USA

^d Department of Urology, Herlev University Hospital, Denmark

^e Division of Cancer Surgery, Peter MacCallum Cancer Centre, St. Andrews Place, East Melbourne, Victoria, Australia

^f Department of Urology, Onze-Lieve-Vrouw Hospital, Aalst, Belgium

^g Vattikuti Urology Institute, Henry Ford Health System, Detroit, MI, USA

^h Department of Urology, Stanford University, Stanford, CA, USA

ⁱ Karolinska University Hospital, Urology, Stockholm, Sweden

^j Department of Urology, Roswell Park Cancer Institute, Buffalo, NY, USA

[†] Contributed equally.

***Corresponding author.** Department of Surgery, Oncology, and Gastroenterology - Urology Clinic, University of Padua, Via Giustiniani 2, 35100 - Padua, Italy. Tel.: +39 049 8211250; Fax: +39 049 8218757.

E-mail: giacomonovara@gmail.com; giacomo.novara@unipd.it

Keywords: Radical cystectomy, Robotics, Robotic radical cystectomy, Laparoscopic radical cystectomy

Abstract

Context: Although open radical cystectomy (ORC) is still the standard approach, laparoscopic radical cystectomy (LRC) and robot-assisted radical cystectomy (RARC) have gained popularity.

Objective: To report a systematic literature review and cumulative analysis of perioperative outcomes and complications of RARC in comparison with ORC and LRC.

Evidence acquisition: Medline, Scopus, and Web of Science databases were searched using a free-text protocol including the terms *robot-assisted radical cystectomy* or *da Vinci radical cystectomy* or *robot* radical cystectomy*. RARC case series and studies comparing RARC with either ORC or LRC were collected. Cumulative analysis was conducted.

Evidence synthesis: The searches retrieved 105 papers. According to the different diversion type, overall mean operative time ranged from 360 to 420 min. Similarly, mean blood loss ranged from 260 to 480 ml. Mean in-hospital stay was about 9 d for all diversion types, with consistently high readmission rates. In series reporting on RARC with either extracorporeal or intracorporeal conduit diversion, overall 90-d complication rates were 59% (high-grade complication: 15%). In series reporting RARC with intracorporeal continent diversion, the overall 30-d complication rate was 45.7% (high-grade complication: 28%). Reported mortality rates were $\leq 3\%$ for all diversion types. Comparing RARC and ORC, cumulative analyses demonstrated shorter operative time for ORC, whereas blood loss and in-hospital stay were better with RARC (all p values < 0.003). Moreover, 90-d complication rates of any-grade and 90-d grade 3 complication rates were lower for RARC (all p values < 0.04), whereas high-grade complication and mortality rates were similar.

Conclusions: RARC can be performed safely with acceptable perioperative outcome, although complications are common. Cumulative analyses demonstrated that operative time was shorter with ORC, whereas RARC may provide some advantages in terms of blood loss

and transfusion rates and, more limitedly, for postoperative complication rates over ORC and LRC.

Patient summary: Although open radical cystectomy (RC) is still regarded as a standard treatment for muscle-invasive bladder cancer, laparoscopic and robot-assisted RC are becoming more popular. Robotic RC can be safely performed with acceptably low risk of blood loss, transfusion, and intraoperative complications; however, as for open RC, the risk of postoperative complications is high, including a substantial risk of major complication and reoperation.

1. Introduction

Radical cystectomy (RC) with regional lymph node dissection is the standard surgical treatment for muscle-invasive and high-risk non–muscle-invasive urothelial carcinoma of the bladder [1]. Although open RC (ORC) is still the most commonly adopted surgical approach [2], minimally invasive techniques have gained popularity such that laparoscopic RC (LRC) and robot-assisted RC (RARC) are routinely performed with promising short- and intermediate-term results [3].

Due to increasing evidence in the field of RARC and in preparation for the Pasadena international consensus meeting on best practice in RARC and urinary diversion, we performed a systematic literature review of perioperative, functional, and oncologic outcomes of RARC in comparison with ORC and LRC. We report the findings of this review with a cumulative analysis of perioperative outcomes and postoperative complications.

2. Evidence acquisition

The systematic literature search was initially performed in September 2013 using the Medline, Scopus, and Web of Science databases. The searches included a free-text protocol using the terms *robot-assisted radical cystectomy* or *da Vinci radical cystectomy* or *robot* radical cystectomy* in all fields of the records for PubMed and Scopus searches and in the *Title* and *Topic* fields for the Web of Science search. No limits were applied. A full update of the searches was done April 28, 2014.

Two authors (G.N. and B.Y.) separately reviewed the records to select RARC case series as well as studies that compared RARC with ORC and RARC with LRC. Discrepancies were resolved by open discussion. Other significant studies cited in the reference lists of the selected papers were evaluated, as were studies published after the systematic search.

All noncomparative studies reporting intraoperative and perioperative data (operative time, blood loss, transfusion rate, in-hospital stay, readmission, complication rates), functional data

(urinary continence, erectile function), and oncologic data (positive surgical margins, lymph node yield, disease-free survival, cancer-specific survival, overall survival) of RARC were collected. The present review included only studies reporting perioperative outcomes and complications.

Studies reporting partial cystectomy, prostate-sparing cystectomy, salvage cystectomy, cystectomy for urachal cancers or benign diseases, single-case reports, or pure laparoscopic (or mixed) series; those focusing on RC with laparoendoscopic single-site or natural orifice transluminal endoscopic surgery; experimental studies on animal models; congress abstracts; review papers; editorials; population-based studies; and book chapters were not included in the review. All data retrieved from the selected studies were recorded in an electronic database.

Papers were categorized according to the Oxford Level of Evidence Working Group 2011 levels of evidence (LOEs) for therapy studies: LOE 1, systematic review of randomized trials or n-of-1 trials; LOE 2, randomized trial or observational study with dramatic effect; LOE 3, nonrandomized controlled cohort or follow-up study; LOE 4, case series, case-control study, or historically controlled study; or LOE 5, mechanism-based reasoning [4]. Papers were categorized according to the IDEAL recommendations [5]. Methodological reporting of complications was evaluated according to the Martin criteria [6]. The systematic review was performed in agreement with the PRISMA statement [7].

2.1. Statistical analysis

Cumulative analysis was conducted using Review Manager v5.2 software designed for composing Cochrane Reviews (Cochrane Collaboration, Oxford, UK). Statistical heterogeneity was tested using the chi-square test. A p value <0.10 was used to indicate heterogeneity. If there was a lack of heterogeneity, fixed-effects models were used for the cumulative analysis. Random-effects models were used in cases of heterogeneity. For

continuous outcomes, the results were expressed as weighted mean differences (WMDs) and standard deviations (SDs); for dichotomous variables, results were given as odds ratios (ORs) and 95% confidence intervals (CIs). Due to limitations in the Review Manager v5.2 software, cumulative analysis of continuous variables was possible only when rough data were presented as mean and SD. Authors of the papers were contacted to provide missing data whenever necessary. For all statistical analyses, two-sided $p < 0.05$ was considered statistically significant.

3. Evidence synthesis

3.1. Quality of the studies and level of evidence

The flow of this systematic review of the literature is shown in Figure 1. In total, 70 surgical series [8–77] and 23 comparative studies [78–100] reported perioperative outcomes and complications of RARC.

Most of the surgical series were retrospective, single-center studies, with the exception of some prospective studies [8,9,29,32,33,48,62,63,67,69,71,73,77] and some multi-institutional collaboration papers [19,24,26,30,38,39,43,45,72] (LOE 4). Only three of the comparative studies were randomized [78–80] (LOE 2b); all other comparative studies were nonrandomized, whether prospective or retrospective (LOE 4).

3.2. Perioperative outcomes after robot-assisted radical cystectomy

Table 1 summarizes mean operative time, blood loss, transfusion rate, intraoperative complication rate, time to flatus, time to bowel movement, in-hospital stay, and readmission rate in the RARC surgical series.

Once duplicate publications and collaborative studies were excluded, weighted mean operative time was 360 min (range: 230–618 min) for RARC with extracorporeal conduit, 420 min (range: 300–496 min) for RARC with extracorporeal neobladder, 340 min (range: 292–660 min) for RARC with intracorporeal conduit, and 420 min (range: 420–450 min) for

RARC with intracorporeal neobladder. Overall mean blood loss was 375 ml (range: 208–763 ml) for RARC with extracorporeal conduit, 390 ml (range: 167–400 ml) for RARC with extracorporeal neobladder, 270 ml (range: 200–1118 ml) for RARC with intracorporeal conduit, and 480 ml (range: 225–500 ml) for RARC with intracorporeal neobladder.

Transfusion rates vary, at 12% for RARC with extracorporeal conduit, 44% for RARC with extracorporeal neobladder, 14.7% for RARC with intracorporeal conduit, and 7% for RARC with intracorporeal neobladder.

The intraoperative complication rate was 3% in the series reporting RARC with extracorporeal conduit, whereas no intraoperative complications were reported in papers evaluating either extracorporeal neobladder or intracorporeal diversions. Sufficient data on time to flatus and bowel movements were available only for the series analyzing RARC with extracorporeal conduit, demonstrating mean time to flatus of 2.5 d (range: 2.1–3.4 d) and mean time to bowel movement of 3.1 d (range: 2.8–4 d).

Length of stay was 8.7 d (range: 3.3–20.7 d) for RARC with extracorporeal conduit, 8.9 d (range: 6.7–9 d) for RARC with extracorporeal neobladder, 8.6 d (range: 4.5–9 d) for RARC with intracorporeal conduit, and 8.5 d (range: 8–9 d) for RARC with intracorporeal neobladder. Readmission rates were consistently high, ranging from 19% for RARC with extracorporeal conduit to 75% in one small study for RARC with intracorporeal neobladder.

3.3. Perioperative outcomes after robot-assisted radical cystectomy and patient characteristics

Two studies analyzed the impact of patient body mass index (BMI) on perioperative outcomes [64,70] (Table 2). Butt et al assessed a cohort of 49 patients receiving RARC and extracorporeal ileal conduit at Roswell Park Cancer Institute (Buffalo, NY, USA) and failed to demonstrate any major significant difference in perioperative outcomes in patients with BMI <25, 25–29, and \geq 30 [70]. More recently, Poch et al reported on 56 consecutive patients

treated at the same institution with RARC and intracorporeal conduit and demonstrated that only blood loss was significantly higher in obese patients [64].

3.4. Aspects of surgery influencing perioperative outcomes after robot-assisted radical cystectomy

Table 3 summarizes the studies assessing the effects of particular surgical aspects on perioperative outcomes. Five papers evaluated the effect of the number of cases previously performed on perioperative outcomes [18,35,65,68,71]. Whereas Pruthi et al failed to demonstrate any significant modification of the perioperative outcomes among the first 50 cases of RARC with extracorporeal diversion [18], Hayn et al found significant improvements in both mean time for RARC (from 180 min in the first 50 cases to 136 min in the last 64 cases; $p < 0.001$) and lymph node yield (from 16 nodes in the first 50 cases to 24 nodes in the last 64 cases; $p < 0.001$) among the first 164 RARC cases performed with extracorporeal urinary diversion [71]. Analyzing their first 60 cases of RARC with extracorporeal urinary diversion, Richards et al demonstrated reduction in overall complication rates from 70% in the first 20 cases to 30% in the second and third 20 cases ($p = 0.013$) [35].

With regard to reporting intracorporeal diversion, in a series of 100 cases receiving RARC with mainly conduit diversion, Azzouni et al found significant reduction in overall diversion time with experience (from 140 min in the first 25 cases to 103 min in the last 25 cases; $p = 0.002$) [65]. Finally, in a series of 45 patients treated with RARC and intracorporeal neobladder, Schumacher et al demonstrated significant improvement in many aspects, including increased adoption of lymph node dissection and reduction in operative time (from 523 min in the first 15 cases to 434 min in the last 15 cases; $p = 0.005$), in-hospital stay (from 22.5 d in the first 15 cases to 9.5 d in the last 15 cases; $p = 0.006$), and >30-d complication rates (from 54% in the first 15 cases to 20% in the last 15 cases; $p = 0.005$) [68].

Hayn et al evaluated the impact of previous experience with robot-assisted radical prostatectomy (RARP) on RARC outcome [72]. Specifically, RARP experience was stratified into four groups: <50, 51–100, 101–150, and >150 cases. RARC operative time, blood loss, and lymph node yield were all significantly associated with prior RARP experience (all p values <0.001), with the most experienced RARP surgeons experiencing lower blood loss but longer operative time and lower lymph node yield [72].

Finally, two studies compared perioperative outcomes in RARC with intracorporeal and extracorporeal diversion [73,74]. Specifically, Guru et al compared the outcomes of 13 patients receiving intracorporeal ileal conduit and 13 receiving extracorporeal ileal conduit at Roswell Park Cancer Institute and failed to demonstrate any significant difference between the two groups [73]. Similarly, Kang et al compared 38 patients receiving RARC with either extracorporeal conduit ($n = 22$) or neobladder ($n = 14$) and 4 patients receiving RARC with intracorporeal diversion (three conduits and one neobladder) and demonstrated shorter operative time for extracorporeal diversions [74]. Both studies had low power to draw definitive conclusions on the issue.

3.5. Postoperative complication rates after robot-assisted radical cystectomy

Table 4 summarizes complication rates in the RARC surgical series stratified by diversion type. In series reporting on RARC with extracorporeal conduit diversion, overall 30- and 90-d complication rates were 44% (range: 26–78%) and 59% (range: 30–77%), respectively.

Low-grade complications were the most prevalent, at 29.4% (range: 8–62%) and 54% (range: 15–79%) at 30 d and 90 d, respectively. High-grade complications at 30 d and 90 d were present in 11.8% (range: 0–35%) and 15% (range: 4–19%), respectively, including high reoperation rates (9.7% at 30 d and 14% at 90 d) and relatively low mortality rates (1.6% at 30 d and 3% at 90 d).

With regard to RARC with extracorporeal continent diversion, virtually all studies reported the experience of City of Hope Comprehensive Cancer Center (Duarte, CA, USA) with an overall 90-d complication rate of up to 77%, including 45% low- and 32% high-grade complications. The 90-d mortality rate was as high as 5% [25,55,56,58,59].

In series reporting on RARC with intracorporeal conduit diversion, the overall complication rates at 30, 30–90, and 90 d were 67% (range: 42–86%), 22% (range: 14–23%), and 59% (range: 30–77%), respectively. Low-grade complication rates were 45% (range: 32–50%), 2% (range: 0–14%), and 66% at 30, 30–90, and 90 d, respectively. High-grade complications were present in 24% (range: 0–54%), 20% (range: 0–23%), and 15% at 30, 30–90, and 90 d, respectively. Reoperation rates were 39% at 30 d, 19% at 30–90 d, and 25% at 90 d. Reported mortality rates were relatively low (0% at 30 d, 1.7% at 30–90 d, and 1.7% at 90 d).

In series reporting RARC with intracorporeal continent diversion, the overall complication rates at 30, 30–90, and 90 d were 45.7% (range: 43–62%), and 30% (range: 12–34%), respectively. Low-grade complications were reported in 19% (range: 12–33%) and 13.5% (range: 13–15%) at 30 d and 30–90 d, respectively. High-grade complications were present in 28% (range: 15–33%) and 18% (range: 12–21%) at 30 d and 30–90 d, respectively.

Reoperation rates were 17% at 30 d, 16% at 30–90 d, and 33% at 90 d. Reported mortality rates were relatively low (1% at 30 d, 1.7% at 30–90 d, and 2.7% at 90 d).

3.6. Patient characteristics and aspects of surgery influencing postoperative complications after robot-assisted radical cystectomy

Table 5 summarizes the studies evaluating the impact of patient characteristics and surgical factors on complication rates in RARC series. Two studies analyzed the impact of patient BMI on complication rates [64,70]. Both papers failed to identify any significant difference in complication rates according to patient BMI. Five papers evaluated the effect of the number of cases performed on postoperative complications [18,35,65,68,71,77]. With regard

to RARC with extracorporeal urinary diversion, Richards et al [35] demonstrated significant improvement in 90-d complication rates among the first 60 cases performed, with overall complication rates decreasing from 70% in the first 20 cases to 30% in the last 20 [35]. Conversely, Hayn et al reported stable 240-d complication rates in a larger series of 164 patients treated at Roswell Park Cancer Institute [71]. With regard to the series reporting intracorporeal conduit diversion, Azzouni et al demonstrated little change in 30- and 90-d complication rates among the first 100 cases performed [65]. Conversely, two series from the Karolinska Institute (Stockholm, Sweden), mainly reporting on RARC with intracorporeal neobladder, showed significant improvements in complication rates at 30 d and 30–90 d [68,77]. Finally, two studies compared postoperative complication rates for RARC with intracorporeal and extracorporeal diversion [73,74]. Both studies reported overlapping complication rates, but small sample size and other methodological limitations prevented any definitive conclusions from being drawn.

Few studies evaluated independent predictors of postoperative complications in a more formal way, including multivariable analyses (Table 6). Specifically, with regard to series reporting on RARC with mainly extracorporeal conduit diversion, three studies reported on predictors of complications [28,39,43] and one reported on predictors of readmission [48]. Specifically, Kauffman et al analyzed 79 patients treated at Weill Cornell Medical Center (New York, NY, USA) and demonstrated that preoperative creatinine level >1.4 mg/dl (OR: 4.2; $p = 0.038$) and intravenous fluids >5000 ml (OR: 4.1; $p = 0.025$) were predictors of any-grade complication, whereas patient age of >65 yr (OR: 12.7; $p = 0.04$), estimated blood loss >500 ml (OR: 9.7; $p = 0.015$), and intravenous fluids >5000 ml (OR: 42.1; $p = 0.003$) were predictors of high-grade complications [28]. In a multicenter series of 279 patients treated at four US institutions, Smith et al demonstrated that younger age of <65 yr (OR: 0.4; $p = 0.230$) and American Society of Anesthesiologists (ASA) score ($p = 0.025$) were associated

with higher risk of complications [39]. In another, larger multi-institutional study involving >900 patients from >20 institutions, Johar et al performed sophisticated analyses evaluating preoperative and intraoperative predictors of any-grade and high-grade complications. Among preoperative variables, age at surgery (OR: 1.34; $p < 0.0001$); BMI (OR: 1.04; $p = 0.006$); and, notably, use of neoadjuvant chemotherapy (OR: 1.71; $p = 0.007$) were associated with any grade of complications, whereas age at surgery (OR: 1.39; $p = 0.02$), BMI (OR: 1.04; $p = 0.024$), use of neoadjuvant chemotherapy (OR: 1.88; $p = 0.006$), and current smoking status (OR: 1.68; $p = 0.018$) were predictive of high-grade complications. Among intraoperative variables, blood transfusion (OR: 1.84; $p = 0.006$) and conduit diversion (OR: 1.44; $p = 0.036$) were predictive of any grade of complications, whereas only blood transfusion (OR: 1.94; $p = 0.009$) was associated with high-grade complications [43]. In the same study, predictors of 90-d mortality were also assessed, with age (OR: 1.62; $p = 0.018$), among the preoperative variables, and blood transfusions (OR: 4.20; $p = 0.001$), among the intraoperative variables, as the only independent predictors [43]. Finally, Al-Daghmin et al reported on readmission rates and demonstrated 30- and 90-d readmission rates of 15% and 25%, respectively. Patient BMI (OR: 1.12; $p = 0.004$) and presence of any grade of complications (OR: 0.09; $p = 0.03$) were predictive of 30-d readmission, whereas male sex (OR: 0.41; $p = 0.014$) and BMI (OR: 1.1; $p = 0.004$) were predictive of 90-d readmission [48].

With regard to series reporting on RARC with mainly extracorporeal continent diversion, two papers reporting the experience of City of Hope Comprehensive Cancer Center assessed predictors [56,58]. In the largest series, reporting on 91 patients receiving orthotopic neobladder, 51 receiving Indiana pouch, and 67 receiving ileal conduit, Nazmy et al demonstrated that ASA score (OR: 7.39; $p = 0.01$), preoperative hematocrit (HCT; OR: 0.85; $p = 0.002$), and diversion type (Indiana pouch vs conduit: OR: 6.59; $p = 0.002$; neobladder vs

conduit: OR: 4.0; $p = 0.007$) were associated with complications of any grade at 90 d, whereas Charlson comorbidity index (OR: 1.44; $p = 0.003$), preoperative HCT (OR: 0.88; $p = 0.0009$), and diversion type (neobladder vs conduit: OR: 4.9; $p = 0.001$) were predictive of high-grade complications at 90 d [58]. Yuh et al also included intraoperative variables and found that operative time (OR: 1.71; $p = 0.006$) and blood loss (OR: 1.0; $p = 0.0003$) were predictive of complications of any grade at 90 d [56].

3.7. Cumulative analysis of studies comparing robot-assisted radical cystectomy with open or laparoscopic radical cystectomy

Table 7 summarizes the comparative studies that report perioperative parameters and intraoperative complication rates after ORC, LRC, and RARC. With regard to the comparison of RARC and ORC, cumulative analyses showed statistically significant differences in terms of rates for operative time (WMD: 83.60; 95% CI, 57.1–110.1; $p < 0.00001$ in favor of ORC), blood loss (WMD: -521; 95% CI, -644 to -399; $p < 0.00001$ in favor of RARC), transfusion (OR: 0.16; 95% CI, 0.1–0.27; $p < 0.00001$ in favor of RARC), and in-hospital stay (WMD: -1.26; 95% CI, -2.08 to -0.43; $p = 0.003$ in favor of RARC), whereas rates for intraoperative complications (OR: 1.34; 95% CI, 0.37–4.77; $p = 0.65$) were similar for RARC and ORC (Fig. 2). Cumulative analysis of mean time to flatus and mean time to bowel movement was not possible. Notably, considering only the few available randomized controlled trials (RCTs), operative time (WMD: 74.7; 95% CI, -30.1 to 179.5; $p = 0.16$) and in-hospital stay (WMD: 0.03; 95% CI, -1.37 to 1.44; $p = 0.96$) were overlapping for the two procedures.

With regard to the comparison of RARC and LRC, cumulative analyses showed statistically significant differences in terms of rates for transfusion (OR: 0.19; 95% CI, 0.07–0.53; $p = 0.001$ in favor of RARC) (Fig.3). Cumulative analysis of the other variables was not possible.

Table 8 summarized the comparative studies that report postoperative complication rates after ORC, LRC, and RARC. With regard to the comparison of RARC and ORC, cumulative analyses showed that rates for any grade of complication at 90 d (OR: 0.44; 95% CI, 0.31–0.61; $p < 0.0001$) and for grade 3 complications at 90 d (OR: 0.55; 95% CI, 0.31–0.98; $p = 0.04$) were in favor of RARC. In contrast, rates at 30 d for any grade of complication (OR: 0.77; 95% CI, 0.56–1.4; $p = 0.09$), for grade 3 complications (OR: 0.70; 95% CI, 0.43–1.13; $p = 0.14$), and for high-grade complications (OR: 0.64; 95% CI, 0.32–1.29; $p = 0.21$); 30-d mortality rates (OR: 0.45; 95% CI, 0.14–1.44; $p = 0.18$); and rates at 90 d for high-grade complications (OR: 0.62; 95% CI, 0.37–1.03; $p = 0.06$) and mortality (OR: 0.45; 95% CI, 0.12–1.66; $p = 0.23$) were similar for RARC and ORC (Fig. 4).

With regard to the comparison of RARC and LRC, cumulative analyses showed that rates at 30 d for any grade of complication (OR: 0.18; 95% CI, 0.08–0.38; $p < 0.0001$) and for grade 3 complications (OR: 0.35; 95% CI, 0.15–0.82; $p = 0.02$) were significantly lower with RARC (Fig. 5).

3.8. Discussion

Following the success of RARP and other robotic procedures, da Vinci technology (Intuitive Surgical, Sunnyvale, CA USA) has been applied to RC, and the number of RARCs performed is increasing; however, according to the most current data available, <20% of RCs are performed robotically in the United States [2]. Our systematic review demonstrated that RARC can be performed safely, with acceptable operative time, relatively little blood loss, and relatively low transfusion rates. Although the risk of intraoperative complications is low, postoperative complications are common, and the rate of readmission is relatively high. Some preoperative patient characteristics, including age, BMI, renal function, and comorbidity, may be associated with the risk of complications. Our cumulative analyses demonstrated that operative time was shorter with ORC, whereas blood loss and transfusion rates were

significantly lower with RARC than with ORC. Conversely, rates for any-grade and grade 3 complication at 90 d were slightly lower with RARC than with ORC. Similarly, transfusion rates were lower with RARC than with LRC, as were any-grade and grade 3 complication rates.

Perioperative outcomes and complication rates are critical issues for complex procedures such as RC. These outcomes have been reported extensively for ORC, including reports using standardized Martin criteria. In general, ORC is associated with a high risk of complications (>60%), including a considerable risk of high-grade complications (13–40% in large series) and mortality (up to 7% in some series) [101–105]. ORC outcomes appear to be associated with hospital and surgeon experience and volumes, with several studies demonstrating improved performance in high-volume centers by high-volume surgeons [106,107]. Our analysis suggests that RARC might provide benefit in terms of reduced blood loss and transfusion rates when compared with ORC, whereas operative time is shorter with ORC. Conversely, complication rates were mostly similar between RARC and ORC and slightly better for RARC in comparisons with LRC. However, many of the perioperative complications following RC may come from the reconstructive part of the procedure. Because most RARC cases reported had extracorporeal reconstruction, it can be hypothesized that this approach mitigated some potential benefit of a totally intracorporeal approach. However, intracorporeal diversion (especially orthotopic neobladder) is a very complex robotic procedure that is currently performed in very few centers and that has complication and readmission rates that appear quite high.

With regard to predictors of complications, in the most comprehensive report on ORC, Shabsigh et al demonstrated that sex, ASA score, and type of urinary diversion were associated with any grade of complications, whereas age at surgery, prior abdominal surgery, and estimated blood loss were associated with high-grade complications [101]. Moreover,

Svatek et al suggested a role for BMI as a predictor of both any-grade and high-grade complications [104]. Very similar results were identified in our systematic review of the literature on RARC, with age, ASA score, Charlson comorbidity index, BMI, and blood transfusion among the most common predictors [28,39,43]. Notably, in a large multi-institutional study involving >900 patients from >20 institutions, Johar et al identified the use of neoadjuvant chemotherapy as a predictor for any-grade and major complication rates [43]. That finding is not in agreement with the literature on ORC [108] and needs to be reconfirmed in larger analyses.

With regard to the comparison of ORC and RARC, in a population-based analysis of the US Nationwide Inpatient Sample, Yu et al reported on >7000 patients receiving ORC and 1100 treated with RARC in 1050 hospitals from 44 states in the United States. Specifically, the authors found that RARC was associated with a lower rate of complications (49% vs 64%), reduced perioperative mortality (0% vs 2.5%), and lower parenteral nutrition use (6.4% vs 13.3%) compared with ORC, whereas blood transfusions and length of stay were similar in the two groups [109]. Conversely, in another population-based study evaluating almost 35 000 patients treated with ORC and 2100 with RARC at 279 hospitals across the United States between 2004 and 2010 and included in the Premier Perspective Database, Leow et al failed to demonstrate significant differences in 90-d postoperative mortality and major complication rates between RARC and ORC, whereas 46% decreased odds of minor complications, mainly due to reduced need for blood transfusion and total parenteral nutrition, were demonstrated [2]. On the whole, the data for our systematic review reconfirmed a lower risk of blood loss and transfusion for RARC compared with ORC and LRC and slightly lower risks of any-grade and high-grade complications at 90 d with RARC, whereas 30-d complication rates and 30- and 90-d mortality rates were similar for ORC and RARC. The reasons for such discrepancies are not clear but could include the well-known limited accuracy of population-

based studies and differences in baseline characteristics of the patients treated with ORC and RARC both in population-based studies and in the comparative studies included in the present systematic review.

Although the conclusions of this systematic review represent the best evidence available in the literature, some drawbacks must be considered. The papers included in the present systematic review contained only three RCTs [78–80], and only one was adequately powered to assess a difference in complications [80]. Unfortunately, at the present time, that paper is published as a letter to the editor in the *New England Journal of Medicine* and reports a limited amount of data; a more detailed report of the study is awaited. Moreover, most of the other low-quality evidence did not adopt accurate methodology for reporting complications. It was almost impossible to evaluate the impact of surgeon ability on the reported results due to the fact that advanced analyses of the RARC learning curve are lacking, and most of the available studies stratifying patient outcomes according to prior experience with RARP or to the number of prior RARC cases performed were small and retrospective. However, two studies from the Karolinska Institute suggested a decrease in complication rates with increasing surgical experience [68,77]. Finally, our comparative analyses were not adjusted for the baseline differences in patient characteristics and surgical experience. Considering that most of the studies included were not RCTs, it is likely that major differences were present between study arms, and this might account for some of the observed findings.

4. Conclusions

RARC can be performed safely with acceptable operative time, little blood loss, and low transfusion rates. The risk of intraoperative complications is low, but postoperative complications and readmission after discharge are common. Cumulative analyses demonstrated that operative time was shorter with ORC, whereas blood loss and transfusion rates were significantly lower with RARC than with ORC. Conversely, rates for any-grade

and grade 3 complications at 90 d were slightly lower with RARC than with ORC. Similarly, transfusion rates were lower with RARC than with LRC, as were any-grade and grade 3 complication rates. The lack of solid, high-quality evidence limits the strength of the data.

Author contributions: Giacomo Novara had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Novara.

Acquisition of data: Novara, Yuh.

Analysis and interpretation of data: Novara.

Drafting of the manuscript: Novara.

Critical revision of the manuscript for important intellectual content: Novara, Catto, Wilson, Annerstedt, Chan, Declan G. Murphy, Motttrie, Peabody, Skinner, Wiklund, Guru, Yuh.

Statistical analysis: Novara.

Obtaining funding: Wilson.

Administrative, technical, or material support: Catto, Wiklund.

Supervision: Catto, Guru, Wiklund, Wilson.

Other (specify): None.

Financial disclosures: Giacomo Novara certifies that all conflicts of interest, including specific financial interests and relationships and affiliations relevant to the subject matter or materials discussed in the manuscript (eg, employment/ affiliation, grants or funding, consultancies, honoraria, stock ownership or options, expert testimony, royalties, or patents filed, received, or pending), are the following: Dr. Annerstedt has had assignments as a proctor for Intuitive Surgical. Dr. Murphy has received travel grants to support conference participation from Device Technologies Australia and has received reimbursement for proctoring robotic surgery. Dr. Wiklund has received proctoring assignments and a research

grant from Intuitive Surgical. Dr. Wilson has been a consultant and speaker for Intuitive Surgical. The other authors have nothing to disclose.

Funding/Support and role of the sponsor: A private donor assisted with travel costs for this review.

References

- [1] Witjes JA, Comperat E, Cowan NC, et al. EAU guidelines on muscle-invasive and metastatic bladder cancer: summary of the 2013 guidelines. *Eur Urol* 2014;65:778–92.
- [2] Leow JJ, Reese SW, Jiang W, et al. Propensity-matched comparison of morbidity and costs of open and robot-assisted radical cystectomies: a contemporary population-based analysis in the United States. *Eur Urol* 2014;66:569–76.
- [3] Challacombe BJ, Bochner BH, Dasgupta P, et al. The role of laparoscopic and robotic cystectomy in the management of muscle-invasive bladder cancer with special emphasis on cancer control and complications. *Eur Urol* 2011;60:767–75.
- [4] Howick J, Chalmers I, Glasziou P, et al. Explanation of the 2011 Oxford Centre for Evidence-Based Medicine (OCEBM) levels of evidence (background document). OCEBM Web site. <http://www.cebm.net/index.aspx?o=5653>.
- [5] McCulloch P, Altman DG, Campbell WB, et al. No surgical innovation without evaluation: the IDEAL recommendations. *Lancet* 2009;374:1105–12.
- [6] Martin RC II, Brennan MF, Jaques DP. Quality of complication reporting in the surgical literature. *Ann Surg* 2002;235:803–13.
- [7] Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151:264–9.

- [8] Guru KA, Nogueira M, Piacente P, Nyquist J, Mohler JL, Kim HL. Robot-assisted anterior exenteration: technique and initial series. *J Endourol* 2007;21:633–9.
- [9] Guru KA, Kim HL, Piacente PM, Mohler JL. Robot-assisted radical cystectomy and pelvic lymph node dissection: initial experience at Roswell Park Cancer Institute. *Urology* 2007;69:469–74.
- [10] Mottrie A, Carpentier P, Schatteman P, et al. Robot-assisted laparoscopic radical cystectomy: initial experience on 27 consecutive patients. *J Robot Surg* 2007;1:197–201.
- [11] Hemal AK, Kolla SB, Wadhwa P. First case series of robotic radical cystoprostatectomy, bilateral pelvic lymphadenectomy, and urinary diversion with the da Vinci S system. *J Robot Surg* 2008;2:35–40.
- [12] Lowentritt BH, Castle EP, Woods M, Davis R, Thomas R. Robot-assisted radical cystectomy in women: technique and initial experience. *J Endourol* 2008;22:709–12.
- [13] Murphy DG, Challacombe BJ, Elhage O, et al. Robotic-assisted laparoscopic radical cystectomy with extracorporeal urinary diversion: initial experience. *Eur Urol* 2008;54:570–80.
- [14] Park SY, Cho KS, Park KK, Park SJ, Ham WS, Rha KH. Robot-assisted laparoscopic radical cystectomy with ileal conduit urinary diversion. *Korean J Urol* 2008;49:506–9.
- [15] Park SY, Jeong W, Choi YD, Chung BH, Hong SJ, Rha KH. Yonsei experience in robotic urologic surgery-application in various urological procedures. *Yonsei Med J* 2008;49:897–900.
- [16] Pruthi RS, Wallen EM. Robotic-assisted laparoscopic radical cystoprostatectomy. *Eur Urol* 2008;53:310–22.
- [17] Pruthi RS, Stefaniak H, Hubbard JS, Wallen EM. Robot-assisted laparoscopic anterior pelvic exenteration for bladder cancer in the female patient. *J Endourol* 2008;22:2397–402.

- [18] Pruthi RS, Smith A, Wallen EM. Evaluating the learning curve for robot-assisted laparoscopic radical cystectomy. *J Endourol* 2008;22:2469–74.
- [19] Woods M, Thomas R, Davis R, et al. Robot-assisted extended pelvic lymphadenectomy. *J Endourol* 2008;22:1297–302.
- [20] Yuh B, Padalino J, Butt ZM, et al. Impact of tumour volume on surgical and pathological outcomes after robot-assisted radical cystectomy. *BJU Int* 2008;102:840–3.
- [21] Gamboa AJ, Young JL, Dash A, Abraham JB, Box GN, Ornstein DK. Pelvic lymph node dissection and outcome of robot-assisted radical cystectomy for bladder carcinoma. *J Robot Surg* 2009;3:7–12.
- [22] Pruthi RS, Stefaniak H, Hubbard JS, Wallen EM. Robotic anterior pelvic exenteration for bladder cancer in the female: outcomes and comparisons to their male counterparts. *J Laparoendosc Adv Surg Tech* 2009;19:23–7.
- [23] Yuh BE, Ciccone J, Chandrasekhar R, et al. Impact of previous abdominal surgery on robot-assisted radical cystectomy. *JSLs* 2009;13:398–405.
- [24] Hayn MH, Hussain A, Mansour AM, et al. The learning curve of robot-assisted radical cystectomy: results from the International Robotic Cystectomy Consortium. *Eur Urol* 2010;58:197–202.
- [25] Josephson DY, Chen JA, Chan KG, Lau CS, Nelson RA, Wilson TG. Robotic-assisted laparoscopic radical cystoprostatectomy and extracorporeal continent urinary diversion: highlight of surgical techniques and outcomes. *Int J Med Robot* 2010;6:315–23.
- [26] Kang SG, Kang SH, Lee YG, et al. Robot-assisted radical cystectomy and pelvic lymph node dissection: a multi-institutional study from Korea. *J Endourol* 2010;24:1435–40.
- [27] Kasraeian A, Barret E, Cathelineau X, et al. Robot-assisted laparoscopic cystoprostatectomy with extended pelvic lymphadenectomy, extracorporeal

enterocystoplasty, and intracorporeal enterourethral anastomosis: initial Montsouris experience. *J Endourol* 2010;24:409–13.

[28] Kauffman EC, Ng CK, Lee MM, et al. Critical analysis of complications after robotic-assisted radical cystectomy with identification of preoperative and operative risk factors. *BJU Int* 2010;105:520–7.

[29] Kwon SY, Kim BS, Kim TH, Yoo ES, Kwon TG. Initial experiences with robot-assisted laparoscopic radical cystectomy. *Korean J Urol* 2010;51:178–82.

[30] Martin AD, Nunez RN, Pacelli A, et al. Robot-assisted radical cystectomy: intermediate survival results at a mean follow-up of 25 months. *BJU Int* 2010;105:1706–9.

[31] Pruthi RS, Nielsen ME, Nix J, Smith A, Schultz H, Wallen EM. Robotic radical cystectomy for bladder cancer: surgical and pathological outcomes in 100 consecutive cases. *J Urol* 2010;183:510–5.

[32] Hayn MH, Hellenthal NJ, Hussain A, Stegemann AP, Guru KA. Defining morbidity of robot-assisted radical cystectomy using a standardized reporting methodology. *Eur Urol* 2011;59:213–8.

[33] Khan MS, Elhage O, Challacombe B, Rimington P, Murphy D, Dasgupta P. Analysis of early complications of robotic-assisted radical cystectomy using a standardized reporting system. *Urology* 2011;77:357–62.

[34] Lavery HJ, Martinez-Suarez HJ, Abaza R. Robotic extended pelvic lymphadenectomy for bladder cancer with increased nodal yield. *BJU Int* 2011;107:1802–5.

[35] Richards KA, Kader K, Pettus JA, Smith JJ, Hemal AK. Does initial learning curve compromise outcomes for robot-assisted radical cystectomy? A critical evaluation of the first 60 cases while establishing a robotics program. *J Endourol* 2011;25:1553–8.

[36] Shah AD, Abaza R. Clinical pathway for 3-day stay after robot-assisted cystectomy. *J Endourol* 2011;25:1253–8.

- [37] Lau CS, Talug J, Williams SB, et al. Robotic-assisted laparoscopic radical cystectomy in the octogenarian. *Int J Med Robot* 2012;8:247–52.
- [38] Mmeje CO, Nunez-Nateras R, Nielsen ME, et al. Oncologic outcomes for lymph node-positive urothelial carcinoma patients treated with robot assisted radical cystectomy: with mean follow-up of 3.5 years. *Urol Oncol* 2013;31:1621–7.
- [39] Smith AB, Raynor M, Amling CL, et al. Multi-institutional analysis of robotic radical cystectomy for bladder cancer: perioperative outcomes and complications in 227 patients. *J Laparoendosc Adv Surg Tech A* 2012;22:17–21.
- [40] Treiyer A, Saar M, Butow Z, Kamradt J, Siemer S, Stockle M. Robotic-assisted laparoscopic radical cystectomy: surgical and oncological outcomes. *Int Braz J Urol* 2012;38:324–9.
- [41] Tsui KH, Chen CL, Lin YH, Hou CP, Chang PL. Robotic assisted laparoscopic radical cystectomy for bladder carcinoma: early experience and oncologic outcomes. *Formosan J Surg* 2012;45:178–82.
- [42] Abbas DN, Kamal JM, El Sheikh SM, Mahmud AM. Early experience in anesthesia of robot assisted cystoprostatectomy. *Egyptian J Anesth* 2013;29:77–81.
- [43] Johar RS, Hayn MH, Stegemann AP, et al. Complications after robot-assisted radical cystectomy: results from the International Robotic Cystectomy Consortium. *Eur Urol* 2013;64:52–7.
- [44] Khan MS, Elhage O, Challacombe B, et al. Long-term outcomes of robot-assisted radical cystectomy for bladder cancer. *Eur Urol* 2013;64:219–24.
- [45] Marshall SJ, Hayn MH, Stegemann AP, et al. Impact of surgeon and volume on extended lymphadenectomy at the time of robot-assisted radical cystectomy: results from the International Robotic Cystectomy Consortium (IRCC). *BJU Int* 2013;111:1075–80.

- [46] Saar M, Ohlmann CH, Siemer S, et al. Fast-track rehabilitation after robot-assisted laparoscopic cystectomy accelerates postoperative recovery. *BJU Int* 2013;112:E99–106.
- [47] Xylinas E, Green DA, Otto B, et al. Robotic-assisted radical cystectomy with extracorporeal urinary diversion for urothelial carcinoma of the bladder: analysis of complications and oncologic outcomes in 175 patients with a median follow-up of 3 years. *Urology* 2013;82:1323–9.
- [48] Al-Daghmin A, Aboumohamed A, Din R, et al. Readmission after robot-assisted radical cystectomy: outcomes and predictors at 90-day follow-up. *Urology* 2014;83:350–6.
- [49] Lin CY, Yang CR, Cheng CL, et al. Application in robotic urologic surgery. *J Chin Med Assoc* 2014;77:242–5.
- [50] Phillips EA, Uberoi V, Tuerk IA. Robot-assisted radical cystectomy in octogenarians. *J Endourol* 2014;28:219–23.
- [51] Snow-Lisy DC, Campbell SC, Gill IS, et al. Robotic and laparoscopic radical cystectomy for bladder cancer: long-term oncologic outcomes. *Eur Urol* 2014;65:193–200.
- [52] Menon M, Hemal AK, Tewari A, et al. Nerve-sparing robot-assisted radical cystoprostatectomy and urinary diversion. *BJU Int* 2003;92:232–6.
- [53] Menon M, Hemal AK, Tewari A, et al. Robot-assisted radical cystectomy and urinary diversion in female patients: technique with preservation of the uterus and vagina. *J Am Coll Surg* 2004;198:386–93.
- [54] Manoharan M, Katkooi D, Kishore TA, Antebie E. Robotic-assisted radical cystectomy and orthotopic ileal neobladder using a modified Pfannenstiel incision. *Urology* 2011;77:491–3.
- [55] Torrey RR, Chan KG, Yip W, et al. Functional outcomes and complications in patients with bladder cancer undergoing robotic-assisted radical cystectomy with extracorporeal Indiana pouch continent cutaneous urinary diversion. *Urology* 2012;79:1073–8.

- [56] Yuh BE, Nazmy M, Ruel NH, et al. Standardized analysis of frequency and severity of complications after robot-assisted radical cystectomy. *Eur Urol* 2012;62:806–13.
- [57] Pham KN, Sack BS, O'Connor RC, et al. V-Loc urethro-intestinal anastomosis during robotic cystectomy with orthotopic urinary diversion. *Can Urol Assoc J* 2013;7:E663–6.
- [58] Nazmy M, Yuh B, Kawachi M, et al. Early and late complications of robot-assisted radical cystectomy: a standardized analysis by urinary diversion type. *J Urol* 2014;191:681–7.
- [59] Yuh B, Torrey RR, Ruel NH, et al. Intermediate-term oncologic outcomes of robot-assisted radical cystectomy for urothelial carcinoma. *J Endourol* 2014;28:939–45.
- [60] Yohannes P, Puri V, Yi B, Khan AK, Sudan R. Laparoscopy-assisted robotic radical cystoprostatectomy with ileal conduit urinary diversion for muscle-invasive bladder cancer: initial two cases. *J Endourol* 2003;17:729–32.
- [61] Pruthi RS, Nix J, McRackan D, et al. Robotic-assisted laparoscopic intracorporeal urinary diversion. *Eur Urol* 2010;57:1013–21.
- [62] Jonsson MN, Adding LC, Hosseini A, et al. Robot-assisted radical cystectomy with intracorporeal urinary diversion in patients with transitional cell carcinoma of the bladder. *Eur Urol* 2011;60:1066–73.
- [63] Goh AC, Gill IS, Lee DJ, et al. Robotic intracorporeal orthotopic ileal neobladder: replicating open surgical principles. *Eur Urol* 2012;62:891–901.
- [64] Poch MA, Stegemann A, Chandrasekhar R, Hayn M, Wilding G, Guru KA. Does body mass index impact the performance of robot-assisted intracorporeal ileal conduit? *J Endourol* 2012;26:857–60.
- [65] Azzouni FS, Din R, Rehman S, et al. The first 100 consecutive, robot-assisted, intracorporeal ileal conduits: evolution of technique and 90-day outcomes. *Eur Urol* 2013;63:637–43.

- [66] Bishop CV, Vasdev N, Boustead G, Adshead JM. Robotic intracorporeal ileal conduit formation: initial experience from a single UK centre. *Adv Urol* 2013;64:2836.
- [67] Collins JW, Tyritzis S, Nyberg T, et al. Robot-assisted radical cystectomy: description of an evolved approach to radical cystectomy. *Eur Urol* 2013;64:654–63.
- [68] Schumacher MC, Jonsson MN, Hosseini A, et al. Surgery-related complications of robot-assisted radical cystectomy with intracorporeal urinary diversion. *Urology* 2011;77:871–6.
- [69] Tyritzis SI, Hosseini A, Collins J, et al. Oncologic, functional, and complications outcomes of robot-assisted radical cystectomy with totally intracorporeal neobladder diversion. *Eur Urol* 2013;64:734–41.
- [70] Butt ZM, Perlmutter AE, Piacente PM, et al. Impact of body mass index on robot-assisted radical cystectomy. *JSL* 2008;12:241–5.
- [71] Hayn MH, Hellenthal NJ, Seixas-Mikelus SA, et al. Is patient outcome compromised during the initial experience with robot-assisted radical cystectomy? Results of 164 consecutive cases. *BJU Int* 2011;108:882–7.
- [72] Hayn MH, Hellenthal NJ, Hussain A, et al. Does previous robot-assisted radical prostatectomy experience affect outcomes at robot-assisted radical cystectomy? Results from the International Robotic Cystectomy Consortium. *Urology* 2010;76:1111–6.
- [73] Guru K, Seixas-Mikelus SA, Hussain A, et al. Robot-assisted intracorporeal ileal conduit: Marionette technique and initial experience at Roswell Park Cancer Institute. *Urology* 2010;76:866–71.
- [74] Kang SG, Ko YH, Jang HA, et al. Initial experience of robot-assisted radical cystectomy with total intracorporeal urinary diversion: comparison with extracorporeal method. *J Laparoendosc Adv Surg Tech A* 2012;22:456–62.

- [75] Akbulut Z, Canda AE, Ozcan MF, Atmaca AF, Ozdemir AT, Balbay MD. Robot-assisted laparoscopic nerve-sparing radical cystoprostatectomy with bilateral extended lymph node dissection and intracorporeal studer pouch construction: outcomes of first 12 cases. *J Endourol* 2011;25:1469–79.
- [76] Canda AE, Atmaca AF, Altinova S, Akbulut Z, Balbay MD. Robot-assisted nerve-sparing radical cystectomy with bilateral extended pelvic lymph node dissection (PLND) and intracorporeal urinary diversion for bladder cancer: initial experience in 27 cases. *BJU Int* 2012;110:434–44.
- [77] Collins JW, Tyritzis S, Nyberg T, et al. Robotic assisted radical cystectomy with intracorporeal neobladder - what is the effect of the learning curve on outcomes? *BJU Int* 2014;113:100–7.
- [78] Nix J, Smith A, Kurpad R, Nielsen ME, Wallen EM, Pruthi RS. Prospective randomized controlled trial of robotic versus open radical cystectomy for bladder cancer: perioperative and pathologic results. *Eur Urol* 2010;57:196–201.
- [79] Parekh DJ, Messer J, Fitzgerald J, Ercole B, Svatek R. Perioperative outcomes and oncologic efficacy from a pilot prospective randomized clinical trial of open versus robotic assisted radical cystectomy. *J Urol* 2013;189:474–9.
- [80] Bochner BH, Sjoberg DD, Laudone VP; Memorial Sloan Kettering Cancer Center Bladder Cancer Surgical Trials G. A randomized trial of robot-assisted laparoscopic radical cystectomy. *N Engl J Med* 2014;371:389–90.
- [81] Rhee JJ, Lebeau S, Smolkin M, Theodorescu D. Radical cystectomy with ileal conduit diversion: early prospective evaluation of the impact of robotic assistance. *BJU Int* 2006;98:1059–63.

- [82] Galich A, Sterrett S, Nazemi T, Pohlman G, Smith L, Balaji KC. Comparative analysis of early perioperative outcomes following radical cystectomy by either the robotic or open method. *JSLs* 2006;10:145–50.
- [83] Pruthi RS, Wallen EM. Robotic assisted laparoscopic radical cystoprostatectomy: operative and pathological outcomes. *J Urol* 2007;178:814–8.
- [84] Ng CK, Kauffman EC, Lee MM, et al. A comparison of postoperative complications in open versus robotic cystectomy. *Eur Urol* 2010;57:274–81.
- [85] Richards KA, Hemal AK, Kader AK, Pettus JA. Robot assisted laparoscopic pelvic lymphadenectomy at the time of radical cystectomy rivals that of open surgery: single institution report. *Urology* 2010;76:1400–4.
- [86] Martin AD, Nunez RN, Castle EP. Robot-assisted radical cystectomy versus open radical cystectomy: a complete cost analysis. *Urology* 2011;77:621–5.
- [87] Gondo T, Yoshioka K, Nakagami Y, et al. Robotic versus open radical cystectomy: prospective comparison of perioperative and pathologic outcomes in Japan. *Jpn J Clin Oncol* 2012;42:625–31.
- [88] Khan MS, Challacombe B, Elhage O, et al. A dual-centre, cohort comparison of open, laparoscopic and robotic-assisted radical cystectomy. *Int J Clin Pract* 2012;66:656–62.
- [89] Styn NR, Montgomery JS, Wood DP, et al. Matched comparison of robotic-assisted and open radical cystectomy. *Urology* 2012;79:1303–8.
- [90] Sung HH, Ahn JS, Seo SI, et al. A comparison of early complications between open and robot-assisted radical cystectomy. *J Endourol* 2012;26:670–5.
- [91] Anderson CB, Morgan TM, Kappa S, et al. Ureteroenteric anastomotic strictures after radical cystectomy-does operative approach matter? *J Urol* 2013;189:541–7.

- [92] Kader AK, Richards KA, Krane LS, Pettus JA, Smith JJ, Hemal AK. Robot-assisted laparoscopic vs open radical cystectomy: comparison of complications and perioperative oncological outcomes in 200 patients. *BJU Int* 2013;112:E290–4.
- [93] Knox ML, El-Galley R, Busby JE. Robotic versus open radical cystectomy: identification of patients who benefit from the robotic approach. *J Endourol* 2013;27:40–4.
- [94] Maes AA, Brunkhorst LW, Gavin PW, Todd SP, Maatman TJ. Comparison of robotic-assisted and open radical cystectomy in a community-based, non-tertiary health care setting. *J Robot Surg* 2013:1–5.
- [95] Musch M, Janowski M, Steves A, et al. Comparison of early postoperative morbidity between robot-assisted and open radical cystectomy - results of a prospective observational study. *BJU Int* 2014;113:458–67.
- [96] Nepple KG, Strobe SA, Grubb RL III, Kibel AS. Early oncologic outcomes of robotic vs. open radical cystectomy for urothelial cancer. *Urol Oncol* 2013;31:894–8.
- [97] Trentman TL, Fassett SL, McGirr D, et al. Comparison of anesthetic management and outcomes of robot-assisted versus open radical cystectomy. *J Robot Surg* 2013;7:273–9.
- [98] Ahdoot M, Almario L, Araya H, Busch J, Conti S, Gonzalgo ML. Oncologic outcomes between open and robotic-assisted radical cystectomy: a propensity score matched analysis. *World J Urol* 2014;32:1441–6.
- [99] Wang GJ, Barocas DA, Raman JD, Scherr DS. Robotic vs open radical cystectomy: prospective comparison of perioperative outcomes and pathological measures of early oncological efficacy. *BJU Int* 2008;101:89–93.
- [100] Abraham JB, Young JL, Box GN, Lee HJ, Deane LA, Ornstein DK. Comparative analysis of laparoscopic and robot-assisted radical cystectomy with ileal conduit urinary diversion. *J Endourol* 2007;21:1473–80.

- [101] Shabsigh A, Korets R, Vora KC, et al. Defining early morbidity of radical cystectomy for patients with bladder cancer using a standardized reporting methodology. *Eur Urol* 2009;55:164–74.
- [102] Novara G, De Marco V, Aragona M, et al. Complications and mortality after radical cystectomy for bladder transitional cell cancer. *J Urol* 2009;182:914–21.
- [103] Stimson CJ, Chang SS, Barocas DA, et al. Early and late perioperative outcomes following radical cystectomy: 90-day readmissions, morbidity and mortality in a contemporary series. *J Urol* 2010;184:1296–300.
- [104] Svatek RS, Fisher MB, Matin SF, et al. Risk factor analysis in a contemporary cystectomy cohort using standardized reporting methodology and adverse event criteria. *J Urol* 2010;183:929–34.
- [105] Aziz A, May M, Burger M, et al. Prediction of 90-day mortality after radical cystectomy for bladder cancer in a prospective European multicenter cohort. *Eur Urol* 2014;66:156–63.
- [106] Birkmeyer JD, Siewers AE, Finlayson EV, et al. Hospital volume and surgical mortality in the United States. *N Engl J Med* 2002;346:1128–37.
- [107] Birkmeyer JD, Stukel TA, Siewers AE, Goodney PP, Wennberg DE, Lucas FL. Surgeon volume and operative mortality in the United States. *N Engl J Med* 2003;349:2117–27.
- [108] Gandaglia G, Popa I, Abdollah F, et al. The effect of neoadjuvant chemotherapy on perioperative outcomes in patients who have bladder cancer treated with radical cystectomy: a population-based study. *Eur Urol* 2014;66:561–8.
- [109] Yu HY, Hevelone ND, Lipsitz SR, et al. Comparative analysis of outcomes and costs following open radical cystectomy versus robot-assisted laparoscopic radical cystectomy: results from the US Nationwide Inpatient Sample. *Eur Urol* 2012;61:1239–44.

Figure legends

Fig. 1 – Flowchart of the systematic review.

Fig. 2 – Comparison of (a) operative time, (b) blood loss, (c) transfusion rates, (d) intraoperative complication rates, and (e) in-hospital stay following robot-assisted radical cystectomy or open radical cystectomy.

CI = confidence interval; M-H = Mantel-Haenszel test; ORC = open radical cystectomy; RARC = robot-assisted radical cystectomy; SD = standard deviation; WMD = weighted mean difference.

Fig. 3 – Comparison of transfusion rates following robot-assisted radical cystectomy or laparoscopic radical cystectomy.

CI = confidence interval; LRC = laparoscopic radical cystectomy; M-H = Mantel-Haenszel test; RARC = robot-assisted radical cystectomy.

Fig. 4 – Comparison of rates for any grade of complication at (a) 30 d and (b) 90 d, (c) grade 3 complications at 30 and 90 d, (d) mortality at 30 and 90 d, and (e) major complication at 30 and 90 d following robot-assisted radical cystectomy or open radical cystectomy.

CI = confidence interval; M-H = Mantel-Haenszel test; ORC = open radical cystectomy; RARC = robot-assisted radical cystectomy.

Fig. 5 – Comparison of rates at 30 d for (a) any grade of complication and (b) grade 3 complications following robot-assisted radical cystectomy or laparoscopic radical cystectomy.

CI = confidence interval; LRC = laparoscopic radical cystectomy; M-H = Mantel-Haenszel test; RARC = robot-assisted radical cystectomy.

Table 1 – Perioperative outcomes in robot-assisted radical cystectomy series stratified by urinary diversion type

Reference	Institution	IDEAL stage	Cases	Study design	Nerve-sparing surgery	Median/mean operative time, min	Median/mean blood loss, ml	Transfusion rate, %	Intraoperative complications, %	Mean time to flatus, d	Mean time to bowel movement, d	In-hospital stay, d	Readmission rate
Mainly extracorporeal conduit diversion													
Guru et al, 2007 [8]	Roswell Park Cancer Institute, Buffalo, NY, USA	1	7	Prospective	–	–	335	–	0	–	–	8	–
Guru et al, 2007 [9]	Roswell Park Cancer Institute, Buffalo, NY, USA	1	20	Prospective	–	442	555	0	0	–	4	10	10%
Mottrie et al, 2007 [10]	O.L.V. Clinic, Aalst, Belgium	2a	27	Retrospective	29%	340	301	7	–	–	–	–	–
Hemal et al, 2008 [11]	All India Institute of Medical Sciences, New Delhi, India	1	6	Retrospective	–	330	200	17	0	–	–	9.2	0
Lowentritt et al, 2008 [12]	Tulane University Health Center, New Orleans, LA, USA	2a	4	Retrospective	–	375	338	0	0	–	–	5	–
Murphy et al, 2008 [13]	Guy's & St Thomas' NHS Foundation Trust, London, UK	2a	23	Retrospective	20%	397	278	4	–	–	–	11.6	–
Park et al, 2008 [14]	Yonsei University College of Medicine, Seoul, Korea	2a	4	Retrospective	–	355	550	–	–	–	–	12	–
Park et al, 2008 [15]	Yonsei University College of Medicine, Seoul, Korea	2a	11	Retrospective	–	309	615	–	–	–	–	–	–
Pruthi et al, 2008 [16]	University of North Carolina, Chapel Hill, NC, USA	2a	20	Retrospective	85%	366	313	–	5	2.1	2.8	4.4	–
Pruthi et al, 2008 [17]	University of North Carolina, Chapel Hill, NC, USA	2a	12 female	Retrospective	0	276	221	–	–	1.9	2.4	4.8	–
Pruthi et al, 2008 [18]	University of North Carolina, Chapel Hill, NC, USA	2b	50	Retrospective	–	306	271	–	–	2	2.6	4.5	–
Woods et al, 2008 [19]	Multicenter	2b	27	Retrospective	–	499	277	11	–	–	–	–	–
Yuh et al, 2008 [20]	Roswell Park Cancer Institute, Buffalo, NY, USA	2a	54	Retrospective	–	–	557	13	0	–	–	9.1	–
Gamboa et al, 2009 [21]	University of California, Irvine, CA, USA	2a	41	Retrospective	–	498	254	44	7	–	4	8	–
Pruthi et al, 2009 [22]	University of North Carolina, Chapel Hill, NC, USA	2b	50	Retrospective	–	302	268	–	–	1.9	2.6	4.5	–
Yuh et al, 2009 [23]	Roswell Park Cancer	2b	73	Retrospective	–	378	573	–	1	–	–	10	–

	Institute, Buffalo, NY, USA												
Hayn et al, 2010 [24]	Multicenter	2b	482	Retrospective	–	385	408	–	–	–	–	–	–
Josephson et al, 2010 [25]	City of Hope Comprehensive Cancer Center, Duarte, CA, USA	2b	58	Retrospective	–	480	450	38	7	–	–	10	–
Kang et al, 2010 [26]	Multicenter	2b	104	Retrospective	–	554	526	–	4	3.4	–	18.4	–
Kasraeian et al, 2010 [27]	Montsouris Institute, Paris, France	2a	9	Retrospective	–	270	400	55	–	–	–	14	–
Kauffman et al, 2010 [28]	Weill Cornell Medical Center, New York, NY, USA	2b	79	Retrospective	–	360	400	3	0	–	–	5	–
Kwon et al, 2010 [29]	Kyungpook National University, Daegu, Korea	2a	17	Prospective	–	379	210	35	0	–	–	20.7	–
Martin et al, 2010 [30]	Multicenter	2b	59	Retrospective	–	–	–	–	–	–	–	–	–
Pruthi et al, 2010 [31]	University of North Carolina, Chapel Hill, NC, USA	2b	100	Retrospective	–	276	250/271	–	2	2.1	2.8	4.9	11% (30 d)
Hayn et al, 2011 [32]	Roswell Park Cancer Institute, Buffalo, NY, USA	2b	156	Prospective	–	577	400	16	–	–	–	8	21%
Khan et al, 2011 [33]	Guy's & St Thomas' NHS Foundation Trust, London, UK	2a	50	Prospective	–	361	340	4	4	–	–	10	18%
Lavery et al, 2011 [34]	Ohio State University, Columbus, OH, USA	2a	15	Retrospective	–	423	160	–	–	–	–	3.4	13% (30 d)
Richards et al, 2011 [35]	Wake Forest University Baptist Medical Center, Winston-Salem, NC, USA	2b	60	Retrospective	–	492	483	–	3	–	–	8.1	–
Shah et al, 2011 [36]	Ohio State University, Columbus, OH, USA	2b	30	Retrospective	–	411	170	3	–	–	–	3.3	20%
Lau et al, 2012 [37]	City of Hope Comprehensive Cancer Center, Duarte, CA, USA	2b	23 aged >80 yr	Retrospective	–	384	300	61	4	–	–	13	–
Mmeje et al, 2013 [38]	Multicenter	2b	50	Retrospective	–	–	–	–	–	–	–	–	–
Smith et al, 2012 [39]	Multicenter	2b	227	Retrospective	–	291/327	200/256	–	–	–	–	5	–
Treyer et al, 2012 [40]	University of Saarland, Homburg/Saar, Germany	2b	91	Retrospective	–	412	294	–	–	2.1	2.9	18.8	11% (30 d)
Tsui et al, 2012 [41]	Chang Gung University College of Medicine, Taoyuan, Taiwan	2a	8	Retrospective	–	430	763	63 (intraoperative)	–	–	–	10.8	–
Abbas et al, 2013 [42]	Cairo University, Cairo, Egypt	2a	25	Retrospective	–	618	700	40	–	–	–	–	–
Johar et al, 2013 [43]	Multicenter	2b	939	Retrospective	–	–	580	15	–	–	–	8	20%

	Cancer Center, Duarte, CA, USA												
Yuh et al, 2014 [59]	City of Hope Comprehensive Cancer Center, Duarte, CA, USA	2b	162	Retrospective	–	438	400	–	–	–	–	–	–
Overall*						420 (range: 300–496)	390 (range: 167–400)	44	0	–	–	8.9 (range: 6.7–9)	39%
Mainly intracorporeal conduit diversion													
Yohannes et al, 2003 [60]	Creighton University, Omaha, NE, USA	1	2	Retrospective	0	660	1118	–	–	–	–	6	–
Pruthi et al, 2010 [61]	University of North Carolina, Chapel Hill, NC, USA	2a	12	Retrospective	–	318	221	–	0	2.2	3.2	4.5	17%
Jonsson et al, 2011 [62]	Karolinska Institute, Stockholm, Sweden	2b	9	Prospective	–	460	350	–	–	–	–	17	–
Goh et al, 2012 [63]	Keck School of Medicine, University of Southern California, Los Angeles, CA, USA	2a	7	Prospective	–	450	200	71	0	–	–	9	43%
Poch et al, 2012 [64]	Roswell Park Cancer Institute, Buffalo, NY	2b	56	Retrospective	–	356	338	–	–	–	–	7.9	29% (30 d)
Azzouni et al, 2013 [65]	Roswell Park Cancer Institute, Buffalo, NY, USA	2b	100	Retrospective	–	352	300	10	–	–	–	9	16% (30 d) /20% (90 d)
Bishop et al, 2013 [66]	Hertfordshire and South Bedfordshire Urological Cancer Centre, Lister Hospital, Stevenage, UK	2a	8	Not reported	–	360	225	25	–	–	–	9	0
Collins et al, 2013 [67]	Karolinska Institute, Stockholm, Sweden	2b	43	Prospective	16%	292	200	–	–	–	–	9	–
Overall*						340 (range: 292–660)	270 (range: 200–1118)	14.7% (range: 10–71%)	0	2.2	3.2	8.6 (range: 4.5–9)	19.7% (range: 0–43%)
Mainly intracorporeal continent diversion													
Jonsson et al, 2011 [62]	Karolinska Institute, Stockholm, Sweden	2b	36	Prospective	–	480	625	–	–	–	–	9	–
Schumacher et al, 2011 [68]	Karolinska Institute, Stockholm, Sweden	2b	45	Retrospective	–	476	669	–	–	–	–	9	–
Goh et al, 2012 [63]	Keck School of Medicine, University of Southern California, Los Angeles, CA, USA	2a	8	Prospective	–	450	225	37	0	–	–	8	75%
Collins et al, 2013 [67]	Karolinska Institute, Stockholm, Sweden	2b	70	Prospective	70%	420	500	–	–	–	–	9	–

Tyritzis et al, 2013 [69]	Karolinska Institute, Stockholm, Sweden	2b	70	Prospective	58% BNS 8% UNS	420	500	4	–	–	–	9	–
Overall*						420 (range: 420–450)	480 (range: 225–500)	7% (range: 4–37%)	0	–	–	8.5 (range: 8–9)	75%

BNS = bilateral nerve sparing; UNS = unilateral nerve sparing.

* Once duplicate publications from the same centers and multicenter papers were excluded.

Table 2 – Impact of patient characteristics on perioperative outcomes in robot-assisted radical cystectomy series

Reference	Institution	IDEAL stage	Cases	Study design	Intracorporeal urinary diversion %	Conduit diversion, %	Median/mean operative time, min	Median/mean blood loss, ml	Transfusion rate, %	Intraoperative complications, %	Mean time to flatus, d	Mean time to bowel movement, d	In-hospital stay, d	Readmission rate
Patient BMI														
Butt et al, 2008 [70]	Roswell Park Cancer Institute, Buffalo, NY, USA	2a	BMI <25: 14	Retrospective	0	100	359	630	–	0	–	–	11.8	–
			BMI 25–29: 18			89	366	496					7.7	
			BMI ≥30: 17			94	371	532					9.1	
Poch et al, 2012 [64]	Roswell Park Cancer Institute, Buffalo, NY, USA	2b	56	Retrospective	100	100	356	338	–	–	–	–	7.9	29% (30 d)
			BMI <25 : 14				349	150					7	23% (30 d)
			BMI 25 to <30: 21				380	300					8	33% (30 d)
			BMI ≥30: 21				349	500					8.5	32% (30 d)

BMI = body mass index.

Table 3 – Impact of surgical factors on perioperative outcomes in robot-assisted radical cystectomy series

Reference	Institution	IDEAL stage	Cases	Study design	Intracorporeal urinary diversion, %	Conduit diversion, %	Median/mean operative time, min	Median/mean blood loss, ml	Transfusion rate, %	Intraoperative complications	Mean time to flatus, d	Mean time to bowel movement, d	In-hospital stay, d	Readmission rate
Case volume														
Pruthi et al, 2008 [18]	University of North Carolina, Chapel Hill, NC, USA	2b	50	Retrospective	0	58	306	271	–	–	2	2.6	4.5	–
			Cases 1–10			70	378	335			2.1	2.8	4.2	
			Cases 11–20			60	342	330			2.2	3	4.6	
			Cases 21–30			50	276	245			1.9	2.4	4.6	
			Cases 31–40			40	270	233			1.7	2.1	4.2	
			Cases 41–50			70	264	210			1.9	2.6	4.9	
Hayn et al, 2011 [71]	Roswell Park Cancer Institute, Buffalo, NY, USA	2a	Cases 1–50	Prospective	–	93	–	566	–	–	–	–	–	–
			Cases 51–100					631						
			Case 101–164					521						
Richards et al, 2011 [35]	Wake Forest University Baptist Medical Center, Winston-Salem, NC, USA	2b	60	Retrospective	0	92	492	483	–	3%	–	–	8.1	–
			Cases 1–20			80	524	511					9.2	
			Cases 21–40			95	503	459					7.8	
			Cases 41–60			100	449	479					7.4	
Schumacher et al, 2011 [68]	Karolinska Institute, Stockholm, Sweden	2b	45	Retrospective	100	20	476	669	–	–	–	–	9	–
			Cases 1–15			33	532	627					12	
			Cases 16–30			20	462	728					8	
			Cases 31–45			7	434	654					8	
Azzouni et al, 2013 [65]	Roswell Park Cancer Institute, Buffalo, NY, USA	2b	100	Retrospective	100	100	352	300	10	–	–	–	9	16% (30 d), 20% (90 d)
			Cases 1–25				366	400	0				7	12% (30 d), 12% (90 d)
			Cases 26–50				349	350	12				9	20% (30 d), 28% (90 d)
			Cases 51–				373	300	4				10	20% (30 d)

			75											d), 24% (90 d)
			Cases 76–100				344	200	24				9	16% (30 d), 20% (90 d)
Prior RARP experience														
Hayn et al, 2010 [72]	Multicenter	2b	482	Retrospective	–	75	385	408	–	–	–	–	–	–
			≤50 previous RARP: 83				421	418						
			51–100 previous RARP: 173				338	286						
			101–150 previous RARP: 168				401	575						
			>150 previous RARP: 48				444*	188*						
Intracorporeal vs extracorporeal diversion														
Guru et al, 2010 [73]	Roswell Park Cancer Institute, Buffalo, NY, USA	2a	13 extracorporeal ileal conduit	Prospective	0	100	387	454	–	0	–	–	8.5	23%
			13 intracorporeal ileal conduit		100		391	315		8			8.8	8%
Kang et al, 2012 [74]	Korea University School of Medicine, Seoul	2a	22 extracorporeal ileal conduit	Retrospective	0	100	420	370	–	–	2.5	–	14.5	–
			3 intracorporeal ileal conduit		100		510	400		14.2				
Kang et al, 2012 [74]	Korea University School of Medicine, Seoul	2a	14 extracorporeal neobladder	Retrospective	0	0	496	390	–	–	2.3	–	16.8	–
			1 intracorporeal neobladder		100		545	500		2.5			14	

RARP = robot-assisted radical prostatectomy.

Table 4 – Complication rates in robot-assisted radical cystectomy series

Reference	Institution	IDE AL stage	Cases	Study design	Martin criteria	Follow-up duration	Overall complication rate, %	Complication rate, %				
								Grade 1	Grade 2	Grade 3	Grade 4	Grade 5
Mainly extracorporeal conduit diversion												
Guru et al, 2007 [8]	Roswell Park Cancer Institute, Buffalo, NY, USA	1	7	Prospective	4	–	14	0	14	0	0	0
Guru et al, 2007 [9]	Roswell Park Cancer Institute, Buffalo, NY, USA	1	20	Prospective	4	–	–	–	–	10	0	5
Hemal et al, 2008 [11]	All India Institute of Medical Sciences, New Delhi, India	1	6	Retrospective	3	–	–	–	–	0		
Murphy et al, 2008 [13]	Guy's & St Thomas' NHS Foundation Trust, London, UK	2a	23	Retrospective	3	–	26	4	4	18	–	–
Park et al, 2008 [14]	Yonsei University College of Medicine, Seoul, Korea	2a	4	Retrospective	–	–	–	–		0		
Pruthi et al, 2008 [16]	University of North Carolina, Chapel Hill, NC, USA	2a	20	Retrospective	5	90 d	30	15		15	–	–
Woods et al, 2008 [19]	Multi-institutional	2b	27	Retrospective	3	–	33	–		–		
Gamboa et al, 2009 [21]	University of California, Irvine, CA, USA	2a	41	Retrospective	4	–	29	17		12		0
Kauffman et al, 2010 [28]	Weill Cornell Medical Center, New York, NY, USA	2b	79	Retrospective	9	30 d	69	58		8	3	0
						90 d	100	37	42	17	4	
Kang et al, 2010 [26]	Multicenter	2b	104	Retrospective	8	–	27	19		6	0	2
Kwon et al, 2010 [29]	Kyungpook National University, Daegu, Korea	2a	17	Prospective	7	–	29	29		0	0	0
Pruthi et al, 2010 [31]	University of North Carolina, Chapel Hill, NC	2b	100	Retrospective	7	30 d	36	28		8		
Hayn et al, 2011 [32]	Roswell Park Cancer Institute, Buffalo, NY, USA	2b	156	Prospective	10	30 d	40	10	17	11	0	2
						90 d	48	14	21	15	0	2
Khan et al, 2011 [33]	Guy's & St Thomas' NHS Foundation Trust, London, UK	2a	50	Prospective	8	90 d	34	6	18	10	0	0
Lau et al, 2012 [37]	City of Hope Comprehensive	2b	23 (aged >80 yr)	Retrospective	8	30 d	78	4	58	31	0	4

	Cancer Center, Duarte, CA, USA											
Saar et al, 2013 [46]	University of Saarland, Homburg/Saar, Germany	2b	62	Retrospective	9	30 d	44	11	23	8	0	2
Smith et al, 2012 [39]	Multicenter	2b	227	Retrospective	5	30 d	30	23		7		0
Treyer et al, 2012 [40]	University of Saarland, Homburg/Saar, Germany	2b	91	Retrospective	6	30 d	49	15	23	7	3	1
Johar et al, 2013 [43]	Multicenter	2b	939	Retrospective	9	30 d	41	-		-	-	1.3
						90 d	48	29		14.8		4.2
Xylinas et al, 2013 [47]	Weill Cornell Medical Center, New York, NY, USA	2b	175	Retrospective	10	30 d	42	8	22	7	3	2
						90 d	45	-	-	-	-	4
Al-Daghmin et al, 2014 [48]	Roswell Park Cancer Institute, Buffalo, NY, USA	2b	272	Prospective	8	30 d	-	-		-		1
						90 d	77	58		14		5
Phillips et al, 2014 [50]	Boston Medical Center, Boston, MA, USA	2b	23 (aged ≥80 yr)	Retrospective	7	90 d	35	0	31	4	0	0
Snow-Lisy et al, 2014 [51]	Cleveland Clinic Lerner College of Medicine, Cleveland, OH, USA	2b	17	Retrospective	5	-	53	12	12	17	12	0
Overall*						30 d	Any-grade complication: 44% (range: 26–78%)	Low-grade complication: 29.4% (range: 8–62%)		High-grade complication: 11.8% (range: 0–35%) Reoperation: 9.7% (range: 0–31%) Mortality: 1.6% (range: 0–4%)		
						90 d	Any-grade complication: 59% (range: 30–77%)	Low-grade complication: 54% (range: 15–79%)		High-grade complication: 15% (range: 4–19%) Reoperation: 14% (range: 4–17%) Mortality: 3% (range: 0–5%)		
Mainly extracorporeal continent diversion												
Josephson et al, 2010 [25]	City of Hope Comprehensive Cancer Center, Duarte, CA, USA	2b	58	Retrospective	4	90 d	69	64		3		2
Kasraeian et al, 2010 [27]	Montsouris Institute, Paris, France	2a	9	Retrospective	5	-	33	0	11	22	0	0
Torrey et al, 2012 [55]	City of Hope Comprehensive Cancer Center, Duarte, CA, USA	2b	34	Retrospective	9	90 d	91	16	69	14	0	1
Yuh et al, 2012 [56]	City of Hope Comprehensive Cancer Center,	2b	196	Retrospective	10	90 d	80	45		31		4

	Duarte, CA, USA											
Nazmy et al, 2014 [58]	City of Hope Comprehensive Cancer Center, Duarte, CA, USA	2b	209	Retrospective	10	90 d	77	45	27	5		
Yuh et al, 2014 [59]	City of Hope Comprehensive Cancer Center, Duarte, CA, USA	2b	162	Retrospective	8	52 mo	82	45	37			
Mainly intracorporeal conduit diversion												
Pruthi et al, 2010 [61]	University of North Carolina, Chapel Hill, NC, USA	2a	12	Retrospective	5	30 d	42	–	–	–	–	–
						90 d	58					
Jonsson et al, 2011 [62]	Karolinska Institute, Stockholm, Sweden	2b	9	Prospective	8	30 d	44	11	0	22	11	0
						After first 30 d	33	22	0	22	0	0
Goh et al, 2012 [63]	Keck School of Medicine, University of Southern California, Los Angeles, CA, USA	2a	7	Prospective	8	30 d	45	45	0			
						After first 30 d	14	14				
Azzouni et al, 2013 [65]	Roswell Park Cancer Institute, Buffalo, NY, USA	2b	100	Retrospective	8	30 d	63	50	13			
						90 d	81	66	15			
Bishop et al, 2013 [66]	Hertfordshire and South Bedfordshire Urological Cancer Centre, Lister Hospital, Stevenage, UK	2a	8	Not reported	6	30 d	75	25	25	25	0	0
Collins et al, 2013 [67]	Karolinska Institute, Stockholm, Sweden	2b	43	Prospective	8	30 d	86	9	23	42	12	0
						After first 30 d	23	0	0	19	2	2
Overall*						30 d	Any-grade complication: 67% (range: 42–86%)	Low-grade complication: 45% (range: 32–50%)		High-grade complication: 24% (range: 0–54%) Reoperation: 39% (range: 25–42%) Mortality: 0%		
						30–90 d	Any-grade complication: 22% (range: 14–23%)	Low-grade complication: 2% (range: 0–14%)		High-grade complication: 20% (range: 0–23%) Reoperation: 19% Mortality: 1.7% (range: 0–2%)		
						90 d	Any-grade complication: 59% (range: 30–77%)	Low-grade complication: 66%		High-grade complication: 15% Reoperation: 25% (range: 14–51%) Mortality: 1.7% (range: 0–2%)		
Mainly intracorporeal continent diversion												
Akbulut et al, 2011 [75]	Ankara Ataturk training and research hospital	2a	12	Not reported	7	30 d	67	25	25	17	0	0
						After first 30 d	41	8	17	8	0	8
Jonsson et al, 2011 [62]	Karolinska Institute, Stockholm, Sweden	2b	36	Prospective	8	30 d	39	14	6	19	0	0
						After first 30 d	33	14	3	16	0	0
Schumacher et al, 2011 [68]	Karolinska Institute, Stockholm, Sweden	2b	45	Retrospective	10	30 d	40	13	4	20	2	0
						After first 30 d	31	11	2	18	0	0

Canda et al, 2012 [76]	Ankara Ataturk training and research hospital	2a	27	Not reported	6	30 d	48	11	22	11	0	4
						After first 30 d	27	4	11	8	0	4
Goh et al, 2012 [63]	Keck School of Medicine, University of Southern California, Los Angeles, CA, USA	2a	8	Prospective	8	30 d	62	37		25		0
						After first 30 d	12	-		12		0
Collins et al, 2013 [67]	Karolinska Institute, Stockholm, Sweden	2b	70	Prospective	8	30 d	43	4	8	20	11	0
						After first 30 d	34	0	13	19	1	1
Overall*						30 d	Any-grade complication: 45.7% (range: 43–62%)	Low-grade complication: 19% (range: 12–33%)		High-grade complication: 28% (range: 15–33%) Reoperation: 17% (range: 11–20%) Mortality: 1% (range: 0–4%)		
						30–90 d	Any-grade complication: 30% (range: 12–34%)	Low-grade complication: 13.5% (range: 13–15%)		High-grade complication: 18% (range: 12–21%) Reoperation :16% (range: 8–19%) Mortality: 1.7% (range: 0–4%)		
						90 d	-	-		Reoperation: 33% (range: 19–39%) Mortality: 2.7% (range: 0–8%)		

* Once duplicate publications from the same centers and multicenter papers were excluded.

Table 5 – Impact of patients characteristics and surgical factors on complication rates in robot-assisted radical cystectomy series

Reference	Institution	IDEAL stage	Cases	Study design	Intracorporeal urinary diversion, %	Conduit diversion, %	Martin criteria	Follow-up duration	Overall complication rate, %	Complication rate, %										
										Grade 1	Grade 2	Grade 3	Grade 4	Grade 5						
Patients BMI																				
Butt et al, 2008 [70]	Roswell Park Cancer Institute, Buffalo, NY, USA	2a	BMI <25: 14	Retrospective	0	100	5	–	21	–	–	–	–	0						
			BMI 25–29: 18			89			33					0						
			BMI ≥30: 17			94			24					6						
Poch et al, 2012 [64]	Roswell Park Cancer Institute, Buffalo, NY, USA	2b	56	Retrospective	100	100	5	90 d	57	–	–	–	–	–						
			BMI <25 : 14						43											
			BMI 25 to <30: 21						67											
			BMI ≥30: 21						57											
Case volume																				
Hayn et al, 2011 [71]	Roswell Park Cancer Institute, Buffalo, NY, USA	2a	Cases 1–50	Prospective	–	93	9	240 d	68	26	18	18	0	6						
			Cases 51–100						62						14	22	24	0	2	
			Case 101–164						62						17	23	21	0	0	
Richards et al, 2011 [35]	Wake Forest University Baptist Medical Center, Winston-Salem, NC, USA	2b	60	Retrospective	0	92	6	90 d	43	3	53	33	10	3						
			Cases 1–20						80						70*	5	45	5	10	5
			Cases 21–40						95						30*	0	20	10	0	0
			Cases 41–60						100						30*	0	15	10	5	0
Schumacher et al, 2011 [68]	Karolinska Institute, Stockholm, Sweden	2b	45	Retrospective	100	20	10	30 d (>30 d)	40 (31) *	13 (11)	4 (2)	20 (18)	2 (0)	0						
			Cases 1–15						33						66 (53) *	27 (7)	7 (7)	27 (40)	7 (0)	
			Cases 16–30						20						27 (20) *	13 (7)	7 (0)	7 (14)	0 (0)	
			Cases 31–45						7						27 (20) *	0 (20)	0 (0)	27 (0)	0 (0)	
Azzouni et al, 2013 [65]	Roswell Park Cancer Institute, Buffalo, NY, USA	2b	100	Retrospective	100	100	8	30 d (90 d)	63 (81)	50 (66)	13 (15)									
			Cases 1–25						52 (72)						32 (48)	20 (24)				
			Cases 26–50						56 (76)						36 (56)	20 (20)				
			Cases 51–75						76 (88)						68 (80)	8 (8)				
			Cases 76–100						68 (88)						64 (80)	4 (8)				
Collins et al, 2014 [77]	Karolinska	2b	Cases 1–10	Prospective	100	0	6	30 d (>30 d)	70 * (60)*	10 (0)	30 (10)	20 (50)	10 (0)	0						
			Cases 11–20						20 * (40)*						10 (0)	0 (10)	0 (20%)	10%		

	Institute, Stockholm, Sweden												(10%)	
			Cases 21–30						20 * (20)*	0 (0)	0 (20)	20 (0)	0 (0)	
			Cases 31–40						30 * (10)*	0 (0)	0	30 (10)	0 (0)	
			Cases 41–47						29 * (29)*	0 (0)	0 (29)	0 (0)	29 (0)	
Intracorporeal vs extracorporeal diversion														
Guru et al, 2010 [73]	Roswell Park Cancer Institute, Buffalo, NY, USA	2a	13 intracorporeal ileal conduit	Prospective	100	100	7	90 d	30	0	15	15	0	0
			13 extracorporeal ileal conduit		0				38		23	15		
Kang et al, 2012 [74]	Korea University School of Medicine, Seoul, Korea	2a	38 extracorporeal diversion	Retrospective	0	58	5	90 d	42	21			21	
			4 intracorporeal diversion		100	75			25	25			0	

BMI = body mass index.

* Statistically significant.

Table 6 – Predictors of complication rates in robot-assisted radical cystectomy series

Reference	Institution	IDEA L stage	Cases	Study design	Martin criteria	Follow-up duration	Overall complication rate, %	Predictors of complications
Mainly extracorporeal conduit diversion								
Kauffman et al, 2010 [28]	Weill Cornell Medical Center, New York, NY, USA	2b	79	Retrospective	9	30 d	69	Any-grade complications: creatinine level >1.4 mg/dl, i.v. fluids >5000 ml High-grade complications: patients aged >65 yr, EBL >500 ml, and i.v. fluids >5000 ml
						90 d	100	
Smith et al, 2012 [39]	Multicenter	2b	227	Retrospective	5	30 d	30	High-grade complications: aged <65 yr, higher ASA score
Johar et al, 2013 [43]	Multicenter	2b	939	Retrospective	9	30 d	41	Any-grade complications: age, BMI, neoadjuvant chemotherapy, receipt of blood transfusion, conduit diversion High-grade complications: age, BMI, neoadjuvant chemotherapy, current smoking, receipt of blood transfusion
						90 d	48	
Al-Daghmin et al, 2014 [48]	Roswell Park Cancer Institute, Buffalo, NY, USA	2b	272	Prospective	8	30 d	–	30-d readmission: BMI 90-d readmission: sex and BMI
						90 d	77	
Mainly extracorporeal continent diversion								
Yuh et al, 2012 [56]	City of Hope Comprehensive Cancer Center, Duarte, CA, USA	2b	196	Retrospective	10	90 d	80	90-d any-grade complications: age, ASA, preop HCT, OR time, EBL, diversion type 90-d high-grade complications: CCI, preop. HCT, orthotopic diversion
Nazmy et al, 2014 [58]	City of Hope Comprehensive Cancer Center, Duarte, CA, USA	2b	209	Retrospective	10	90 d	77	90-d any-grade complications: ASA, preop HCT, diversion type 90-d high-grade complications: CCI, HCT, diversion type

ASA = American Society of Anesthesiologists; BMI = body mass index; CCI = Charlson comorbidity index; EBL = estimated blood loss; HCT = hematocrit; i.v. = intravenous; OR = operating room; preop = preoperative.

Table 7 – Perioperative parameters and intraoperative complication rates after open, laparoscopic, and robot-assisted radical cystectomy

Comparison	LOE	Reference	No. of cases	Median/mean operative time, min	Median/mean blood loss, ml	Transfusion rate, %	Intraoperative complications, %	Mean time to flatus, d	Mean time to bowel movement, d	In-hospital stay, d
RARC vs ORC	2b									
		Nix et al, 2010 [78]	21 RARC 20 ORC	252 211	258 575	– –	– –	2.3 3.2	3.2 4.3	5.1 6.0
		Parekh et al, 2013 [79]	20 RARC 20 ORC	308 ± 77 288 ± 60	627 ± 554 1113 ± 935	40 50	– –	– –	– –	9.2 ± 7.8 8.9 ± 5.6
		Bochner et al, 2014 [80]	60 RARC 58 ORC	456 ± 82 329 ± 77	– –	– –	– –	– –	– –	8 ± 3 8 ± 5
	4									
		Rhee et al, 2006 [81]	7 RARC 23 ORC	638 ± 46 507 ± 110	479 ± 551 1109 ± 398	57 87	– –	– –	– –	11 ± 2 13 ± 3
		Galich et al, 2006 [82]	13 RARC 24 ORC	697 395	500 1250	54 75	– –	– –	– –	– –
		Pruthi et al, 2007 [83]	20 RARC 24 ORC	366 222	313 588	– –	5 0	2.1 2.9	2.8 3.8	4.4 5.3
		Ng et al, 2010 [84]	83 RARC 104 ORC	375 ± 90 357 ± 132	460 ± 299 1172 ± 916	– –	– –	– –	– –	5.5 8
		Richards et al, 2010 [85]	35 RARC 35 ORC	530 240	350 1000	17 71	– –	– –	– –	7 8
		Martin et al, 2011 [86]	19 RARC 14 ORC	280 320	255 696	– –	– –	– –	– –	5 10
		Gondo et al, 2012 [87]	11 RARC 15 ORC	408.5 ± 55.886 363 ± 111.255	656.9 ± 452.02 1788.7 ± 972.13	0 40	9 0	– –	– –	40.2 ± 9.282 37 ± 9.921
		Khan et al, 2012 [88]	48 RARC 52 ORC	386 320	337 1351	4 58	– –	– –	– –	9.9 19.2

		Styn et al, 2012 [89]	50 RARC 100 ORC	455 ± 100 349 ± 87	350 475	4 24	– –	– –	– –	9.5 ± 8.8 10.2 ± 8.4
		Sung et al, 2012 [90]	35 RARC 104 ORC	578 ± 153 501 ± 110	448.0 ± 231.6 1063.4 ± 892.7	11 57	1 0	– –	– –	28.9 ± 11.9 27.1 ± 13.4
		Anderson et al, 2013 [91]	103 RARC 375 ORC	403 ± 93 281 ± 77	411 ± 271 806 ± 660	–	–	–	–	–
		Kader et al, 2013 [92]	100 RARC 100 ORC	451 393	420 983	15 47	–	–	–	7.8 12.2
		Knox et al, 2013 [93]	58 RARC 84 ORC	468 396	276 1522	5 80	–	4.3 5.9	–	6.3 10.8
		Maes et al, 2013 [94]	14 RARC 14 ORC	383 268	470 942	7 29	–	–	–	11.2 11.4
		Musch et al, 2014 [95]	100 RARC 42 ORC	410 ± 68 351 ± 92	351 ± 170 810 ± 621	27 60	3 5	–	2.3 ± 1.5 2.3 ± 1.1	17.1 ± 7.6 19.9 ± 12
		Nepple et al, 2013 [96]	36 RARC 29 ORC	410 345	675 1497	39 83	–	–	–	7.9 9.6
		Trentman et al, 2013 [97]	96 RARC 102 ORC	372 ± 73 259 ± 70	257.7 ± 164.3 601.8 ± 491.4	31 60	–	–	–	7.1 ± 5.8 9.8 ± 5
		Ahdoot et al, 2014 [98]	51 RARC 51 ORC	346 369	300 900	22 33	–	–	–	7 7
RARC vs LRC	4									
		Abraham et al, 2007 [100]	14 RARC 20 LRC	410 419	212 653	42 70	7 15	–	–	5.8 ± 0.9 9.4 ± 7.4
		Khan et al, 2012 [88]	48 RARC 58 LRC	386 316	337 480	4 26	–	–	–	9.9 16

LOE = level of evidence; LRC = laparoscopic radical cystectomy; ORC = open radical cystectomy; RARC = robot-assisted radical cystectomy.

Table 8 – Comparative studies evaluating complication rates after open, laparoscopic, and robot-assisted radical cystectomy

Comparison	LOE	Reference	No. of cases	Intracorporeal urinary diversion, %	Conduit diversion, %	Martin criteria	Follow-up	Overall complication rate, %	Complications rate, %				
									Grade 1	Grade 2	Grade 3	Grade 4	Grade 5
RARC vs ORC	2b	Nix et al, 2010 [78]	21 RARC	0	66	3	–	33	–	–	–	–	–
			20 ORC		70			50					
		Parekh et al, 2013 [79]	20 RARC 20 ORC	0	–	4	–	–	–	25 25			
		Bochner et al, 2014 [80]	60 RARC 58 ORC	0				–	–	–	22 21		
	4												
		Galich et al, 2006 [82]	13 RARC 24 ORC	0	100	6	–	15 17	–	–	–	–	0 4
		Wang et al, 2008 [99]	33 RARC 21 ORC	0	53 52	5	–	21 24	–	–	–		
		Pruthi et al, 2007 [83]	20 RARC 24 ORC	0	50	5	90 d	30 33	–	–	–	–	–
		Ng et al, 2010 [84]	83 RARC 104 ORC	0	57 49	10	30 d	41 59	12 7	19 22	8 19	1 6	0 5
	90 d						48 61	13 8	17 23	16 20	1 6	0 6	
		Richards et al, 2010 [85]	35 RARC 35 ORC	0	86	6	30 d	60 66	3 11	37 29	11 14	6 11	3 0
		Gondo et al, 2012 [87]	11 RARC 15 ORC	0	63 60	5	30 d	54 73	18 40	36 27	0 7	0 0	0 0
		Khan et al, 2012 [88]	48 RARC 52 ORC	0	87 90	7	–	42 71	25 40		17 27	0 2	0 2
		Styn et al, U2012 [89]	50 RARC 100 ORC	0	72 72	8	30 d	66 62	72 79		28 21		
		Sung et al, 2012 [90]	35 RARC 104 ORC	0	37 82	8	90 d	63 74	26 5	29 45	6 16	0 4	3 3
		Kader et al, 2013	100	0	97	6	90 d	35	1	25	6	3	1

		[92]	RARC 100 ORC		83			57	7	30	11	11	0			
		Knox et al, 2013 [93]	58 RARC 84 ORC	0	91 89	8	30 d	24	5	12	21	3	2			
							90 d	45 78								
							>90 d	73 88								
		Maes et al, 2013 [94]	14 RARC 14 ORC	0	100 –	4	–	57 78		36 64		21 14				
		Musch et al, 2014 [95]	100 RARC 42 ORC	0	76 –	10	90 d (60 d for ORC)	59 93		35 51		24 43				
RARC vs LRC	4															
		Abraham et al, 2007 [100]	14 RARC 20 LRC	0 0	100 100	5	–	21 55	– –	14 35	7 20	– –	– –			
		Khan et al, 2012 [88]	48 RARC 58 ORC	0	87 96	7	–	42 81		25 41	17 36	0 0	0 3			

LOE = level of evidence; LRC = laparoscopic radical cystectomy; ORC = open radical cystectomy; RARC = robot-assisted radical cystectomy.

Figure 1






Figure 2


(A)




(B)



(C)



(D)



(E)




Figure 3






Figure 4


(A)




(B)



(C)



(D)



(E)






Figure 5

(A)



(B)



Typesetter instructions re EURUROL-D-14-01778

Figure 1

--Change Pubmed to: Medline

--Change 17th to 17: April 17, 2014

Figure 2a–2e

--Change Favours to: Favors

--Delete hyphen: Nonrandomized

--Format p values as: $p < 0.00001$, $p = 0.002$

--Change Chi^2 to: χ^2

Figure 3

--Change Favours to: Favors

--Format p values as: $p < 0.00001$, $p = 0.002$

--Change Chi^2 to: χ^2

Figure 4a–4e

--Change Favours to: Favors

--Delete hyphen: Nonrandomized

--Format p values as: $p < 0.00001$, $p = 0.002$

--Change Chi^2 to: χ^2