



UNIVERSITY OF LEEDS

This is a repository copy of *Barley Hv CIRCADIAN CLOCK ASSOCIATED 1 and Hv PHOTOPERIOD H1 Are Circadian Regulators That Can Affect Circadian Rhythms in Arabidopsis*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/88926/>

Version: Published Version

---

**Article:**

Kusakina, J, Rutterford, Z, Cotter, S et al. (5 more authors) (2015) Barley Hv CIRCADIAN CLOCK ASSOCIATED 1 and Hv PHOTOPERIOD H1 Are Circadian Regulators That Can Affect Circadian Rhythms in Arabidopsis. PLoS One, 10 (6). e0127449. ISSN 1932-6203

<https://doi.org/10.1371/journal.pone.0127449>

---

**Reuse**

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

RESEARCH ARTICLE

# Barley *Hv CIRCADIAN CLOCK ASSOCIATED 1* and *Hv PHOTOPERIOD H1* Are Circadian Regulators That Can Affect Circadian Rhythms in Arabidopsis

Jelena Kusakina<sup>1☯✉</sup>, Zoe Rutterford<sup>2,3☯</sup>, Sean Cotter<sup>1☯</sup>, María C. Martí<sup>2</sup>, David A. Laurie<sup>4</sup>, Andy J. Greenland<sup>3</sup>, Anthony Hall<sup>1</sup>, Alex A. R. Webb<sup>2,\*</sup>

**1** Institute of Integrative Biology, University of Liverpool, Crown Street, Liverpool, United Kingdom,

**2** Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge, United Kingdom,

**3** National Institute of Agricultural Botany, Cambridge, United Kingdom, **4** John Innes Centre, Coney Lane, Norwich, United Kingdom

☯ These authors contributed equally to this work.

✉ Current address: Centre for Plant Sciences, University of Leeds, Leeds, LS2 9JT, United Kingdom

\* [alex.webb@plantsci.cam.ac.uk](mailto:alex.webb@plantsci.cam.ac.uk)



**OPEN ACCESS**

**Citation:** Kusakina J, Rutterford Z, Cotter S, Martí MC, Laurie DA, Greenland AJ, et al. (2015) Barley *Hv CIRCADIAN CLOCK ASSOCIATED 1* and *Hv PHOTOPERIOD H1* Are Circadian Regulators That Can Affect Circadian Rhythms in Arabidopsis. PLoS ONE 10(6): e0127449. doi:10.1371/journal.pone.0127449

**Academic Editor:** David E. Somers, Ohio State University, UNITED STATES

**Received:** December 22, 2014

**Accepted:** April 15, 2015

**Published:** June 15, 2015

**Copyright:** © 2015 Kusakina et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** Relevant data are available at: <https://www.repository.cam.ac.uk/handle/1810/247449>.

**Funding:** ZR is grateful to the National Institute of Agricultural Botany for the award of a Scholarship. The authors acknowledge funding from a Marie Curie Early Stage Training project MEST-CT-2005-020526 for JK, and the Biology and the Biological Sciences Research Council Doctoral Training Programme for SC. AARW and MCM are grateful to the BBSRC for the award of BBSRC Grant BB/M006212/1, which

## Abstract

Circadian clocks regulate many aspects of plant physiology and development that contribute to essential agronomic traits. Circadian clocks contain transcriptional feedback loops that are thought to generate circadian timing. There is considerable similarity in the genes that comprise the transcriptional and translational feedback loops of the circadian clock in the plant Kingdom. Functional characterisation of circadian clock genes has been restricted to a few model species. Here we provide a functional characterisation of the *Hordeum vulgare* (barley) circadian clock genes *Hv CIRCADIAN CLOCK ASSOCIATED 1* (*HvCCA1*) and *Hv PHOTOPERIODH1*, which are respectively most similar to *Arabidopsis thaliana CIRCADIAN CLOCK ASSOCIATED 1* (*AtCCA1*) and *PSEUDO RESPONSE REGULATOR 7* (*AtPRR7*). This provides insight into the circadian regulation of one of the major crop species of Northern Europe. Through a combination of physiological assays of circadian rhythms in barley and heterologous expression in wild type and mutant strains of *A. thaliana* we demonstrate that *HvCCA1* has a conserved function to *AtCCA1*. We find that *Hv PHOTOPERIOD H1* has *AtPRR7*-like functionality in *A. thaliana* and that the effects of the *Hv photoperiod h1* mutation on photoperiodism and circadian rhythms are genetically separable.

## Introduction

Circadian clocks are timing mechanisms that are an adaptation to the Earth's rotation. These circadian clocks are entrained to the day/night cycle by sensing environmental cues such as light and temperature and act as master regulators to synchronise biological events to specific

supported aspects of the study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

times of the day [1]. In plants, the circadian clock has been shown to control approximately one third of the genome and it regulates a wide range of key processes such as hypocotyl elongation, leaf movement, stomatal opening and flowering [1]. Furthermore, accurate and robust circadian function enhances yield, water use efficiency and overall plant performance [2,3].

While the molecular components of the clock are not conserved across taxonomic groups, most circadian oscillators comprise a number of coupled transcription/translation feedback loops [4]. In the model dicotyledonous plant *Arabidopsis*, the central clock is proposed to consist of several interlocking loops [5]. A loop of morning active genes is formed by the partially redundant MYB-transcription factors *AtCCA1* (*CIRCADIAN CLOCK ASSOCIATED 1*) and *AtLHY* (*LATE ELONGATED HYPOCOTYL*) and two members of the pseudo-response regulators family, *AtPRR7* (*PSEUDO-RESPONSE REGULATOR 7*), and *AtPRR9* (*PSEUDO-RESPONSE REGULATOR 9*). *AtCCA1/LHY* activate *AtPRR7/9* and in turn *AtPRR7/9* repress *AtCCA1/LHY*. A loop of evening-active genes is formed by *AtGI* (*GIGANTEA*) and *AtPPR1/TOC1* (*PSEUDO-RESPONSE REGULATOR 1/TIMING OF CAB EXPRESSION*) which both negatively regulate each other. The loops are coupled by the repressive effects of *AtCCA1/LHY* on *AtTOC1* and *AtTOC1* on *AtCCA1/LHY* [6,7]. Another loop of mutual repression exists between the morning loop and an “evening complex” of *AtELF3*, *AtELF4* and *LUX* (*EARLY FLOWERING 3, 4* and *LUXARRHYTHMIO*) through recruitment of the transcriptional repressor *LUX* by *ELF3* to the *PRR9* promoter [8].

Monocot circadian clocks appear to be comprised of a similar structure. For example, in *Lemna gibba*, a floating monocotyledon plant, orthologues for *AtCCA1*, *AtLHY*, *AtGI*, *AtELF3* and several genes similar to the *AtPRR* family have been identified and their function in the *Lemna* circadian clock characterised [9]. In rice, another model monocot plant, several orthologues of central clock genes have also been identified, including *OsGI*, *OsZTL* (*ZEITLUPE*), *OsCCA1/LHY* and members of the *OsPRR* family [10]. These components are under circadian control with similar expression patterns reported for homologous clock genes in *Arabidopsis*.

Most monocots contain only a single orthologues of *AtCCA1/LHY*. For example, rice has a single orthologues of *AtCCA1/LHY* (*OsLHY*) which functions in a similar manner to the *Arabidopsis*, acting as a repressor of the rice *AtTOC1* orthologues [11]. However, *OsLHY* does not homodimerize due to the loss of a key phosphorylation site, which demonstrates divergence between the rice and *Arabidopsis* clocks [11].

Barley contains orthologues of *Arabidopsis* circadian clock genes, with transcripts that are rhythmic suggesting that *HvCCA1*, *HvGI* and *HvPRR1* have a functions similar to their *Arabidopsis* counterparts [12]. For some of the *PRR* homologues it has proven difficult to determine which barley gene is orthologous to *Arabidopsis* members of the family and therefore the designation reflects the uncertainty. *HvPRR37*, *HvPRR73*, describe two genes both of which are orthologues to either *AtPRR3* or *AtPRR7*, and *HvPRR59* and *HvPRR95* are orthologous to either *AtPRR5* or *AtPRR9* [12]. Other circadian clock genes are thought to be important for agronomic traits such as *HvLUX1* and *HvELF3* underlying the *EARLY MATURITY 10* [13] and 8 [14] loci respectively.

*HvPRR37* is also known as *Hv PHOTOPERIOD H 1* (*HvPpd-H1*), which controls sensitivity to photoperiod. The *Ppd-H1* region contains a single *PRR* gene originally thought to be most similar to *AtPRR7* [15]. Landraces from south-west Asia, southern Europe, and the Mediterranean basin have the *Ppd-H1* allele that confers early flowering in long days of the spring. Whereas the photoperiodic-insensitive *ppd-H1* allele is present in landraces from central and northern Europe (this is known as spring barley, because it is planted in the spring and flowers at the end of summer). The reduced response to photoperiod of *ppd-H1* allows spring-sown plants to extend the period of vegetative growth and accumulate additional biomass that supports higher yields [15]. The late-flowering *ppd-H1* allele is recessive, suggesting that reduced

response results from a mutation that impairs gene function [15]. The modern variants of barley used to map *Ppd-H1* were Igri which contains the photoperiod-sensitive allele (*Ppd-H1*) and Triumph which is late flowering (*ppd-H1*). The central role of *HvPpd-H1* (and *TaPpd-D1a*) in flowering is well characterised, but it is not known whether *HvPpd-H1* is a circadian clock gene in barley, or whether it is functionally equivalent to *AtPRR7* or *AtPRR3* because natural variation at the *HvPpd1-H1* has no effect on circadian rhythms of circadian clock gene transcripts but does affect the timing of expression of genes associated with flowering time [12, 14].

Barley represents a challenging system for studying circadian behaviour and it does not have the wide resource tools for molecular genetics developed for the model *Arabidopsis*, therefore we have used a heterologous approach to examine barley clock gene function by mis-expression in *Arabidopsis*. We provide evidence that the barley circadian clock genes have conserved functions with their counterparts in *Arabidopsis*. We demonstrate that *HvCCA1* has *AtCCA1*-like activity and *HvPpd-H1* has *AtPRR7*-like functionality in *Arabidopsis*. We suggest that the effects of *HvPpd-H1* on photoperiodism and circadian rhythms are separable. Our data suggests that flowering time in crops can be manipulated by appropriate breeding strategies without compromising circadian clock function.

## Methods

### Cloning of *HvCCA1* into *Arabidopsis*

The entire *HvCCA1* transcript was amplified using KOD TAQ polymerase (Novagen). *HvCCA1* was cloned into GATEWAY pCR8/GW/TOPO entry vector using pCR8/GW/TOPO TA cloning kit (Invitrogen) following the manufacturer's instructions. *HvCCA1* was then sub-cloned into the binary plasmid PMDC32 using the Gateway LR recombination reaction (Invitrogen). The binary plasmid PMDC32 confers hygromycin resistance in plants and contains a double 35S cauliflower mosaic virus promoter for constitutive expression of the inserted gene [16]. The CaMV35S:*HvCCA1* was introduced into GV3101 *Agrobacterium tumefaciens* and subsequently into *Arabidopsis thaliana* Ws-2 via the floral dip method [17]. Out of 21 primary transformants, nine were identified as homozygous lines carrying an insert at a single locus. *HvCCA1* overexpression was confirmed by RT-PCR (S1 Fig). Homozygous lines were used for all experiments.

### RNA extraction and transcript analysis

For circadian experiments seedlings were grown at 22°C in 12 h:12 h L:D for 7 days before being transferred to continuous light for 54 h. Samples were taken every 6 h until 72 h. Plant tissue was disrupted in the TissueLyser (QIAGEN) and total RNA was extracted using RNeasy plant kit (Qiagen). cDNA was synthesised from 0.5 µg of template RNA using the QuantiTect Reverse Transcription kit (Qiagen). For quantitative PCR, transcript abundance was measured using Power SYBR Green PCR master mix (Applied Biosystems, USA) in the Applied Biosystems 7500 Fast Real-Time PCR System. Transcript abundance of *AtCCA1*, *AtLHY* and *AtTOC1* was normalised to *UBIQUITIN10* (UBQ). The primers for q-PCR reactions are presented in S1 Table. All reactions were carried out in triplicate. Normalization and transcript abundance calculations were automated and performed using the Applied Biosystems 7500 Fast Real-Time PCR System software.

## Non-homologous Complementation of Arabidopsis *PRR* Mutants

Arabidopsis *PRR7* T-DNA insertion mutant (*prp7-11*) was complemented with *HvPpd-H1* and *Hvppd-H1* to test for similarity of function between *AtPRR7* and *HvPpd-H1*. pENTR<sub>PpdH1</sub> (Igri) cDNA and of pENTR<sub>ppdH1</sub> (Triumph) were generated, each containing a fragment comprising the coding region of the *HvPpdH1* or *HvppdH1* cDNA from Cultivar “Igri” or “Triumph” respectively, cloned directionally into a pENTR-D plasmid (Invitrogen, UK). The genomic *AtPRR7* (AT5G02810) promoter (p) sequence was used to drive the *HvPpd-H1* cDNA sequences extending from -871 to +1 bases where the first nucleotide of the inferred ATG initiation codon. The promoter was amplified (S1 Table) from *pAtPRR7:LUC*, a luciferase binary vector supplied by C. Robertson McClung, originally derived from pZPomegaLUCplus [18] by replacement of the gentamicin resistance cassette with the *BASTA* resistance gene from p35SBarn [19]. The resulting vector, pZPBAR, was then made Gateway compatible (Invitrogen, Carlsbad, CA) by inserting the PCR-amplified attR-flanked destination cassette from pK7WG2D [20] at the *Bam*HI and *Hind*III sites upstream of *LUC* to create pZPBAR-DONR. The *AtPRR7* promoter was inserted into pZPBAR-DONR. The binary vector used was pBGW, supplied by the Functional Genomics unit of the Department of Plant Systems Biology (VIB-Ghent University). The 894 bp *pAtPRR7* product and pENTR-D Topo Ppd-H1 and pENTR-D Topo ppd-H1 recipient plasmids were digested with SacII restriction enzyme. The digested pENTR-D Topo PpdH1 and pENTR-D Topo ppdH1 samples were dephosphorylated by adding Shrimp Alkaline Phosphatase (SAP) buffer (Fermentas, UK). The *pAtPRR7* insert DNA and pENTR-D Topo PpdH1 / ppdH1 vector backbones were mixed in a ratio of 3:1 (insert:backbone) with ligase (New England BioLabs, UK) and incubated at 4°C overnight. pENTR-D Topo PpdH1 or pENTR-D Topo ppdH1 with the *AtPRR7* promoter were introduced into chemically competent DH5α *E. coli* cells (Invitrogen, UK). Colony PCR was used to identify bacteria carrying the pENTR-D Topo PpdH1 or pENTR-D Topo ppdH1 with the *AtPRR7* promoter in the correct orientation. Plasmid DNA was isolated using Promega mini-prep Wizard kits (Promega, UK). The insertion and orientation of *AtPRR7* into the pENTR-D Topo backbones was verified again by restriction digestion alongside the parent plasmids using SacI and SacII. Plasmids carrying the *AtPRR7* promoter in the correct insertion were fully sequenced through the *AtPRR7* promoter and the Igri or Triumph coding regions. *A. tumefaciens* strain GV3101 was transformed with *pBGW-PRR7:PpdH1* or *pBGW-PRR7:ppdH1* and then used to transform Arabidopsis by floral dip [17]. Positive transformants were selected by spray with Finale glufosinate ammonium (0.31% v/v; Bayer, Germany). For physiological assays the F2 transgenic plants were used.

To ensure functionality of the promoter construct tissue was collected from 12-day-old seedlings grown in 12 h light–12 h dark cycles (100 μmol m<sup>-2</sup> s<sup>-1</sup>) six h after dawn. Total RNA was extracted from frozen tissue of three biological replicates of at least five pooled plants each, using the RNeasy Plant Mini Kit (QIAGEN) and RNase-Free DNase on-column treatment (QIAGEN). cDNA was synthesized from 500 ng RNA with the RevertAid First Strand cDNA Synthesis Kit (Thermo Scientific) using oligo(dT) primers. The gene-specific products were amplified using the Rotor-Gene SYBR Green PCR Kit on a Rotor-Gene 6000 Real-Time PCR machine (QIAGEN). The primer sequences were: PP2a-F, 5'-TAACGTGGCCAAAATG ATGC-3'; PP2a-R, 5'-GTTCTCCACAACCGCTTGGT-3' and *Hvppd-H1*-F, 5'-GATGGATT CAAAGGCAAGGA-3'; *Hvppd-H1*-R, 5'-GAACAATTGGCTCCTCCAAA-3' [12]. Relative transcript levels were determined by incorporating PCR efficiencies [21].

## Circadian rhythms of leaf movement, delayed fluorescence and analysis of hypocotyl elongation and flowering time in Arabidopsis

For leaf movement, delayed fluorescence and hypocotyl elongation experiments Arabidopsis seed was surface-sterilized in 70% ethanol for 1 min, followed by 50% (w/v) bleach for 10 min and 3 rinses with distilled water. Seeds were resuspended in 0.15% (w/v) agar and stratified at 4°C for 3 days prior to sowing on Murashige and Skoog medium 1.5% (w/v) agar plates. For leaf movement, ten day old seedlings entrained to 12h light 12 h dark cycles were moved to Sanyo MLR350 plant growth chambers and imaged under constant light at 22°C [22].

Delayed chlorophyll fluorescence was imaged in a Sanyo MIR-553 cooled incubator (Sanyo Gallenkamp, UK) using an ORCA-II-BT 1024 16-bit camera (Hamamatsu Photonics, Japan) cooled to -80°C [20]. Images from leaf movement and delayed fluorescence were processed using Metamorph 6.0 image-analysis software (Molecular Devices, UK). Period estimates and relative amplitude errors (RAE) were calculated in BRASS (available from <http://www.amillar.org/downloads.html>) by running fast Fourier transformed non-linear least-square analysis [23]. RAE is a measure of rhythm robustness that ranges from 0 (a perfect fit to the wave) to 1 (no fit).

Hypocotyl length was measured in ten day old seedlings grown at 22°C in 12h light 12 h dark of 100  $\mu\text{mol m}^{-2} \text{s}^{-1}$ . Seedlings were harvested and laid flat on agar plates for imaging. Images of Arabidopsis seedlings were analysed in Metamorph 6.0. Student's *t*-test was used to determine statistical significance between the controls and mutants.

Flowering time assays were performed on Arabidopsis seedlings grown in a controlled environment under 16h light 8 h dark at 22°C. The flowering time was expressed as a number of days and rosette leaves when a bolt of 0.5 cm long was observed.

## Infra-Red Gas Exchange Analyser (IRGA) Assay of Circadian Rhythms in Arabidopsis and barley

Circadian rhythms of gas exchange in Arabidopsis plants was measured as described in [24]. For measurement of circadian gas exchange in individual barley plants an IRGA system based on the portable LI-COR LI-6400XT (LI-COR, USA) was developed. Humid CO<sub>2</sub> free air was obtained by pumping atmospheric air through two 4.5 cm x 30 cm cylinders in series filled with soda lime (Fisher Scientific, UK) and a flask of water. Relative humidity was limited to a maximum of 85% through the control of the IRGA. CO<sub>2</sub> (BOC, UK) was provided at between 180 and 220 PSIG and a flow of 500  $\mu\text{mol s}^{-1}$ . The [CO<sub>2</sub>] in the chamber was controlled at 380 PPM. A 6 cm<sup>2</sup> area of mature barley (4–8 weeks old) leaf was clamped in the cuvette with a clear glass screen and therefore received illumination from the cabinet equivalent to that of the rest of the plant. Gaseous exchange was recorded every 10 min.

## Results

### Over-expression of *HvCCA1* causes arrhythmia of the Arabidopsis circadian clock

Over-expression of *AtCCA1* in Arabidopsis abolishes circadian rhythms in multiple outputs of the circadian clock [25]. Here we investigated whether *HvCCA1* is functionally orthologous to *AtCCA1* by assaying circadian regulation of leaf movement, delayed chlorophyll fluorescence, transcript abundance and gas exchange in Arabidopsis overexpressing *HvCCA1* driven by 2x 35S cauliflower mosaic virus promoters (*HvCCA1*-ox; (S1 Fig). In continuous light (LL), *Ws*-2 and *Col*-0 wildtypes had robust circadian rhythms of delayed fluorescence (*Ws*-2 = 23.4 h; *Col*-0 = 23.6 h; Table 1, Fig 1) and leaf movement (*Ws*-2 = 24.1 h; *Col*-0 = 23.6 h; Table 1, Fig




**Table 1. Summary of mean circadian period estimates, standard error (SE) and percent rhythmic (%Rh) for leaf movement (LM) and delayed fluorescence (DF).**

Line	LM data				DF Data			
	Period (h)	SE	n	%Rh	Period (h)	SE	n	%Rh
Col	23.6	0.1	30	93	23.6	0.1	8	100
CCA1-ox	26.6	1.0	30	70	24.9	0.9	8	38
WS	24.1	0.1	30	100	23.4	0.1	8	100
8-3	20.0	0.6	30	83	25.7	0.6	8	75
14-2	20.4	1.1	30	80	N/A	N/A	N/A	N/A
16-4	24.2	0.1	30	87	27.8	1.6	8	75
17-7	24.1	0.2	30	87	21.2	1.1	8	38
18-1	24.2	0.3	30	77	25.9	1.7	8	75
19-4	24.8	0.1	30	90	24.7	1.3	8	75

N/A—no data available. Col indicates Col-0 and WS indicates Ws-2 ecotypes.

doi:10.1371/journal.pone.0127449.t001

1). By contrast, in *AtCCA1-ox* [25] and *HvCCA1-ox* lines circadian rhythms of both delayed fluorescence and leaf movement were severely disrupted (Fig 1B, 1C and 1D) and there was a lower percentage of rhythmic plants (defined as RAE > 0.5; Table 1). Similarly, rhythms of stomatal conductance and photosynthesis were abolished by *HvCCA1-ox* and *AtCCA1-ox*



**Fig 1. Over expression of *HvCCA1* causes circadian arrhythmia in *Arabidopsis*.** Normalised delayed chlorophyll fluorescence (a) and period estimates vs R.A.E (b) for Ws-2, Col-0, *AtCCA1-ox* (*AtCCA1-ox* 038) and two independent *HvCCA1-ox* transgenic lines ( $n = 8$ ). Period estimates vs R.A.E for leaf movement in LL or individual leaves Col-0 and *AtCCA1-ox* (*AtCCA1-ox* 038) (c) and Ws-2 and two independent transgenic lines of *HvCCA1-ox* (d).  $n = 30$ . All experiments were independently repeated at least twice. Ws-2 (closed squares), Col-0 (closed triangles) *AtCCA1-ox* (open triangles) and two independent *HvCCA1-ox* transgenic lines (8-3 and 18-1) (open squares and diamonds).

doi:10.1371/journal.pone.0127449.g001

**Table 2. The effect of *HvCCA1-ox* on circadian rhythms of photosynthesis, transpiration rate and stomatal conductance in Arabidopsis.**

Line	Photosynthesis				Transpiration rate				Stomatal conductance			
	Period (h)	SE	n	%Rh	Period (h)	SE	n	%Rh	Period (h)	SE	n	%Rh
Col	23.5	0.4	5	100	24.6	0.5	5	60	24.7	2.0	5	80
<i>CCA1-ox</i>	26.4	8.5	5	45	26	2.2	5	60	29.2	0.4	5	60
WS	23.4	0.4	8	75	23.5	0.2	8	86	23.5	0.3	8	86
8-3	28.7	5.8	5	40	26.0	1.6	5	40	-	-	5	20
17.7	-	-	2	0	-	-	2	0	-	-	2	50

%Rh—percentage of rhythmic plants. “-” no rhythm could be assigned during data analysis. Col indicates Col-0 and WS indicates Ws-2 ecotypes.

doi:10.1371/journal.pone.0127449.t002

(Table 2). Of seven *HvCCA1-ox* plants, only one plant had detectable rhythms of both CO<sub>2</sub> assimilation (34.5 h, 0.4 RAE) and transpiration (24.4 h, 0.3 RAE). One other plant had rhythmic CO<sub>2</sub> assimilation in LL (22.9 h, 0.5 RAE). In contrast, leaf gas exchange of the Ws-2 wildtype background line had robust circadian rhythms. Of eight Ws-2 plants, five had circadian rhythms of CO<sub>2</sub> assimilation (23.7 ± 0.3 h, 0.3 ± 0.1 RAE) and four Ws-2 plants, had detectable circadian rhythms of transpiration (23.2 ± 0.3 h, 0.3 ± 0.1 RAE).

Since analysis of circadian rhythms of delayed fluorescence, leaf movement, stomatal movements and photosynthesis all indicated that heterologous overexpression of barley *HvCCA1* made Arabidopsis plants circadian arrhythmic, we next investigated the effects of *HvCCA1-ox* on the expression of the circadian oscillator genes *AtCCA1*, *AtLHY* and *AtTOC1*. Over-expression of *HvCCA1* or *AtCCA1* resulted in arrhythmic or reduced amplitude expression of the endogenous *AtCCA1*, *AtTOC1* and *AtLHY* in LL (Fig 2).

### Overexpression of *HvCCA1* causes morphological and developmental defects in Arabidopsis


Rhythmic regulation by *AtCCA1* has profound effects on Arabidopsis development, such as the elongation of the hypocotyl, and the transition from vegetative to reproductive growth. We next investigated whether *HvCCA1* regulates development similar to *AtCCA1*. Over expression of both *AtCCA1* and *HvCCA1* resulted in longer hypocotyls, with those of *HvCCA1-ox* lines being significantly longer (3.5 to 5 mm) than the Col-0 and Ws-2 controls (1.5 mm and 2 mm; p<0.001; Fig 3A).

When grown under 16 h light 8 h dark, *AtCCA1-ox* delayed bolting and increased the number of leaves at bolt and this was phenocopied by over expression of *HvCCA1*. *HvCCA1-ox* lines flowered at least 20 days later than the wild type Ws-2 (p<0.001; Fig 3B) consistent with the *CCA1-ox-038* line, which bolted approximately 30 days later than the control Col-0. The number of rosette leaves at bolting was also significantly higher in *HvCCA1-ox* plants compared to Ws-2 (p<0.001; Fig 3B).

### Analysis of the function of *HvPpd-H1* in barley

We next investigated whether *HvPpd-H1* is similar in function to the Arabidopsis morning loop gene *AtPRR7*, with which *HvPpd-H1* shares most similarity. We analysed the function of the photoperiod responsive *HvPpd-H1* and photoperiod unresponsive *Hvppd-H1* variants to determine whether this locus can function in circadian regulation and whether natural mutations that alter photoperiodic sensitivity can affect circadian function. Circadian rhythms of stomatal conductance and CO<sub>2</sub> assimilation were measured in *Ppd-H1* and *ppd-H1* *H. vulgare* plants to determine if the natural mutant form *ppd-H1* had a consequence for circadian






**Fig 2. Overexpression of *HvCCA1* or *AtCCA1* (*AtCCA1-ox 038*) in *Arabidopsis* abolishes rhythms of transcript abundance of (a) *AtCCA1*, (b) *AtLHY* and (c) *AtTOC1*.** *Ws-2* (white diamonds), *HvCCA1-ox* (line 8–3 (filled circles), *Col-0* (black squares) and *AtCCA1-ox 038* (white triangle). Transcript abundance was normalised to *UBIQUITIN10*.



doi:10.1371/journal.pone.0127449.g002

rhythms in barley. No significant difference between *Ppd-H1* and *ppd-H1* in the period of circadian rhythms of CO<sub>2</sub> assimilation and stomatal conductance was detected (Fig 4). Measurement of gas exchange in barley is a low throughput assay in which only one individual at a time can be recorded. We obtained 15 measurements in white light representing >75 days of



**Fig 3. Over expression of *HvCCA1* in *Arabidopsis* causes elongated hypocotyls and delayed flowering.** (a) Hypocotyl length was measured in WS and Col-0 background (white bars) or *AtCCA1-ox* (*AtCCA1-ox 038*) or *HvCCA1-ox* (indicated by the independent transgenic line numbers). Stars indicate significant difference in the hypocotyl experiments. (b) Bolting time in days (black bars) and numbers of leaves at bolting (white) in *Ws-2* (WS) and *Col-0* background or *AtCCA1-ox* (*CCA1-ox*) or *HvCCA1-ox* (indicated by the independent transgenic line numbers) when grown in 16h light, 8 h dark.

doi:10.1371/journal.pone.0127449.g003



**Fig 4. Estimates of period of circadian rhythms of (a) photosynthesis and (b) stomatal conductance in *HvPpd-H1* (filled circles) and *Hvppd-H1* (open circles) in constant white light of 100  $\mu\text{mol m}^{-2} \text{s}^{-1}$  (black outlined symbols) or red light (red symbols). Each data point is derived from one individual seedling. Triangles represent the mean values.**

doi:10.1371/journal.pone.0127449.g004

continuous measurement. To ensure that the assay, despite its low-throughput nature had sufficient sensitivity to report changes in circadian period, we also investigated the effects of red light on circadian period. As expected the circadian period in equal intensity illumination was shorter under red than white light and this was most clear when assaying rhythms of stomatal conductance (Fig 4B). This demonstrated that whilst measurements of circadian rhythms of assimilation were not robust, the assay of stomatal conductance had sufficient sensitivity to detect period alteration when present (Fig 4A and 4B). The effect of both *Ppd-H1* and *ppd-H1* and also was monitored in red light, because it has previously been demonstrated that in Arabidopsis circadian period phenotypes in *prrr7-11* were more pronounced in red, rather than white light [26]. However, even in red light there was no significant difference between the period of the rhythms in *Ppd-H1* and *ppd-H1* (Fig 4).

### *HvPpd-H1* complements *AtPRR7-11*

The lack of a detectable effect of *Hvppd-H1* on circadian rhythms in barley might suggest that the mutation which regulates photoperiodic sensitivity does not affect the function of the circadian clock. Additionally, this finding raises the possibility that *HvPpd-H1* is not a circadian clock gene in barley, despite the sequence similarity to *AtPRR7*. We proceeded to use heterologous expression of barley genes in Arabidopsis to test whether *HvPpd-H1* can function as a circadian clock gene. Arabidopsis Col-0 and the *Atprrr7-11* loss of function mutant were transformed with *HvPpd-H1* and *Hvppd-H1* under the control of the *AtPRR7* promoter resulting in elevated expression of the transgenes (S2 Fig). We reasoned that complementation of the long circadian period phenotype of *prrr7-11* by heterologous expression of *HvPpd-H1* would provide evidence for *HvPpd-H1* acting in the circadian clock of barley.

The mean period of leaf movement rhythms of *prrr7-11* plants was significantly longer than that of the wildtype (*prrr7-11* 26.6  $\pm$  0.3 h compared to Col-0 24.9  $\pm$  0.2 h  $P = 0.0046$ ). The *prrr7-11* mutation was complemented by one of two lines of *pAtPRR7::Ppd-H1* and both lines of *pAtPRR7::ppd-H1*, with the complemented lines not being significantly different from wild type, except in the case of line *prrr7-11-pAtPRR7:Ppd-H1-c* (Table 3). Since three of the four lines complemented with natural variants of *HvPpd-H1* were restored to wild type circadian period, we conclude that *HvPpd-H1* in barley is likely to have a role orthologous to *AtPRR7* in

**Table 3. Summary of circadian period estimates for leaf movement in Col-0, *prrr7-11* and *prrr7-11* transformed with either *pPRR7::Ppd-H1* or *pPRR7::ppd-H1*.**

Line	Period (h)	SEM	RAE	n	%Rh
Col-0	24.9	0.2	0.2	32	75
<i>prrr7-11</i> *	26.6	0.3	0.2	29	59
<i>prrr7-11 pAtPRR7:Ppd-H1-c</i>	25.8	0.3	0.2	35	69
<i>prrr7-11 pAtPRR7:Ppd-H1-d*</i>	25.0	0.3	0.2	25	64
<i>prrr7-11 pAtPRR7:ppd-H1-e*</i>	25.3	0.3	0.2	21	86
<i>prrr7-11 pAtPRR7:ppd-H1-f*</i>	25.6	0.3	0.2	29	59

\* indicates significant difference at 5% level compared to background.

The background for *prrr7-11* is Col-0. The background for the complemented lines is *prrr7-11*.

SEM = standard error of the mean. %Rh = Percentage of rhythmic seedlings.

doi:10.1371/journal.pone.0127449.t003

Arabidopsis and further supports the finding of the gas exchange analyses that *Hvppd-H1* mutation is without effect on circadian rhythms.

## Discussion

We provide experimental evidence that *HvCCA1* and *HvPpd-H1* have functionality similar to *AtCCA1* and *AtPRR7* in Arabidopsis. Heterologous expression in Arabidopsis of *HvCCA1-ox* results in circadian arrhythmia and morphological changes consistent with *HvCCA1* being functionally equivalent to *AtCCA1*. Our data show that constitutive expression of *HvCCA1* in Arabidopsis causes arrhythmia in leaf movement, delayed fluorescence and several circadian clock genes. It also causes significant hypocotyl elongation and late flowering in *HvCCA1* transformants. These results are consistent with *AtCCA1-ox* Arabidopsis data and provide a strong support that *CCA1* and its function are highly conserved throughout the plant kingdom. Expression of either *HvPpd-H1* or *Hvppd-H1* under control of the endogenous *AtPRR7* promoter partially restores the wildtype period of circadian leaf movement rhythms in an *AtPRR7* tDNA insertion mutant (*prrr7-11*) suggesting that the function of *Hv-Ppd1* is as a circadian clock gene. A similar experiment in rice provided evidence that the *OsPRR37* and *AtPRR7* genes are partially interchangeable [27]. Together these findings show that the clock-associated functions of *CCA1* and *PRR7* are conserved between Arabidopsis, rice and barley.

Since both the photoperiodic-sensitive and -insensitive alleles of *Hv-Ppd1* were able to complement the *prrr7-11* phenotype, this suggested that the causative SNP affects photoperiodism independent of circadian rhythms. To test this directly, circadian rhythms of gas exchange from a Triumph (*ppd-H1*) *H. vulgare* line were compared to those from the same line into which the Igrri *Ppd-H1* allele had been introgressed and it was found that the circadian period was not significantly altered between barley plants carrying the *Hv-Ppd1* and *Hv-ppd1* alleles. Whilst mutation in *HvPpd-H1* did not produce detectable effects on circadian function either in barley or through complementation tests in Arabidopsis *Atprrr-11*, the *Hvppd-H1* allele does abolish circadian rhythms of photoperiodic-response and vernalisation-response genes *HvCO1*, *HvCO2* and *Vrn-H1* [12]. These findings coupled with our demonstration that both *HvPpd1-H1* and *Hvppd-H1* can complement *AtPrr7-11* suggest that *HvPpd-H1* has at least two regulatory roles, one required for proper functioning of the circadian system, and one required for generation of rhythmic outputs, and it is the latter that is compromised in *Hvppd-H1*.

The analysis of circadian rhythms in barley and the other grain crops is of critical importance, since domestication has required breeders and early farmers to select for varieties with altered photoperiodic sensitivity and adaptations for cold tolerance. These are both traits that are regulated by the circadian clock [1, 28]. We have described the use of gas exchange and heterologous expression in Arabidopsis circadian mutants as new tools for the dissection of the barley circadian signalling network and determined that *HvPpd-H1* is functionally orthologous to *AtPRR7*.

## Supporting Information

**S1 Fig. Transcript abundance of *AtCCA1* (black) and 5'UTR *AtCCA1* (endogenous) in *AtCCA1-ox* (white) and *HvCCA1* in *HvCCA1-ox* plants (grey).** Seedlings were entrained for one week in 12 light/12hr dark cycles at 22°C then transferred to constant light. Samples were taken 3 h after dawn. Lines used were Ws-2, Col-0, *AtCCA1-ox* 038 and *HvCCA1-ox* (8–3 and 18–1). Transcript abundance was normalised to Ubiquitin10. (PNG)

**S2 Fig. Transcript levels of *HvPpd-H1* and *Hvppd-H1* relative to *PP2a* expression in *prp7-11* Arabidopsis seedlings transformed with *HvPpd-H1* and *Hvppd-H1* under the control of the *AtPRR7* promoter (mean  $\pm$  s.d.;  $n = 3$ ).**

(PNG)

**S1 Table. Sequences of the primers used in the course of this study.** Forward (F) and reverse orientation (R) primers.

(PDF)

## Acknowledgments

ZR is grateful to the National Institute of Agricultural Botany for the award of a Scholarship. We acknowledge funding from a Marie Curie Early Stage Training project MEST-CT-2005-020526 for JK and the BBSRC-DTP for funding SC. AARW and MCM are grateful to the BBSRC for the award of BBSRC Grant BB/M006212/1, which supported aspects of the study.

## Author Contributions

Conceived and designed the experiments: JK ZR SC AG AH AARW. Performed the experiments: JK ZR SC MCM. Analyzed the data: JK ZR SC AH AARW. Contributed reagents/materials/analysis tools: DAL. Wrote the paper: JK ZR SC AG AH AARW.

## References

1. Harmer SL (2009) The circadian system in higher plants. *Ann. Rev. Plant Biol.* 60: 357–377.
2. Dodd AN, Salathia N, Hall A, Kévei E, Tóth R, Nagy F, et al. (2005) Plant circadian clocks improve growth, competitive advantage and survival. *Science* 309: 630–633. PMID: [16040710](#)
3. Green RM, Tingay S, Wang ZY, Tobin EM (2002) Circadian rhythms confer a higher level of fitness to Arabidopsis plants. *Plant Phys.* 129: 576–584.
4. Johnson CH, Kyriacou CP (2007) Clock evolution and adaptation: Whence and whither? *Ann. Plant Rev.* 21: 237–260.
5. Locke JC, Kozma-Bognár L, Gould PD, Fehér B, Kévei E, Nagy F, et al. (2006) Experimental validation of a predicted feedback loop in the multi-oscillator clock of Arabidopsis thaliana. *Mol. Sys Biol.*, 2(1).
6. Gendron JM, Pruneda-Paz JL, Doherty CJ, Gross AM, Kang SE, Kay SA (2012) Arabidopsis circadian clock protein, TOC1, is a DNA-binding transcription factor. *Proceedings of the National Academy of Sciences*, 109, 3167–3172.
7. Huang W, Perez-Garcia P, Pokhilko A, Millar AJ, Antoshechkin I, Riechmann JL, et al. (2012) Mapping the core of the Arabidopsis circadian clock defines the network structure of the oscillator. *Science*, 336: 75–79. doi: [10.1126/science.1219075](#) PMID: [22403178](#)
8. Herrero E, Kolmos E, Bujdosó N, Yuan Y, Wang M, Berns M, et al. (2012) EARLY FLOWERING4 recruitment of EARLY FLOWERING3 in the nucleus sustains the Arabidopsis circadian clock. *Plant Cell*, 24: 428–443. doi: [10.1105/tpc.111.093807](#) PMID: [22327739](#)
9. Serikawa M, Miwa K, Kondo T, Oyama T (2008) Functional conservation of clock-related genes in flowering plants: Overexpression and RNA interference analyses of the circadian rhythm in the monocotyledon *Lemna gibba*. *Plant Phys.* 146: 1952–1963.
10. Matsushika A, Makino S, Kojima M, Mizuno T (2000) Circadian waves of expression of the PRR1/TOC1 family of Pseudo-response regulators of *A.thaliana*: insight into the plant circadian clock. *Plant Cell Physiol.* 41: 1002–1012. PMID: [11100772](#)
11. Ogiso E, Takahashi Y, Sasaki T, Yano M, Izawa T (2010) The role of casein kinase II in flowering time regulation has diversified during evolution. *Plant Phys.* 152: 808–820.
12. Campoli C, Shtaya M, Davis SJ, von Korff M (2012) Expression conservation within the circadian clock of a monocot: natural variation at barley Ppd-H1 affects circadian expression of flowering time genes, but not clock orthologs. *BMC Plant Biol.* 12: 97. doi: [10.1186/1471-2229-12-97](#) PMID: [22720803](#)
13. Campoli C, Pankin A, Drosse B, Casao CM, Davis SJ, Korff M (2013) HvLUX1 is a candidate gene underlying the early maturity 10 locus in barley: phylogeny, diversity, and interactions with the circadian



- clock and photoperiodic pathways. *New Phytologist* 199: 1045–1059. doi: [10.1111/nph.12346](https://doi.org/10.1111/nph.12346) PMID: [23731278](https://pubmed.ncbi.nlm.nih.gov/23731278/)
14. Faure S, Turner AS, Gruszka D, Christodoulou V, Davis SJ, Korff M, et al. (2012) Mutation at the circadian clock gene EARLY MATURITY 8 adapts domesticated barley (*Hordeum vulgare*) to short growing seasons. *Proc. Natl. Acad. Sci. (USA)* 109: 8328–8333. doi: [10.1073/pnas.1120496109](https://doi.org/10.1073/pnas.1120496109) PMID: [22566625](https://pubmed.ncbi.nlm.nih.gov/22566625/)
  15. Turner A, Beales J, Faure S, Dunford RP, Laurie DA (2005) The Pseudo-Response Regulator Ppd-H1 Provides Adaptation to Photoperiod in Barley. *Science* 310: 1031–1034. PMID: [16284181](https://pubmed.ncbi.nlm.nih.gov/16284181/)
  16. Curtis MD, Grossniklaus U (2003) A gateway cloning vector set for high-throughput functional analysis of genes *in planta*. *Plant Physiology* 133: 462–469. PMID: [14555774](https://pubmed.ncbi.nlm.nih.gov/14555774/)
  17. Clough SJ, Bent AF (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J* 16: 735–743. PMID: [10069079](https://pubmed.ncbi.nlm.nih.gov/10069079/)
  18. Schultz TF, Kiyosue T, Yanovsky M, Wada M, Kay SA (2001) A Role for LKP2 in the Circadian Clock of *Arabidopsis*. *Plant Cell* 13: 2659–2670. PMID: [11752379](https://pubmed.ncbi.nlm.nih.gov/11752379/)
  19. LeClerc S, Bartel B (2001) A library of *Arabidopsis* 35S-cDNA lines for identifying novel mutants. *Plant Mol. Biol.* 46: 695–703. PMID: [11575724](https://pubmed.ncbi.nlm.nih.gov/11575724/)
  20. Karimi M, Inze D, Depicker A (2002) GATEWAY vectors for *Agrobacterium* mediated plant transformation. *Trends Plant Sci.* 7: 193–195. PMID: [11992820](https://pubmed.ncbi.nlm.nih.gov/11992820/)
  21. Ramakers C, Ruijter JM, Lekanne Deperz RH, Moorman AFM (2003) Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neuroscience Letters* 339: 62–66. PMID: [12618301](https://pubmed.ncbi.nlm.nih.gov/12618301/)
  22. Gould PD, Diaz P, Hogben C, Kusakina J, Salem R, Hartwell J, et al. (2009) Delayed fluorescence as a universal tool for the measurement of circadian rhythms in higher plants. *Plant J.* 58: 893–901. doi: [10.1111/j.1365-313X.2009.03819.x](https://doi.org/10.1111/j.1365-313X.2009.03819.x) PMID: [19638147](https://pubmed.ncbi.nlm.nih.gov/19638147/)
  23. Plautz JD, Straume M, Stanewsky R, Jamison CF, Brandes C, Dowse HB, et al. (1997) Quantitative analysis of *Drosophila* period gene transcription in living animals. *J. Biol. Rhyth.* 12: 204–217. PMID: [9181432](https://pubmed.ncbi.nlm.nih.gov/9181432/)
  24. Dodd AN, Parkinson K, Webb AAR (2004) Independent circadian regulation of assimilation and stomatal conductance in the *ztl-1* mutant of *Arabidopsis*. *New Phyt.* 162: 63–70.
  25. Wang ZY, Tobin EM (1998) Constitutive expression of the *CIRCADIAN CLOCK ASSOCIATED 1* (*CCA1*) gene disrupts circadian rhythms and suppresses its own expression. *Cell* 93: 1207–1217. PMID: [9657153](https://pubmed.ncbi.nlm.nih.gov/9657153/)
  26. Farré EM, Harmer SL, Harmon FG, Yanovsky MJ, Kay SA (2005) Overlapping and distinct roles of PRR7 and PRR9 in the *Arabidopsis* circadian clock. *Curr. Biol.* 15: 47–54. PMID: [15649364](https://pubmed.ncbi.nlm.nih.gov/15649364/)
  27. Murakami M, Tago Y, Yamashino T, Mizuno T (2007) Characterization of the rice circadian clock-associated pseudo-response regulators in *Arabidopsis thaliana*. *Bioscience, Biotechnology, and Biochemistry.* 71: 1107–1110. PMID: [17420570](https://pubmed.ncbi.nlm.nih.gov/17420570/)
  28. Dodd AN, Kyed Jakobsen M, Baker AJ, Telzerow A, Hou S-W, Laplaze L, et al. (2006) Time of day modulation of Ca<sup>2+</sup> signals in *Arabidopsis*. *Plant J.* 48: 962–973. PMID: [17227550](https://pubmed.ncbi.nlm.nih.gov/17227550/)