White Rose Research Online

eprints@whiterose.ac.uk
https://eprints.whiterose.ac.uk
Universities of Leeds, Sheffield and York

Deposited via The University of Leeds.

White Rose Research Online URL for this paper:
https://eprints.whiterose.ac.uk/id/eprint/89204/

Version: Published Version

Article:
Walker, HM, Stone, D, Ingham, T et al. (2015) Night-time measurements of HOx during the RONOCO project and analysis of the sources of HO2. Atmospheric Chemistry and Physics, 15 (14). 8179 - 8200. ISSN: 1680-7316

https://doi.org/10.5194/acp-15-8179-2015

Reuse
Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown
If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

UNIVERSITY OF LEEDS
University of Sheffield
UNIVERSITY of York

Atmos. Chem. Phys., 15, 8179–8200, 2015
www.atmos-chem-phys.net/15/8179/2015/
doi:10.5194/acp-15-8179-2015
© Author(s) 2015. CC Attribution 3.0 License.

Atmospheric Chemistry and Physics
Open Access

Night-time measurements of HO_x during the RONOCO project and analysis of the sources of HO_2

H. M. Walker[1], D. Stone[1], T. Ingham[1,2], S. Vaughan[1], M. Cain[3,4], R. L. Jones[3], O. J. Kennedy[3], M. McLeod[3], B. Ouyang[3], J. Pyle[3,4], S. Bauguitte[5,6], B. Bandy[7,8], G. Forster[7,8], M. J. Evans[9,10], J. F. Hamilton[9], J. R. Hopkins[9,10], J. D. Lee[9,10], A. C. Lewis[9,10], R. T. Lidster[9], S. Punjabi[9,10], W. T. Morgan[11,12], and D. E. Heard[1,2]

[1]School of Chemistry, University of Leeds, Leeds, UK
[2]National Centre for Atmospheric Science, University of Leeds, Leeds, UK
[3]Department of Chemistry, University of Cambridge, Cambridge, UK
[4]National Centre for Atmospheric Science, University of Cambridge, Cambridge, UK
[5]Facility for Airborne Atmospheric Measurements (FAAM), Cranfield University, Cranfield, UK
[6]National Centre for Atmospheric Science, FAAM, Cranfield University, Cranfield, UK
[7]Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East Anglia, Norwich, UK
[8]National Centre for Atmospheric Science, University of East Anglia, Norwich, UK
[9]Wolfson Atmospheric Chemistry Laboratories, Department of Chemistry, University of York, York, UK
[10]National Centre for Atmospheric Science, University of York, York, UK
[11]School of Earth, Atmospheric and Environmental Sciences, University of Manchester, Manchester, UK
[12]National Centre for Atmospheric Science, University of Manchester, Manchester, UK

Correspondence to: D. E. Heard (d.e.heard@leeds.ac.uk)

Received: 6 November 2014 – Published in Atmos. Chem. Phys. Discuss.: 30 January 2015
Revised: 12 May 2015 – Accepted: 24 May 2015 – Published: 23 July 2015

Abstract. Measurements of the radical species OH and HO_2 were made using the fluorescence assay by gas expansion (FAGE) technique during a series of night-time and daytime flights over the UK in summer 2010 and winter 2011. OH was not detected above the instrument's 1σ limit of detection during any of the night-time flights or during the winter daytime flights, placing upper limits on [OH] of 1.8×10^6 molecule cm^{-3} and 6.4×10^5 molecule cm^{-3} for the summer and winter flights, respectively. HO_2 reached a maximum concentration of 3.2×10^8 molecule cm^{-3} (13.6 pptv) during a night-time flight on 20 July 2010, when the highest concentrations of NO_3 and O_3 were also recorded. An analysis of the rates of reaction of OH, O_3, and the NO_3 radical with measured alkenes indicates that the summer night-time troposphere can be as important for the processing of volatile organic compounds (VOCs) as the winter daytime troposphere. An analysis of the instantaneous rate of production of HO_2 from the reactions of O_3 and NO_3 with alkenes has shown that, on average, reactions of NO_3 dominated the night-time production of HO_2 during summer and reactions of O_3 dominated the night-time HO_2 production during winter.

1 Introduction

Trace gases emitted into the atmosphere, including pollutants and greenhouse gases, are removed primarily by oxidation. The hydroxyl radical, OH, is the most important oxidising species in the daytime troposphere, reacting with numerous species, including volatile organic compounds (VOCs), CO, SO_2, and long-lived anthropogenic halogenated compounds. During the day, primary production of OH (i.e. initialisation of the radical chain) occurs predominantly via photolysis of ozone at $\lambda \leq 340$ nm, followed by the reaction of the resulting electronically excited oxygen atom, $O(^1D)$, with water vapour. The OH-initiated oxidation of VOCs leads to the production of the hydroperoxy radical, HO_2, and together the

two radicals form the HO_x family. A key reaction in the conversion of OH to HO_2 is the reaction with CO:

$$OH + CO \rightarrow H + CO_2 \quad (R1)$$

$$H + O_2 + M \rightarrow HO_2 + M. \quad (R2)$$

The reaction of OH with VOCs results in the production of organic peroxy radicals, RO_2:

$$OH + RH \rightarrow H_2O + R \quad (R3)$$

$$R + O_2 + M \rightarrow RO_2 + M \quad (R4)$$

Reactions of HO_2 and RO_2 with NO propagate the HO_x radical chain, regenerating OH:

$$RO_2 + NO \rightarrow RO + NO_2 \quad (R5)$$

$$RO + O_2 \rightarrow R'O + HO_2 \quad (R6)$$

$$HO_2 + NO \rightarrow OH + NO_2 \quad (R7)$$

The production of OH through the photolysis of ozone (and other species at longer wavelengths) is limited to daylight hours, and the oxidation of trace gases at night proceeds through alternative mechanisms. Two mechanisms are known to initiate HO_x radical chemistry and oxidation chemistry at night: ozonolysis of alkenes and reactions of the nitrate radical, NO_3, with alkenes.

Reactions of ozone with alkenes occur via the addition of ozone to the double bond to form a five-membered ring called a primary ozonide. The primary ozonide decomposes to form one of two possible pairs of products, each pair consisting of a carbonyl compound and a vibrationally and rotationally excited carbonyl oxide termed a Criegee intermediate (CI). The simplest gas-phase CI, CH_2OO, and the alkyl-substituted CH_3CHOO have been observed directly by photoionisation mass spectrometry (Taatjes et al., 2008, 2012, 2013; Beames et al., 2012, 2013; Welz et al., 2012; Stone et al., 2014a), by infrared absorption spectroscopy (Su et al., 2013), and by microwave spectroscopy (Nakajima and Endo, 2013, 2014). Excited CIs may be stabilised by collision with surrounding molecules (Donahue et al., 2011; Drozd and Donahue, 2011) or may undergo isomerisation or decomposition to yield products including OH, H, and subsequently HO_2 (Paulson and Orlando, 1996; Kroll et al., 2001a, b, 2002; Johnson and Marston, 2008). Stabilised CIs (SCIs) are known to react with a variety of compounds, including H_2O, NO_2, SO_2, and a variety of organic compounds (e.g. Mauldin III et al., 2012; Taatjes et al., 2012, 2013, 2014; Ouyang et al., 2013; Stone et al., 2014a). There is experimental evidence for the formation of OH from the thermal decomposition of SCIs, on a much longer timescale than the decomposition or isomerisation of excited CIs (Kroll et al., 2001a, b). The OH produced through these ozonolysis mechanisms will proceed to oxidise other VOC species. Criegee intermediates formed in the ozonolysis of alkenes are known to be an important source of HO_x during the day and at night (Paulson and Orlando, 1996; Donahue et al., 1998; Kanaya et al., 1999; Salisbury et al., 2001; Geyer et al., 2003; Ren et al., 2003a, 2006; Heard et al., 2004; Harrison et al., 2006; Sommariva et al., 2007). The gas-phase ozonolysis of unsaturated VOCs, and in particular the role and subsequent chemistry of the Criegee intermediate, have been reviewed in detail by Johnson and Marston (2008), Donahue et al. (2011), Vereecken and Francisco (2012), and Taatjes et al. (2014).

Another key night-time oxidant, NO_3, is formed primarily by the reaction of NO_2 with ozone. NO_3 reacts with a range of species in the troposphere, and its reaction with alkenes is known to be an important night-time oxidation mechanism (Salisbury et al., 2001; Geyer et al., 2003; Sommariva et al., 2007; Emmerson and Carslaw, 2009; Brown et al., 2011). The reaction between NO_3 and an alkene proceeds primarily via addition to a double bond to form a nitrooxyalkyl radical, R–ONO_2. At atmospheric pressure, the main fate of the nitrooxyalkyl radical is reaction with O_2 (Berndt and Böge, 1994) to produce a nitrooxyalkyl peroxy radical, O_2–R–ONO_2. The nitrooxyalkyl peroxy radical can react with NO_2, HO_2, RO_2, NO, and NO_3, of which the latter two reactions lead to the formation of the nitrooxyalkoxy radical, O–R–ONO_2. The nitrooxyalkoxy radical can undergo isomerisation, decomposition, or reaction with O_2. Reaction with O_2, analogous to the reaction of organic alkoxy radicals, yields HO_2:

$$O–R–ONO_2 + O_2 \rightarrow O–R'–ONO_2 + HO_2. \quad (R8)$$

Thus, the night-time oxidation of hydrocarbons by NO_3 leads to the production of HO_2. The reaction of HO_2 with NO (Reaction R7), O_3, and NO_3 can generate OH:

$$HO_2 + O_3 \rightarrow OH + 2O_2 \quad (R9)$$

$$HO_2 + NO_3 \rightarrow OH + NO_2 + O_2. \quad (R10)$$

Atkinson and Arey (2003) published a detailed review of the tropospheric degradation of VOCs, including reaction with O_3 and NO_3. A comprehensive review of night-time radical chemistry is given by Brown and Stutz (2012).

The oxidising capacity of the nocturnal troposphere is thought to be controlled by the reactions described above, with a limited role for OH and HO_2 due to the absence of their photolytic sources. The oxidation of VOCs at night can have significant effects on daytime air quality and tropospheric ozone production (Brown et al., 2004, 2006, 2011; Wong and Stutz, 2010). Several field measurement campaigns have involved night-time measurements of OH, HO_2, RO_2, and NO_3 (see Table 1) and have highlighted the importance of the vertical profile of night-time radical concentrations and chemistry (Geyer and Stutz, 2004a, b; Stutz et al., 2004; Volkamer et al., 2010), but prior to the current work, there had been no aircraft-based studies of night-time chemistry involving measurements of both NO_3 and HO_2 to enable the vertical profiling of the lower atmosphere and a full evaluation of the nocturnal radical budget. Table 1 gives details of some previous measurements and modelling of night-time

Table 1. Examples of modelling studies and observations of HO_x radicals and VOC oxidation at night. PERCA: peroxy radical chemical amplification; LIF: laser-induced fluorescence; DOAS: differential optical absorption spectroscopy; MCM: Master Chemical Mechanism; MIESR: matrix isolation electron spin resonance; RACM: Regional Atmospheric Chemistry Mechanism; CRDS: cavity ring-down spectroscopy; CIMS: chemical ionisation mass spectrometry; GC: gas chromatography; PTRMS: proton transfer reaction mass spectrometry; FTIR: Fourier transform infrared spectroscopy; DUALER: DUAl channel peroxy radical chemical amplifier; OA-CRD: off-axis cavity ring-down spectroscopy; CRM-PTR-MS: comparative reactivity method proton transfer mass spectrometry.

Location, campaign, date	Methods	Results	Reference
Mace Head, Ireland, Eastern Atlantic Spring Experiment (EASE97), 1997	Measurements: [HO_2+RO_2] measured by PERCA; HO_x measured by LIF; NO_3 measured by DOAS. Modelling: campaign-tailored box model constrained to measurements, based on MCM.	Two nights of HO_x measurements: HO_2 = 1–2 and 0.5–0.7 pptv; OH not detected above limit of detection ($\sim 2.5 \times 10^5$ cm^{-3}). NO_3 dominated radical production in westerly (clean) air masses; O_3 dominated in NE, SE, and SW air masses and dominated radical production overall during the campaign.	Salisbury et al. (2001); Creasey et al. (2002)
Pabstthum, Germany, Berlin Ozone Experiment (BERLIOZ), 1998	Measurements: HO_x measured by LIF; NO_3 measured by DOAS and MIESR. Modelling: zero-dimensional model using lumped VOC reactivity, constrained to measured species.	Night-time OH = 1.85×10^5 cm^{-3}, compared to modelled value of 4.1×10^5 cm^{-3}. Night-time HO_2 = 3×10^7 cm^{-3}, model results in agreement. NO_3 chemistry responsible for 53 % of HO_2 and 36 % of OH during the night. O_3+ alkene responsible for 47 % of HO_2 and 64 % of OH during the night.	Geyer et al. (2003); Holland et al. (2003)
Birmingham, Pollution of the Urban Midlands Atmosphere (PUMA), 1999 and 2000	Measurements: HO_x measured by LIF. Modelling: photochemical box model constrained to measurements, based on MCM.	Daytime OH initiation dominated by O_3 + alkenes, HONO photolysis, and $O(^1D) + H_2O$ during summer. O_3 + alkenes dominated in winter. O_3 + alkenes main radical source at night.	Emmerson et al. (2005); Harrison et al. (2006)
New York, $PM_{2.5}$ Technology Assessment and Characteristics Study-New York (PMTACS-NY), 2001	Measurements: HO_x measured by LIF.	Night-time OH $\sim 7 \times 10^5$ cm^{-3} and night-time HO_2 $\sim 8 \times 10^6$ cm^{-3}. Increase in HO_x after midnight attributed to increase in O_3 due to transport. O_3 + alkenes main source of night-time HO_x.	Ren et al. (2003a, b)
Mace Head, North Atlantic Marine Boundary Layer Experiment (NAMBLEX), 2002	Measurements: HO_x measured by LIF; NO_3 measured by DOAS. Modelling: zero-dimensional box models constrained to measured species, based on MCM.	Night-time HO_2 = 2–3 $\times 10^7$ cm^{-3}; OH below detection limit (6×10^4 cm^{-3}). Model overestimated HO_2. On average, O_3+ alkene reactions contributed 59 % and NO_3+ alkene reactions contributed 41 % to RO_2 production at night, but NO_3 and RO_2 concentrations were always higher in semi-polluted air masses than in clean marine air masses and NO_3 reactions dominated in these conditions.	Fleming et al. (2006); Smith et al. (2006); Sommariva et al. (2007)
Writtle, London, Tropospheric ORganic CHemistry experiment (TORCH), 2003	Measurements: HO_x measured by LIF, RO_2 measured by PERCA, during a heatwave or pollution episode. Modelling: zero-dimensional box model constrained to measured species.	OH and HO_2 observed above the limit of detection on several nights. OH peaked at 8.5×10^5 cm^{-3}; HO_2 peaked at 1×10^8 cm^{-3}. Model overpredicted night-time OH and HO_2 on average by 24 % and 7 % and underpredicted [HO_2 + ΣRO_2] by 22 %.	Lee et al. (2006); Emmerson et al. (2007); Emmerson and Carslaw (2009)
Mexico City Metropolitan Area (MCMA 2003)	Measurements: HO_x measured by LIF, NO_3 measured by DOAS. Modelling: zero-dimensional model based on MCM v3.1, constrained to measured species.	Polluted city location characterised by high levels of NO, NO_2, and O_3. Maximum night-time OH $\sim 1 \times 10^6$ cm^{-3}; maximum night-time HO_2 $\sim$ 6 pptv. Night-time production of radicals dominated by O_3+ alkene reactions (76–92 %); NO_3+ alkene plays a minor role. Daytime radical production $\sim$ 25 times higher than night.	Shirley et al. (2006); Sheehy et al. (2010); Volkamer et al. (2010)
New York City, PMTACS-NY winter 2004	Measurements: HO_x measured by LIF. Modelling: zero-dimensional model based on RACM and constrained by measurements.	Mean maximum OH = 0.05 pptv; mean maximum HO_2 = 0.7 pptv. Model underprediction of HO_2 was pronounced when NO was high. O_3+ alkene reactions were dominant night-time source.	Ren et al. (2006)

Table 1. Continued.

Gulf of Maine, Northeast United States, New England Air Quality Study (NEAQS), 2004	Measurements: NO_3 and N_2O_5 measured by CRDS. Modelling: zero-dimensional model based on MCM v3.1, constrained to measured species. No measurements of OH, HO_2, or RO_2.	Ship-based measurements onboard *RV Ronald H. Brown* in the Gulf of Maine, influenced by unpolluted marine air masses and polluted air masses from the USA and Canada. Maximum modelled night-time $HO_2 = 7.0 \times 10^8$ cm^{-3}. Base model overestimated NO_3 and NO_2 observations by 30–50 %. In anthropogenic air masses, reaction with VOCs and RO_2 each accounted for 40 % of modelled NO_3 loss.	Sommariva et al. (2009)
Houston, Texas, Texas Air Quality Study (TexAQS), 2006	Measurements: NO_3 and N_2O_5 measured by CRDS; VOCs measured by CIMS, GC, and PTRMS. No direct measurements of OH, HO_2, or RO_2.	Loss rates and budgets of NO_3 and highly reactive VOCs calculated. NO_3 primarily lost through reaction with VOCs. VOC oxidation dominated by NO_3, which was 3–5 times more important than O_3.	Brown et al. (2011)
Pearl River Delta, China, Program of Regional Integrated Experiments of Pearl River Delta Region (PRIDE-PRD), 2006	Measurements: HO_x measured by LIF; OH reactivity measured by laser-flash photolysis and LIF; VOCs measured by FTIR and GC. Modelling: box model based on RACM and the Mainz Isoprene Mechanism and constrained by measurements.	Rural site 60 km downwind of large urban region (Guangzhou), with low local wind speeds favouring accumulation of air pollutants. Maximum night-time OH (hourly average) $= 5 \times 10^6$ cm^{-3}; maximum night-time HO_2 (hourly average) $= 1 \times 10^9$ cm^{-3}. Unknown recycling mechanism required for the model to reproduce measured night-time values. OH reactivity peaked at night. Missing night-time reactivity attributed to unmeasured secondary organic compounds.	Lou et al. (2010); Lu et al. (2012, 2013)
Beijing, Campaigns of Air Quality Research in Beijing and Surrounding Regions (CAREBEIJING2006), 2006	Measurements: HO_x measured by LIF; OH lifetime measured by laser-flash photolysis and LIF; VOCs measured by GC. Modelling: box model based on RACM and the Mainz Isoprene Mechanism and constrained by measurements.	Suburban rural site south of Beijing, under the influence of slowly moving, aged polluted air from the south. OH reactivity peaked at night. Model generally underestimated observed night-time OH concentrations.	Lu et al. (2013, 2014)
Cape Verde, Reactive Halogens in the Marine Boundary Layer (RHaMBLe), 2007	Measurements: HO_x measured by LIF. Modelling: box model based on MCM with added halogen chemistry scheme, constrained to measurements of long-lived species.	Clean tropical Atlantic measurement site with occasional continental influence. OH was not measured at night. HO_2 was detected on two nights, up to 2.5×10^7 cm^{-3}. Model underprediction of HO_2 was significantly reduced by constraining the model to 100 pptv of peroxy acetyl nitrate (PAN) at night.	Whalley et al. (2010)
Huelva, Spain, Diel Oxidant Mechanisms in relation to Nitrogen Oxides (DOMINO), 2008	Measurements: $[HO_2+RO_2]$ measured by DUALER; HO_x measured by LIF; NO_3 and N_2O_5 measured by OA-CRD; OH reactivity measured by CRM-PTR-MS. No measurements of anthropogenic VOCs.	Coastal forested site with strong urban-industrial and weak biogenic influences. Maxima in $[HO_2+RO_2]$ and $[HO_2]$ were observed around noon and midnight. Enhanced night-time $[HO_2+RO_2]$ (up to 80 pptv) was observed in air masses from the urban-industrial region. Maximum night-time $HO_2 = 8$ pptv. Measured NO_3 was generally below LOD; calculated NO_3 up to 20 pptv. Calculated production of RO_2 from NO_3 + alkenes accounts for 47–54 % of observed $[HO_2+RO_2]$. Ozonolysis of unmeasured alkenes could account for remaining $[HO_2+RO_2]$.	Andrés-Hernández et al. (2013)

HO_x concentrations in polluted or semi-polluted environments. Highlights from these studies are discussed here, with particular attention paid to those involving measurements of HO_x, NO_3, and O_3 and to those in which the contributions made by O_3 and NO_3 to night-time radical chemistry have been considered.

Geyer et al. (2003) report radical measurements and modelling from the 1998 Berlin Ozone Experiment (BERLIOZ). Measurements of NO_3, RO_2, HO_2, and OH were made by matrix isolation electron spin resonance (MIESR), chemical amplification (CA), and laser-induced fluorescence (LIF) spectroscopy at a site approximately 50 km from Berlin. HO_2 was detected at night, with concentrations frequently as high as 5×10^7 molecule cm^{-3} (approximately 2 pptv) and an average concentration of 1×10^8 molecule cm^{-3} over 1 h (02:00 to 03:00) of nocturnal measurements during an intensive period of the study (Holland et al., 2003). OH was usually below the limit of detection of the LIF instrument (3.5×10^5 molecule cm^{-3}). Modelling revealed that nitrate radical reactions with terpenes were responsible for producing 53 % of HO_2 and 36 % of OH radicals in the night, with ozonolysis accounting for the production of the remaining

47 % of HO_2 and 64 % of OH radicals. A positive linear correlation between RO_2 and NO_3 was observed and was reproduced by the model.

Reactions of O_3 with alkenes were found to be responsible for the majority of the formation of OH during the winter PUMA (Pollution of the Urban Midlands Atmosphere) campaign (a low-photolysis urban environment) (Heard et al., 2004; Emmerson et al., 2005; Harrison et al., 2006). Measurements of OH, HO_2, and RO_2 were unavailable at night, but model-predicted values of these radicals were used to calculate that 90 % of night-time initiation via HO_2 was from O_3 reactions. Without measurements of NO_3 during the campaign, there was no estimate of its contribution to radical initiation.

Modelling results from the MCMA-2003 (Mexico City) field campaign (Volkamer et al., 2010) indicate that night-time radical production at roof-top level (approximately 16 m above the ground) was dominated by ozonolysis of alkenes, and that reactions of NO_3 with alkenes played only a minor role. The measurement site was located in a polluted urban environment, with high levels of NO, NO_2, and O_3. NO_3 was observed at a maximum concentration of 50 pptv during the night at a mean height above the ground of 70 m. Roof-top level concentrations of NO_3 were estimated using a linear scaling factor, calculated from the observed O_3 vertical gradient, and were found to be, on average, 3 times lower than the concentrations measured at 70 m. This predicted vertical gradient accounts for the relative unimportance of NO_3 reactions in radical initiation at roof-top level. The propagation of RO_2 radicals to HO_2 and OH, by reaction with NO_3, was found to be negligible.

The 2006 Texas Air Quality Study (TexAQS) involved a series of night-time flights onboard the NOAA P-3 aircraft over Houston, Texas, and along the Gulf Coast (Brown et al., 2011). Loss rates and budgets of NO_3 and highly reactive VOCs were calculated, but there were no measurements of OH, HO_2, and RO_2 during the flights. Budgets for NO_3 show that it was lost primarily through reactions with unsaturated VOCs, but the contribution to NO_3 loss through reaction with peroxy radicals was uncertain because of the lack of direct measurements of RO_2 during the flights. NO_3 dominated VOC oxidation, being 3 to 5 times more important than O_3.

In summary, NO_3 and O_3 have both been found to dominate radical initiation in the night-time troposphere, and in some situations the two mechanisms were found to be equally important. The relative importance of O_3- and NO_3-initiated oxidation depends on the availability of NO_3, which is determined by the amount of NO_x present in the atmosphere and the ratio of NO to NO_2, and on the concentration and species distribution of VOCs (Bey et al., 2001; Geyer et al., 2003). A modelling study by Bey et al. (2001) suggests that nocturnal radical initiation is driven by alkene ozonolysis in urban environments or in environments with low NO_x concentrations, while both O_3 and NO_3 contribute to radical initiation in rural environments with moderate NO_x levels. It is expected that NO_3 dominates nocturnal radical initiation in air masses containing sufficient NO_2 and O_3 for NO_3 production while being deprived of NO (e.g. air masses downwind of urban areas). Geyer and Stutz (2004b) have found that the effects of suppressed mixing in the nocturnal boundary layer can also control whether NO_3 or O_3 dominates night-time radical chemistry.

In this paper we report airborne measurements of OH and HO_2 made during the RONOCO (ROle of Nighttime chemistry in controlling the Oxidising Capacity of the atmOsphere) and SeptEx (September Experiment) projects in 2010 and 2011. The rates of reaction between O_3, NO_3, and OH with the alkenes measured during the flights are investigated. The analysis of radical production from the night-time reactions of O_3 and NO_3 with alkenes is also given. Comparisons are made between the daytime and night-time chemistry studied and between the summer and winter measurement periods. Details and results of a box modelling study, and a comparison to the observations, are given by Stone et al. (2014b).

2 Details of the RONOCO and SeptEx fieldwork

RONOCO is a Natural Environment Research Council (NERC)-funded consortium project aimed at improving our understanding of the mechanisms and impact of nocturnal oxidation chemistry over the UK. The RONOCO fieldwork consisted of two measurement campaigns, in July 2010 and January 2011. Additional fieldwork, SeptEx, was conducted in September 2010. The RONOCO and SeptEx flights were conducted onboard the BAe-146 research aircraft operated by the Facility for Airborne Atmospheric Measurements (FAAM). Both field measurement campaigns were based at East Midlands Airport (52.8° N, 1.3° W) in the UK. During RONOCO the majority of the flying took place at night, with occasional flights beginning or ending in daylight hours to study chemical behaviour at dusk and dawn. Flights during SeptEx were mainly during the day, providing a useful comparison to the nocturnal chemistry.

Flights were conducted between altitudes of 50 and 6400 m, above the UK and the North Sea. Figure 1 shows the flight tracks during the summer, SeptEx, and winter measurements coloured by altitude. Measurements of OH and HO_2 were made using the University of Leeds aircraft-based fluorescence assay by gas expansion (FAGE) instrument. A suite of supporting measurements, including CO, O_3, NO, H_2O, VOCs, NO_3, and HCHO, were made during the flights and have been used in the current work. Table 2 summarises the techniques used to measure these species.

Air mass histories for each flight have been calculated using the UK Met Office Numerical Atmospheric-dispersion Modelling Environment (NAME). NAME is a three-dimensional Lagrangian particle dispersion model (Jones et al., 2007), which is run here using the UK Meteoro-



Figure 1. Flight tracks for **(a)** summer RONOCO, **(b)** SeptEx, and **(c)** winter RONOCO measurement campaigns, coloured by altitude.

logical Office's Unified Model meteorological fields. Model "particles", restricted to a 300 m deep layer from the surface, were released along the flight path and were tracked backwards through the modelled atmosphere. Model particle densities were integrated over 24 h periods, beginning at 24, 48, 72, and 96 h before each flight. The resulting "footprint" maps show the regions where the measured air has been in contact with the surface over the 4 days preceding a flight. An example is shown in Fig. 2, which shows model particle densities integrated over the 24 h period beginning 48 h prior to flight B535. The majority of the summer flights were characterised by air masses originating from the west and south-west of the UK, having Atlantic or continental European influences. The SeptEx flights were predominantly influenced by air masses from the north-east, east, and south-east of the UK, with northern European influences. The winter flights were mainly characterised by air masses arriving from the west of the UK, bringing Atlantic influences.



Figure 2. Footprint map for flight B535 on 17 July 2010, showing model particle densities ($g\,s\,m^{-3}$) in a 300 m deep layer from the surface, integrated over a 24 h period beginning 48 h prior to the flight.

Table 3 gives mean and maximum mixing ratios of CO, O_3, NO, and NO_2 measured during RONOCO and SeptEx. The mean mixing ratios of NO measured during the summer RONOCO flights are much lower than ground-based night-time measurements (e.g. 1.0 ppbv during the Tropospheric ORganic CHemistry experiment (TORCH) (Emmerson and Carslaw, 2009), 0–20 ppbv during the $PM_{2.5}$ Technology Assessment and Characteristics Study-New York (PMTACS-NY); Ren et al., 2006) but are comparable with previous airborne night-time measurements (e.g. < 30 pptv during the Texas Air Quality Study (TexAQS; Brown et al., 2011). Mean values of NO up to 14 pptv were reported by Salisbury et al. (2001) for semi-polluted air masses sampled at Mace Head. These comparisons indicate that the RONOCO and SeptEx flights enabled the sampling of air masses generally removed from the influence of NO in fresh surface emissions. Table 3 also highlights the unusual chemical conditions encountered during flight B537 on 20 July 2010, discussed further in Sect. 4.1. Night-time altitude profiles of NO_3, O_3, *trans*-2-butene, and propene (the latter two being illustrative of the alkenes measured) are given in Fig. 3.

3 Experimental

3.1 The Leeds FAGE aircraft instrument

The University of Leeds aircraft FAGE instrument has been described in detail by Commane et al. (2010). A brief description is given here. The instrument, which was designed specifically for use onboard the FAAM BAe-146 research aircraft (Floquet, 2006), is housed in two double-width 19 in. aircraft racks, with the inlet, detection cells, and pump set being separate from the two racks. Ambient air is sampled through a 0.7 mm diameter "pinhole" into a cylindrical inlet (length: 50 cm; diameter: 5 cm) which extends through a window blank on the starboard side of the aircraft.

Table 2. Details of supporting measurements.

Species	Instrument, technique	Time resolution; limit of detection (LOD)	References
CO	Aero Laser AL5002 Fast Carbon Monoxide Monitor. Excitation and fast-response fluorescence at $\lambda = 150$ nm.	1 s; 3.5 ppbv	Gerbig et al. (1999)
O_3	Thermo Scientific TEi49C ozone analyser. Absorption spectroscopy at $\lambda = 254$ nm.	1 s; 0.6 ppbv	Hewitt et al. (2010)
NO, NO_2, NO_x ($NO + NO_2$)	Air Quality Design dual-channel fast-response NO_x instrument. Chemiluminescence from $NO + O_3$ reaction. Conversion of NO_2 to NO by photolysis.	10 s; 3 pptv for NO, 15 pptv for NO_2	Stewart et al. (2008)
NO_2, ΣANs, ΣPNs	TD-LIF (thermal dissociation laser-induced fluorescence). Detection of NO_2 by laser-induced fluorescence. Thermal decomposition of ΣANs (total alkyl nitrate) and ΣPNs (total peroxy nitrate) to NO_2.	1 s; 9.8 pptv for NO_2, 28.1 pptv for ΣANs, 18.4 pptv for ΣPNs	Dari-Salisburgo et al. (2009); Di Carlo et al. (2013)
Alkenes	Whole air samples (WAS) analysed by laboratory-based gas chromatography with flame ionisation detection (GC-FID).	Typically 30 s; variable limits of detection	Hopkins et al. (2003)
NO_3, N_2O_5	BBCEAS (broadband cavity-enhanced absorption spectroscopy) of NO_3 at $\lambda = 642$–672 nm. N_2O_5 measured following thermal dissociation to $NO_3 + NO_2$.	1 s; 1.1 pptv for NO_3, 2.4 pptv for $NO_3 + N_2O_5$	Kennedy et al. (2011)
HCHO	Hantzsch technique: liquid-phase reaction of formaldehyde followed by excitation and fluorescence of resulting adduct at $\lambda = 510$ nm.	60 s; 81 pptv	Still et al. (2006)

Table 3. Mean mixing ratios of selected gas-phase species, and air temperature, measured during RONOCO and SeptEx. The flight and season during which the maximum values were measured are given in parentheses. NO_2 data are from the TD-LIF instrument. Zero values indicate measurements below the limit of detection.

Species	Summer RONOCO	SeptEx	Winter RONOCO	Maximum
CO/ppbv	102.3	117.1	139.3	256.0 (B537, summer)
O_3/ppbv	39.6	40.4	38.6	89.8 (B537, summer)
NO_3/pptv	21.1	0	6.2	176.9 (B537, summer)
NO/ppbv	0.05	0	0	18.9 (B539, summer)
NO_2/ppbv	1.6	1.7	2.3	18.6 (B568, winter)
Temperature/K	286.5	286.2	276.4	297.5 (B537, summer)



Figure 3. Night-time altitude profiles of **(a)** NO_3; **(b)** O_3; **(c)** *trans*-2-butene; and **(d)** propene, showing 60 s data (grey points) and mean values in 500 m altitude bins (black lines).

Downstream of the inlet are two low-pressure fluorescence cells positioned in series, the first for the detection of OH and the second for the detection of HO_2. During the RONOCO and SeptEx flights, the pressure inside the cells ranged from 1.9 Torr at ground level to 1.2 Torr at 6 km.

Laser light at $\lambda \sim 308$ nm is generated by a diode-pumped Nd:YAG-pumped tunable Ti:sapphire laser (Photonics Industries DS-532-10 and TU-UV-308 nm) and delivered to the fluorescence cells via optical fibres, on an axis perpendicular to the gas flow. A small fraction of the Ti:sapphire second harmonic ($\lambda = 462$ nm) is directed to the probe of a wavemeter to enable measurement of the laser wavelength to within 0.001 nm. A UV photodiode is positioned opposite the laser input arm on each fluorescence cell to measure laser power.

The sampled air forms a supersonic gas expansion beam in which the rate of collision between OH radicals and ambient air molecules is reduced. The OH fluorescence lifetime is

therefore extended to several hundred nanoseconds, significantly longer than the laser pulse, so that the measured signal can be temporally discriminated from laser scattered light. OH is excited from its ground state, $X^2\Pi_i\left(v''=0\right)$, to its first electronically excited state, $A^2\Sigma^+\left(v'=0\right)$, at $\lambda \sim 308$ nm. The resulting on-resonance fluorescence is detected by a UV-sensitive channel photomultiplier tube on an axis perpendicular to both the gas flow and the laser light. HO_2 is detected by titration with an excess of NO (Reaction R7), the resulting OH being detected as described.

The FAGE instrument was calibrated prior to and following each field measurement period, using a well-established method (Edwards et al., 2003; Faloona et al., 2004; Commane et al., 2010). Light at $\lambda = 184.9$ nm from a mercury pen-ray lamp photolyses water vapour in a flow of synthetic air inside an aluminium flow tube, generating OH and HO_2 at known concentrations. The aircraft FAGE instrument's limit of detection (LOD) for OH and HO_2 is determined by the instrument's sensitivity and the standard deviation of the background signal. During the RONOCO and SeptEx fieldwork, the 1σ LOD for a 5 min averaging period ranged between 0.64 and 1.8×10^6 molecule cm^{-3} for OH and between 5.9 and 6.9×10^5 molecule cm^{-3} for HO_2.

3.2 RO_2-based interference in FAGE measurements of HO_2

It has recently been shown that the reaction of alkene-derived β-hydroxyalkyl peroxy radicals, RO_2, with NO inside the HO_2 detection cell can lead to interference in FAGE HO_2 measurements (Fuchs et al., 2011; Whalley et al., 2013). The magnitude of the interference depends on the parent alkene, the residence time and mean temperature inside the cell, and the amount of NO injected. The interference therefore depends on the chemical environment and differs between FAGE instruments. In view of this, the University of Leeds ground-based and aircraft FAGE instruments have been tested for RO_2 interference. Thorough descriptions of the ground-based experimental method and results, and the results of a modelling study, are given by Whalley et al. (2013). The strongest interference in the aircraft instrument measurements was observed for ethene-derived RO_2, amounting to an increase of 39.7 ± 4.8 % in the observed HO_2 signal, with a cell pressure of 1.8 Torr, an estimated detection cell temperature of 255 K (obtained from rotational excitation spectra performed previously), and $[NO]_{cell} = 10^{14}$ molecule cm^{-3}.

Whalley et al. (2013) show that the chemistry responsible for the observed interferences is well known and that a model using the Master Chemical Mechanism (MCM, version 3.2: Jenkin et al., 1997; Saunders et al., 2003; Bloss et al., 2005, via http://mcm.leeds.ac.uk/MCM) can reproduce the interferences once tuned to the conversion efficiency of HO_2 to OH in the FAGE detection cell. Accordingly, Stone et al. (2014b) have applied the results of the ethene-derived RO_2 interference testing in a modelling study to assess the effect of the interference on the HO_2 measurements made during the RONOCO and SeptEx campaigns. A box model using a detailed MCM scheme was used to calculate a total potential interference in the RONOCO HO_2 measurements. The model was constrained to the conditions in the detection cell (1.8 Torr, 255 K, [NO] $\sim 10^{14}$ molecule cm^{-3}). Equal concentrations of HO_2 and $\sum RO_2$ (sum of all peroxy radicals in the MCM generated from the parent hydrocarbon) were used to initialise the model. The model run time was varied until the model-predicted interference from ethene-derived RO_2 radicals was equal to the experimentally determined interference, thereby tuning the model to the conversion efficiency of HO_2 to OH. An interference factor, f, was calculated for each RO_2 in the MCM as follows:

$$f = \frac{[OH]_{HO_2+RO_2} - [OH]_{HO_2}}{[OH]_{HO_2}}, \quad (1)$$

where $[OH]_{HO_2+RO_2}$ and $[OH]_{HO_2}$ are the modelled concentrations of OH produced from the reactions of RO_2 and HO_2 and the concentration from HO_2 alone, respectively. The greatest interference was calculated to come from isoprene-derived peroxy radicals, followed by aromatic compounds and C_2 to C_5 alkenes. The smallest modelled interference is from the C_1 to C_3 alkanes. The interference factors were applied to model-predicted RO_2 speciation and concentrations for the RONOCO flights. Model-predicted RO_2 species were dominated by CH_3O_2 (33 %; $f = 1.1$ %) and HO_2 (24 %; $f = 0.0$ %), with smaller contributions from RO_2 derived from *iso*-butene (12 %; $f = 0.5$ %), *cis*-2-butene and *trans*-2-butene (10 %; $f = 0.05$ %), and isoprene (2 %; $f = 7.6$ %). RO_2 species with high interference factors were a minor component of the total RO_2. A modelled value of HO_2 including the total potential interference, $HO_2{}^*$, was calculated using

$$\left[HO_2{}^*\right] = [HO_2]_{mod} + \sum_i f_i \left[RO_{2,i}\right]_{mod}. \quad (2)$$

Direct comparison between modelled values of $[HO_2{}^*]$ and the FAGE-measured values of $[HO_2]$ was therefore made possible. The model-predicted interference during the RONOCO campaign is described by $\left[HO_2{}^*\right] = 1.15\,[HO_2] + 2 \times 10^5$ molecule cm^{-3}. The average model-predicted interference in the HO_2 measurements is 14 %. The HO_2 measurements made during RONOCO and SeptEx were not adjusted since speciated RO_2 measurements were not available. The measurements are hereafter referred to as $HO_2{}^*$.

The magnitude of the RO_2 interference can be reduced by lessening the concentration of NO in the detection cell. This also reduces the instrument sensitivity to HO_2. Since the conversion of RO_2 to OH requires at least two NO molecules, while the conversion of HO_2 requires only one molecule, the ratio of HO_2 signal to RO_2 signal can be made favourable by reducing [NO] (Whalley et al., 2013). This effect has been investigated for the ground-based instrument and will

be investigated for the aircraft instrument prior to future HO_x measurement campaigns. An overview of the laboratory and computational studies of the interference in different FAGE instruments is given in a recent review by Stone et al. (2012).

3.3 BBCEAS measurements of NO_3 and N_2O_5

NO_3 and N_2O_5 were measured by the University of Cambridge broadband cavity-enhanced absorption spectroscopy (BBCEAS) instrument. The instrument was designed and built specifically for the RONOCO project and is described in detail in Kennedy et al. (2011). A brief description is given here.

The instrument consists of three 94 cm long high-finesse optical cavities formed by pairs of highly reflecting mirrors. The cavities are irradiated by incoherent broadband continuous wave light sources. Two of the cavities, for the detection of N_2O_5 and NO_3, are irradiated by red light-emitting diodes (LEDs) centred at 660 nm. The third cavity, for the detection of NO_2, is irradiated by a blue LED centred at 460 nm. The light from the LEDs is collimated using optical fibres and a focussing lens at the input of each cavity. A spectrometer, consisting of a spectrograph and charge couple device (CCD), is positioned at the end of each cavity to measure the wavelength-dependent intensity of transmitted light.

Ambient air is sampled through a rear-facing inlet on the aircraft fuselage, positioned approximately 4 m from the aircraft nose and 10 cm from the aircraft body. The air from the inlet is divided into two flows. The flow directed to the N_2O_5 cavity is heated to 120 °C to ensure near complete (> 99.6 %) thermal dissociation of N_2O_5 to NO_2 and NO_3. The cavity itself is heated to 80 °C and is used to measure the sum of the concentrations of ambient NO_3 plus NO_3 from thermal decomposition of N_2O_5. The second flow is unheated and is directed first through the NO_3 cavity and then through the NO_2 cavity. Background spectra are recorded at half-hour intervals during flights by halting the flow of ambient air and purging the cavities with nitrogen.

NO_3 is detected by its strong $B^2E' - X^2A'_2$ electronic transition centred at 662 nm. The concentration of NO_3 is determined by separating the finely structured NO_3 absorption features from the broad features caused by Rayleigh and Mie scattering using a fitting technique analogous to that employed in differential optical absorption spectroscopy (DOAS). A strong water vapour absorption feature that spectrally overlaps with NO_3 absorption around 662 nm is simulated for the pressure and temperature measured in the cavity and is removed from the measured absorption spectrum. The concentration of N_2O_5 is determined by subtracting the concentration of ambient NO_3 measured in the unheated cavity from the sum of the concentrations of ambient and dissociated NO_3 measured in the heated cavity.

Contributions to uncertainties in ambient measurements of NO_3 and N_2O_5, including wall losses of NO_3 and N_2O_5, temperature- and pressure-dependent absorption cross sections of NO_3 and H_2O, and the length of the cavity occupied by the sample, have been thoroughly investigated in laboratory experiments or addressed in the data analysis routine. In addition, wall losses of NO_3 and N_2O_5 were determined before and after each flight to account for changes in the surface properties of the inlet and detection cell walls, which were found to be negligible. The total uncertainty in the measured concentration of ambient NO_3 was 11 %. The uncertainty in the measured concentration of ambient N_2O_5 is determined for each individual ambient measurement, being dependent on the NO_3 / N_2O_5 ratio, and was on the order of 15 %. During RONOCO flights, the 1σ limits of detection for NO_3 and the sum of $NO_3 + N_2O_5$ were 1.1 and 2.4 pptv, respectively, for a 1 s integration time.

4 Overview of OH and HO_2* measurements

FAGE measurements were made on 16 flights during RONOCO and 9 flights during SeptEx. There was insufficient laser power during flights B534 to B536 in the summer campaign to measure both OH and HO_2* by dividing the laser light between the two cells. OH was therefore not measured during these flights. Low laser power throughout the summer fieldwork caused relatively high fluctuations in laser power overall and therefore higher background variability. This resulted in higher limits of detection for OH (1.8×10^6 molecule cm^{-3}) and HO_2* (6.9×10^5 molecule cm^{-3}).

Table 4 summarises the OH and HO_2* measurements during RONOCO and SeptEx and gives the instrument's average 1σ limit of detection for a 5 min averaging period. OH was not detected above the limit of detection during the summer or winter RONOCO flights, resulting in upper limits of 1.8×10^6 and 6.4×10^5 molecule cm^{-3} for mean summer and winter concentrations, respectively. These upper limit values are similar to previously reported night-time OH measurements (Geyer et al., 2003; Holland et al., 2003; Ren et al., 2003b; Emmerson and Carslaw, 2009). The mean daytime OH concentration during SeptEx was 1.8×10^6 molecule cm^{-3}, which was above the limit of detection. The mean HO_2* mixing ratio was highest during SeptEx (2.9 pptv) and was higher during summer (1.6 pptv) than during winter (0.7 pptv). The OH and HO_2* data sets for RONOCO and SeptEx are shown as altitude profiles in Figs. 4 and 5, respectively.

Table 5 gives the mean and maximum HO_2* mixing ratios at different times of day during summer, SeptEx, and winter. Dawn, day, dusk, and night are defined by the solar zenith angle as follows: dawn and dusk are between 90 and 102° and are distinguished by the time of day; day is between 0 and 90°; night is between 102 and 180°.

The mean dusk HO_2* mixing ratio in summer was higher than the mean night-time mixing ratio, suggesting that photochemical production was still active at dusk in summer. The reverse was true for the winter data, with the highest mean

Table 4. Combined daytime and night-time mean concentrations of OH and mean mixing ratios of HO_2* with the FAGE instrument's average 1σ limits of detection for a 5 min averaging period during the RONOCO and SeptEx fieldwork.

	OH/molecule cm^{-3}		HO_2*/pptv	
	Mean concentration	Limit of detection	Mean mixing ratio	Limit of detection
Summer		1.8×10^6	1.6	0.03
SeptEx	1.8×10^6	1.2×10^6	2.9	0.02
Winter		6.4×10^5	0.7	0.02



Figure 4. Altitude profile of OH measured during SeptEx showing 60 s data (grey points) and mean values in 500 m altitude bins (black lines). Error bars are 1 SD.

HO_2* mixing ratio being at night. This suggests that when photochemical production was suppressed in the winter daytime due to low photolysis rates, production via reactions of NO_3 and O_3 with alkenes was an important route to radical initiation. The RONOCO HO_2* measurements are similar to night-time, ground-based, urban measurements. For example, during the TORCH campaign, [HO_2] peaked at 1×10^8 molecule cm^{-3} at night (Emmerson et al., 2007), and during the PMTACS-NY 2001 field campaign, night-time HO_2 concentrations of 8×10^6 molecule cm^{-3} were measured (Ren et al., 2003b).

4.1 Case study flight B537: high night-time HO_2* concentrations

The highest HO_2* concentration (3.2×10^7 molecule cm^{-3}; 13.7 pptv) was measured during night-time flight B537 on 20 July 2010. Take-off from East Midlands Airport was at 22:00 local time (21:00 UTC, sunset at 20:18 UTC). The flight track, coloured by altitude, is shown in Fig. 6. The flight involved a profile descent from 3350 to 460 m down the Norfolk coast and a missed approach at Southend Airport (51.6° N, 0.70° E). Plumes from European continental outflow (see Fig. 7) were intersected by a series of runs at altitudes between 460 m and the upper boundary of the polluted layer.

Table 5. Mean and, in parentheses, maximum HO_2* mixing ratios measured during RONOCO and SeptEx.

	Mean (maximum) HO_2* mixing ratio/pptv		
	Summer	SeptEx	Winter
Dawn	0.74 (1.19)		0.54 (1.81)
Day		3.78 (11.79)	0.49 (1.68)
Dusk	2.73 (9.97)		0.32 (0.97)
Night	1.86 (13.58)		0.98 (2.02)



Figure 5. Altitude profiles of HO_2* measured in RONOCO and SeptEx during **(a)** dawn; **(b)** day; **(c)** dusk; and **(d)** night, showing 60 s data (grey points) and mean values in 500 m altitude bins (black lines). Error bars are 1 SD.

Flight B537 is an unusual flight within the RONOCO data set, with high concentrations of CO, O_3, NO_3, and high temperatures compared to the values measured during other night-time flights (see Table 3). The ambient aerosol surface area was significantly higher during B537 (nearly 800 µ m^2 cm^{-3}) than during other flights (between 100 and 400 µ m^2 cm^{-3}), and the organic aerosol concentration was significantly enhanced (Morgan et al., 2015). Footprint maps for flight B537, indicating regions where the sampled air was in contact with the surface prior to the flight, are shown in Fig. 7. The air sampled during the flight originated primarily over northern France, Belgium, and Germany.

A region of high surface pressure was positioned over the UK on the 20 July, with a mean air pressure of 1012.6 hPa over the 24 h prior to the flight. The mean air temperature 24 h prior to the flight (22:00, 19 July 2010, to 22:00, 20 July 2010), measured at a number of Met Office weather stations in Greater London, was 22.6 °C, and reached a maximum value of 28.6 °C. Wind speeds prior to the flight were low, with an average value of 4.7 knots (2.4 m s^{-1}). No rainfall was recorded at any of the Greater London weather sta-



Figure 6. Flight track of flight B537 on 20 July 2010, coloured by altitude.



Figure 7. Footprint maps for flight B537 on 20 July 2010, showing model particle densities (g s m^{-3}) in a 300 m deep layer from the surface, integrated over 24 h periods beginning **(a)** 24 h, **(b)** 48 h, **(c)** 72 h, and **(d)** 96 h prior to the flight.

tions during the 24 h prior to the flight. At Heathrow Airport (51.5° N, 0.45° W), 12.4 h of sunshine were recorded on the 20 July. High temperatures, combined with low wind speed, exposure to solar radiation, and little precipitation promote the formation of ozone as a result of photochemical processing of VOCs emitted at the surface (e.g. Lee et al., 2006) and offer an explanation for the high ozone mixing ratios measured during flight B537. Peak surface daytime ozone concentrations measured in Teddington, London, on 20 July



Figure 8. Time series of altitude (top panel, blue), HO_2* (middle panel, black, with grey shading representing the uncertainty in the measurements), O_3 (bottom panel, red), and NO_3 (bottom panel, green) during night-time flight B537 on 20 July 2010.

were on the order of 2.0×10^{12} molecule cm^{-3} ($\sim$ 78 ppbv) (data available at http://www.airquality.co.uk). Similar levels were recorded at a number of locations within Greater London.

Figure 8 shows a time series of altitude, HO_2*, O_3, and NO_3 mixing ratios during the flight, demonstrating very similar behaviour between the two radical species. During the missed approach at Southend Airport the mixing ratios of HO_2* and NO_3 increased with decreasing altitude, to reach values of 4.5 and 35 pptv, respectively, at 50 m above the ground. The maximum HO_2* and NO_3 mixing ratios were measured over the North Sea east of Ipswich (52.16° N, 2.34° E) at an altitude of 509 m, in the outflow of the London plume. Figure 9 shows scatter plots of HO_2* against NO_3 and O_3 during flight B537 and during the other night-time flights during RONOCO. Strong positive correlation is evident between HO_2* and NO_3 during B537 ($r = 0.97$), while during the remaining night flights there is still a significant, though weaker, correlation ($r = 0.58$). Moderate negative correlation is evident between HO_2* and O_3 during B537 ($r = -0.46$), with weak positive correlation existing for the other night-time flights ($r = 0.19$). The data suggest that NO_3 was an important initiator of HO_x radicals during flight B537 and that O_3 played a limited role overall during the night-time flights. Further investigation of the roles of NO_3 and O_3 in alkene oxidation and radical initiation at night is described in Sect. 5.

5 Oxidation of alkenes and production of HO_2: method of analysis

Following the work of Salisbury et al. (2001), the total rates of reaction, Φ, of O_3 and NO_3 with the alkenes measured



Figure 9. HO_2^* vs. **(a)** NO_3 and **(b)** O_3 during flight B537 (blue, filled circles) and during all other night-time flights (black, open circles). The lines are lines of best fit to the data.



Figure 10. Generalised reaction scheme for production of RO_2 and HO_2 following reaction of NO_3 with an alkene.

during RONOCO and SeptEx have been calculated:

$$\Phi_{O_3} = \sum_{i}^{\text{alkene}} k_{O_3+\text{alk}_i} [O_3][\text{alkene}_i] \quad (3)$$

$$\Phi_{NO_3} = \sum_{i}^{\text{alkene}} k_{NO_3+\text{alk}_i} [NO_3][\text{alkene}_i]. \quad (4)$$

The reactions of O_3 and NO_3 with alkenes yield OH, HO_2, and RO_2 radicals. A consideration of the reaction mechanisms of NO_3 and O_3 enables the calculation of the rate of instantaneous production of HO_2 (P_{HO_2}) from the reactions of NO_3 and O_3 with the alkenes measured during RONOCO, using the chemistry scheme, rate constants, and branching ratios in the MCM (Jenkin et al., 1997; Saunders et al., 2003).

Figure 10 shows a generalised reaction scheme for the reaction of NO_3 with an alkene. The reaction between NO_3 and an alkene proceeds via the addition of NO_3 to the double bond to form a nitrooxyalkyl radical, followed by rapid reaction with oxygen to yield a nitrooxyalkyl peroxy radical, RO_2 (shown as a single step in Fig. 10). The RO_2 radical can react with a number of species, of which NO, NO_3, and RO_2 lead to the production of an alkoxy radical (RO). Radical termination occurs via the reaction of RO_2 with HO_2 to yield a peroxide (ROOH) or with RO_2 to yield carbonyl ($RC(O)CH_3$) and alcohol (RCH_2OH) products. Reaction of RO with oxygen proceeds via the abstraction of a hydrogen atom to yield HO_2 or an aldehyde (RCHO). This generalised scheme can be applied to the reactions of NO_3 with all the alkenes measured. The rate of instantaneous production of HO_2 is found by first calculating the fraction of RO_2 that reacts to produce RO (F_{RO}) and the fraction of RO that reacts to produce HO_2 (F_{HO_2}):

$$F_{RO} = \frac{k_3[NO] + k_4[NO_3] + 0.6k_5[RO_2]}{k_2[HO_2] + k_3[NO] + k_4[NO_3] + k_5[RO_2]} \quad (5)$$

$$F_{HO_2} = \frac{k_6[O_2]}{k_7 + k_6[O_2]}, \quad (6)$$

where RO_2 represents all peroxy radicals. Average values of F_{RO} for the NO_3+ alkene reactions range between 0.50 for *trans*-2-pentene- and 1-pentene-derived RO_2 species and 0.61 for ethene-derived RO_2 species. F_{HO_2} varies between 0 and 1 for the alkenes studied. Overall, the rate of production of HO_2 (P_{HO_2}) from reactions of NO_3 with alkenes is then given by

$$P_{HO_2} = k_i[NO_3][\text{alkene}_i] \times F_{RO} \times F_{HO_2}. \quad (7)$$

The reaction scheme for the reaction of O_3 with alkenes is more complicated because the number and type of radicals produced in the O_3 + alkene reaction depends on the structure of the alkene. The simplest case is the reaction of ozone with ethene. Ozone adds to the double bond to form a five-membered ring called a primary ozonide. Decomposition of the ozonide yields an excited Criegee intermediate (CH_2OO^*) and a carbonyl compound (in this case formaldehyde, HCHO). The energy-rich Criegee intermediate can be stabilised by collision with a third body or undergo decomposition to yield products including OH, CO, and HO_2. The primary ozonide produced in the O_3+ propene reaction (see Fig. 11) can decompose via two channels, yielding carbonyls and Criegee intermediates with different structures and different products, including RO_2. Reaction of RO_2 with NO, NO_3, and RO_2 (all peroxy radicals) yields RO, which in turn yields HO_2.

The rates of production of HO_2 from reactions of O_3 with alkenes (P_{HO_2}) have been calculated as follows:

$$P_{HO_2,\text{Direct}} = k_i[O_3][\text{alkene}_i] \times \alpha_{HO_2} \quad (8)$$

$$P_{HO_2,RO_2} = k_i[O_3][\text{alkene}_i] \times \alpha_{RO_2} \times F_{RO} \times F_{HO_2} \quad (9)$$

$$P_{HO_2} = P_{HO_2,\text{Direct}} + P_{HO_2,RO_2}, \quad (10)$$

where $P_{HO_2,\text{Direct}}$ is the rate of direct HO_2 production from Criegee intermediate decomposition, α_{HO_2} is the branching ratio to HO_2-producing channels from the Criegee intermediate, P_{HO_2,RO_2} is the rate of HO_2 production from RO_2 radicals produced in the O_3 + alkene reaction, α_{RO_2} is the branching ratio to RO_2-producing channels from the Criegee intermediate, F_{RO} is the fraction of RO_2 radicals that react to produce RO radicals, and F_{HO_2} is the fraction of RO radicals that react to produce HO_2 radicals, which is equal to 1 for all the alkenes studied. Average values of F_{RO} for



Figure 11. Reaction scheme for O_3 + propene, showing production of HO_2 and the methyl peroxy radical, CH_3O_2.



Figure 12. Average night-time rates of reaction between O_3, and NO_3 with alkenes during **(a)** summer and **(b)** winter RONOCO flights. Error bars represent the combined uncertainty in the measurements.

the O_3+ alkene reactions range between 0.54 for 1-pentene-derived RO_2 species and 0.64 for 1-butene- and *trans*-2-pentene-derived RO_2 species.

The reactions of RO_2 with NO to form $RONO_2$ have been omitted from the calculations because the branching ratio is small (0.001 to 0.02) for the radicals studied (Carter and Atkinson, 1989; Lightfoot et al., 1992). The reaction of CH_3O_2 with NO_2 to form $CH_3O_2NO_2$ has been omitted from the calculations, since the reverse reaction is much faster than the forward direction ($k_f = 6.4 \times 10^{-12}\,s^{-1}$; $k_{rev} = 1.08\,s^{-1}$ at a mean temperature of 286.5 K during RONOCO).

The primary aims of the analysis presented here are threefold: (1) to calculate the total rate of initiation through reactions of NO_3 and O_3 with alkenes; (2) to determine the relative importance of NO_3 and O_3 in night-time HO_2 production; (3) to investigate differences in radical production between different seasons and different times of day. The correlation between $[HO_2]^*$ and $[NO_3]$, especially during flight B537, will be investigated.

P_{HO_2} has been calculated for each alkene measured for every 60 s data point where all the requisite data were available and where $HO_2{}^*$ was above the limit of detection of the FAGE instrument. Concentrations of RO_2 were calculated by scaling the observed $HO_2{}^*$ concentrations with the $RO_2 / HO_2{}^*$ ratio calculated using a box model constrained to the concentrations of long-lived species measured during the flights (Stone et al., 2014b), i.e. $RO_{2,obs} = HO_2{}^*{}_{,obs} \times RO_{2,mod} / HO_2{}^*{}_{,mod}$. The rates of reaction and rates of production of HO_2 presented hereafter are average values for individual flights, seasons, or times of day.

6 Results

6.1 Night-time oxidation of alkenes

Figure 12 shows histograms of the rate of reaction between O_3 and NO_3 with individual alkenes during summer and winter, for the night-time data only. The reactivity of measured alkenes ($\Phi_{O_3} + \Phi_{NO_3}$) was greater by a factor of 2.2 during summer flights than during winter flights. The reactions of NO_3 are largely responsible for this seasonal difference, since the contribution from O_3 + alkene reactions varies little between summer (4.1×10^4 molecule $cm^{-3}\,s^{-1}$) and winter (3.9×10^4 molecule $cm^{-3}\,s^{-1}$). The factor of 4.1 difference between the rate of NO_3 reactions in summer (9.8×10^4 molecule $cm^{-3}\,s^{-1}$) and winter (2.4×10^4 molecule $cm^{-3}\,s^{-1}$) can be attributed to the higher mean concentration of NO_3 in summer (5.8×10^8 molecule cm^{-3}) compared to winter (2.0×10^8 molecule cm^{-3}). This seasonal difference in NO_3 concentrations is attributable to the lower mean night-time temperature in winter (277.7 K) compared to summer (286.7 K), which disfavours NO_3 in the thermal equilibrium N_2O_5 $NO_3 + NO_2$. $K_{eq}[NO_2]$, which determines $[N_2O_5] / [NO_3]$, is calculated to be 4.8 in summer and 29.6 in winter. At night in summer, Φ_{NO_3} was greater than Φ_{O_3} by a factor of 2.4, but in winter Φ_{O_3} was a factor of 1.6 greater than Φ_{NO_3}. Figure 12 illustrates the importance of the butene isomers (within the VOCs measured) in the reactions of O_3 and NO_3 and therefore radical initiation and propagation. Reactions with *iso*-butene dominated NO_3 reactivity in summer (42 %) and winter (53 %), with *trans*-2-butene also contributing significantly (28 % in summer and 32 % in winter). Reactions of O_3 were dominated by *trans*-2-butene (42 % in summer and 34 % in winter) and propene (26 % in summer and 38 % in winter). The importance of these alkenes is attributed to their relatively high abundances compared to the other alkenes measured,

Table 6. Average rates of instantaneous production of HO_2 from reactions of O_3 and NO_3 with alkenes.

	HO_2 production rate (ΣP_{HO_2})/ 10^4 molecule cm^{-3} s^{-1}			
	Dawn	Day	Dusk	Night
Summer				
NO_3	0		2.8	3.8
O_3	0.5		2.2	1.7
Total	0.5		5.0	5.5
Winter				
NO_3	0.4	0.4	0.4	0.5
O_3	1.4	1.5	1.2	1.2
Total	1.8	1.9	1.6	1.7

during both summer and winter, combined with their fast rates of reaction with O_3 and NO_3.

For comparison with the reactions of O_3 and NO_3, the total rate of reaction of measured alkenes with OH has been calculated using upper limits on OH concentrations of 1.8×10^6 molecule cm^{-3} and 6.4×10^5 molecule cm^{-3} for the summer and winter flights, respectively, based on the FAGE instrument's limit of detection. The high upper limits make the total rate of reaction of OH with alkenes, Φ_{OH}, unrealistically high for both summer (1.6×10^5 molecule cm^{-3} s^{-1}) and winter (7.8×10^4 molecule cm^{-3} s^{-1}). However, the OH reactivity will likely be considerably lower than the values calculated using the OH upper limits. A box model constrained to concentrations of long-lived species measured during the flights (Stone et al., 2014b) predicts a mean OH concentration of 2.4×10^4 molecule cm^{-3}, significantly lower than the upper limits given by the instrument's limit of detection. Using the mean modelled value for OH gives $\Phi_{OH} = 2.1 \times 10^3$ molecule cm^{-3} s^{-1} for summer and $\Phi_{OH} = 2.9 \times 10^3$ molecule cm^{-3} s^{-1} for winter, indicating a diminished role for OH in alkene oxidation at night, in agreement with previous studies (e.g. Geyer et al., 2003; Emmerson et al., 2005).

6.2 Daytime oxidation of alkenes

Figure 13 shows histograms of rates of reaction of O_3 and OH with alkenes during SeptEx and of O_3 and NO_3 with alkenes during winter RONOCO flights, for daytime data only. OH was detected above the limit of detection (1.2×10^6 molecule cm^{-3}) during the SeptEx flights, so the FAGE OH data were included in the calculations, using a reaction scheme analogous to the one shown in Fig. 10. NO_3 was not detected during the day in SeptEx. NO_3 is not expected to be present at measurable concentrations during daylight hours due to photolysis, but a mean concentration of 8.3×10^7 molecule cm^{-3} (3.3 pptv) was measured during the day in the winter RONOCO flights. These measurements of low mixing ratios of NO_3 may be partly caused by interference from other daytime species as observed by Brown et al. (2005) or by the variability in the instrument baseline, which can be on the order of 1–2 pptv during vertical profiles on the aircraft (Kennedy et al., 2011). This variability is small compared to the range of NO_3 values typically observed during RONOCO flights (0–50 pptv during summer; 0–10 pptv during winter). During SeptEx, Φ_{OH} exceeded Φ_{O_3} by a factor of 8. Ethene and propene were the two most abundant alkenes measured during SeptEx and contributed significantly to OH reactivity. O_3 reactivity with alkenes was dominated by propene and *trans*-2-butene (the six most abundant alkenes measured during SeptEx). NO_3 reactivity with alkenes was dominated by *trans*-2-butene and isobutene (the three most abundant alkene measured during winter daytime flights). The total rate of reaction of O_3 and OH with alkenes during daytime SeptEx flights (3.7×10^5 molecule cm^{-3} s^{-1}) exceeded the total rate of reaction of O_3 and NO_3 during daytime winter RONOCO flights (6.6×10^4 molecule cm^{-3} s^{-1}) by a factor of 6 and was more than double the total rate of reaction of O_3 and NO_3 with alkenes during night-time summer flights (1.4×10^5 molecule cm^{-3} s^{-1}). In winter daytime flights, Φ_{O_3} was greater than Φ_{NO_3} by a factor of 2.4.



Figure 13. Average daytime rates of reaction of **(a)** O_3 and OH with alkenes during SeptEx and **(b)** O_3 and NO_3 with alkenes during winter. Note the different scales. NO_3 was not detected during daytime SeptEx flights (LOD = 1.1 pptv); OH was not detected during daytime winter flights (LOD = 6.4×10^5 molecule cm^{-3}). Error bars represent the combined uncertainties in the measurements.

Figures 12b and 13b reveal that reactions of O_3 dominated alkene reactivity during both daytime and night-time winter RONOCO flights. The concentrations of alkenes were generally higher at night, with the total alkene concentration (sum of concentrations of alkenes measured) being 2.1×10^9 molecule cm^{-3} in the day and 3.4×10^9 molecule cm^{-3} at night. The total measured alkene reactivity ($\Phi_{O_3} + \Phi_{NO_3}$) was marginally higher during the



Figure 14. Average rates of instantaneous production of HO_2 from reactions of O_3 and NO_3 with alkenes during **(a)** summer and **(b)** winter RONOCO flights. Error bars represent the combined uncertainty in the measurements.

day, by a factor of 1.04. This difference is attributable mainly to the change in Φ_{O_3}.

Comparison of Figs. 12a and 13b reveals that the total measured alkene reactivity ($\Phi_{O_3} + \Phi_{NO_3}$) was higher during the summer night-time flights (1.4×10^5 molecule cm^{-3} s^{-1}) than during the winter daytime flights (6.6×10^4 molecule cm^{-3} s^{-1}), indicating a low oxidising environment during winter daytime. The additional contribution to measured alkene reactivity from reactions with OH has been calculated using the OH upper limits as described in Sect. 6.1. Even with this additional, upper-limit OH reactivity (1.6×10^5 and 1.1×10^5 molecule cm^{-3} s^{-1} for summer night-time and winter daytime, respectively), the total summer night-time alkene reactivity remains higher than that during winter daytime, confirming the importance of the summer nocturnal troposphere for the oxidation of the measured alkenes.

6.3 Night-time production of HO_2 from reactions of O_3 and NO_3 with alkenes

Table 6 gives the total rates (ΣP_{HO_2}) of instantaneous production of HO_2 from the reactions of O_3 and NO_3 with alkenes. NO_3 was not detected during the dawn summer RONOCO flights, and there were no daytime RONOCO flights during summer. NO_3 dominated HO_2 production during dusk and night (68 %), in agreement with Geyer et al. (2003), who found that NO_3 was responsible for 53 % of HO_2 production at night in the BERLIOZ campaign. During winter, O_3 dominated HO_2 production at all times, with a night-time contribution of 70 %. This is in agreement with the results from the winter PMTACS-NY 2004 field campaign (Ren et al., 2006).

The total rate of instantaneous production of HO_2 at night was 3.3 times greater in summer than in winter, with production from O_3 decreasing by a factor of 1.5 and production from NO_3 decreasing by a factor of 7.8 between summer and winter. The mean temperature difference of 9 K between summer and winter is thought to be responsible for the lower NO_3 concentrations in winter (2.0×10^8 molecule cm^{-3}, 8.2 pptv, compared to 5.8×10^8 molecule cm^{-3}, 24.5 pptv, in summer), owing to the increased thermal stability of N_2O_5, and for the reduced rate of temperature-dependent reactions between NO_3 and alkenes and subsequent reactions. There was very little difference between summer and winter mean O_3 mixing concentrations (9.6×10^{11} molecule cm^{-3}, 39.6 ppbv, and 9.4×10^{11} molecule cm^{-3}, 38.6 ppbv, respectively).

The production of HO_2 via reactions of NO_3 and O_3 with alkenes is now examined in more detail. The rate of production from individual alkenes was calculated and plotted in a histogram, as shown in Fig. 14 for the summer and winter night-time data. During both summer and winter, reactions of O_3 and NO_3 with *trans*-2-butene were important sources of HO_2, contributing on average 62 % to O_3-initiated HO_2 production and 36 % to NO_3-initiated production during the summer and winter flights. Reactions of NO_3 with isoprene were important during summer, contributing 28 % to NO_3-initiated production. The importance of *trans*-2-butene, despite its relatively low abundance during summer and winter night-time RONOCO flights (1.8 and 1.7 pptv, respectively, compared to ethene mixing ratios of 55.0 and 104.5 pptv), is attributed to its fast rates of reaction with both O_3 and NO_3 compared to the other alkenes measured. The importance of the isoprene + NO_3 reactions during the summer RONOCO flights is similarly attributed to its fast rate of reaction with NO_3 compared to the other alkenes measured. In addition there is no aldehyde-forming channel from the isoprene-derived RO radical (k_7 in Fig. 10), so that the yield of HO_2 from RO is equal to 1. The reaction of isobutene with NO_3 can proceed via one of two channels to produce two different RO_2 radicals but only one channel, with a branching ratio of 0.2, produces HO_2. Isobutene is therefore not a dominant contributor to HO_2 production, despite being the single largest contributor to NO_3 reactivity during daytime and night-time RONOCO flights (Figs. 12 and 13). Figure 14 highlights the small change in total production from O_3 between summer and winter and the dramatic change in total production from NO_3 between summer and winter.

Reactions of formaldehyde with NO_3 were included in the analysis where formaldehyde data were available (mean HCHO = 955 pptv). The NO_3 + HCHO reaction contributed a further 5.5×10^3 molecule cm^{-3} s^{-1} (15 %) to HO_2 production from NO_3 reactions, so that production from NO_3 contributed 79 % of the total production.

6.4 Production of HO_2 during flight B537

Flight B537, on 20 July 2010, has been identified as an interesting flight, with high concentrations of HO_2* (3.2×10^8 molecule cm^{-3}; 13.6 pptv), ozone (peaking at



Figure 15. Average rates of instantaneous production of HO_2 from reactions of O_3 and NO_3 with alkenes during flight B537. Error bars represent the combined uncertainty in the measurements.



Figure 16. [HO_2*] vs. total rate of instantaneous production of HO_2 from reactions of **(a)** O_3 and **(b)** NO_3 during flight B537. Correlation coefficients (r) are given in each plot. The lines are lines of best fit to the data.

1.8×10^{12} molecule cm^{-3}; 89.9 ppbv), and NO_3 (peaking at 4.1×10^9 molecule cm^{-3}; 176.9 pptv), and a strong positive correlation between HO_2* and NO_3 ($r = 0.97$; see Fig. 9). NO, NO_2, and aerosol surface area were also elevated in-flight during flight B537 compared to their mean summer values. The highest concentration of ethene (1.43×10^{10} molecule cm^{-3}; 0.61 ppbv) during the summer RONOCO flights was measured during B537. ΣP_{HO_2} from O_3 + alkene reactions (2.6×10^4 molecule cm^{-3} s^{-1}) was higher in flight B537 than in all the other summer flights, contributing 42 % of HO_2 production, with NO_3 + alkene reactions contributing 3.6×10^4 molecule cm^{-3} (58 %). The total rate of HO_2 production from O_3 and NO_3 reactions during flight B537 was 6.2×10^4 molecule cm^{-3} s^{-1}. While this is higher than the average value of ΣP_{HO_2} for the summer flights (5.4×10^4 molecule cm^{-3} s^{-1}), it is not the highest rate of production during the summer flights. During B534 unusually high concentrations of isoprene, *cis*-2-butene, and 1,3-butadiene contributed to a total rate of HO_2 production of 7.9×10^4 molecule cm^{-3}, which is the highest calculated value.

Figure 15 shows that the reactions of O_3 and NO_3 with *trans*-2-butene are once again important, contributing 74 % of $\Sigma P_{HO_2,O_3}$ and 45 % of $\Sigma P_{HO_2,NO_3}$. The correlation between HO_2* and NO_3 is attributed to the production of HO_2 by reactions of NO_3 with alkenes, especially *trans*-2-butene. Figure 16 shows HO_2* vs. the total instantaneous rate of production from the reactions of O_3 and NO_3 with alkenes during flight B537 at each 60 s data point during the flight for which the requisite data were available. Note that the rates plotted in Fig. 16 are higher than those shown in Fig. 15, where the rates of production of HO_2 from each alkene have been averaged across the whole flight. A strong positive correlation exists between HO_2* and both $\Sigma P_{HO_2,O_3}$ ($r = 0.6$) and $\Sigma P_{HO_2,NO_3}$ ($r = 0.8$), indicating the importance of these reactions for the production of HO_2 during this flight.

7 Comparison with model results

The observations of OH, HO_2*, NO_3, and N_2O_5 have been interpreted in the context of night-time oxidation chemistry using a box model constrained to observations of VOCs, NO_x, O_3, CO, and other long-lived species measured during the RONOCO flights (Stone et al., 2014b). The Dynamically Simple Model of Atmospheric Chemical Complexity (DSMACC) (Emmerson and Evans, 2009; Stone et al., 2010, 2014b) was initiated with concentrations of measured species, using a chemistry scheme based on the Master Chemical Mechanism (MCM, version 3.2: Jenkin et al., 1997, 2003; Saunders et al., 2003; Bloss et al., 2005, via http://mcm.leeds.ac.uk/MCM), and was allowed to run to diurnal steady state. The model output includes concentrations of OH, HO_2, NO_3, RO_2, and other species. Data from daytime flights, or during dawn or dusk periods, were not included in the model analysis. Data from flight B537 were also excluded, owing to the atypical observations of HO_2*, NO_3, O_3, and other chemical species made during this flight. The modelling study and results are described in more detail by Stone et al. (2014b).

The model predicts a mean OH concentration of 2.4×10^4 molecule cm^{-3} for the summer flights, which is consistent with the measured OH concentrations for which the instrument's limit of detection is an upper limit only. The base model underpredicts HO_2* by around 200 % and overpredicts NO_3 and N_2O_5 by 80 and 50 %, respectively. These discrepancies were investigated by determining the processes controlling radical production and loss in the model and using those results to improve model performance. Model production of HO_2 is dominated by reactions of RO + O_2 (42 %), with a significant contribution from OH + CO (31 %) despite low OH concentrations at night. RO_x (= RO + RO_2 + OH + HO_2) radical initiation in the model is dominated by reactions of NO_3 with unsaturated VOCs (80 %), with a much smaller contribution (18 %) from

alkene ozonolysis. Modelled HO_2 loss is dominated by its reactions with NO_3 (45 %) and O_3 (27 %), both of which are radical propagating routes and which are the dominant routes to OH production in the model. In fact NO_3 was found to control both radical initiation and propagation in the model.

These results are in general agreement with the results of the analysis presented in Sect. 6.1, though the model predicts a more important role for NO_3 (80 % of RO_x radical production, which is 7.2 times the contribution from O_3+ alkenes) than is predicted by the analysis based on the observations alone (69 % of HO_2 radical production during summer, which is 2.1 times the contribution from O_3+ alkenes). The model predicts a relatively small role for O_3 in both summer and winter. The model is constrained to measured values of O_3 but overpredicts NO_3. The mean measured NO_3 night-time mixing ratio was 24.5 pptv in the summer and 8.2 pptv in the winter. The mean modelled summer and winter values are 37.4 and 20.7 pptv, respectively. This discrepancy between modelled and measured NO_3 helps to explain the model overprediction of the role of NO_3 in HO_x radical initiation during the RONOCO flights. Modelled NO_3 reactivity was dominated by *iso*-butene (36 %) and *trans*-2-butene (27 %), and modelled O_3 reactivity was dominated by *trans*-2-butene (51 %), in agreement with the night-time alkene reactivities presented in Sect. 6.1.

An improvement to the model predictions of NO_3, N_2O_5, and HO_2^* was made by increasing the concentration of unsaturated VOCs in the model. Increasing the total observed alkene concentration by 4 times resulted in a modelled-to-observed ratio of 1.0 for HO_2^* and of ~ 1.2 for NO_3 and N_2O_5. Two-dimensional gas chromatography (GC × GC) analysis of the whole-air samples taken during RONOCO has revealed a large number of VOCs extra to those routinely measured (Lidster et al., 2014). Calibration standards for the majority of these species are not yet available, and so the quantification of their concentrations is not possible, but their detection confirms that the model overprediction of NO_3 and underprediction of HO_2^* are attributable to reactions of NO_3 with unquantified unsaturated hydrocarbons.

The presence of unquantified unsaturated VOCs during the RONOCO campaign, suggested by the model and confirmed by the two-dimensional GC analysis, has implications for the conclusions drawn from the analysis based on the observations. The relative contributions of NO_3 and O_3 to night-time radical initiation will change with the composition of unsaturated VOCs in the sampled air, due to the different rates of reaction of NO_3 and O_3 with different VOC species and the rates of production of HO_2 following these reactions. The model results indicate that the reaction of NO_3 with the unquantified VOCs leads to increased production of HO_2. The role of NO_3 in night-time radical production would therefore be enhanced by the inclusion of the unquantified VOCs in the observational analysis.

8 Conclusions

Night-time radical chemistry has been studied as part of the RONOCO and SeptEx campaigns onboard the BAe-146 research aircraft during summer 2010 and winter 2011. NO_3, N_2O_5, OH, and HO_2^* were measured simultaneously for the first time from an aircraft, with OH and HO_2^* being measured by the University of Leeds aircraft FAGE instrument. OH was detected above the limit of detection during the daytime SeptEx flights only, with a mean concentration of 1.8×10^6 molecule cm^{-3}. Upper limits of 1.8×10^6 and 6.4×10^5 molecule cm^{-3} are placed on mean OH concentrations for the summer and winter RONOCO (night, dawn, and dusk) measurement campaigns, respectively. HO_2^* was detected above the limit of detection during the summer and winter RONOCO flights and during SeptEx, with a maximum mixing ratio of 13.6 pptv measured during the night-time flight B537 on 20 July 2010. Mean night-time HO_2^* mixing ratios were significantly higher in summer than in winter. Significant concentrations (up to 176.9 pptv) of NO_3 were measured during night-time flights, since the air masses sampled were sufficiently removed from the surface that the loss of NO_3 by reaction with NO was minimised. The RONOCO flights were therefore an excellent opportunity to study the role of NO_3 in nocturnal oxidation and radical initiation.

The rates of reaction of O_3 and NO_3 with the alkenes measured have been calculated. At night during summer, NO_3 dominated alkene reactivity. Several previous night-time studies have also found NO_3 to be the dominant nocturnal oxidant (e.g. Geyer et al., 2003; Brown et al., 2011). During night-time winter RONOCO flights the total rate of reaction of NO_3 with alkenes was much reduced, but the rate of reaction of O_3 with alkenes was similar to that in summer. During day and night in winter, O_3+ alkene reactions were faster than NO_3+ alkene reactions. Overall, during RONOCO, the combined rate of alkene oxidation by O_3 and NO_3 was highest at night during summer.

The calculation of the rates of the instantaneous production of HO_2 from reactions of O_3 and NO_3 with alkenes, using measurements made during the flights, has revealed that night-time production was dominated by NO_3 in summer and by O_3 in winter. The rate of instantaneous production of HO_2 from reactions of NO_3 with alkenes decreased significantly from summer to winter (87 %), whereas production from O_3+ alkene reactions was similar in summer and winter, decreasing by just 31 %. Strong positive correlation between HO_2^* and NO_3, especially during flight B537, is attributed to the production of HO_2 from reactions of NO_3 with alkenes, particularly *trans*-2-butene and other isomers of butene.

Significant concentrations of HO_2^* were detected at night, with the highest HO_2^* concentration (13.6 pptv) being measured during a summer night-time flight, indicating that HO_x radical chemistry remains active at night under the right con-

ditions. The role of HO_x is diminished in the low photolysis winter daytime atmosphere, with alkene ozonolysis being primarily responsible for oxidation and radical initiation, in agreement with previous studies (e.g. Heard et al., 2004; Emmerson et al., 2005). Both the analysis presented here and the results of the box modelling study by Stone et al. (2014b) indicate that in air masses removed from sources of NO, NO_3 plays an important role in the oxidation of alkenes and radical initiation at night, in agreement with previous studies (e.g. Brown et al., 2011). Alkene ozonolysis also plays a significant role in nocturnal oxidation in agreement with Salisbury et al. (2001), Geyer et al. (2003), Ren et al. (2003a, 2006), Emmerson et al. (2005), and Volkamer et al. (2010). The balance between the roles of NO_3 and O_3 was controlled in part by [NO_3], with colder winter temperatures forcing the NO_3–N_2O_5 equilibrium towards N_2O_5.

The total rate of reaction of O_3 and NO_3 with alkenes during night-time summer flights (1.4×10^5 molecule cm^{-3} s^{-1}) was higher by a factor of 2.1 than during daytime winter flights (6.6×10^4 molecule cm^{-3} s^{-1}). Whilst it should be remembered that measurements at different times of day and in different seasons reflect composition changes in air masses (such as the abundance of reactive alkenes), this result supports the hypothesis that oxidation of certain VOCs, in particular the reactive alkenes, in the nocturnal summer atmosphere can be as rapid as in the winter daytime atmosphere.

A box model of night-time chemistry constrained to measurements of long-lived species has been used to investigate the night-time chemistry sampled during RONOCO (Stone et al., 2014b). The base model underpredicts HO_2* and overpredicts NO_3. These discrepancies were minimised by increasing the concentration of alkenes in the model, thereby increasing the reaction of NO_3 with alkenes and the production of HO_2. The presence of unquantified unsaturated VOCs has been confirmed by 2D-GC analysis, though the exact nature and concentrations of the 'missing' species are unclear. The inclusion of these species in the analysis presented in this paper would likely increase the role of NO_3 in the oxidation of alkenes and production of HO_2 at night.

Acknowledgements. This work was funded by the UK Natural Environment Research Council (NE/F004664/1). The authors would like to thank ground staff, engineers, scientists, and pilots involved in RONOCO for making this project a success. Airborne data were obtained using the BAe 146-301 Atmospheric Research Aircraft (ARA) flown by Directflight Ltd. and managed by the Facility for Airborne Atmospheric Measurements (FAAM), which is a joint entity of the Natural Environment Research Council (NERC) and the Met Office.

Edited by: S. Brown

References

Andrés-Hernández, M. D., Kartal, D., Crowley, J. N., Sinha, V., Regelin, E., Martínez-Harder, M., Nenakhov, V., Williams, J., Harder, H., Bozem, H., Song, W., Thieser, J., Tang, M. J., Hosaynali Beigi, Z., and Burrows, J. P.: Diel peroxy radicals in a semi-industrial coastal area: nighttime formation of free radicals, Atmos. Chem. Phys., 13, 5731–5749, doi:10.5194/acp-13-5731-2013, 2013.

Atkinson, R. and Arey, J.: Gas-phase tropospheric chemistry of biogenic volatile organic compounds: a review, Atmos. Environ., 37, 197–219, 2003.

Beames, J. M., Liu, F., Lu, L., and Lester, M. I.: Ultraviolet Spectrum and Photochemistry of the Simplest Criegee Intermediate CH_2OO, J. Am. Chem. Soc., 134, 20045–20048, 2012.

Beames, J. M., Liu, F., Lu, L., and Lester, M. I.: UV spectroscopic characterization of an alkyl substituted Criegee intermediate CH_3CHOO, J. Chem. Phys., 138, 244307, doi:10.1063/1.4810865, 2013.

Berndt, T. and Böge, O.: Kinetics of Oxirane Formation in the Reaction of Nitrate Radicals with Tetramethylethylene, Berichte der Bunsengesellschaft für physikalische Chemie, 98, 869–871, 1994.

Bey, I., Aumont, B., and Toupance, G.: A modeling study of the nighttime radical chemistry in the lower continental troposphere, J. Geophys. Res.-Atmos., 106, 9991–10001, 2001.

Bloss, C., Wagner, V., Bonzanini, A., Jenkin, M. E., Wirtz, K., Martin-Reviejo, M., and Pilling, M. J.: Evaluation of detailed aromatic mechanisms (MCMv3 and MCMv3.1) against environmental chamber data, Atmos. Chem. Phys., 5, 623–639, doi:10.5194/acp-5-623-2005, 2005.

Brown, S. S. and Stutz, J.: Nighttime radical observations and chemistry, Chem. Soc. Rev., 41, 6405–6447, 2012.

Brown, S. S., Dibb, J. E., Stark, H., Aldener, M., Vozella, M., Whitlow, S., Williams, E. J., Lerner, B. M., Jakoubek, R., Middlebrook, A. M., DeGouw, J. A., Warneke, C., Goldan, P. D., Kuster, W. C., Angevine, W. M., Sueper, D. T., Quinn, P. K., Bates, T. S., Meagher, J. F., Fehsenfeld, F. C., and Ravishankara, A. R.: Nighttime removal of NO_x in the summer marine boundary layer, Geophys. Res. Lett., 31, L07108–L07113, 2004.

Brown, S. S., Osthoff, H. D., Stark, H., Dubé, W. P., Ryerson, T. B., Warneke, C., de Gouw, J. A., Wollny, A. G., Parrish, D. D., Fehsenfeld, F. C., Ravishankara, A. R.: Aircraft observations of daytime NO_3 and N_2O_5 and their implications for tropospheric chemistry, J. Photoch. Photobio. A, 176, 270–278, 2005.

Brown, S. S., Ryerson, T. B., Wollny, A. G., Brock, C. A., Peltier, R., Sullivan, A. P., Weber, R. J., Dubé, W. P., Trainer, M., Meagher, J. F., Fehsenfeld, F. C., and Ravishankara, A. R.: Variability in Nocturnal Nitrogen Oxide Processing and Its Role in Regional Air Quality, Science, 311, 67–70, 2006.

Brown, S. S., Dubé, W. P., Peischl, J., Ryerson, T., Atlas, E., Warneke, C., de Gouw, J. A., te Lintel Hekkert, S., Brock, C. A., Flocke, F., Trainer, M., Parrish, D. D., Feshenfeld, F. C., and Ravishankara, A.: Budgets for nocturnal VOC oxidation by nitrate radicals aloft during the 2006 Texas Air Quality Study, J. Geophys. Res.-Atmos., 116, D24305–D24320, 2011.

Carter, W. P. L. and Atkinson, R.: Alkyl Nitrate Formation from the Atmospheric Photooxidation of Alkanes; a Revised Estimation Method, J. Atmos. Chem., 8, 165–173, 1989.

Commane, R., Floquet, C. F. A., Ingham, T., Stone, D., Evans, M. J., and Heard, D. E.: Observations of OH and HO_2 radicals over West Africa, Atmos. Chem. Phys., 10, 8783–8801, doi:10.5194/acp-10-8783-2010, 2010.

Creasey, D. J., Heard, D. E., and Lee, J. D.: Eastern Atlantic Spring Experiment 1997 (EASE97) 1. Measurements of OH and HO_2 concentrations at Mace Head, Ireland, J. Geophys. Res.-Atmos., 107, 4091–4106, 2002.

Dari-Salisburgo, C., Di Carlo, P., Giammaria, F., Kajii, Y., and D'Altorio, A.: Laser induced fluorescence instrument for NO_2 measurements: Observations at a central Italy background site, Atmos. Environ., 43, 970–977, 2009.

Di Carlo, P., Aruffo, E., Busilacchio, M., Giammaria, F., Dari-Salisburgo, C., Biancofiore, F., Visconti, G., Lee, J., Moller, S., Reeves, C. E., Bauguitte, S., Forster, G., Jones, R. L., and Ouyang, B.: Aircraft based four-channel thermal dissociation laser induced fluorescence instrument for simultaneous measurements of NO_2, total peroxy nitrate, total alkyl nitrate, and HNO_3, Atmos. Meas. Tech., 6, 971–980, doi:10.5194/amt-6-971-2013, 2013.

Donahue, N. M., Kroll, J. H., Anderson, J. G., and Demerjian, K. L.: Direct observation of OH production from the ozonolysis of olefins, Geophys. Res. Lett., 25, 59–62, 1998.

Donahue, N. M., Drozd, G. T., Epstein, S. A., Presto, A. A., and Kroll, J. H.: Adventures in ozoneland: down the rabbit-hole, Phys. Chem. Chem. Phys., 13, 10848–10857, 2011.

Drozd, G. T. and Donahue, N. M.: Pressure Dependence of Stabilized Criegee Intermediate Formation from a Sequence of Alkenes, J. Phys. Chem. A, 115, 4381–4387, 2011.

Edwards, G. D., Cantrell, C., Stephens, S., Hill, B., Goyea, O., Shetter, R., Mauldin, R. L., Kosciuch, E., Tanner, D., and Eisele, F.: Chemical Ionization Mass Spectrometer Instrument for the Measurement of Tropospheric HO_2 and RO_2, Anal. Chem., 75, 5317–5327, 2003.

Emmerson, K. M. and Carslaw, N.: Night-time radical chemistry during the TORCH campaign, Atmos. Environ., 43, 3220–3226, 2009.

Emmerson, K. M. and Evans, M. J.: Comparison of tropospheric gas-phase chemistry schemes for use within global models, Atmos. Chem. Phys., 9, 1831–1845, doi:10.5194/acp-9-1831-2009, 2009.

Emmerson, K. M., Carslaw, N., and Pilling, M. J.: Urban Atmospheric Chemistry During the PUMA Campaign 2: Radical Budgets for OH, HO_2 and RO_2, J. Atmos. Chem., 52, 165–183, 2005.

Emmerson, K. M., Carslaw, N., Carslaw, D. C., Lee, J. D., McFiggans, G., Bloss, W. J., Gravestock, T., Heard, D. E., Hopkins, J., Ingham, T., Pilling, M. J., Smith, S. C., Jacob, M., and Monks, P. S.: Free radical modelling studies during the UK TORCH Campaign in Summer 2003, Atmos. Chem. Phys., 7, 167–181, doi:10.5194/acp-7-167-2007, 2007.

Faloona, I. C., Tan, D., Lesher, R. L., Hazen, N. L., Frame, C. L., Simpas, J. B., Harder, H., Martinez, M., di Carlo, P., Ren, X., and Brune, W. H.: A Laser-Induced Fluorescence Instrument for Detecting Troposhperic OH and HO_2: Characteristics and Calibration, J. Atmos. Chem., 47, 139–167, 2004.

Fleming, Z. L., Monks, P. S., Rickard, A. R., Heard, D. E., Bloss, W. J., Seakins, P. W., Still, T. J., Sommariva, R., Pilling, M. J., Morgan, R., Green, T. J., Brough, N., Mills, G. P., Penkett, S. A., Lewis, A. C., Lee, J. D., Saiz-Lopez, A., and Plane, J. M. C.: Peroxy radical chemistry and the control of ozone photochemistry at Mace Head, Ireland during the summer of 2002, Atmos. Chem. Phys., 6, 2193–2214, doi:10.5194/acp-6-2193-2006, 2006.

Floquet, C. F. A.: Airborne Measurements of Hydroxyl Radicals by Fluorescence Assay by Gas Expansion, PhD, School of Chemistry, University of Leeds, Leeds, 155 pp., 2006.

Fuchs, H., Bohn, B., Hofzumahaus, A., Holland, F., Lu, K. D., Nehr, S., Rohrer, F., and Wahner, A.: Detection of HO_2 by laser-induced fluorescence: calibration and interferences from RO_2 radicals, Atmos. Meas. Tech., 4, 1209–1225, doi:10.5194/amt-4-1209-2011, 2011.

Gerbig, C., Schmitgen, S., Kley, D., and Volz-Thomas, A.: An improved fast-response vacuum-UV resonance fluorescence CO instrument, J. Geophys. Res.-Atmos., 104, 1699–1704, 1999.

Geyer, A. and Stutz, J.: The vertical structure of OH-HO_2-RO_2 chemistry in the nocturnal boundary layer: A one-dimensional model study, J. Geophys. Res.-Atmos., 109, D16301–D16316, 2004a.

Geyer, A. and Stutz, J.: Vertical profiles of NO_3, N_2O_5, O_3, and NO_x in the nocturnal boundary layer: 2. Model studies on the altitude dependence of composition and chemistry, J. Geophys. Res.-Atmos., 109, D12307, doi:10.1029/2003JD004211, 2004b.

Geyer, A., Bächmann, K., Hofzumahaus, A., Holland, F., Konrad, S., Klüpfel, T., Pätz, H.-W., Perner, D., Mihelcic, D., Schäfer, H.-J., Volz-Thomas, A., and Platt, U.: Nighttime formation of peroxy and hydroxyl radicals during the BERLIOZ campaign: Observations and modeling studies, J. Geophys. Res.-Atmos., 108, 8249–8263, 2003.

Harrison, R. M., Yin, J., Tilling, R. M., Cai, X., Seakins, P. W., Hopkins, J. R., Lansley, D. L., Lewis, A. C., Hunter, M. C., Heard, D. E., Carpenter, L. J., Creasey, D. J., Lee, J. D., Pilling, M. J., Carslaw, N., Emmerson, K. M., Redington, A., Derwent, R. G., Ryall, D., Mills, G., and Penkett, S. A.: Measurement and modelling of air pollution and atmospheric chemistry in the U.K. West Midlands conurbation: Overview of the PUMA Consortium project, Sci. Total Environ., 360, 5–25, 2006.

Heard, D. E.: Atmospheric field measurements of the hydroxyl radical using laser-induced fluorescence spectroscopy, Annu. Rev. Phys. Chem., 57, 191–216, 2006.

Heard, D. E., Carpenter, L. J., Creasey, D. J., Hopkins, J. R., Lee, J. D., Lewis, A. C., Pilling, M. J., Seakins, P. W., Carslaw, N., and Emmerson, K. M.: High levels of the hydroxyl radical in the winter urban troposphere, Geophys. Res. Lett., 31, L18112–L18117, 10.1029/2004gl020544, 2004.

Hewitt, C. N., Lee, J. D., MacKenzie, A. R., Barkley, M. P., Carslaw, N., Carver, G. D., Chappell, N. A., Coe, H., Collier, C., Commane, R., Davies, F., Davison, B., DiCarlo, P., Di Marco, C. F., Dorsey, J. R., Edwards, P. M., Evans, M. J., Fowler, D., Furneaux, K. L., Gallagher, M., Guenther, A., Heard, D. E., Helfter, C., Hopkins, J., Ingham, T., Irwin, M., Jones, C., Karunaharan, A., Langford, B., Lewis, A. C., Lim, S. F., MacDonald, S. M., Mahajan, A. S., Malpass, S., McFiggans, G., Mills, G., Misztal, P., Moller, S., Monks, P. S., Nemitz, E., Nicolas-Perea, V., Oetjen, H., Oram, D. E., Palmer, P. I., Phillips, G. J., Pike, R., Plane, J. M. C., Pugh, T., Pyle, J. A., Reeves, C. E., Robinson, N. H., Stewart, D., Stone, D., Whalley, L. K., and Yin, X.: Overview: oxidant and particle photochemical processes above a south-east Asian tropical rainforest (the OP3 project): introduction, ratio-

nale, location characteristics and tools, Atmos. Chem. Phys., 10, 169–199, doi:10.5194/acp-10-169-2010, 2010.

Holland, F., Hofzumahaus, A., Schäfer, J., Kraus, A., and Pätz, H.-W.: Measurements of OH and HO_2 radical concentrations and photolysis frequencies during BERLIOZ, J. Geophys. Res.-Atmos., 108, 8246–8261, 2003.

Hopkins, J. R., Lewis, A. C., and Read, K. A.: A two-column method for long-term monitoring of non-methane hydrocarbons (NMHCs) and oxygenated volatile organic compounds (*o*-VOCs), J. Environ. Monitor., 5, 8–13, 2003.

Jenkin, M. E., Saunders, S. M., and Pilling, M. J.: The tropospheric degradation of volatile organic compounds: A protocol for mechanism development, Atmos. Environ., 31, 81–104, 1997.

Jenkin, M. E., Saunders, S. M., Wagner, V., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic compounds, Atmos. Chem. Phys., 3, 181–193, doi:10.5194/acp-3-181-2003, 2003.

Johnson, D. and Marston, G.: The gas-phase ozonolysis of unsaturated volatile organic compounds in the troposphere, Chem. Soc. Rev., 37, 699–716, 2008.

Jones, A. R., Thomson, D. J., Hort, M., and Devenish, B.: The U.K. Met Office's next-generation atmospheric dispersion model, NAME III, in: Air Pollution Modeling and its Application XVII (Proceedings of the 27th NATO/CCMS International Technical Meeting on Air Pollution Modelling and its Application), edited by: Borrego, C. and Norman, A.-L., Springer, Dordrecht, the Netherlands, 580–589, 2007.

Kanaya, Y., Sadanaga, Y., Matsumoto, J., Sharma, U. K., Hirokawa, J., Kajii, Y., and Akimoto, H.: Nighttime observation of the HO_2 radical by an LIF instrument at Oki Island, Japan, and its possible origins, Geophys. Res. Lett., 26, 2179–2182, 1999.

Kennedy, O. J., Ouyang, B., Langridge, J. M., Daniels, M. J. S., Bauguitte, S., Freshwater, R., McLeod, M. W., Ironmonger, C., Sendall, J., Norris, O., Nightingale, R., Ball, S. M., and Jones, R. L.: An aircraft based three channel broadband cavity enhanced absorption spectrometer for simultaneous measurements of NO_3, N_2O_5 and NO_2, Atmos. Meas. Tech., 4, 1759–1776, doi:10.5194/amt-4-1759-2011, 2011.

Kroll, J. H., Clarke, J. S., Donahue, N. M., Anderson, J. G., and Demerjian, K. L.: Mechanism of HO_x Formation in the Gas-Phase Ozone-Alkene reaction. 1. Direct, Pressure-Dependent Measurements of Prompt OH Yields, J. Phys. Chem. A, 105, 1554–1560, 2001a.

Kroll, J. H., Sahay, S. R., Anderson, J. G., Demerjian, K. L., and Donahue, N. M.: Mechanism of HO_x Formation in the Gas-Phase Ozone-Alkene Reaction. 2. Prompt versus Thermal Dissociation of Carbonyl Oxides to Form OH, J. Phys. Chem. A, 105, 4446–4457 2001b.

Kroll, J. H., Donahue, N. M., Cee, V. J., Demerjian, K. L., and Anderson, J. G.: Gas-Phase Ozonolysis of Alkenes: Formation of OH from Anti Carbonyl Oxides, J. Am. Chem. Soc., 124, 8518–8519, 2002.

Lee, J. D., Lewis, A. C., Monks, P. S., Jacob, M., Hamilton, J. F., Hopkins, J. R., Watson, N. M., Saxton, J. E., Ennis, C., Carpenter, L. J., Carslaw, N., Fleming, Z. L., Bandy, A., Oram, D. E., Penkett, S. A., Slemr, J., Norton, E. G., Rickard, A. R., Whalley, L. K., Heard, D. E., Bloss, W. J., Gravestock, T., Smith, S. C., Stanton, J., Pilling, M. J., and Jenkin, M. E.: Ozone photochemistry and elevated isoprene during the UK heatwave of August 2003, Atmos. Environ., 40, 7598–7613, 2006.

Lidster, R. T., Hamilton, J. F., Lee, J. D., Lewis, A. C., Hopkins, J. R., Punjabi, S., Rickard, A. R., and Young, J. C.: The impact of monoaromatic hydrocarbons on OH reactivity in the coastal UK boundary layer and free troposphere, Atmos. Chem. Phys., 14, 6677–6693, doi:10.5194/acp-14-6677-2014, 2014.

Lightfoot, P. D., Cox, R. A., Crowley, J. N., Destriau, M., Hayman, G. D., Jenkin, M. E., Moortgat, G. K., and Zabel, F.: Organic Peroxy Radicals: Kinetics, Spectroscopy and Tropospheric Chemistry, Atmos. Environ., 26A, 1805–1961, 1992.

Lou, S., Holland, F., Rohrer, F., Lu, K., Bohn, B., Brauers, T., Chang, C.C., Fuchs, H., Häseler, R., Kita, K., Kondo, Y., Li, X., Shao, M., Zeng, L., Wahner, A., Zhang, Y., Wang, W., and Hofzumahaus, A.: Atmospheric OH reactivities in the Pearl River Delta – China in summer 2006: measurement and model results, Atmos. Chem. Phys., 10, 11243–11260, doi:10.5194/acp-10-11243-2010, 2010.

Lu, K. D., Rohrer, F., Holland, F., Fuchs, H., Bohn, B., Brauers, T., Chang, C. C., Häseler, R., Hu, M., Kita, K., Kondo, Y., Li, X., Lou, S. R., Nehr, S., Shao, M., Zeng, L. M., Wahner, A., Zhang, Y. H., and Hofzumahaus, A.: Observation and modelling of OH and HO_2 concentrations in the Pearl River Delta 2006: a missing OH source in a VOC rich atmosphere, Atmos. Chem. Phys., 12, 1541–1569, doi:10.5194/acp-12-1541-2012, 2012.

Lu, K. D., Hofzumahaus, A., Holland, F., Bohn, B., Brauers, T., Fuchs, H., Hu, M., Häseler, R., Kita, K., Kondo, Y., Li, X., Lou, S. R., Oebel, A., Shao, M., Zeng, L. M., Wahner, A., Zhu, T., Zhang, Y. H., and Rohrer, F.: Missing OH source in a suburban environment near Beijing: observed and modelled OH and HO_2 concentrations in summer 2006, Atmos. Chem. Phys., 13, 1057–1080, doi:10.5194/acp-13-1057-2013, 2013.

Lu, K. D., Rohrer, F., Holland, F., Fuchs, H., Brauers, T., Oebel, A., Dlugi, R., Hu, M., Li, X., Lou, S. R., Shao, M., Zhu, T., Wahner, A., Zhang, Y. H., and Hofzumahaus, A.: Nighttime observation and chemistry of HO_x in the Pearl River Delta and Beijing in summer 2006, Atmos. Chem. Phys., 14, 4979–4999, doi:10.5194/acp-14-4979-2014, 2014.

Mauldin III, R. L., Berndt, T., Sipilä, M., Paasonen, P., Petäjä, T., Kim, S., Kurtén, A., Stratmann, F., Kerminen, V.-M., and Kulmala, M.: A new atmospherically relevent oxidant of sulphur dioxide, Nature, 488, 193–196, 2012.

Morgan, W. T., Ouyang, B., Allan, J. D., Aruffo, E., Di Carlo, P., Kennedy, O. J., Lowe, D., Flynn, M. J., Rosenberg, P. D., Williams, P. I., Jones, R., McFiggans, G. B., and Coe, H.: Influence of aerosol chemical composition on N_2O_5 uptake: airborne regional measurements in northwestern Europe, Atmos. Chem. Phys., 15, 973–990, doi:10.5194/acp-15-973-2015, 2015.

Nakajima, M. and Endo, Y.: Determination of the molecular structure of the simplest Criegee intermediate CH_2OO, J. Chem. Phys., 139, 101103, doi:10.1063/1.4821165, 2013.

Nakajima, M. and Endo, Y.: Spectroscopic characterization of an alkyl substituted Criegee intermediate *syn*-CH_3CHOO through pure rotational transitions, J. Chem. Phys., 140, 011101, doi:10.1063/1.4861494, 2014.

Ouyang, B., McLeod, M. W., Jones, R. L., and Bloss, W. J.: NO_3 radical production from the reaction between the Criegee intermediate CH_2OO and NO_2, Phys. Chem. Chem. Phys., 15, 17070–17075, 2013.

Paulson, S. E., and Orlando, J. J.: The reactions of ozone with alkenes: An important source of HO_x in the boundary layer, Geophys. Res. Lett., 23, 3727–3730, 1996.

Ren, X., Harder, H., Martinez, M., Lesher, R., Oliger, A., Simpas, J. B., Brune, W. H., Schwab, J. J., Demerjian, K. L., He, Y., Zhou, X., and Gao, H.: OH and HO_2 chemistry in the urban atmosphere of New York City, Atmos. Environ., 37, 3639–3651, 2003a.

Ren, X., Harder, H., Martinez, M., Lesher, R. L., Oliger, A., Shirley, T., Adams, J., Simpas, J. B., and Brune, W. H.: HO_x concentrations and OH reactivity observations in New York City during PMTACS-NY2001, Atmos. Environ., 37, 3627–3637, 2003b.

Ren, X., Brune, W. H., Mao, J., Mitchell, M. J., Lesher, R., Simpas, J. B., Metcalf, A. R., Schwab, J. J., Cai, C., Li, Y., Demerjian, K. L., Felton, H. D., Boynton, G., Adams, A., Perry, J., He, Y., Zhou, X., and Hou, J.: Behavior of OH and HO_2 in the winter atmosphere in New York City, Atmos. Environ., 40, 252–263, 2006.

Salisbury, G., Rickard, A. R., Monks, P. S., Allan, B. J., Bauguitte, S. J. B., Penkett, S. A., Carslaw, N., Lewis, A. C., Creasey, D. J., Heard, D. E., Jacobs, P. J., and Lee, J. D.: Production of peroxy radicals at night via reactions of ozone and the nitrate radical in the marine boundary layer, J. Geophys. Res.-Atmos., 106, 12669–12687, 2001.

Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, Atmos. Chem. Phys., 3, 161–180, doi:10.5194/acp-3-161-2003, 2003.

Sheehy, P. M., Volkamer, R., Molina, L. T., and Molina, M. J.: Oxidative capacity of the Mexico City atmosphere – Part 2: A RO_x radical cycling perspective, Atmos. Chem. Phys., 10, 6993–7008, doi:10.5194/acp-10-6993-2010, 2010.

Shirley, T. R., Brune, W. H., Ren, X., Mao, J., Lesher, R., Cardenas, B., Volkamer, R., Molina, L. T., Molina, M. J., Lamb, B., Velasco, E., Jobson, T., and Alexander, M.: Atmospheric oxidation in the Mexico City Metropolitan Area (MCMA) during April 2003, Atmos. Chem. Phys., 6, 2753–2765, doi:10.5194/acp-6-2753-2006, 2006.

Smith, S. C., Lee, J. D., Bloss, W. J., Johnson, G. P., Ingham, T., and Heard, D. E.: Concentrations of OH and HO_2 radicals during NAMBLEX: measurements and steady state analysis, Atmos. Chem. Phys., 6, 1435–1453, doi:10.5194/acp-6-1435-2006, 2006.

Sommariva, R., Pilling, M. J., Bloss, W. J., Heard, D. E., Lee, J. D., Fleming, Z. L., Monks, P. S., Plane, J. M. C., Saiz-Lopez, A., Ball, S. M., Bitter, M., Jones, R. L., Brough, N., Penkett, S. A., Hopkins, J. R., Lewis, A. C., and Read, K. A.: Night-time radical chemistry during the NAMBLEX campaign, Atmos. Chem. Phys., 7, 587–598, doi:10.5194/acp-7-587-2007, 2007.

Sommariva, R., Osthoff, H. D., Brown, S. S., Bates, T. S., Baynard, T., Coffman, D., de Gouw, J. A., Goldan, P. D., Kuster, W. C., Lerner, B. M., Stark, H., Warneke, C., Williams, E. J., Fehsenfeld, F. C., Ravishankara, A. R., and Trainer, M.: Radicals in the marine boundary layer during NEAQS 2004: a model study of day-time and night-time sources and sinks, Atmos. Chem. Phys., 9, 3075–3093, doi:10.5194/acp-9-3075-2009, 2009.

Stewart, D. J., Taylor, C. M., Reeves, C. E., and McQuaid, J. B.: Biogenic nitrogen oxide emissions from soils: impact on NO_x and ozone over west Africa during AMMA (African Monsoon Multidisciplinary Analysis): observational study, Atmos. Chem. Phys., 8, 2285–2297, doi:10.5194/acp-8-2285-2008, 2008.

Still, T. J., Al-Haider, S., Seakins, P. W., Sommariva, R., Stanton, J. C., Mills, G., and Penkett, S. A.: Ambient formaldehyde measurements made at a remote marine boundary layer site during the NAMBLEX campaign – a comparison of data from chromatographic and modified Hantzsch techniques, Atmos. Chem. Phys., 6, 2711–2726, doi:10.5194/acp-6-2711-2006, 2006.

Stone, D., Evans, M. J., Commane, R., Ingham, T., Floquet, C. F. A., McQuaid, J. B., Brookes, D. M., Monks, P. S., Purvis, R., Hamilton, J. F., Hopkins, J., Lee, J., Lewis, A. C., Stewart, D., Murphy, J. G., Mills, G., Oram, D., Reeves, C. E., and Heard, D. E.: HO_x observations over West Africa during AMMA: impact of isoprene and NO_x, Atmos. Chem. Phys., 10, 9415–9429, doi:10.5194/acp-10-9415-2010, 2010.

Stone, D., Whalley, L. K., and Heard, D. E.: Tropospheric OH and HO_2 radicals: field measurements and model comparisons, Chem. Soc. Rev., 41, 6348–6404, 2012.

Stone, D., Blitz, M., Daubney, L., Howes, N. U. M., and Seakins, P.: Kinetics of CH_2OO reactions with SO_2, NO_2, NO, H_2O and CH_3CHO as a function of pressure, Phys. Chem. Chem. Phys., 16, 1139–1149, 2014a.

Stone, D., Evans, M. J., Walker, H., Ingham, T., Vaughan, S., Ouyang, B., Kennedy, O. J., McLeod, M. W., Jones, R. L., Hopkins, J., Punjabi, S., Lidster, R., Hamilton, J. F., Lee, J. D., Lewis, A. C., Carpenter, L. J., Forster, G., Oram, D. E., Reeves, C. E., Bauguitte, S., Morgan, W., Coe, H., Aruffo, E., Dari-Salisburgo, C., Giammaria, F., Di Carlo, P., and Heard, D. E.: Radical chemistry at night: comparisons between observed and modelled HO_x, NO_3 and N_2O_5 during the RONOCO project, Atmos. Chem. Phys., 14, 1299–1321, doi:10.5194/acp-14-1299-2014, 2014b.

Stutz, J., Alicke, B., Ackermann, R., Geyer, A., White, A., and Williams, E.: Vertical profiles of NO_3, N_2O_5, O_3, and NO_x in the nocturnal boundary layer: 1. Observations during the Texas Air Quality Study 2000, J. Geophys. Res.-Atmos., 109, D12306–D12320, 2004.

Su, Y.-T., Huang, Y.-H., Witek, H. A., and Lee, Y.-P.: Infrared Absorption Spectrum of the Simplest Criegee Intermediate CH_2OO, Science, 340, 174–176, 2013.

Taatjes, C. A., Meloni, G., Selby, T. M., Trevitt, A. J., Osborn, D. L., Percival, C. J., and Shallcross, D. E.: Direct Observation of the Gas-Phase Criegee Intermediate (CH_2OO), J. Am. Chem. Soc., 130, 11883–11885, 2008.

Taatjes, C. A., Welz, O., Eskola, A. J., Savee, J. D., Osborn, D. L., Lee, E. P. F., Dyke, J. M., Mok, D. W. K., Shallcross, D. E., and Percival, C. J.: Direct measurement of Criegee intermediate (CH_2OO) reactions with acetone, acetaldehyde, and hexafluoroacetone, Phys. Chem. Chem. Phys., 14, 10391–10400, 2012.

Taatjes, C. A., Welz, O., Eskola, A. J., Savee, J. D., Scheer, A. M., Shallcross, D. E., Rotavera, B., Lee, E. P. F., Dyke, J. M., Mok, D. W. K., Osborn, D. L., and Percival, C. J.: Direct Measurements of Conformer-Dependent Reactivity of the Criegee Intermediate CH_3CHOO, Science, 340, 177–180, 2013.

Taatjes, C. A., Shallcross, D. E., and Percival, C. J.: Research frontiers in the chemistry of Criegee intermediates and tropospheric ozonolysis, Phys. Chem. Chem. Phys., 16, 1689–2180, 2014.

Vereecken, L. and Francisco, J. S.: Theoretical studies of atmospheric reaction mechanisms in the troposphere, Chem. Soc. Rev., 41, 6217–6708, 2012.

Volkamer, R., Sheehy, P., Molina, L. T., and Molina, M. J.: Oxidative capacity of the Mexico City atmosphere – Part 1: A radical source perspective, Atmos. Chem. Phys., 10, 6969–6991, doi:10.5194/acp-10-6969-2010, 2010.

Welz, O., Savee, J. D., Osborn, D. L., Vasu, S. S., Percival, C. J., Shallcross, D. E., and Taatjes, C. A.: Direct Kinetic Measurements of Criegee Intermediate (CH_2OO) Formed by Reaction of CH_2I with O_2, Science, 335, 204–207, 2012.

Whalley, L. K., Furneaux, K. L., Goddard, A., Lee, J. D., Mahajan, A., Oetjen, H., Read, K. A., Kaaden, N., Carpenter, L. J., Lewis, A. C., Plane, J. M. C., Saltzman, E. S., Wiedensohler, A., and Heard, D. E.: The chemistry of OH and HO_2 radicals in the boundary layer over the tropical Atlantic Ocean, Atmos. Chem. Phys., 10, 1555–1576, doi:10.5194/acp-10-1555-2010, 2010.

Whalley, L. K., Blitz, M. A., Desservettaz, M., Seakins, P. W., and Heard, D. E.: Reporting the sensitivity of laser-induced fluorescence instruments used for HO_2 detection to an interference from RO_2 radicals and introducing a novel approach that enables HO_2 and certain RO_2 types to be selectively measured, Atmos. Meas. Tech., 6, 3425–3440, doi:10.5194/amt-6-3425-2013, 2013.

Wong, K. W. and Stutz, J.: Influence of nocturnal vertical stability on daytime chemistry: A one-dimensional model study, Atmos. Environ., 44, 3753–3760, 2010.