



Deposited via The University of Sheffield.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/id/eprint/89716/>

Version: Accepted Version

---

**Article:**

Bauso, D. and Pesenti, R. (2012) Team Theory and Person-by-Person Optimization with Binary Decisions. *SIAM Journal on Control and Optimization (SICON)*, 50 (5). 3011 - 3028. ISSN: 0363-0129

<https://doi.org/10.1137/090769533>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# TEAM THEORY AND PERSON-BY-PERSON OPTIMIZATION WITH BINARY DECISIONS\*

D. BAUSO<sup>†</sup> AND R. PESENTI<sup>‡</sup>

**Abstract.** In this paper, we extend the notion of person by person optimization to binary decision spaces. The novelty of our approach is the adaptation to a dynamic team context of notions borrowed from the pseudo-boolean optimization field as completely local-global or unimodal functions and sub-modularity. We also generalize the concept of pbp optimization to the case where the Decision Makers (DMs) make decisions sequentially in groups of  $m$ , we call it *mbm optimization*. The main contribution are certain sufficient conditions, verifiable in polynomial time, under which a pbp or an *mbm* optimization algorithm leads to the team-optimum. As a second contribution, we present a *local* and *greedy* algorithm that allows the DMs to select a small neighborhood which guarantees them to behave as if they had complete information. As a last contribution, we also show that there exists a subclass of sub-modular team problems, recognizable in polynomial time, for which the convergence is guaranteed if the pbp algorithm is opportunely initialized.

**Key words.** team theory, person-by-person optimality, approximation algorithms

**1. Introduction.** Most fundamental results in team theory concern linear quadratic Gaussian problems or, in general, problems with continuous decision spaces, where the cost is somehow convex in the strategies and the information structure is a “nice” one (see, e.g., partial nested structures) [12, 16]. In such particular cases, it is well known that a simple solution idea consisting in a sequential optimization on the part of the Decision Makers (DMs), called *person by person optimization* (pbp), leads to the team-optimum [12], namely the argument minimizing the team objective function.

In this paper, on the same line of [8, 9], we restrict our attention to boolean decision spaces. The novelty of our approach is the adaptation to a dynamic team context of notions borrowed from *pseudo-boolean optimization* [5], as Completely Local-Global (CLG) functions, Completely Unimodal (CU) functions (also known as acyclic unique sink orientations and abstract objective functions [15]) and sub-modular functions [6, 11].

Boolean decision spaces can be found in finite-alphabet control and in particular on-off control problems [2, 10], impulsively-controlled systems (activate the impulse or not) [7], or switching control (switches between active and passive modes) [17]. Boolean decisions are encountered in many applications as inventory with set up costs (reordering or not from a warehouse in order to meet a demand) [3, 4], distributed computer systems (processing or not the assigned task) [9], in air-conditioning systems control, in economics and finance (see, e.g., [5] and references therein).

As first contribution, we generalize the concept of pbp optimization to the case where the Decision Makers (DMs) make decisions sequentially in groups of  $m$ , we call it *mbm optimization*.

---

\*A conference version [1] of this paper has been published in the Proc. of the IEEE American Control Conference, Seattle, Washington, USA, June 2008. Corresponding author D. Bauso Tel. +39-320-4648142. Research supported by MURST-PRIN 2007ZMVK5T “Decisional model for the design and the management of logistics networks characterized by high interoperability and information integration”.

<sup>†</sup>D. Bauso is with DINFO, Università di Palermo, Italy [dario.bauso@unipa.it](mailto:dario.bauso@unipa.it)

<sup>‡</sup>R. Pesenti is with DMA, Università “Ca’ Foscari” di Venezia, Italy [pesenti@unive.it](mailto:pesenti@unive.it)

The main contribution of this paper consists in providing certain sufficient conditions, verifiable in polynomial time, for the optimality of such pbp (respectively *mbm*) optimization algorithms based on the knowledge of the agents' states. Then we can frame our results in the literature on person by person algorithms in team theory, which has drawn the attention of the control audience since the '70s (see, e.g., [12]).

A second contribution, which makes this paper to differ from the conference version [1], is the *local* and *greedy* algorithm of Section 5. This algorithm allows the DMs to select a small neighborhood which guarantees them to behave as if they had complete information, i.e., they knew all the other agents' states. "Small" neighborhood means that each DM is not required to communicate with all the other DMs but only with a restricted, possibly the minimal, number of them. "Local" means that the algorithm is implemented by the same DM who has to make a decision without any centralized mechanism. "Greedy" means that the DM who has to make a decision implements an iterative algorithm that at each iteration picks the smallest number of DMs whose state knowledge may be as informative as the knowledge of all the other agents' states.

As a last contribution, we have paid special attention to problems with sub-modular team objective function (*sub-modular team problems*). Though sub-modularity alone does not guarantee the convergence of any pbp optimization algorithm, we show that there exists a special class of sub-modular team problems, recognizable in polynomial time, for which the convergence is guaranteed when the algorithm is opportunely initialized. This class is characterized by so-called *threshold strategies*.

This paper is organized as follows. In Section 2, we introduce some notions from team theory [12] and pseudo-boolean optimization [5]. In Section 3, we introduce the class of completely local-global functions and completely unimodal functions [6], and [11]. In Section 4, we address the *mbm* optimization. In Section 5, we present the greedy algorithm. In Section 6, we focus on sub-modular team problems. In Section 7 we provide numerical examples. Finally, in Section 8, we discuss how to extend the obtained results.

**2. Definitions and Problem Statement.** Consider a set  $N$  of  $n$  DMs making decisions  $x$  from a discrete hypercube  $\mathbb{B}^n = \{0, 1\}^n$ . Decisions are made in order to optimize a common team objective function,  $J(x) : \mathbb{B}^n \mapsto \mathbb{Z}$ , where  $\mathbb{Z}$  is the set of integer numbers.

ASSUMPTION 2.1. *The team objective function  $J(x)$  is injective and has the following quadratic form*

$$(2.1) \quad J(x) = \sum_{i=1}^n b_i x_i + \sum_{i \in N} \sum_{j \in N} a_{ij} x_i x_j$$

with  $a_{ij}$  and  $b_i$  integer (this causes  $J(x)$  assuming only integer values).

The following definitions are slightly modified from [9].

DEFINITION 2.1. (Team-optimum) *A point  $x^*$  is a team-optimum if*

$$x^* = \arg \min_{x \in \mathbb{B}^n} J(x).$$

As the set  $\mathbb{B}^n$  is finite, a team optimum  $x^*$  always exists. Furthermore, as  $J(x)$  is injective, the team optimum is unique.

DEFINITION 2.2. (pbp optimum) *The point  $x^*$  is a pbp optimum if for any DM  $i$  the following condition holds*

$$(2.2) \quad J(x_i^*, x_{-i}^*) < J(x_i, x_{-i}^*), \quad \forall x_i \neq x_i^*$$

where  $x_i \in \mathbb{B}$  is the decision of DM  $i$  and  $x_{-i} = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)^T \in \mathbb{B}^{n-1}$  is a vector collecting decisions of all other DMs. From the above definitions we have that a team-optimum always implies pbp optimality but not vice versa.

Let  $S$  any subset of  $N$  with  $m$  elements. We indicate this with  $S \subseteq N$  with  $|S| = m$ , where  $|S|$  means cardinality of  $S$ . Let  $x_S \in \mathbb{B}^m$  be a vector collecting the decisions of all the DMs belonging to  $S$ , namely,  $x_S = (x_i : i \in S)$ . Analogously, let  $x_{-S} \in \mathbb{B}^{n-m}$  be a vector collecting the decisions of all the other DMs,  $x_{-S} = (x_i : i \in N \setminus S)$ .

DEFINITION 2.3. (*mbm optimum*) *The point  $x^*$  is an mbm optimum if, for any subset  $S \subseteq N$  with  $|S| = m$ , the following condition holds*

$$(2.3) \quad J(x_S^*, x_{-S}^*) < J(x_S, x_{-S}), \quad \forall x_S \neq x_S^*.$$

All the results stated in the following hold true for any value of the parameter  $m$  from 1 to  $n$ .

For each subset  $S \subseteq N$ , we isolate from the team objective function (2.1) the only terms in  $x_i$  with  $i \in S$  as follows

$$J_S(x) = \sum_{i \in S} b_i x_i + \sum_{i \in S} \sum_{j \in N} a_{ij} x_i x_j$$

and denote this last function as the  $S$ -projection of  $J(x)$ .

We observe that for any  $S \subseteq N$ ,

$$\arg \min_{\tilde{x}_S \in \{0,1\}} J(\tilde{x}_S, x_{-S}) = \arg \min_{\tilde{x}_S \in \{0,1\}} J_S(\tilde{x}_S, x_{-S}).$$

Moreover, assume that DMs in  $S$  know the decisions of the only DMs in a *neighborhood*  $\Gamma_S$ , with  $\Gamma_S \subseteq N \setminus S$ .

Then, for each  $\Gamma_S \subseteq N \setminus S$  we can also define as  $\Gamma_S$ -approximation of  $J_S(x)$  the following function

$$(2.4) \quad \hat{J}_{S, \Gamma_S}(x) = \sum_{i \in S} b_i x_i + \sum_{i \in S} \sum_{j \in S \cup \Gamma_S} a_{ij} x_i x_j + \sum_{i \in S} \sum_{i \in N \setminus (S \cup \Gamma_S)} a_{ij} x_i \hat{x}_j$$

where

$$\hat{x}_j = \begin{cases} 1 & \text{if } a_{ij} > 0 \\ 0 & \text{otherwise} \end{cases}.$$

The above definition implies that  $\hat{J}_{S, \Gamma_S}(x)$  approximates from above  $J_S(x)$ , i.e.,  $\hat{J}_{S, \Gamma_S}(x) \geq J_S(x)$  for all  $x \in \mathbb{B}^n$  and all  $\Gamma_S \subseteq N \setminus S$ .

Hereafter, for the sake of notation, we use the notation  $J_i(x)$ ,  $\hat{J}_i(x)$ , and  $\Gamma_i$ , and  $\Gamma_{\{i\}}$ , in state of  $J_{\{i\}}(x)$ ,  $\hat{J}_{\{i\}}(x)$ , and  $\Gamma_{\{i\}}$  respectively.

We are ready to generalize the concept of pbp strategy, introduced in [9] and [12], as follows.

DEFINITION 2.4. *A strategy  $\mu_i : \mathbb{B}^{n-1} \mapsto \mathbb{B}$  is pbp strict for DM  $i$  if, for any  $x_{-i} \in \mathbb{B}^{n-1}$ , we have*

$$\mu_i(x_{-i}) = \arg \min_{\tilde{x}_i \in \{0,1\}} J(\tilde{x}_i, x_{-i}) = \arg \min_{\tilde{x}_i \in \{0,1\}} J_i(\tilde{x}_i, x_{-i}).$$

A strategy  $\hat{\mu}_i : \mathbb{B}^{n-1} \mapsto \mathbb{B}$  is a  $\Gamma_i$ -approximation, for some  $\Gamma_i \subseteq N \setminus \{i\}$ , of a pbp strict strategy  $\mu_i(x_{-i})$  if, for any  $x_{-i} \in \mathbb{B}^{n-1}$ , we have

$$\hat{\mu}_i(x_{-i}) = \arg \min_{\tilde{x}_i \in \{0,1\}} \hat{J}_{i,\Gamma_i}(\tilde{x}_i, x_{-i}).$$

As  $J(x)$  is injective, the above equations have a unique solution. Then, under a strict pbp strategy  $\mu_i(\cdot)$ , DM  $i$  changes decision from zero to one or vice versa only if such a change lets the  $\{i\}$ -projection  $J_i(\cdot, \cdot)$ , and the team objective function  $J(\cdot, \cdot)$  as well, decrease for fixed decisions of all other DMs  $j \neq i$ .

We can repeat the same argument for the  $\Gamma_i$ -approximate strict pbp strategy  $\hat{\mu}_i(\cdot)$  with respect to the  $\Gamma_i$ -approximation  $\hat{J}_{i,\Gamma_i}(\cdot, \cdot)$ .

DEFINITION 2.5. A strategy  $\mu_S : \mathbb{B}^{n-m} \mapsto \mathbb{B}^m$  is mbm strict for DMs in  $S$  where  $S \subseteq N$  with cardinality  $|S| = m$  if, for any  $x_{-S} \in \mathbb{B}^{n-m}$ , we have

$$\mu_S(x_{-S}) = \arg \min_{\tilde{x}_S \in \mathbb{B}^m} J(\tilde{x}_S, x_{-S}) = \arg \min_{\tilde{x}_S \in \mathbb{B}^m} J_S(\tilde{x}_S, x_{-S}).$$

A strategy  $\hat{\mu}_S : \mathbb{B}^{n-m} \mapsto \mathbb{B}^m$  is a  $\Gamma_S$ -approximation, for some  $\Gamma_S \subseteq N \setminus S$ , of a mbm strict strategy  $\mu_S(x_{-S})$  if, for any  $x_{-S} \in \mathbb{B}^{n-m}$ , we have

$$\hat{\mu}_S(x_{-S}) = \arg \min_{\tilde{x}_S \in \{0,1\}} \hat{J}_{S,\Gamma_S}(\tilde{x}_S, x_{-S}).$$

The above definition of strict mbm strategy has the following geometric interpretation. For any  $x \in \mathbb{B}^n$  and  $S \subseteq N$ , denote by  $\Pi_S(x)$  as the the corresponding  $m$ -dimensional face  $\{\tilde{x} = (\tilde{x}_S, x_{-S}) \in \mathbb{B}^n : x_{-S} \text{ fixed}\}$  of hypercube  $\mathbb{B}^n$ . Then, a strict mbm strategy means that either  $(x_S, x_{-S})$  is the optimal vertex in  $\Pi_S(x)$  or the DMs in  $S$  coordinate their decisions to find an optimal vertex in  $\Pi_S(x)$ .

With the above definitions in mind, we call *pbp optimization algorithm*, any algorithm that returns a sequence of decisions  $x(0) \rightarrow x(1) \rightarrow \dots$  where, for each iteration  $t$ , we denote by  $x(t) = \{x_1(t) \dots x_n(t)\}$  and  $x_i(t)$  the vector of decisions and the decision of DM  $i$  respectively. We also require that each decision  $x(t)$  is obtained from  $x(t-1)$  by a unilateral improvement on the part of a single DM  $i = \sigma(t)$ , i.e.,  $x(t) = [\mu_i(x_{-i}(t-1)), x_{-i}(t-1)]$ , where  $\sigma : \mathbb{N} \mapsto N$ , is a periodic surjective function, with period  $n$ , that returns a DM for each iteration  $t$ . For instance,  $\sigma(1) = 2, \sigma(2) = 5 \dots$  means that at iteration 1, DM 2 plays the strict pbp strategy for fixed decisions of all other DMs, and similarly for DM 5 at iteration 2. We define an *mbm optimization algorithm* in a similar manner. Here, the function  $\sigma$  becomes  $\sigma : \mathbb{N} \mapsto \mathcal{Q}$ , with period  $|\mathcal{Q}|$ , where  $\mathcal{Q}$  is the set of all subsets  $S \subseteq N$  with  $|S| = m$ , and the vector of decisions at iteration  $t$  becomes  $x(t) = [\mu_S(x_{-S}(t-1)), x_{-S}(t-1)]$ . We define an algorithm approximate when it uses approximate strategies  $\hat{\mu}_i(\cdot)$  or  $\hat{\mu}_S(\cdot)$ .

We can now state the problem of interest.

PROBLEM 1. Find conditions under which any pbp (respectively mbm) optimization algorithm converges to the team-optimum. Furthermore, design local information mechanisms under which approximate algorithms return the same decisions as in the complete information case.

Throughout the paper, convergence means “from any generic  $x(0)$ ”, unless specified differently.

REMARK 2.1. Any strict pbp (respectively mbm) optimization algorithm converges to a pbp (mbm) optimum  $x_{pbp}^*$  (respectively  $x_{mbm}^*$ ) in a finite number of iterations. Actually, the set  $\mathbb{B}^n$  is finite and at each iteration  $t$  of the algorithm the value of objective function  $J(x(t))$  decreases.

There is a vast literature on functions  $f(x) : \mathbb{B}^n \mapsto \mathbb{Z}$  that map from a discrete hypercube  $\mathbb{B}^n$  to the ordered field  $\mathbb{Z}$  of integer numbers. They are usually referred to as *pseudo-boolean functions* [5].

In the following, we recall some notions and optimality conditions in the context of pseudo-boolean optimization that we use to prepare and motivate the results of the next sections.

Let us now associate to a binary vector  $x \in \mathbb{B}^n$  its neighborhood  $N_r(x)$  of *radius*  $r$ , defined as  $N_r(x) = \{y : \rho_H(x, y) \leq r\}$ , where  $\rho_H(x, y)$  denotes the *Hamming distance* of the vectors  $x$  and  $y$ , defined as the number of components in which these two vectors differ. According to this definition, the neighborhood of radius  $n$  of each  $x \in \mathbb{B}^n$  is equal to  $\mathbb{B}^n$ , that is  $N_n(x) = \mathbb{B}^n$ .

A vector  $x$  is a *local minimum* of a pseudo-boolean  $f(\cdot)$  if  $f(y) \geq f(x)$  for all neighboring vectors  $y \in N_1(x)$ . It is a *global minimum* if  $f(y) \geq f(x)$  for all vectors  $y \in \mathbb{B}^n$ .

Local minima can be determined by means of local search algorithms. In particular, [6] defines as a *single switch algorithm* any algorithm that at each iteration proceeds to a better neighbor of the current iterate, by changing one coordinate at a time, until a local optimum is found. Similarly, they define as a *multiple switch algorithm* of order  $m$  any algorithm that at each iteration proceeds to a next better iterate that differs from the current vertex in at most  $m$  coordinates.

REMARK 2.2. *The following statements hold true:*

- i) *The team objective function  $J(x)$  is a pseudo-boolean function.*
- ii) *Any pbp (respectively mbm) optimum is a local optimum in a neighborhood of radius one (respectively  $m$ ).*
- iii) *The team-optimum is a global optimum.*
- iv) *Strict pbp (respectively mbm) strategies are single (respectively multiple) switch algorithms.*

There is a large variety of techniques applied in the literature for solving problems that can be modelled by quadratic pseudo-Boolean functions optimization. As this last problem is NP-hard, many of the published algorithms are implicitly enumerative. However, specialized optimization algorithms have been developed for increasing or decreasing pseudo-Boolean functions.

We can associate to a pseudo-boolean function its first order  $i$ th *derivative*

$$\frac{\partial f}{\partial x_i}(x) = f(x_1, \dots, x_{i-1}, 1, x_{i+1}, \dots, x_n) - f(x_1, \dots, x_{i-1}, 0, x_{i+1}, \dots, x_n),$$

which will be used later on. If  $f(\cdot)$  is injective,  $\frac{\partial f}{\partial x_i}(x) \neq 0$  for all  $x \in \mathbb{B}^n$ , for all  $i \in N$ . Let us finally introduce the following operation.

DEFINITION 2.6. *Given a function  $f : \mathbb{B}^n \mapsto \mathbb{R}$ , for any subset  $S \subseteq N$ , define restriction of  $f$  into  $S$ ,  $\mathcal{R}_S f(x) : \mathbb{B}^n \mapsto \mathbb{R}$  the function*

$$\mathcal{R}_S f(x) = \sum_{i \in S} b_i + \sum_{i \in S} \sum_{j \in S} a_{ij} + \sum_{k \notin S} \sum_{i \in S} a_{ik} x_k.$$

The above definition has the following geometric interpretation. Consider the face  $\Pi_S(x) : \{x = (x_S, x_{-S}) \in \mathbb{B}^n : x_{-S} \text{ fixed}\}$  of  $\mathbb{B}^n$  and extract two points  $\bar{x} = (\mathbf{1}, x_{-S})$  and  $\underline{x} = (\mathbf{0}, x_{-S})$  from it. Note that, for fixed  $x_{-S}$ , in  $\bar{x}$  all DMs  $i \in S$  set  $x_i = 1$  while in  $\underline{x}$  all DMs  $i \in S$  set  $x_i = 0$ . Then, the restriction is the difference  $J(\bar{x}) - J(\underline{x})$  of the team objective function computed on the two points. Also, note that for a singleton,  $S = \{i\}$ , then  $\mathcal{R}_S f(x) = \frac{\partial f}{\partial x_i}(x)$ .

**3. Person by Person Optimization.** In this section, we present sufficient conditions, verifiable in polynomial time, for the convergence of any pbp algorithm to the team-optimum.

DEFINITION 3.1. (*CLG-functions [11]*) An injective function  $f : \mathbb{B}^n \mapsto \mathbb{Z}$  is Completely Local-Global (CLG) if in  $\mathbb{B}^n$  there is a unique local minimum.

LEMMA 3.1. Any pbp optimization algorithm guarantees convergence to the team-optimum  $x^*$  if and only if  $J(x)$  is a CLG-function.

*Proof.* (sufficiency) If  $J(\cdot)$  is a CLG-function then there is a unique pbp optimum which is also team-optimum. Any pbp optimization algorithm guarantees convergence to it.

(necessity) If  $J(\cdot)$  is not a CLG-function then there is a second pbp optimum  $\bar{x}$  which is not team-optimum. Any pbp optimization algorithm starting at  $\bar{x}$  cannot deviate from it and therefore does not reach the global optimum.  $\square$

The class of CLG-functions includes the class of completely unimodal functions.

DEFINITION 3.2. (*CU-functions*) An injective function  $f : \mathbb{B}^n \mapsto \mathbb{Z}$  is Completely Unimodal (CU) if  $f$  has a unique local minimum on every face of  $\mathbb{B}^n$ .

We can derive the following corollary from the above lemma.

COROLLARY 3.1. Any pbp optimization algorithm converges to the team-optimum  $x^*$  if  $J(x)$  is a CU-function.

To the best of author's knowledge, recognizing CU-functions or CLG-functions is, in general, a difficult task. Actually, it involves an exponential number of conditions as shown next. Furthermore, even if  $f$  is a CLG or CU-function, strict pbp strategies may converge in exponential time.

To see why complete unimodality involves an exponential number of conditions consider that for the existence of two local minima on a 2-face containing  $x_i$  and  $x_j$ , it must hold

$$(3.1) \quad \left. \frac{\partial f(x)}{\partial x_i} \right|_{x_j=0} \cdot \left. \frac{\partial f(x)}{\partial x_i} \right|_{x_j=1} < 0$$

$$(3.2) \quad \left. \frac{\partial f(x)}{\partial x_j} \right|_{x_i=0} \cdot \left. \frac{\partial f(x)}{\partial x_j} \right|_{x_i=1} < 0.$$

Then for  $f$  to be CU it is necessary that, on each 2-face and for all  $x$ , the above conditions are not satisfied, which implies an exponential number of verifications.

EXAMPLE 3.1. Consider the set  $\mathbb{B}^3 = \{0, 1\}^3$  and the team objective function  $J(x) : \mathbb{B}^3 \mapsto \mathbb{Z}$ , taking on the values displayed in Fig. 3.1.a. The explicit expression of the function  $J$  according to the formula (2.1) is

$$J(x) = \underbrace{4x_1^2 + 4x_2^2 - 8x_1x_2 + 2x_2}_{\mathcal{J}(x_1, x_2)} - 10x_3 - 10x_1x_3 + 3x_2x_3,$$

where we denote by  $\mathcal{J}(x_1, x_2)$  the function obtained considering terms only in  $x_1$  and  $x_2$ . In Fig. 3.1.a, the oriented arcs indicate the decreasing directions for the team objective function  $J(x)$ . Function  $J(x)$  is a CLG-function as it has a unique local (global) minimum in  $\mathbb{B}^3$  which is  $x = (1, 0, 1)$  (point C in the figure). However note that  $\mathcal{J}(x_1, x_2)$  is not a CLG-function as it has two local minima in  $\mathbb{B}^2$ . For instance, see the 2-face  $x_1$ - $x_2$  with  $x_3 = 0$  which has two local minima in  $x = (0, 0, 0)$  and  $x = (1, 1, 0)$  (point A and B). We complete the example by considering a different




FIG. 3.1. Unit 3-dimensional cubes: oriented arcs indicate decreasing directions for  $J(x)$  when (a)  $J(x)$  is CLG-function or (b)  $J(x)$  is CU-function. Solutions  $x = (0, 0, 0)$  and  $x = (1, 1, 0)$  (point A and B in (a)) are two local minima for the 2-face  $x_1$ - $x_2$  with  $x_3 = 0$ . In both cases, the global minimum is  $x = (1, 0, 1)$  (point C).

function  $\hat{J}(x) : \mathbb{B}^3 \mapsto \mathbb{Z}$ , taking on the values displayed in Fig. 3.1.b. The explicit expression is

$$\hat{J}(x) = \underbrace{x_1^2 + 4x_2^2 - 5x_1x_2 + 2x_2}_{\hat{J}(x_1, x_2)} - 10x_3 - 10x_1x_3 + 3x_2x_3,$$

where again  $\hat{J}(x_1, x_2)$  is obtained considering terms only in  $x_1$  and  $x_2$ . In Fig. 3.1.b, the unique global minimum is again  $x = (1, 0, 1)$  (point C in the figure) but differently from before function  $J(x)$  is a CU-function in  $\mathbb{B}^3$  as it has a unique local minimum on each 2-face. In correspondence to such a situation we also have that  $\hat{J}(x_1, x_2)$  is a CLG-function on  $\mathbb{B}^2$  as it has a unique local minimum in  $\mathbb{B}^2$  (see the 2-face  $x_1$ - $x_2$  with  $x_3 = 0$  which has a local minimum in  $x = (0, 0, 0)$  (point A)).

A special case of completely unimodality is when  $f(\cdot)$  is monotonic along any single direction, which corresponds to being both left hand side of (3.1) and (3.2) positive. Now,  $f(\cdot)$  is monotonic along any single direction, when for all  $i = 1 \dots, n$ , one of the following mutually exclusive conditions holds true

$$(3.3) \quad \max_{x \in \mathbb{B}^n} \frac{\partial J(x)}{\partial x_i} < 0$$

$$(3.4) \quad \min_{x \in \mathbb{B}^n} \frac{\partial J(x)}{\partial x_i} > 0.$$

We can specialize Corollary 3.1 to such a particular case.

LEMMA 3.2. (Sufficient conditions) If the team objective function  $J(x)$  is such that, for all  $i \in N$ , either (3.3) or (3.4) hold, then

1. the team optimum is

$$x_i^* = \begin{cases} 1 & \text{if } \max_{x \in \mathbb{B}^n} \frac{\partial J(x)}{\partial x_i} < 0 \\ 0 & \text{if } \min_{x \in \mathbb{B}^n} \frac{\partial J(x)}{\partial x_i} > 0 \end{cases}$$

2. the team optimum  $x^*$  is also the unique pbp optimum,

3. any pbp optimization algorithm converges to the team optimum  $x^*$  in at most  $n$  iterations.

*Proof.* Item 3 is straightforward from item 2. To prove item 1 and 2 consider that if  $\max \frac{\partial J(x)}{\partial x_i} < 0$ , then  $\frac{\partial J(x)}{\partial x_i} < 0$  for all  $x$ . Analogously, if  $\min \frac{\partial J(x)}{\partial x_i} > 0$  then  $\frac{\partial J(x)}{\partial x_i} > 0$  for all  $x$ .  $\square$

Let us finally observe that verifying whether (3.3) or (3.4) holds is easy (polynomial in  $n$ ), as we just have to find the maxima, respectively the minima, of the  $n$  functions  $\frac{\partial J(x)}{\partial x_i}$  linear in  $x \in \mathbb{B}^n$ .

**4. Generalization to mbm Optimization.** Let us now generalize the results established in the preceding section to the case where DMs make decisions sequentially in groups of  $m$ .

**THEOREM 4.1.** *(Sufficient conditions) Let  $x^* = \mathbf{1}$  be an  $(m-1)b(m-1)$  optimum, if the team objective function  $J(\cdot)$  is such that for all  $S \subseteq N$  with  $|S| = m$  it holds*

$$(4.1) \quad \max_{x \in \mathbb{B}^n} \mathcal{R}_S J(x) < 0$$

then

1.  $x^*$  is the team-optimum
2.  $x^*$  is also the unique mbm optimum,
3. any mbm optimization algorithm converges to the team-optimum  $x^*$ .

*Proof.* Item 3 is straightforward from item 2. To prove item 1 and 2, let us assume by contradiction that there exists a team optimum value  $x^* \neq \mathbf{1}$ . Let  $V = \{i : x_i^* = 0\}$ . The cardinality of  $V$  cannot be greater than or equal to  $m$ . Indeed consider  $S \subseteq V$  with  $|S| = m$ , since  $\mathcal{R}_S J(x^*) < 0$  implies  $J(x^\circ) < J(x^*)$ , where  $x^\circ \in \mathbb{B}^n$  differs from  $x^*$  only for the components in  $S$ , i.e.,  $x_i^\circ = 0$  if  $i \in V \setminus S$ ,  $x_i^\circ = 1$  otherwise. Then  $x^*$  should be within an Hamming distance strictly less than  $m$  from  $\mathbf{1}$ , but this situation cannot occur since  $\mathbf{1}$  by definition is optimum within its neighborhood of radius  $m-1$ .  $\square$

**EXAMPLE 4.1.** Consider the team objective function  $J(x) = x_1 + x_2 - 3x_3 - 5x_1x_2 + x_1x_3 + x_2x_3$ . The solution  $x^* = \mathbf{1}$  is a pbp optimum as, for all  $i$ ,  $b_i + \sum_{k \neq i} a_{ik} < 0$ . Since for all  $S$ , with  $|S| = 2$  condition (4.1) holds (for  $i = 1$  and  $j = 2$ , we have  $b_1 + b_2 + a_{12} + \max_{x \in \mathbb{B}^n} (a_{13} + a_{23})x_3 = -1$ ), then  $x^* = \mathbf{1}$  is also team-optimum.

**REMARK 4.1.** In the above lemma, the assumption  $x^* = \mathbf{1}$  is without loss of generality. Actually, if the team problem has a unique team optimum  $x^* \neq \mathbf{1}$  then the following transformation can be applied to the decision space such that the new team optimum is  $\hat{x}^* = \mathbf{1}$ :

$$(4.2) \quad \hat{x}_i = \begin{cases} x_i & \text{if } x_i^* = 1 \\ 1 - x_i & \text{if } x_i^* = 0. \end{cases}$$

Let us finally observe that verifying whether (4.1) holds is, for fixed  $m$ , polynomial in  $n$  although exponential in  $m$ , as we just have to find the maxima of the  $\binom{n}{m}$  functions  $\mathcal{R}_S J(x)$  linear in  $x \in \mathbb{B}^n$ .

**5. A greedy algorithm to find a small set  $\Gamma_{\sigma(t)}$ .** Assume that at time  $t$  DM  $\sigma(t)$  may choose its neighborhood  $\Gamma_{\sigma(t)}$ . In this context, we present a *local* (implemented by DM  $\sigma(t)$ ) and *greedy* algorithm to find a small neighborhood  $\Gamma_{\sigma(t)}$  for

which  $\Gamma_i$ -approximate, respectively  $\Gamma_S$ -approximate, strategies  $\hat{\mu}_i(\cdot)$  or  $\hat{\mu}_S(\cdot)$  reproduce exactly the behavior of strict pbp strategies  $\hat{\mu}_i(\cdot)$ , or *mbm* strategies  $\hat{\mu}_S(\cdot)$ .

Given an initial point  $x(0)$ , consider the pbp optimization algorithm. Let  $i = \sigma(t)$  be the DM picked by the algorithm at step  $t$ . The DM  $i$  strategy provides the following result

$$x_i(t) = \begin{cases} 1 & \text{if } \frac{\partial J_i}{\partial x} \Big|_{x(t-1)} = b_i + \sum_{j \in N} a_{ij} x_j(t-1) \leq 0 \\ 0 & \text{otherwise} \end{cases}.$$

Here note that the value of  $x_i(t)$  depends only on the sign of  $\frac{\partial J_i}{\partial x} \Big|_{x(t-1)}$  and not on its exact value. In general, the strategy  $\mu_i(x_{-i}(t-1))$  may not need to know the values of all the  $x_j(t-1)$ , for  $j \in N$  to determine the sign of  $\frac{\partial J_i}{\partial x} \Big|_{x(t-1)}$  and, hence, to return the value of  $x_i(t)$ . As a consequence, an approximate pbp optimization algorithm certainly converges to  $x_{pbp}^*$  if, at each step  $t$ , the DM  $i$  chooses a  $\Gamma_i(t)$  so that the sign of  $\frac{\partial \hat{J}_{i, \Gamma_i}}{\partial x}$  is equal to the sign of  $\frac{\partial J_i}{\partial x}$  and, hence,  $\hat{\mu}_i(x_{-i}(t-1)) = \mu_i(x_{-i}(t-1))$ .

In particular, the DM  $i$  may choose the set  $\Gamma_i(t)$  by iteratively solving the following problem:

$$(5.1) \quad \begin{aligned} z &= \min \sum_{j \in N} y_j \\ b_i + \sum_{j \in D_i} a_{ij} x_j(t-1) + \sum_{j \in A_i^+} a_{ij} (1 - y_j) + \sum_{j \in A_i^-} a_{ij} y_j &\leq 0 \\ y_j &\in \{0, 1\} \quad \forall j \in N \end{aligned}$$

where: the binary variables  $y_j$ , for  $j \in N$ , are defined as follows

$$y_j = \begin{cases} 1 & \text{if } j \in \Gamma_i(t) \\ 0 & \text{otherwise} \end{cases};$$

the set  $D_i$  is the set of DMs whose values  $x_j(t-1)$  are known in advance by DM  $i$ ; finally, the sets  $A_i^+$  and  $A_i^-$  are such that  $A_i^+ = \{j \in N \setminus D_i : a_{ij} > 0\}$  and  $A_i^- = \{j \in N \setminus D_i : a_{ij} \leq 0\}$ .

Let  $y^* = \{y_j^* : j \in N\}$  be an optimal solution (if it exists) for (5.1) and  $\Gamma_i(t) = \{j \in N : y_j^* = 1\} \cup D_i$  the corresponding neighborhood of  $i$ . Problem (5.1) determines a minimal set  $\Gamma_i(t)$  of DMs that must be considered by DM  $i$  to be sure that  $\hat{\mu}_i(x_{-i}(t-1)) = \mu_i(x_{-i}(t-1)) = 1$ , given the knowledge of  $x_j(t-1)$ , for  $j \in D_i$ . Indeed, if the following conditions hold

$$(5.2) \quad \begin{aligned} x_j(t-1) &= 0, \text{ for each } j \text{ such that } y_j^* = 1 \text{ and } j \in A_i^+, \\ x_j(t-1) &= 1, \text{ for each } j \text{ such that } y_j^* = 1 \text{ and } j \in A_i^-, \end{aligned}$$

then

$$\frac{\partial J_i}{\partial x} \Big|_{x(t-1)} \leq b_i + \sum_{j \in D_i} a_{ij} x_j(t-1) + \sum_{j \in A_i^+} a_{ij} (1 - y_j^*) + \sum_{j \in A_i^-} a_{ij} y_j^* = \frac{\partial \hat{J}_{i, \Gamma_i}}{\partial x} \Big|_{x(t-1)} \leq 0.$$

Now, let us observe that three situations may occur:

- i) problem (5.1) has no feasible solution, that is  $\Gamma_i(t) = D_i$ ;
- ii) problem (5.1) has an optimal solution  $y^*$  and conditions (5.2) hold;

*iii*) problem (5.1) has an optimal solution  $y^*$  and conditions (5.2) do not hold.

If situation *i*) occurs, then both  $\frac{\partial \hat{J}_{i,\Gamma_i}}{\partial x} \Big|_{x(t-1)}$  and  $\frac{\partial J_i}{\partial x} \Big|_{x(t-1)}$  are strictly positive. Hence,  $\hat{\mu}_i(x_{-i}(t-1)) = \mu_i(x_{-i}(t-1)) = 0$ .

If situation *ii*) occurs, then both  $\frac{\partial \hat{J}_{i,\Gamma_i}}{\partial x} \Big|_{x(t-1)}$  and  $\frac{\partial J_i}{\partial x} \Big|_{x(t-1)}$  are non positive. Hence,  $\hat{\mu}_i(x_{-i}(t-1)) = \mu_i(x_{-i}(t-1)) = 1$ .

Differently, if situation *iii*) occurs, DM  $i$  cannot conclude that  $\frac{\partial \hat{J}_{i,\Gamma_i}}{\partial x} \Big|_{x(t-1)} = \frac{\partial J_i}{\partial x} \Big|_{x(t-1)}$ . In this last situation, DM  $i$  must enlarge the set  $D_i$  by including even the indexes of DMs whose values  $x_j(t-1)$  have been interrogated by DM  $i$  before realizing that conditions (5.2) do not hold. Then, DM  $i$  determines a further tentative set  $\Gamma_i(t)$  by solving (5.1) again.

The above procedure is iterated, starting from  $D_i = \emptyset$ , until either situation *i*) or *ii*) occurs. At each iteration the cardinality of  $D_i$  increases at least by one unit. Then, in the worst case, after at maximum  $n-1$  iterations, the procedure stops as  $\Gamma_i(t)$  has become equal to  $N \setminus \{i\}$  and hence  $\frac{\partial \hat{J}_{i,\Gamma_i}}{\partial x} \Big|_{x(t-1)} = \frac{\partial \hat{J}_{i,N \setminus \{i\}}}{\partial x} \Big|_{x(t-1)} = \frac{\partial J_i}{\partial x} \Big|_{x(t-1)}$  as  $\hat{J}_{i,N \setminus \{i\}} = J_i$ .

Let us finally observe that Problem (5.1) can be solved in polynomial time. Indeed, rewrite Problem (5.1) as

$$(5.3) \quad \begin{aligned} z &= \min \sum_{j \in N} y_j \\ &\sum_{j \in A_i^+ \cup A_i^-} \hat{a}_{ij} y_j \geq \hat{b}_i \\ y_j &\in \{0, 1\} \quad \forall j \in N \end{aligned}$$

where  $\hat{b}_i = b_i + \sum_{j \in D_i} a_{ij} x_j(t-1) + \sum_{j \in A_i^+} a_{ij}$  and  $\hat{a}_{ij} = \begin{cases} a_{ij} & \text{if } a_{ij} \in A_i^+ \\ -a_{ij} & \text{if } a_{ij} \in A_i^- \end{cases}$ .

Problem (5.3) is a relaxed (polynomial time) version of the change making problem [14] and its optimal solution can be trivially determined. Initially, re-denominate the DMs in  $A_i^+ \cup A_i^-$  so that  $\hat{a}_{ij} \geq \hat{a}_{ik}$  if  $j < k$ , then set

$$y_j^* = \begin{cases} 0 & \text{if } \sum_{k \in A_i^+ \cup A_i^-, k < j} \hat{a}_{ik} \geq \hat{b}_i \\ 1 & \text{otherwise} \end{cases}.$$

Similarly to (5.1), the DMs  $i \in S$  may determine whether  $\mathcal{R}_S f(x) \leq 0$  choosing a set  $\Gamma_S(t)$  by iteratively solving the following problem:

$$(5.4) \quad \begin{aligned} z &= \min \sum_{j \in N \setminus S} y_j \\ &\sum_{i \in S} b_i + \sum_{i \in S} \sum_{j \in S} a_{ij} + \sum_{i \in S} \sum_{k \in (N \setminus S) \cap D_S} a_{ik} x_k(t-1) + \\ &+ \sum_{i \in S} \sum_{k \in (N \setminus S) \cap A_S^+} a_{ik} (1 - y_j) + \sum_{i \in S} \sum_{k \in (N \setminus S) \cap A_S^-} a_{ij} y_j \leq 0 \\ y_j &\in \{0, 1\} \quad \forall j \in N \end{aligned}$$

where: the binary variables  $y_j$ , for  $j \in N \setminus S$ , are defined as follows

$$y_j = \begin{cases} 1 & \text{if } j \in \Gamma_S(t) \\ 0 & \text{otherwise} \end{cases} ;$$

the set  $D_S$  is the set of DMs whose values  $x_j(t-1)$  are known in advance by DMs  $i \in S$ ; finally, the sets  $A_S^+$  and  $A_S^-$  are such that  $A_S^+ = \{j \in N \setminus D_S : a_{ij} > 0\}$  and  $A_S^- = \{j \in N \setminus D_S : a_{ij} \leq 0\}$ .

A little more difficult is for the DMs  $i \in S$  to choose a  $\Gamma_S(t)$  so that  $\hat{\mu}_i(x_{-S}(t-1)) = \mu_i(x_{-S}(t-1))$ .

Actually,  $x_S^* = \mu_i(x_{-S}(t-1))$  if, for all  $\tilde{x}_S \in \mathbb{B}^m$ ,

$$\sum_{i \in S} b_i(x_i^* - \tilde{x}_i) + \sum_{i \in S} \sum_{j \in S} a_{ij}(x_i^* x_j^* - \tilde{x}_i \tilde{x}_j) + \sum_{i \in S} \sum_{k \in N \setminus S} a_{ik}(x_i^* - \tilde{x}_i)x_k(t-1) \leq 0$$

Then, the DMs  $i \in S$  may determine whether a tentative  $\hat{x}_S \in \mathbb{B}^m$  is equal to  $\mu_i(x_{-S}(t-1))$  choosing a set  $\Gamma_S(t)$  by iteratively solving the following problem:

$$\begin{aligned} z = \min & \sum_{j \in N \setminus S} y_j \\ & \sum_{i \in S} b_i(\hat{x}_i - \tilde{x}_i) + \sum_{i \in S} \sum_{j \in S} a_{ij}(\hat{x}_i \hat{x}_j - \tilde{x}_i \tilde{x}_j) + \sum_{i \in S} \sum_{k \in (N \setminus S) \cap D_S} a_{ik}(\hat{x}_i - \tilde{x}_i)x_k(t-1) + \\ (5.5) & + \sum_{i \in S} \sum_{k \in (N \setminus S) \cap A_S^+} a_{ik}(\hat{x}_i - \tilde{x}_i)(1 - y_j) + \\ & + \sum_{i \in S} \sum_{k \in (N \setminus S) \cap A_S^-} a_{ij}(\hat{x}_i - \tilde{x}_i)y_j \leq 0 \quad \forall \tilde{x}_S \in \mathbb{B}^m \\ & y_j \in \{0, 1\} \quad \forall j \in N \end{aligned}$$

where the binary variables  $y_j$ , for  $j \in N \setminus S$ , and the sets  $D_S$ ,  $A_S^+$  and  $A_S^-$  are defined as for (5.4).

Now, let us observe that three situations may occur:

- i)* problem (5.5) has no feasible solution;
- ii)* problem (5.5) has an optimal solution  $y^*$  and conditions (5.2) hold;
- iii)* problem (5.5) has an optimal solution  $y^*$  and conditions (5.2) do not hold.

If situation *i)* occurs, then  $\hat{x}_S$  is not  $\mu_i(x_{-S}(t-1))$ , a different  $x_S \in \mathbb{B}^m$  must be considered as tentative  $\mu_i(x_{-S}(t-1))$ .

If situation *ii)* occurs, then  $\hat{x}_S = \mu_i(x_{-S}(t-1))$ .

Differently, if situation *iii)* occurs, DMs  $i \in S$  cannot conclude neither  $\hat{x}_S = \mu_i(x_{-S}(t-1))$  nor  $\hat{x}_S \neq \mu_i(x_{-S}(t-1))$ . In this situation, DMs  $i \in S$  must enlarge the set  $D_S$  by including even the indexes of DMs whose values  $x_j(t-1)$  have been interrogated by DM  $i \in S$  before realizing that conditions (5.2) do not hold. Then, DMs  $i \in S$  solves (5.5) again.

The above procedure is iterated, starting from  $D_S = \emptyset$ , until either situation *i)* or *ii)* occurs. At each iteration the cardinality of  $D_S$  increases at least by one unit. Then, in the worst case, after at maximum  $n - m$  iterations, the procedure stops as  $\Gamma_S(t)$  has become equal to  $N \setminus S$ .

**6. Sub-modular Team Problems.** In the past sections we have provided conditions for the convergence from any initial state  $x(0)$ . Now, we show that we can

TABLE 6.1

Sequence of DMs' decisions: blocks on the left, middle and right describe the first, second and third round of optimization.

DM	$x$	$J(x)$	DM	$x$	$J(x)$	DM	$x$	$J(x)$
1	(1,0,0,0,0)	-1	1	(0,1,1,0,0)	-4	1	(1,1,1,1,1)	-7
2	(1,1,0,0,0)	-2	2	(0,1,1,0,0)	-4	2	(1,1,1,1,1)	-7
3	(1,1,1,0,0)	-3	3	(0,1,1,0,0)	-4	3	(1,1,1,1,1)	-7
4	(1,1,1,0,0)	-3	4	(0,1,1,1,0)	-5	4	(1,1,1,1,1)	-7
5	(1,1,1,0,0)	-3	5	(0,1,1,1,1)	-6	5	(1,1,1,1,1)	-7

recognize in polynomial time a special class of sub-modular team problems, which do not meet the aforementioned conditions and for which the convergence is guaranteed at least when the pbp algorithm is opportunely initialized. This class is characterized by so-called *threshold strategies*.

Let us call *sub-modular team problems*, all team problems with sub-modular team objective function. From [5], we remind from that i) a pseudo-Boolean function  $f(\cdot)$  is *sub-modular* if  $f(v) + f(w) \leq f(vw) + f(v \vee w)$  ii) a quadratic pseudo-Boolean function  $f(\cdot)$  is submodular iff its quadratic terms are nonpositive. However, from the following example, it is apparent that sub-modularity alone does not guarantee the convergence of any pbp optimization algorithm.

EXAMPLE 6.1. Consider the sub-modular team objective function  $J(x) = x_1 + x_2 - 3x_1x_2$  and take  $x(0) = (0, 0)$ . The team optimum is  $(1, 1)$  but observe that at iteration 1, no DM alone benefits from changing its decision from 0 to 1. Hence the pbp optimization algorithm starts and terminates in  $(0, 0)$ .

We can generalize the above reasoning to show that sub-modularity alone does not guarantee the convergence of any mbm optimization algorithm. On this purpose, note that if the team objective function is sub-modular, then condition (4.1) reduces to

$$(6.1) \quad \sum_{i \in S} b_i + \sum_{i \in S} \sum_{j \in S} a_{ij} < 0, \quad \text{for all } S, \text{ with } |S| = m.$$

We derive the above result by reminding that all quadratic terms are nonpositive and therefore  $\max_x \sum_{k \neq i, j} (a_{ik} + a_{jk})x_k \leq 0$  with the equality verified in  $x = 0$ .

EXAMPLE 6.2. Consider the sub-modular team objective function  $J(x) = 2x_1 + 2x_2 + 2x_3 - 3x_1x_2 - 3x_1x_3 - 3x_2x_3$  and take  $x(0) = (0, 0)$ . The team optimum is again  $(1, 1)$  but observe that at iteration 1, no pairs  $i$  and  $j$  of DMs alone benefits from changing their decisions from 0 to 1. Note that condition (6.1) for  $m = 2$  becomes  $b_i + b_j + a_{ij} < 0$  and there is no pair  $i$  and  $j$  that satisfies such a condition. Hence the mbm optimization algorithm starts and terminates in  $(0, 0)$ .

**6.1. A Special Class with Threshold Strategies.** Threshold strategy means that a DM  $i$  chooses  $x_i = 1$  if and only if at least other  $l_i$  DMs do the same. The following simple example shows that when DMs have threshold strategies the team objective function is sub-modular. The team objective function is as in (2.1). We say that DM  $i$  has a threshold strategy with threshold  $l_i = k$ , if its strict pbp strategy is

$$(6.2) \quad \mu_i(x_{-i}) = \begin{cases} 1 & \text{if } \|x_{-i}\|_1 \geq k \\ 0 & \text{otherwise.} \end{cases}$$

TABLE 6.2

Sequence of decisions: first and second round of pbp optimization (left and middle blocks), 2b2 optimization (right block).

DM	$x$	$J(x)$	DM	$x$	$J(x)$	DM	$x$	$J(x)$
1	(0,0,0,0,0)	0	1	(0,0,1,0,0)	-3	1-2	(1,1,0,0,0)	-3
2	(0,0,0,0,0)	0	2	(0,0,1,0,0)	-3	3-4	(1,1,1,1,0)	-11
3	(0,0,1,0,0)	-3	3	(0,0,1,0,0)	-3	5-1	(1,1,1,1,1)	-23
4	(0,0,1,0,0)	-3	4	(0,0,1,0,0)	-3	2-3	(1,1,1,1,1)	-23
5	(0,0,1,0,0)	-3	5	(0,0,1,0,0)	-3	4-5	(1,1,1,1,1)	-23

LEMMA 6.1. *If all DMs have threshold strategies then the team objective function  $J(x)$  must be sub-modular.*

*Proof.* Observe that DM  $i$  has a threshold strategy with  $l_i = k$ . Denote by  $\mathcal{S}(k)$  the set of all subsets of  $N$ , which do not contain DM  $i$  and have cardinality less than  $k$ . Now, for a generic subset  $S \in \mathcal{S}(k)$ , take  $x_{-i}$  such that  $x_j = 1$  for all  $j \in S$  and  $x_j = 0$  for all  $j \in N \setminus (S \cup \{i\})$  and observe that from (6.2) it must hold that  $\mu_i(x_{-i}) = 0$ . But this means that the following condition holds true

$$(6.3) \quad b_i + \sum_{j \in S} a_{ij} \geq 0 \text{ for all } S \in \mathcal{S}(k).$$

Repeat the same reasoning considering a generic subset  $S \subseteq N \setminus \mathcal{S}(k)$ , and take  $x_{-i}$  such that  $x_j = 1$  for all  $j \in S$  with  $j \neq i$  and  $x_j = 0$  for all  $j \in N \setminus S$ . Observe that from (6.2) it must hold that  $\mu_i(x_{-i}) = 1$  which implies that the following condition hold true

$$(6.4) \quad b_i + \sum_{j \in S} a_{ij} < 0 \text{ for all } S \subseteq N \setminus \mathcal{S}(k).$$

Now, consider two sets  $S^1 \in \mathcal{S}(k)$  with  $|S^1| = k - 1$  and  $S^2 = S^1 \cup \{j\} \in N \setminus \mathcal{S}(k)$ . Observe that  $S^2$  has cardinality  $|S^2| = k$  as it is obtained from  $S^1$  by adding a single DM  $j$ . We complete the proof by observing that for (6.3) and (6.4) to be valid it must be  $a_{ij} < 0$  for all  $i$  and  $j$ . Then  $J(\cdot)$  has all quadratic terms negative which proves that  $J(\cdot)$  is sub-modular. □

This special class of sub-modular team problems is interesting as i) threshold structures can be recognized in polynomial time and ii) any pbp optimization algorithm initialized at  $x(0) = \mathbf{1}$  converges to the team-optimum  $x^*$ , in general different from  $\mathbf{1}$ , as established in the following theorem.

THEOREM 6.1. *There exists a polynomial algorithm that verifies conditions (6.3) and (6.4) in  $O(n^2 \log n)$ . In case of positive answer, any pbp optimization algorithm initialized at  $x(0) = \mathbf{1}$  converges to the team-optimum.*

*Proof.* (Complexity) Given a DM  $i$ , consider all DMs except  $i$  in the order  $\sigma(1), \dots, \sigma(n)$  with  $a_{i\sigma(1)} \leq \dots \leq a_{i\sigma(n)}$ . We remind here that the ordering process has a complexity  $O(n \log n)$ . Now, conditions (6.3) and (6.4) are verified if and only if  $b_i + a_{i\sigma(1)} + \dots + a_{i\sigma(k-1)} \geq 0$  and  $b_i + a_{i\sigma(n-k)} + \dots + a_{i\sigma(n)} < 0$ . We can limit ourselves to verify the latter two conditions for any possible value of the threshold  $l_i$  from 1 to  $n$ . Such a procedure is carried out via a dicotomic search and has a complexity of  $O(\log n)$ . Then, for fixed  $i$  the total complexity is  $O(n \log n) + O(\log n)$ ,

and as  $O(n \log n)$  dominates (is always greater than)  $O(\log n)$  the total complexity simply reduces to the cost of the ordering process  $O(n \log n)$ . We conclude our proof by noticing that the ordering process must be repeated  $n$  times (one for all DM  $i$ ) and therefore the resulting complexity is  $O(n^2 \log n)$ .

(Convergence of pbp) Assume DMs ordered by increasing thresholds, i.e.,  $l_1 \leq \dots \leq l_n$ . Starting at  $x(0) = \mathbf{1}$  any pbp optimization algorithm converges to the pbp optimum nearest to  $\mathbf{1}$  (in terms of Hamming distance), call it  $\hat{x}$ . In other words  $\hat{x} = \arg \min\{\|x - \mathbf{1}\| : x \text{ is pbp-opt.}\}$ . We must show that  $\hat{x}$  is also the team-optimum. To prove this fact corresponds to proving that, if there exists a second pbp optimum, call it  $\tilde{x}$ , it must hold

$$\begin{aligned} J(\hat{x}) - J(\tilde{x}) &= \mathcal{R}_S J(\tilde{x}) = \\ &= \sum_{i \in S} b_i + \sum_{i \in S} \sum_{j \in S} a_{ij} + \sum_{r \notin S} \sum_{i \in S} a_{ir} \tilde{x}_r \leq 0, \end{aligned}$$

where  $S$  is the set of components which are zero in  $\tilde{x}$  and one in  $\hat{x}$ . Now note that  $\sum_{i \in S} \sum_{j \in S} a_{ij} + \sum_{r \notin S} \sum_{i \in S} a_{ir} \tilde{x}_r = \sum_{i \in S} \sum_{r \in N} a_{ir} \hat{x}_r$  and therefore we can rewrite the above inequality as

$$(6.5) \quad J(\hat{x}) - J(\tilde{x}) = \sum_{i \in S} (b_i + \sum_{r \in N} a_{ir} \hat{x}_r) = \sum_{i \in S} (b_i + \sum_{r \in \bar{S}} a_{ir}) \leq 0,$$

where we denote by  $\bar{S}$  the set of components which are one in  $\hat{x}$ . Then we need to prove the validity of (6.5). Now, note that if DMs are ordered by increasing thresholds, it must hold  $\tilde{x} \leq \hat{x}$  component-wise. Hence, as  $\hat{x}$  is a pbp optimum then each  $i \in S$  has threshold  $l_i < \|\hat{x} - \mathbf{0}\| = \|\hat{x}\|$  which in turns implies that  $\sum_{i \in S} (b_i + \sum_{r \in \bar{S}} a_{ir}) \leq 0$  and therefore (6.5) hold true.  $\square$

**REMARK 6.1.** *Threshold strategies simplify the search for a small neighborhood  $\Gamma_{\sigma(t)}$ . DMs may implement a random selection of neighbors based only on their threshold. So, if DM  $i$  has threshold 4, then, it will start selecting randomly four neighbor DMs, and still randomly increase their number until it can certainly affirm or exclude that at least four of them play 1.*

**7. Numerical example.** In this first example we simulate a pbp optimization and show that the algorithm converges to the team optimum. Consider the following team objective function

$$\begin{aligned} J(x) &= -x_1 + x_2 + x_3 + x_4 + 5x_5 - 2x_1x_2 + 4x_1x_3 + \\ &\quad + 2x_1x_4 - 4x_1x_5 - 6x_2x_3 - 2x_2x_4 - 7x_4x_5 \end{aligned}$$

By direct verification, it can be proved that the above function is a CLG-function as it has a unique local minimum in  $(1, 1, 1, 1, 1)$ . Similarly, we can see that it is not a CU-function as, for instance, on the 2-face  $x_1$ - $x_3$  with  $x_2 = x_4 = x_5 = 0$ , conditions (3.1)-(3.2) are both verified. The function is not submodular because of the presence of positive quadratic terms.

Start from the decision vector  $x = 0$  and assume that the DMs make their decision in the following order: DM 1, DM 2, ..., DM 5. Table 6.1 reports the sequence of DMs' decisions (decisions are starred when they change with respect to the previous round). Blocks on the left describe the first and second round of optimization while block on the right describes the third round of optimization.

If we consider only the vectors  $x$  that change from a decision to another one we obtain the sequence

$$\begin{aligned} \sigma = & (1, 0, 0, 0, 0), (1, 1, 0, 0, 0), (1, 1, 1, 0, 0), (0, 1, 1, 0, 0), \\ & (0, 1, 1, 1, 0), (0, 1, 1, 1, 1), (1, 1, 1, 1, 1). \end{aligned}$$

In this second example we simulate the pbp and the 2b2 optimization for the following team objective function and show that only in the second case we converge to the team optimum:

$$\begin{aligned} J(x) = & x_1 + x_2 - 3x_3 + x_4 + x_5 - 5x_1x_2 + x_1x_3 + x_2x_3 + \\ & -4x_1x_4 - 4x_1x_5 - 4x_2x_4 - 4x_2x_5 - 5x_4x_5. \end{aligned}$$

First observe that the solution  $x^* = \mathbf{1}$  is a pbp optimum as, for all  $i$ ,  $b_i + \sum_{k \neq i} a_{ik} < 0$ . Furthermore, since for all  $S$ , with  $|S| = 2$  condition (4.1) holds, then  $x^* = \mathbf{1}$  is also team-optimum. The pbp optimization is carried out as in the previous example and decisions are reported in Table 6.2 (left blocks describe the first and second round). Convergence is on  $x = (0, 0, 1, 0, 0) \neq x^*$ . Differently, the 2b2 optimization converges to  $x^*$  as evident from the sequence of decisions listed in the right block.

**8. Concluding Remarks.** In future works, we wish to extend the obtained results to consensus problems. Actually, consensus problems have been recently reinterpreted as special potential games [13]. For these games there exist algorithms, very similar in spirit to pbp algorithms and called *best response path algorithm*, that guarantee the distributed convergence to Nash equilibria.

A second line of research aims at providing a parallel between *mbm* and self organizing/Kohonen maps, since both are optimization methods that can be applied to boolean spaces with decreasing goal functions that in each iteration modify a subset of decision variables.

#### REFERENCES

- [1] D. Bauso, R. Pesenti, "Generalized person-by-person optimization in team problems with binary decisions", *Proc. of the American Control Conference 2008*, Seattle, USA, 2008, pp. 717–722.
- [2] D. Bauso, "Boolean-controlled systems via receding horizon and linear programming", *Journal of Mathematics of Control Signals and Systems*, vol. 21, no. 1, 2009, pp. 69–91.
- [3] D. Bauso, L. Giarrè, R. Pesenti, "Consensus in Noncooperative Dynamic Games: a Multi-Retailer Inventory Application", *IEEE Transactions on Automatic Control*, vol. 53, no. 4, 2008, pp. 998–1003.
- [4] D. Bertsimas, A. Thiele, "A Robust Optimization Approach to Inventory Theory", *Operations Research*, vol. 54, no. 1, 2006, pp. 150–168.
- [5] E. Boros and P. L. Hammer, "Pseudo-Boolean optimization", *Discrete Applied Mathematics*, vol. 123, pp. 155–225, 2002.
- [6] H. Björklund and S. Sandberg and S. Vorobyov, "Optimization on completely unimodal hypercubes", *Technical Report 2002-018*, Department of Information Technology, Uppsala University, May 2002.
- [7] M. S. Branicky, V. S. Borkar and S. K. Mitter, "A Unified Framework for Hybrid Control: Model and Optimal Control Theory", *IEEE Trans. on Automatic Control*, vol. 43, no. 1, 1998, pp. 31–45.
- [8] R. Cogill and S. Lall, "An approximation algorithm for the discrete team decision problem", *SIAM Journal on Control and Optimization*, vol. 45, no. 4, 2006, pp. 1359–1368.
- [9] P. R. De Waal and J. H. Van Schuppen, "A class of team problems with discrete action spaces: optimality conditions based on multimodularity", *SIAM Journal on Control and Optimization*, vol. 38, 2000, pp. 875–892.

- [10] G.C. Goodwin and D.E. Quevedo, “Finite alphabet control and estimation”, *International Journal of Control, Automation, and Systems*, vol.1, no.4, 2003, pp. 412–430.
- [11] P. L. Hammer and B. Simeone and T. Lieblich and D. de Werra, “From linear separability to unimodality: a hierarchy of pseudo-boolean functions”, *SIAM Journal of Discrete Mathematics*, vol. 1, 1988, pp. 174–184.
- [12] Y.-C. Ho, “Team decision theory and information structures”, *Proceedings IEEE*, 1980, vol. 68, pp. 644–654.
- [13] J.R. Marden, G. Arslan, and J.S. Shamma, “Connections between cooperative control and potential games illustrated on the consensus problem”, *European Control Conference*, Kos, Greece, July 2-5, 2007, pp. 4604-4611.
- [14] S. Martello, and P. Toth, *Knapsack Problems: Algorithms and Computer Implementations*, John Wiley & Sons, Hoboken, NJ, 1990.
- [15] J. Matousek, “The number of unique-sink orientations of hypercube”, *Combinatorica*, vol. 26, no. 1, 2006, pp. 91–99.
- [16] A. Rantzer, “Linear quadratic team theory revisited”, *Proceedings of the 2006 American Control Conference* Minneapolis, Minnesota, USA, June 14-16, 2006, pp. 1637–1641.
- [17] D. C. Tarraf, A. Megretski and M. A. Dahleh, “A Framework for Robust Stability of Systems Over Finite Alphabets”, *IEEE Transactions on Automatic Control*, vol. 53, no. 5, 2008, pp. 1133–1146.