

This is a repository copy of Rasch Analysis of the Hospital Anxiety and Depression Scale in Psoriatic Arthritis: Results from the Presta Study.

White Rose Research Online URL for this paper: http://eprints.whiterose.ac.uk/91604/

Version: Accepted Version

Proceedings Paper:

Ndosi, ME, Hsu, M, Cappelleri, J et al. (3 more authors) (2015) Rasch Analysis of the Hospital Anxiety and Depression Scale in Psoriatic Arthritis: Results from the Presta Study. In: Arthritis and Rheumatology. 2015 ACR/ARHP Annual Meeting, 06-11 Nov 2015, San Francisco, CA. Wiley.

https://doi.org/10.1002/art.39448

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Rasch Analysis of the Hospital Anxiety and Depression Scale in Psoriatic Arthritis: Results from the PRESTA Study.

Mwidimi Ndosi¹, Ming-Anne Hsu², J. Cappelleri³, Heather Jones⁴, Amit Chhabra⁵ and Philip S. Helliwell⁶,

¹School of Healthcare, University of Leeds, Leeds, United Kingdom, ²445 Eastern Point Road, Pfizer Inc, Groton, CT, ³Statistics, Pfizer Inc, New London, CT, ⁴Inflammation Global Medical Affairs, Pfizer, Collegeville, PA, 5Pfizer Inc, Collegeville, PA, ⁶Section of Musculoskeletal Disease, University of Leeds, Leeds, United Kingdom

Background/Purpose

The Hospital Anxiety and Depression Scale (HADs) is a generic measure of psychological status comprising anxiety and depression subscales. The aim of this study was to assess the psychometric properties of HADs in psoriatic arthritis (PsA), to calibrate the scale, and to provide interval-level scale for use in parametric analyses when required.

Methods

We used HADS data from patients with PsA recruited in PRESTA trial. [1] The data was subjected to Rasch analysis to determine fit to the Rasch model (implying construct validity and unidimensionality), reliability and targeting in subjects with PsA.

Results

The number of evaluable subjects was 740 at baseline, 701 at week 12, and 653 at week 24. Both the anxiety subscale and the depression subscale satisfied the expectation of the Rasch model (table 1). The overall scale was shown to fit the Rasch model (item-by-severity interaction Chi-Square = 15.878, p = 0.601) and had excellent reliability (person separation index = 0.888). Validity and reliability of HADS were confirmed at baseline and both follow-up visits.

Figure 1 presents person location relative to all items (logarithmically transformed scores) along the same scale (logits). The top plot representing 'persons severity'; those with higher scores (impaired psychological status) on the right of the scale and those with lower scores (better psychological status) on the left. The bottom plot presents relative 'difficulty' of the items. HADS appears to be well targeted across all 'severity' levels, providing for calibration of the scale by transforming raw scores into interval-level (Rasch-transformed) scores.

Conclusions

The validity and reliability of the HADS are confirmed in PsA and continues to be a useful psychological status instrument to use in PsA clinical studies. Raw scores can be Rasch-transformed into interval scores for use alongside other outcomes in parametric analyses.

Table 1: Fit Statistics for the Anxiety and Depression Subscales in PsA (Baseline).

Subscale	Location	SE	Item Fit Residuals	Chi-Square	DF	p-value
Anxiety	-0.120	0.021	-0.074	6.494	9	0.690
Depression	0.120	0.022	0.356	9.384	9	0.403

SE, Standard error; DF, Degrees of freedom, Non-significant p-value for Chi-Square suggests fit to Rasch model

Person-Item Threshold Distribution (Grouping Set to Interval Length of 0.50 making 18 Groups) PERSONS 150 20.2% No. Mean [741] -0.986 1.277 Total 125 r e q u e n c y 75 10.1% 50 25 0 0.0% Location (logits) ITEMS 0 0.0% 23.8%

Figure 1: Person-item Threshold Distribution Showing Targeting of the HADS (Baseline Data)

Reference

1. Sterry W, et al. Comparison of two etanercept regimens for treatment of psoriasis and psoriatic arthritis: PRESTA randomised double blind multicentre trial. BMJ 2010;340(c147)