



UNIVERSITY OF LEEDS

This is a repository copy of *Optimising primary care research participation: a comparison of three recruitment methods in data-sharing studies*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/94719/>

Version: Accepted Version

---

**Article:**

Lord, P, Willis, T, Carder, P et al. (2 more authors) (2016) Optimising primary care research participation: a comparison of three recruitment methods in data-sharing studies. *Family Practice*, 33 (2). pp. 200-204. ISSN 0263-2136

<https://doi.org/10.1093/fampra/cmw003>

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

Table 2. Comparison of characteristics and quality measures between recruited and non-recruited practices for each recruitment method

	Opt-in			Mixed opt-in and opt-out			Opt-out		
	Not-recruited	Recruited	p	Not-recruited	Recruited	p	Not-recruited	Recruited	p
Practices, n	47 (42%)	64* (58%)		47 (30%)	110* (70%)		22 (20%)	89 (80%)	
List Size (Patients)	<b>4722</b>	<b>7153</b>	<b>0.03</b>	<b>5857</b>	<b>7091</b>	<b>0.043</b>	5534	6455	0.121
Number of GPs in the practice (mean)	<b>4.2</b>	<b>5.3</b>	<b>0.033</b>	<b>4</b>	<b>5</b>	<b>0.04</b>	3.6	5	0.055
Male GPs	2.3	2.6	0.301	<b>2</b>	<b>2.6</b>	<b>0.018</b>	2.2	2.6	0.17
IMD (proportionally applied across practice population)	22.5	25.8	0.997	33.1	33.7	0.769	30.6	33.1	0.25
Clinical Quality Measure									
Diabetic patients with HbA1c<=59mmol/mol (%)	57.2	56.6	0.562	<b>53.9</b>	<b>58.4</b>	<b>&lt;0.01</b>	57.7	57.1	0.921
Patient Experience Measure									
Rating of GP treating you with care and concern (good/very good) (%)	84.8	84.7	0.673	80.5	81.4	0.76	84.4	83.3	0.595
Local Service Impact Measure									
Emergency Admissions / 1000	111.9	101.1	0.19	112.3	105.3	0.673	102.7	104.9	0.947

Averages are median unless otherwise stated. Comparison with Wilcoxon rank sum test with Monte Carlo permutation sampling (10,000 permutations).

\*One practice closed between the end of the original studies and this analysis therefore full data were not available for comparison here.