



This is a repository copy of *Prevalence of visits to five types of complementary and alternative medicine practitioners by the general population: A systematic review.*

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/96913/>

Version: Accepted Version

---

**Article:**

Cooper, K.L., Harris, P.E., Relton, C. et al. (1 more author) (2013) Prevalence of visits to five types of complementary and alternative medicine practitioners by the general population: A systematic review. *Complementary Therapies in Clinical Practice* , 19 (4). pp. 214-220. ISSN 1744-3881

<https://doi.org/10.1016/j.ctcp.2013.06.006>

---

**Reuse**

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## **Prevalence of visits to five types of complementary and alternative medicine practitioners by the general population: a systematic review**

**Katy L Cooper<sup>a</sup>, Philip E Harris<sup>b</sup>, Clare Relton<sup>c</sup>, Kate J Thomas<sup>d</sup>.**

<sup>a</sup>School of Health and Related Research, University of Sheffield, Regent Court, 30 Regent Street, Sheffield, S1 4DA, UK. Email: [k.l.cooper@sheffield.ac.uk](mailto:k.l.cooper@sheffield.ac.uk)

<sup>b</sup>Cardiff School of Health Sciences, Cardiff Metropolitan University, PO Box 377, Western Avenue, Cardiff, CF5 2YB, UK. Email: [peharris@cardiffmet.ac.uk](mailto:peharris@cardiffmet.ac.uk)

<sup>c</sup>School of Health and Related Research, University of Sheffield, Regent Court, 30 Regent Street, Sheffield, S1 4DA, UK. Email: [c.relton@sheffield.ac.uk](mailto:c.relton@sheffield.ac.uk)

<sup>d</sup>School of Health and Related Research, University of Sheffield, Regent Court, 30 Regent Street, Sheffield, S1 4DA, UK. Email: [k.thomas@sheffield.ac.uk](mailto:k.thomas@sheffield.ac.uk)

### **Corresponding author:**

Katy Cooper, Research Fellow

School of Health and Related Research, University of Sheffield, Regent Court, 30 Regent Street, Sheffield, S1 4DA, UK. Email: [k.l.cooper@sheffield.ac.uk](mailto:k.l.cooper@sheffield.ac.uk) Tel: +44 (0)114 2220773.

**Keywords:** systematic review; prevalence; complementary medicine; acupuncture; homeopathy; osteopathy; chiropractic; herbal medicine

**Disclosures:** None.

## **ABSTRACT**

**Objective:** To systematically review surveys of 12-month prevalence of visits to complementary and alternative medicine (CAM) practitioners for five therapies: acupuncture, homeopathy, osteopathy, chiropractic, and medical herbalism.

**Methods:** Studies were identified via database searches to 2011. Study quality was assessed using a six-item tool.

**Results:** Forty-one surveys across 12 countries were included. Twenty-five (61%) met four of six quality criteria. Prevalence of visits by adults were (median, range): acupuncturists 1.4% (0.2-7.5%, N=27 surveys), homeopaths 1.5% (0.2-2.9%, N=20 surveys), osteopaths 1.9% (0.2-4.4%, N=9 surveys), chiropractors 7.5% (0.3-16.7, N=33 surveys), medical herbalists 0.9% (0.3-4.7%, N=14 surveys). Estimates were slightly lower for children and higher for older adults. There was little change over the past 15-20 years.

**Conclusions:** This review summarises 12-month prevalence of visits to CAM practitioners in Europe, North America, Australia, East Asia, Saudi Arabia and Israel. A small but significant percentage of these general populations visit CAM practitioners each year.

## **INTRODUCTION**

We recently published results of a broad-scale systematic review assessing prevalence of use of complementary and alternative medicine (CAM) within general populations across 15 countries.<sup>1</sup> Estimates of 12-month prevalence of use of any CAM ranged from 9.8% to 76% (based on 32 studies), while estimates of 12-month prevalence of visits to CAM practitioners ranged from range 1.8% to 48.7% (based on 33 studies). Though these ranges were wide, estimates of 12-month prevalence of any CAM use (excluding prayer) from surveys using consistent measurement methods showed remarkable stability within some countries, such as Australia (49%, 52% and 52% in 1993, 2000 and 2004 respectively) and USA (36% and 38% in 2002 and 2007).

The focus of the study reported here is to systematically review the subset of these general population studies that reported 12-month prevalence of visits to practitioners for any one of five types of complementary and alternative medicine (CAM): acupuncture, homeopathy, osteopathy, chiropractic, and medical herbalism. Each of these therapies has established training and governance practices in the countries surveyed, and each claims to utilise a specific diagnostic approach which is independent of Western medical practice.<sup>2</sup>

Brief definitions of the five therapies are as follows.<sup>2</sup> Acupuncturists insert small needles into various points in the body. Traditional Chinese acupuncturists use the idea of 'qi' (vital energy) travelling around the body along 'meridians', while Western acupuncturists prefer to think of needle insertion as affecting nerve impulses and the central nervous system. Homeopaths use the principle of treating 'like with like'. Homeopathic remedies are highly diluted and serially succussed substances that if given to a healthy person would produce the symptoms that the remedies are being given to treat. Osteopaths use a system of diagnosis and treatment, usually by manipulation, which mainly focuses on musculo-skeletal problems, though some branches aim to treat a wider spectrum of disorders. Chiropractors treat musculo-skeletal complaints through adjusting muscles, tendons and joints and using manipulation and massage

techniques. Medical herbalists use remedies derived from plants and plant extracts to treat disorders and maintain good health.

This review focusses on visits to practitioners rather than self-treatment using over-the-counter products (for example for homeopathic remedies and herbal preparations). This decision was made on the basis that practitioner visits represent significant examples of health behaviour, and estimates for this behaviour are likely to be better-defined and less prone to recall bias than estimates for self-treatment.

## **METHODS**

### Search strategy

The systematic review followed the recommendations in the PRISMA statement.<sup>3</sup> The following databases were searched in February 2011: MEDLINE, Medline in Process, EMBASE, Cochrane Database of Systematic Reviews, Cochrane CENTRAL Register of Controlled Trials, HTA database, Science Citation Index, AMED, and PsycINFO. The search strategy combined terms for: i) complementary and alternative medicines, ii) prevalence, surveys or patterns of use, and iii) population-level or national-level data. The full search strategy is provided in our previous article on prevalence of use of any CAM.<sup>1</sup> The search was restricted to studies published from 1998 onwards. Studies published prior to 1998 were identified from two previous systematic reviews of CAM prevalence.<sup>4,5</sup> Bibliographies of included papers were checked for further relevant studies.

### Inclusion and exclusion criteria

Studies were included if they reported 12-month prevalence of overall CAM use and/or visits to CAM practitioners, in addition to prevalence of visits to at least one of five types of CAM practitioner: acupuncturists, homeopaths, osteopaths, chiropractors, and medical herbalists. Prevalence had to be reported over a 12-month retrospective period within a representative

general population sample of a nation or a defined geographical area. Surveys of clearly-defined age groups (such as adults, children or older adults) were included. Included studies used survey methods such as structured interviews or self-complete questionnaires. Studies were excluded if they did not report 12-month prevalence, or were not written in English. Studies were also excluded if they were not based on representative samples of the general population; for example, surveys of sub-populations with specific clinical conditions or socio-demographic characteristics (other than age).

#### Study selection and data extraction

Study titles retrieved by the search were assessed for inclusion by one reviewer and a sample of excluded titles was checked by a second reviewer. Potentially relevant abstracts and full texts were assessed by two reviewers and any discrepancies resolved through discussion. Data were extracted by one reviewer and checked by a second.

#### Quality assessment

There is no agreed set of criteria for assessing the quality of health-related surveys. As part of our wider systematic review on prevalence of overall CAM use, we devised a six-item, literature-based quality assessment tool comprising important and assessable criteria of methodological quality.<sup>1</sup> This was applied to each of the included studies. The criteria covered by the quality assessment tool include 1) whether CAM-use questions were clearly described and number of therapies/questions reported; 2) whether the survey was piloted (this was assumed for government surveys); 3) whether the sample size was  $\geq 1,000$  and/or a CAM-specific sample size calculation was reported; 4) whether the reported response rate was  $\geq 60\%$ ; 5) whether data were weighted to population characteristics (where appropriate) to reduce non-response bias; and 6) whether a 95% confidence interval or standard error were reported for the main prevalence estimates.

## **RESULTS**

### **Number of surveys included**

The search identified 2312 unique citations. Of these, 2208 were excluded at the title and abstract stage, while the full texts of 104 references were examined. Forty-four references were included in the review. These covered 49 reports (for different age groups) from 41 independent surveys conducted in 12 countries: USA, Canada, Australia, UK, Germany, Denmark, Italy, Israel, Singapore, Japan, South Korea and Saudi Arabia. In terms of surveys of adults or all ages, there were 27 surveys reporting visits to acupuncturists, 20 for homeopaths, 9 for osteopaths, 33 for chiropractors, and 14 for medical herbalists (Table 1). The number of surveys reporting practitioner visits by children ranged from 0 to 4 for the different therapies, while the number reporting visits by older adults ranged from 0 to 8.

Of the 49 surveys included in our previous publication on any CAM use,<sup>1</sup> 8 did not report data on visits to any of the five types of practitioner covered here, and so were excluded. Data on individual practitioner visits from the remaining 41 surveys are included here.

### **Prevalence of visits to five types of CAM practitioner**

Table 1 outlines the 12-month prevalence of visits to each of the five types of practitioner (as well as to any CAM practitioner) for the 41 surveys across 12 countries. Survey data is ordered by country, then survey type (government, other national, or sub-national), then year of survey. Data are grouped by age: adults or all ages; children; and older adults. Further detail (sampling and data collection methods for each survey) is provided in our earlier publication.<sup>1</sup>

Table 2 provides a summary of the median and range for prevalence of visits to each type of practitioner, by age group. Estimates from surveys of adults or all ages were as follows (median and range): acupuncturists 1.4% (0.2 to 7.5%, N=27 surveys), homeopaths 1.5% (0.2 to 2.9%, N=20 surveys), osteopaths 1.9% (0.2 to 4.4%, N=9 surveys), chiropractors 7.5% (0.3 to 16.7, N=33 surveys), and medical herbalists 0.9% (0.3 to 4.7%, N=14 surveys). Estimates for

prevalence of visits by children were slightly lower, ranging from 0.2% for acupuncturists to 2.4% for chiropractors (based on 4 surveys each), while estimates for older adults were similar or slightly higher, ranging from 0.0% for homeopaths (based on 1 survey) to 8.4% for chiropractors (based on 8 surveys). Where possible, the following narrative identifies key data on prevalence of visits to CAM practitioners by national populations from data obtained using consistent methodologies.

#### *Visits to acupuncturists*

The five US government surveys estimated that between 0.6% and 1.4% of the adult (or all ages) population had visited an acupuncturist in the previous 12 months. Rates were similar over the years surveyed (1995-2007). Rates for other government surveys were similar: 1.6% for the UK (2001), 1.0% to 2.3% for Canada (1994-2005), and 2.0% to 2.8% for Australia (1993-2004). Estimates for visits by children were lower (0.1% to 0.3% across US and Australian surveys), while estimates for older adults tended to be higher, although many of these were from smaller or non-government sources.

#### *Visits to homeopaths*

Three US government surveys (1996-2007) estimated that between 0.4% and 1.8% of the adult (or all ages) population had visited a homeopath in the previous 12 months. Rates for other government surveys were similar: 1.7% to 1.9% for the UK (2001-2005), 2.0% to 2.3% for Canada (1994-2005), and 0.5% to 1.2% for Australia (1993-2004). Estimates for visits by children were 0.03% to 1.3% in US government surveys, while surveys of older adults generally did not report data on visits to homeopaths.

#### *Visits to osteopaths*

Data on osteopathy were less well reported than those for other therapies, with no US or Canadian surveys reporting these data. Two UK government surveys (2001-2005) estimated that between 1.9% and 2.7% of the adult (or all ages) population had visited an osteopath in the



previous 12 months, while estimates for Australia ranged from 0.2% to 0.4% (1993-2004).

These data were not well reported for children or older adults.

#### *Visits to chiropractors*

The five US government surveys estimated that between 3.3% and 10.9% of the adult (or all ages) population had visited a chiropractor in the previous 12 months. Rates were similar over the years surveyed (1995-2007). Estimates from other government surveys showed a similarly wide range: 1.6% to 2.2% for the UK (2001-2005), 1.4% to 11.0% for Canada (1994-2005), and 15.0% to 16.7% for Australia (1993-2004). Estimates for visits by children were lower but still significant (0.8% to 6.4% across US and Australian surveys), while estimates for older adults tended to be towards the higher end of the range of estimates for all adults.

#### *Visits to medical herbalists*

These data were not particularly well-reported, since many surveys simply reported use of medical herbs including over-the-counter products and did not specify visits to practitioners which is the focus of our review. A US government survey from 1996 estimated visits for adults (or all ages) at 1.8%, while rates for other government surveys were similar: 0.8% to 1.8% for the UK (2001-2005), 0.6% for Canada (2001-2005), and 0.4% to 1.9% for Australia (1993-2004). Visits by children were poorly reported (0.2% in one 1996 US survey), and these data were not reported for older adults.

#### *Trends over time*

Where data on trends over time are available, these appear to indicate little change in prevalence of visits per therapy over the past 15-20 years. Visits to acupuncturists and chiropractors had the most data points available. For visits to acupuncturists, the US National Health Interview Survey estimated fairly constant rates between 1.1-1.4% for all surveyed years between 1999-2007, while UK surveys gave similar rates of 1.6% in 1998-2001, and rates from Australian government surveys ranged from 2.0-2.8% for all surveyed years between 1993-2004. For visits to chiropractors, the US NHIS estimated rates of 7.5-8.6% for the years 1999-

2007 (the estimate of 8.6% included osteopathy), while UK surveys gave slightly lower rates ranging between 0.5-3.6% for all surveyed years between 1998-2005, and Australian surveys gave slightly higher but again constant rates of 15.0-16.7 for the years 1993-2004.

### Study quality

Table 3 provides a summary of the quality of included survey reports. Full details for each survey are reported in our earlier publication.<sup>1</sup> The proportion of all survey reports achieving each of our criteria ranged from 49% to 83%. The criteria least likely to be met were (5) data weighting to population characteristics to reduce non-response bias, and (6) reporting of 95% confidence interval or standard error for key prevalence estimates. Of all surveys, 61% met four or more quality criteria; these percentages were 79% for government-sponsored surveys and 45% for other surveys. However, although there was a trend towards more government surveys meeting each individual criterion, the only marked difference observed was for the piloting criterion, where we made the assumption that all government-sponsored surveys were piloted.

## **DISCUSSION**

This report provides a comprehensive and systematic review of surveys reporting 12-month prevalence of visits by general populations to five key types of CAM practitioner. This complements our previous report which systematically reviewed prevalence of any CAM use and visits to any CAM practitioner by general populations.<sup>1</sup> The data reported here include estimates from 41 surveys across 12 countries. Data were well reported for visits to acupuncturists, homeopaths and chiropractors (reported in 27, 20 and 33 of the 36 surveys of adults or all ages, respectively). Data on visits to osteopaths and medical herbalists were slightly less well-reported (within 9 and 14 surveys of adults or all ages).

The survey data indicated that a small but significant percentage of the general population (adult or all ages) had visited each type of CAM practitioner over the previous 12 months, with median estimates of 1.4% for acupuncturists, 1.5% for homeopaths, 1.9% for osteopaths, 7.5%

for chiropractors and 0.9% for medical herbalists. Practitioner visits by children were less frequent but still substantial for some therapies (median estimate of 2.4% for chiropractors) while estimates of visits by older adults were similar to or slightly higher than estimates for adults or all ages.

Data were obtained from surveys which also reported overall 12-month prevalence of any CAM use and/or visits to any CAM practitioner. Therefore, surveys only reporting visits to one of the included types of therapist but not reporting overall CAM use or visits were not included in this review. This was due to the design of this review which focussed on studies reporting this overall data, and is a limitation of this review. Conversely, data from the types of survey included here (many of which were government surveys or large population surveys) may be expected to be of higher quality than data from surveys of a single therapy. A further limitation is the exclusion of studies not reported in English, although English-language reports of surveys from any country were included.

The quality of methodological reporting was variable; 25 of 41 surveys (61%) met four or more of six quality criteria. These rates were higher for government surveys, though this was mainly due to our assumption that all government surveys were piloted. Our earlier report showed wide variation in estimates of 12-month prevalence of any CAM use (range 9.8% to 76%) and visits to CAM practitioners (range 1.8% to 48.7%), which was likely due in part to the variation in the way CAM is defined for data collection. Conversely, data reported here on visits to individual types of practitioner were more consistent with narrower ranges. This formed part of our rationale for focussing on prevalence of visits to practitioners rather than self-treatment; we did not include estimates of homeopathy or medical herbs which included use of over-the-counter remedies (or which were ambiguous on this point) since we felt that data on practitioner visits would be better-defined and less prone to recall bias.

Periodic surveys of general populations are important to monitor changing patterns in CAM use. Where data on trends over time are available, these appear to indicate little change in

prevalence of visits per therapy over the past 15-20 years for these five types of CAM. This, and the relative consistency of estimates across the different countries despite differences in local legislation and access, suggest a pattern of consistent healthcare seeking behaviour, rather than a response to fashions or trends.<sup>6</sup> Recent reports in Australia, UK, and USA<sup>7-9</sup> emphasised the need to improve communication between physicians and patients about their use of CAM; openness and non-judgemental communication is needed to determine the risks of drug interactions and other potential complications.<sup>7</sup> In summary, this review provides a comprehensive overview of prevalence of visits to key types of CAM practitioners by general populations in Europe, North America, Australia, East Asia, Saudi Arabia and Israel.

**Acknowledgements:** None

**Conflict of interest statement:** None.

**Author contributions:** All authors contributed to the design of the review, extraction and compiling of the data, drafting and critical revision of the manuscript.

## **REFERENCES**

- (1) Harris PE, Cooper KL, Relton C, Thomas KJ. Prevalence of complementary and alternative medicine (CAM) use by the general population: a systematic review and update. *Int J Clin Pract* 2012; 66(10):924-939.
- (2) House of Lords Select Committee on Science and Technology. Sixth Report: Complementary and Alternative Medicine. <http://www.publications.parliament.uk/pa/ld199900/ldselect/ldsctech/123/12302.htm>. 2000.
- (3) Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009; 151(4):264-269.
- (4) Ernst E. Prevalence of use of complementary/alternative medicine: a systematic review. *Bulletin of the World Health Organization* 2000; 78(2):252-257.

- (5) Harris P, Rees R. The prevalence of complementary and alternative medicine use among the general population: a systematic review of the literature. *Complementary Therapies in Medicine* 2000; 8(2):88-96.
- (6) Kelner M, Wellman B. Complementary and alternative medicine: challenge and change. London and New York: Taylor & Francis; 2000.
- (7) MacLennan AH, Myers SP, Taylor AW. The continuing use of complementary and alternative medicine in South Australia: costs and beliefs in 2004. *Medical Journal of Australia* 2006; 184(1):27-31.
- (8) Hunt KJ, Coelho HF, Wider B, Perry R, Hung SK, Terry R et al. Complementary and alternative medicine use in England: results from a national survey. *International Journal of Clinical Practice* 2010; 64(11):1496-1502.
- (9) Barnes PM, Bloom B, Nahin RL. Complementary and alternative medicine use among adults and children: United States, 2007. *National health statistics reports* 2008;(12):1-23.
- (10) Barnes PM, Powell-Griner E, McFann K, Nahin RL. Complementary and alternative medicine use among adults: United States, 2002. *Advance Data* 2004;(343):1-19.
- (11) Ni H, Simile C, Hardy AM. Utilization of complementary and alternative medicine by United States adults: results from the 1999 national health interview survey. *Medical Care* 2002; 40(4):353-358.
- (12) Druss BG, Rosenheck RA. Association between use of unconventional therapies and conventional medical services. *JAMA* 1999; 282(7):651-656.
- (13) Honda K, Jacobson JS. Use of complementary and alternative medicine among United States adults: the influences of personality, coping strategies, and social support. *Preventive Medicine* 2005; 40(1):46-53.
- (14) Eisenberg DM, Davis RB, Ettner SL, Appel S, Wilkey S, Van Rompay M. et al. Trends in alternative medicine use in the United States, 1990-1997: results of a follow-up national survey. *JAMA* 1998; 280(18):1569-1575.
- (15) Landmark Healthcare. The Landmark report on public perceptions of alternative care. 1998. Sacramento, Landmark Healthcare.
- (16) Paramore LC. Use of alternative therapies: Estimates from the 1994 Robert Wood Johnson Foundation National Access to Care Survey. *Journal of Pain and Symptom Management* 1997; 13(2):83-89.
- (17) Eisenberg DM, Kessler RC, Foster C, Norlock FE, Calkins DR, Delbanco TL. Unconventional medicine in the United States - Prevalence, costs, and patterns of use. *New England Journal of Medicine* 1993; 328(4):246-252.
- (18) Astin JA. Why patients use alternative medicine: results of a national study. *JAMA* 1998; 279(19):1548-1553.
- (19) Arcury TA, Preisser JS, Gesler WM, Sherman JE. Complementary and Alternative Medicine Use Among Rural Residents in Western North Carolina. *Complementary Health Practice Review* 2004; 9(2):93-102.
- (20) Oldendick R, Coker AL, Wieland D, Raymond JI, Probst JC, Schell BJ et al. Population-based survey of complementary and alternative medicine usage, patient satisfaction,

- and physician involvement. South Carolina Complementary Medicine Program Baseline Research Team. *Southern Medical Journal* 2000; 93(4):375-381.
- (21) Thomas K, Coleman P. Use of complementary or alternative medicine in a general population in Great Britain. Results from the National Omnibus survey. *Journal of Public Health* 2004; 26(2):152-157.
  - (22) Ernst E, White A. The BBC survey of complementary medicine use in the UK. *Complementary Therapies in Medicine* 2000; 8(1):32-36.
  - (23) Thomas KJ, Nicholl JP, Coleman P. Use and expenditure on complementary medicine in England: a population based survey. *Complementary Therapies in Medicine* 2001; 9(1):2-11.
  - (24) Thomas KJ, Fall M, Nicholl J, Williams B. Methodological study to investigate the feasibility of conducting a population-based survey of the use of complementary health care. 1993. University of Sheffield: SchARR.
  - (25) Yung B, Lewis P, Charny M, Farrow S. Complementary medicine: Some population-based data. *Complementary Medical Research* 1988; 3(1):23-28.
  - (26) Metcalfe A, Williams J, McChesney J, Patten SB, Jette N. Use of complementary and alternative medicine by those with a chronic disease and the general population--results of a national population based survey. *BMC Complementary & Alternative Medicine* 2010; 10:58.
  - (27) Millar WJ. Patterns of use--alternative health care practitioners. *Health Reports* 2001; 13(1):9-21.
  - (28) Millar WJ. Use of alternative health care practitioners by Canadians. *Canadian Journal of Public Health* 1997; 88(3):154-158.
  - (29) Northcott HC, Bachynsky JA. Concurrent utilization of chiropractic, prescription medicines, nonprescription medicines and alternative health care. *Social Science and Medicine* 1993; 37(3):431-435.
  - (30) Xue CC, Zhang AL, Lin V, Da CC, Story DF. Complementary and alternative medicine use in Australia: a national population-based survey. *Journal of Alternative & Complementary Medicine* 2007; 13(6):643-650.
  - (31) MacLennan AH, Wilson DH, Taylor AW. The escalating cost and prevalence of alternative medicine. *Preventive Medicine* 2002; 35(2):166-173.
  - (32) MacLennan AH, Wilson DH, Taylor AW. Prevalence and cost of alternative medicine in Australia. *Lancet* 1996; 347(9001):569-573.
  - (33) Niskar AS, Peled-Leviatan T, Garty-Sandalon N. Who uses complementary and alternative medicine in Israel? *Journal of Alternative & Complementary Medicine* 2007; 13(9):989-995.
  - (34) Shmueli A, Shuval J. Use of complementary and alternative medicine in Israel: 2000 vs. 1993. *Israel Medical Association Journal: Imaj* 2004; 6(1):3-8.
  - (35) Rasmussen NK, Morgall JM. The use of alternative treatments in the Danish adult population. *Complementary Medical Research* 1990; 4:16-22.

- (36) Lim MK, Sadarangani P, Chan HL, Heng JY. Complementary and alternative medicine use in multiracial Singapore. *Complementary Therapies in Medicine* 2005; 13(1):16-24.
- (37) Schwarz S, Messerschmidt H, Volzke H, Hoffmann W, Lucht M, Doren M. Use of complementary medicinal therapies in West Pomerania: A population-based study. *Climacteric* 2008; 11(2):124-134.
- (38) Yamashita H, Tsukayama H, Sugishita C. Popularity of complementary and alternative medicine in Japan: a telephone survey. *Complementary Therapies in Medicine* 2002; 10(2):84-93.
- (39) Al-Faris EA, Al-Rowais N, Mohamed AG, Al-Rukban MO, Al-Kurdi A, Balla Al-Noor MA et al. Prevalence and pattern of alternative medicine use: the results of a household survey. *Annals of Saudi Medicine* 2008; 28(1):4-10.
- (40) Ock SM, Choi JY, Cha YS, Lee J, Chun MS, Huh CH et al. The use of complementary and alternative medicine in a general population in South Korea: results from a national survey in 2006. *Journal of Korean Medical Science* 2009; 24(1):1-6.
- (41) Davis MP, Darden PM. Use of complementary and alternative medicine by children in the United States. *Archives of Pediatrics & Adolescent Medicine* 2003; 157(4):393-396.
- (42) Yussman SM, Ryan SA, Auinger P, Weitzman M. Visits to complementary and alternative medicine providers by children and adolescents in the United States. *Ambulatory Pediatrics* 2004; 4(5):429-435.
- (43) Smith C, Eckert K. Prevalence of complementary and alternative medicine and use among children in South Australia. *Journal of Paediatrics & Child Health* 2006; 42(9):538-543.
- (44) McMahan S, Lutz R. Alternative therapy use among the young-old (Ages 65 to 74): An evaluation of the MIDUS database. *Journal of Applied Gerontology* 2004; 23(2):91-103.
- (45) Astin JA, Pelletier KR, Marie A, Haskell WL. Complementary and alternative medicine use among elderly persons: one-year analysis of a Blue Shield Medicare supplement. *Journals of Gerontology Series A-Biological Sciences & Medical Sciences* 2000; 55(1):M4-M9.
- (46) Cheung CK, Wyman JF, Halcon LL. Use of complementary and alternative therapies in community-dwelling older adults. *Journal of Alternative and Complementary Medicine* 2007; 13(9):997-1006.
- (47) Shreffler-Grant J, Weinert C, Nichols E, Ide B. Complementary therapy use among older rural adults. *Public Health Nursing* 2005; 22(4):323-331.
- (48) Zhang AL, Xue CC, Lin V, Story DF. Complementary and alternative medicine use by older Australians. *Annals of the New York Academy of Sciences* 2007; 1114:204-215.
- (49) Feng L, Chiam PC, Kua EH, Ng TP. Use of complementary and alternative medicines and mental disorders in community-living Asian older adults. *Archives of Gerontology & Geriatrics* 2010; 50(3):243-249.
- (50) Dello Buono M, Urciuoli O, Marietta P, Padoani W, De Leo D. Alternative medicine in a sample of 655 community-dwelling elderly. *Journal of Psychosomatic Research* 2001; 50(3):147-154.

**Table 1: Prevalence of visits to five types of CAM practitioners across 12 countries**

Country	Survey type	Year of survey	Name of survey <sup>†</sup>	Sample size	Sample ages (% males)	Meets ≥4 quality criteria	Visited acupuncturist (%)	Visited homeopath (%)	Visited osteopath (%)	Visited chiropractor (%)	Visited medical herbalist (%)	Visited any CAM practitioner (%)	Reference
<b>Adult or all ages</b>													
USA	Government national	2007	NHIS	23,393	18+ (NR)	Yes	1.4	1.8	-	8.6*	-	16.2	Barnes (2008) <sup>9</sup> Barnes (2004) <sup>10</sup> Ni (2002) <sup>11</sup> Druss (1999) <sup>12</sup> Honda (2005) <sup>13</sup>
		2002	NHIS	31,044	18+ (NR)	Yes	1.1	1.7	-	7.5	-	12.5	
		1999	NHIS	30,801	18+ (NR)	Yes	1.4	-	-	7.6	-	-	
		1996	MEPS	16,068	18+ (47)	Yes	0.6	0.4	-	3.3	1.8	8.3	
		1995-6	MIDUS	4,242	25-74 (43)	Yes	1.1	-	-	10.9	-	-	
USA	Other national	1997		2,055	18+ (48)	Yes	0.9	0.6	-	9.9	1.8	19.5	Eisenburg (1998) <sup>14</sup> Landmark (1998) <sup>15</sup> Paramore (1997) <sup>16</sup> Eisenberg (1993) <sup>17</sup> Astin (1998) <sup>18</sup>
		1997		1,500	18+ (NR)	-	2.0	-	-	16.0	-	-	
		1994		2,056	18-64 (49)	Yes	0.4	-	-	9.3	-	9.4**	
		1990		1,539	18+ (52)	Yes	0.4	0.2	-	7.1	0.3	12.3	
		NR		1,035	18+ (49)	-	-	-	-	15.7	-	-	
USA	Sub-national	1999		1,059	18+ (NR)	-	1.2	0.7	-	6.7	0.8	8.6	Arcury (2004) <sup>19</sup> Oldendick (2000) <sup>20</sup>
		1998		1,584	18+ (38)	Yes	-	-	-	8.7	-	-	
UK	Government national	2005	HSE	7,630	16+ (45)	Yes	-	1.7	2.7	2.2	1.8	12.1	Hunt (2010) <sup>8</sup> Thomas (2004) <sup>21</sup>
		2001	NOS	1,794	16+ (47)	Yes	1.6	1.9	1.9	1.6	0.8	10.0	
UK	Other national	1999		1,204	18+ (45)	-	-	-	0.7	0.5	-	-	Ernst (2000) <sup>22</sup> Thomas (2001) <sup>23</sup> Thomas (1993) <sup>24</sup>
		1998		2,669	18+ (43)	Yes	1.6	1.2	4.3	3.6	0.9	13.6	
		1993		676	18+ (47)	Yes	0.5	1.7	4.4	2.1	0.9	8.5	
UK	Government sub-national	1986	CHS	4,268	18+ (NR)	Yes	0.2	-	-	0.8*	-	2.6	Yung (1988) <sup>25</sup>
Canada	Government national	2001-5	CCHS	400,055	12+ (49)	Yes	2.3	2.3	-	1.4	0.6	12.4	Metcalf (2010) <sup>26</sup> Millar (2001) <sup>27</sup> Millar (1997) <sup>28</sup>
		1998-9	NPHS	14,150	18+ (46)	Yes	-	-	-	11.0	-	17.0	
		1994-5	NPHS	17,626	15+ (NR)	Yes	1.0	2.0	-	11.0	-	15.0	
Canada	Government sub-national	1988	AEAS	464	18+ (49)	-	-	-	-	10.8	-	-	Northcott (1993a) <sup>29</sup> Northcott (1993b) <sup>29</sup>
		1979	AEAS	439	18+ (47)	-	-	-	-	8.2	-	-	
Australia	Other national	2005		1,067	18+ (49)	Yes	7.5	2.9	3.5	14.6	4.7	44.1	Xue (2007) <sup>30</sup>
Australia	Government sub-national	2004	SAHOS	3,015	15+ (49)	Yes	2.1	0.5	0.4	16.7	1.9	26.5	MacLennan (2006) <sup>7</sup> MacLennan (2002) <sup>31</sup> MacLennan (1996) <sup>32</sup>
		2000	SAHOS	3,027	15+ (49)	Yes	2.8	1.2	0.4	16.7	0.9	23.3	
		1993	SAHOS	3,004	15+ (49)	Yes	2.0	1.2	0.2	15.0	0.4	20.3	
Israel	Government national	2003-4	INHIS	2,365	21+ (44)	-	1.6	1.3	-	0.3*	-	5.8	Niskar (2007) <sup>33</sup>
Israel	Sub-national	2000		2,505	45-75 (47)	-	2.9	2.9	-	1.3	-	9.8	Shmueli (2004a) <sup>34</sup> Shmueli (2004b) <sup>34</sup>
		1993		2,003	45-75 (48)	-	1.3	1.8	-	0.4	-	6.1	
Denmark	Government national	1987	DICE	4,753	16+ (NR)	-	1.5	-	-	-	-	10.0	Rasmussen (1990) <sup>35</sup>
Singapore	Sub-national	2002		468	18+ (46)	-	5.2	-	-	-	-	-	Lim (2005) <sup>36</sup>
Germany	Sub-national	1997-2001		4,291	20-79 (49)	-	-	1.0	-	0.9	2.4	6.0	Schwarz (2008) <sup>37</sup>
Japan	Other national	2001		1,000	20-79 (49)	Yes	6.7	-	-	7.1*	-	-	Yamashita (2002) <sup>38</sup>



Country	Survey type	Year of survey	Name of survey <sup>†</sup>	Sample size	Sample ages (% males)	Meets ≥4 quality criteria	Visited acupuncturist (%)	Visited homeopath (%)	Visited osteopath (%)	Visited chiropractor (%)	Visited medical herbalist (%)	Visited any CAM practitioner (%)	Reference
Saudi Arabia	Sub-national	2003		1,408	Mean 36; SD 14 (39)	Yes	0.3	-	-	-	-	23.9	Al-Faris (2008) <sup>39</sup>
South Korea	Other national	2006		3,000	30-69 (50)	-	-	-	-	0.5	-	-	Ock (2009) <sup>40</sup>
<b>Children</b>													
USA	Government national	2007 1996	NHIS MEPS	9,417 6,262	0-17 (NR) 0-17 (52)	Yes Yes	0.2 0.1	1.3 0.03	- -	2.8* 0.8	- 0.2	- 1.8	Barnes (2008) <sup>9</sup> Davis (2003), Yussman (2004) <sup>41,42</sup>
USA	Other national	1994		980	1-17 (NS)	Yes	0.1	-	-	1.9	-	-	Paramore (1997) <sup>16</sup>
Australia	Government sub-national	2004	SAHOS	911	0-15 (46)	Yes	0.3	-	-	6.4	-	-	Smith (2006) <sup>43</sup>
<b>Older adults</b>													
USA	Government national	1995-6	MIDUS	335	65-74 (48)	Yes	1.2	-	-	8.5	-	-	Honda (2005), McMahan (2004) <sup>13,44</sup>
USA	Other national	1994		414	65+ (NS)	Yes	0.4	-	-	6.5	-	-	Paramore (1997) <sup>16</sup>
USA	Sub-national	1997-8 NR NR		728 445 325	65+ (45) 65-94 (45) 60+ (51)	- - -	5.5 4.0 -	- - -	- - -	8.3 17.8 14.2	- - -	- - 17.5	Astin (2000) <sup>45</sup> Cheung (2007) <sup>46</sup> Shreffler-Grant (2005) <sup>47</sup>
Australia	Other national	2005		178	65+ (43)	Yes	9.2	-	5.9	15.7	-	34.9	Xue (2007), Zhang (2007) <sup>30,48</sup>
Singapore	Government national	2003-4	NMHSE	1,092	60+ (44)	Yes	3.7	0.0	0.0	0.04	-	-	Feng (2010) <sup>49</sup>
Italy	Sub-national	1996-7		655	65+ (37)	-	9.6	-	-	-	-	-	Dello Buono (2001) <sup>50</sup>
Japan	Other national	2001		1,000	60-79 (46)	Yes	5.9	-	-	5.9*	-	-	Yamashita (2002) <sup>38</sup>

\*Combined data for visits to chiropractor or osteopath. \*\*Estimate for all ages. <sup>†</sup>Survey names are provided where reported for government surveys: AEAS = Annual Edmonton Area Survey; CCHS = Canadian Community Health Survey; CHS = Cardiff Health Survey; DICE = Danish Institute for Clinical Epidemiology; HSE = Health Survey for England; INHIS = Israeli National Health Interview Survey; MEPS = Medical Expenditure Panel Survey; MIDUS = Midlife Development in the US; NHIS = National Health Interview Survey; NOS = National Omnibus Survey; NMHSE = National Mental Health Survey of the Elderly; NPHS = National Population Health Survey; SAHOS = South Australian Health Omnibus Survey.

**Table 2: Summary of prevalence of visits to five types of CAM practitioners**

Age group	Visited acupuncturist (%)		Visited homeopath (%)		Visited osteopath (%)		Visited chiropractor (%)		Visited medical herbalist (%)		Visited any CAM practitioner (%)	
	N surveys	Median % (range)	N surveys	Median % (range)	N surveys	Median % (range)	N surveys	Median % (range)	N surveys	Median % (range)	N surveys	Median % (range)
<b>Adults or all ages</b>	27	1.4 (0.2-7.5)	20	1.5 (0.2-2.9)	9	1.9 (0.2-4.4)	33	7.5 (0.3-16.7)	14	0.9 (0.3-4.7)	25	12.3 (2.6-44.1)
<b>Children</b>	4	0.2 (0.1-0.3)	2	0.7 (0.0-1.3)	0	-	4	2.4 (0.8-6.4)	1	0.2 (-)	1	1.8 (-)
<b>Older adults</b>	8	4.8 (0.4-9.6)	1	0.0 (-)	2	3.0 (0.0-5.9)	8	8.4 (0.0-17.8)	0	-	2	26.2 (17.5-34.9)

**Table 3: Summary of the methodological quality of surveys**

Quality criterion	All survey reports N = 41		Govt. sponsored survey reports N = 19		Other CAM survey reports N = 22	
	n	%	n	%	n	%
1. CAM-use questions clearly described and number of therapies/questions reported	27	66	13	68	14	64
2. Piloting of survey reported (or assumed for government surveys)	29	71	19	100 <i>(assumed)</i>	10	45
3. Sample size ≥1,000 and/or CAM-specific sample size calculation reported	34	83	17	89	17	77
4. Reported survey response rate ≥60%	26	63	14	74	12	55
5. Data weighted to population characteristics (where appropriate) to reduce non-response bias	23	56	12	63	11*	50
6. 95% confidence interval or standard error reported for main prevalence estimates	20	49	9	47	11	50
<b>Four or more criteria met</b>	<b>25</b>	<b>61</b>	<b>15</b>	<b>79**</b>	<b>10</b>	<b>45</b>

\*Including one survey (Al-Faris et al 2008) with 95% response reported where weighting was deemed unnecessary.

\*\*This includes the assumption that the piloting criterion is met by 100% of government surveys.