



UNIVERSITY OF LEEDS

This is a repository copy of *Tethered N-Heterocyclic Carbene-Carboranes: Unique Ligands that Exhibit Unprecedented and Versatile Coordination Modes at Rhodium*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/98345/>

Version: Accepted Version

Article:

Holmes, J, Pask, CM, Fox, MA et al. (1 more author) (2016) Tethered N-Heterocyclic Carbene-Carboranes: Unique Ligands that Exhibit Unprecedented and Versatile Coordination Modes at Rhodium. *Chemical Communications*, 52 (38). pp. 6443-6446. ISSN 1359-7345

<https://doi.org/10.1039/C6CC01650B>

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Tethered N-Heterocyclic Carbene-Carboranes: Unique Ligands that Exhibit Unprecedented and Versatile Coordination Modes at Rhodium

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Jordan Holmes,^a Christopher M. Pask,^a Mark A. Fox^b and Charlotte E. Willans^{a*}

www.rsc.org/

Four brand new hybrid ligands combining an N-heterocyclic carbene tethered with two isomeric *nido*-dicarbadecaborane dianions, a neutral *closo*-dicarbadodecaborane or a *closo*-dicarbadodecaborane anion are described. Versatile coordination of the ligands to Rh^I is demonstrated, in which both NHC and carborane moieties covalently coordinate a metal centre.

Ligand design is central to the development of new organometallic and coordination compounds as they control the overall properties, the activities and the reactivities of a metal centre. This leads to potential new applications in materials, biomedicine and catalysis.^{1–6} Whilst modifying current ligands may induce incremental changes in the chemical behaviour of a metal complex, novel ligand architectures can lead to more diverse variations. N-Heterocyclic carbenes (NHCs) are excellent two electron donor ligands, with their steric and electronic properties being controlled and tuned through alteration of the N-substituents and backbone substituents respectively (Figure 1A. R¹–R⁴).^{7,8}

Carborane anions are also important classes of ligands, having very different properties to neutral NHCs. While there are many known classes of carboranes, the well-known *ortho*-carborane of type 1,2-R⁵,R⁶-1,2-C₂B₁₀H₁₀ (Figure 1B) and the *nido*-carborane dianion of type [7,8-R⁵,R⁶-7,8-C₂B₉H₉]²⁻ (Figure 1D) are explored here. They can coordinate to a metal through either a boron atom or a carbon atom of a *closo*-carborane anion (e.g. Figure 1C),⁹ or through the open face of a *nido*-carborane dianion (e.g. Figure 1D) to form a metallacarborane,¹⁰ in analogy to the widely used cyclopentadienyl ligand. *Ortho*-carborane precursors can be easily modified by substituting the acidic cage hydrogen atom(s) at the cluster carbons with groups that have desirable electronic and steric effects. This includes the addition of tethering substituents such as amino^{11,12} or cyclopentadienyl¹³

groups. Herein, we report unprecedented tethered NHC-carborane ligands (E–H) in which both the carborane and the NHC moieties are available for metal binding.





Fig. 1 Representation of an NHC (A), a neutral *ortho*-carborane (B), an *ortho*-carborane anion (C), a *nido*-carborane dianion (D) and novel hybrid ligands described in this study (E–H).

Slow addition of one equivalent of ^tbutylimidazole to bromoethyl-*ortho*-carborane in toluene was carried out at 75 °C for 18 hours (Scheme 1). The product, isolated in 72 % yield, was characterised as an imidazolium salt linked to a *closo*-carborane *via* an ethylene tether (1). Addition of excess ^tbutylimidazole to bromoethyl-*ortho*-carborane induced deboronation, with the ¹¹B NMR data for the resulting product exhibiting characteristic resonances for a *nido*-carborane monoanion of type [7,8-R⁵,R⁶-7,8-C₂B₉H₁₀]. The product, isolated in 65 % yield, was identified as an unusual imidazolium *nido*-carborane zwitterion 2. This compound could also be prepared from imidazolium 1 with two equivalents of ^tbutylimidazole. Compounds 1 and 2 were characterised using multinuclear NMR spectroscopy, mass spectrometry, elemental analysis and X-ray crystallography (see Supporting Information).

^a School of Chemistry, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT, UK. E-mail: c.e.willans@leeds.ac.uk

^b Department of Chemistry, Durham University, South Road, Durham, DH1 3LE, UK. Electronic Supplementary Information (ESI) available: Detailed experimental procedures, spectroscopic and crystallographic data. See DOI: 10.1039/x0xx00000x



Scheme 1 Synthesis of new ligand precursors **1** and **2**. Deprotonation and variable coordination of the ligands to Rh^I (unlabelled vertices = BH).

A common method to prepare metallocarboranes involves deprotonation of the *nido*-carborane of type $[7,8\text{-R}^5, \text{R}^6\text{-}7,8\text{-C}_2\text{B}_9\text{H}_{10}]^-$ using NaH with subsequent coordination of the dianion of type $[7,8\text{-R}^5, \text{R}^6\text{-}7,8\text{-C}_2\text{B}_9\text{H}_9]^{2-}$ (Figure 1D) to a metal. As NaH may also be used to generate free carbenes from imidazolium salts, zwitterion **2** was reacted with NaH to deprotonate both sites (Scheme 1). Subsequent treatment with $[\text{Rh}(\text{COD})\text{Cl}]_2$ for 3 hours at room temperature resulted in the presence of two new Rh-NHC species, **3**·MeCN and **4**·MeCN, in the ^1H NMR spectrum (CD_3CN), with the $^{11}\text{B}\{^1\text{H}\}$ NMR spectrum showing resonances characteristic of *closo* MC_2B_9 metalladiboraboranes. Crystals grown from the product mixture were identified as complex **3**·MeCN (Figure 2), a bimetallic complex featuring two linked Rh^I centres in very different chemical environments, formally $[\text{Rh}^I(\text{NHC})(\text{COD})(\text{MeCN})]^+$ and $[\text{Rh}^I(\text{carborane})(\text{COD})]^-$. While derivatives of the $[\text{Rh}^I(\text{carborane})(\text{COD})]^-$ anion have been known for decades,^{14, 15} this is the first solid-state structural elucidation of its kind.

Heating of the reaction mixture of **3**·MeCN and **4**·MeCN in MeCN for 18 hours gave only the thermodynamically stable product **4**·MeCN in 67 % yield. Similarly to **3**·MeCN, crystals of **4**·MeCN revealed $[\text{Rh}^I(\text{NHC})(\text{COD})(\text{MeCN})]^+$ and $[\text{Rh}^I(\text{carborane})(\text{COD})]^-$ moieties, but with the 2,1,8- MC_2B_9 metalladiboraborane isomer present, where one cluster carbon atom is not bonded to Rh. The cluster carbon atoms in both **3**·MeCN and **4**·MeCN were identified using the Vertex-to-Centroid Distance (VCD) Method.¹⁶ The observed 3,1,2 to 2,1,8 RhC_2B_9 cluster rearrangement is known, and attributed to relief of steric crowding.¹⁷ In addition, Stone and co-workers reported a series of Pd^{II} -metalladiboraboranes, and found that polytopal rearrangement occurs readily in the presence of a COD co-ligand, thought to be promoted by dissociation of the COD.¹⁸ Rearrangement of complex **3**·MeCN to **4**·MeCN occurs under relatively mild conditions, with both steric crowding and the presence of a COD co-ligand being contributing factors.





Fig. 2 Molecular structure of Rh^I complexes **3**-MeCN (left) and **4**-MeCN (right). H atoms are omitted for clarity and ellipsoids are plotted at 50 % probability level. Selected bond lengths (Å): **3**-MeCN Rh1-C7, 2.061(4); C12-C13, 1.380(6); C16-C17, 1.389(6); Rh2-C1, 2.231(4); Rh2-C2, 2.306(4); C1-C2, 1.608(5); C25-C26, 1.410(5); C22-C29, 1.399(6). **4**-MeCN Rh1-C7, 2.048(3); C12-C13, 1.394(4); C16-C17, 1.371(5); Rh2-C1, 2.290(3); C22-23, 1.404(5); C26-C27, 1.393(5).

Recrystallisation of **4**-MeCN with a mixed solvent system (MeCN/Et₂O/hexane) resulted in the unexpected formation of a dimeric solid-state structure (**4**₂) with MeCN ligands absent. The cationic [Rh^I(NHC)(COD)]⁺ moiety appears to be stabilised by the anionic [Rh^I(carborane)(COD)]⁻ moiety of a second molecule *via* close Rh...H-B links and vice versa, resulting in the formation of a dimer. Rhodacarborane clusters containing Rh...H-B bridges have previously been reported,¹⁹ with the complexes usually bimetallic. The dimer **4**₂ does not appear to persist in solution, as the coordinated acetonitrile ligand is observed in the ¹H NMR spectrum (acetone-d₆). However, the demonstrated lability of the MeCN ligand and the contrasting Rh^I environments renders complex **4**-MeCN interesting for development in, for example, tandem catalysis without the need for a hetero-bimetallic system.^{20, 21}



Oxidative addition of a carborane C-H or B-H bond to the Rh^I centre^{25, 26} appears to be resisted, and is likely due to the steric bulk surrounding the Rh^I centre.

In conclusion, we have developed new ligand systems which combine soft and hard ligands, namely NHCs and carborane anions. Examination of the ligands has revealed unique and versatile coordination to Rh^I through both the NHC and either a *nido*-dicarbaundecaborane dianion or a *closo*-dicarbadodecaborane anion. The *nido*-carborane ligands form homo-bimetallic complexes, with two Rh^I centres in considerably different chemical environments. The *closo*-carborane ligand forms a remarkable 7-membered metallacycle coordinating through both the NHC and the carbon atom of the carborane. Expansion of these highly tailorable ligand classes through substituent modifications, varying tethers and different metals is currently underway in our laboratory, with the possibilities of extending to pincer ligands²⁷ and introducing chiral centres into the ligand architectures.²⁸ Our novel ligands will be of broad interest for development in catalytic, materials and biomedical applications, with the NHC and carborane moieties working synergistically to tune a particular system.

Notes and references

1. L. Zhang, Y.-J. Lin, Z.-H. Li and G.-X. Jin, *J. Am. Chem. Soc.*, 2015, **137**, 13670-13678.
2. R. H. Morris, *Acc. Chem. Res.*, 2015, **48**, 1494-1502.
3. H. A. Younus, N. Ahmad, W. Su and F. Verpoort, *Coord. Chem. Rev.*, 2014, **276**, 112-152.
4. R. D. Hancock, *Chem. Soc. Rev.*, 2013, **42**, 1500-1524.
5. F. A. A. Paz, J. Klinowski, S. M. F. Vilela, J. P. C. Tome, J. A. S. Cavaleiro and J. Rocha, *Chem. Soc. Rev.*, 2012, **41**, 1088-1110.
6. Z. Liu and P. J. Sadler, *Acc. Chem. Res.*, 2014, **47**, 1174-1185.
7. M. N. Hopkinson, C. Richter, M. Schedler and F. Glorius, *Nature*, 2014, **510**, 485-496.
8. S. Diez-Gonzalez, N. Marion and S. P. Nolan, *Chem. Rev.*, 2009, **109**, 3612-3676.
9. Z. J. Yao, W. B. Yu, Y. J. Lin, S. L. Huang, Z. H. Li and G. X. Jin, *J. Am. Chem. Soc.*, 2014, **136**, 2825-2832.
10. R. N. Grimes, *Carboranes (Second Edition)*, Academic Press, Oxford, 2011, pp. 773-1014.
11. L. Xiang, K. Mashima and Z. Xie, *Chem. Commun.*, 2013, **49**, 9039-9041.
12. D.-H. Kim, J. H. Won, S.-J. Kim, J. Ko, S. H. Kim, S. Cho and S. O. Kang, *Organometallics*, 2001, **20**, 4298-4300.
13. D. Liu, Z. Qiu, H.-S. Chan and Z. Xie, *Can. J. Chem.* 2011, **90**, 108-117.
14. D. M. Speckman, C. B. Knobler and M. F. Hawthorne, *Organometallics*, 1985, **4**, 426-428.
15. R. Núñez, O. Tutusaus, F. Teixidor, C. Viñas, R. Sillanpää and R. Kivekäs, *Organometallics*, 2004, **23**, 2273-2280.
16. A. McAnaw, G. Scott, L. Elrick, G. M. Rosair and A. J. Welch, *Dalton Trans.*, 2013, **42**, 645-664.
17. J. A. Long, T. B. Marder, P. E. Behnken and M. F. Hawthorne, *J. Am. Chem. Soc.*, 1984, **106**, 2979-2989.
18. K. A. Fallis, D. F. Mullica, E. L. Sappenfield and F. G. A. Stone, *Inorg. Chem.*, 1994, **33**, 4927-4933.
19. P. E. Behnken, T. B. Marder, R. T. Baker, C. B. Knobler, M. R. Thompson and M. F. Hawthorne, *J. Am. Chem. Soc.*, 1985, **107**, 932-940.
20. J. A. Mata, F. E. Hahn and E. Peris, *Chem. Sci.*, 2014, **5**, 1723-1732.
21. S. Proch and R. Kempe, *Angew. Chem.-Int. Edit.*, 2007, **46**, 3135-3138.
22. C. E. Willans, C. A. Kilner and M. A. Fox, *Chem.-Eur. J.*, 2010, **16**, 10644-10648.
23. Z.-J. Yao, Y.-J. Lin, B. Xu and G.-X. Jin, *Dalton Trans.*, 2014, **43**, 4938-4940.
24. S. Bresadola, B. Longato and F. Morandini, *Coord. Chem. Rev.*, 1975, **16**, 19-26.
25. J. Estrada, S. E. Lee, S. G. McArthur, A. El-Hellani, F. S. Tham and V. Lavallo, *J. Organomet. Chem.*, 2015, **798**, 214-217.
26. X. L. Zhang, Z. W. Zhou and H. Yan, *Chem. Commun.*, 2014, **50**, 13077-13080.
27. A. M. Spokoiny, M. G. Reuter, C. L. Stern, M. A. Ratner, T. Seideman and C. A. Mirkin, *J. Am. Chem. Soc.*, 2009, **131**, 9482-9483.
28. A. P. da Costa, R. Lopes, J. M. S. Cardoso, J. A. Mata, E. Peris and B. Royo, *Organometallics*, 2011, **30**, 4437-4442.